## Order Lagomorpha

Pikas, hares, and rabbits range from 100-3000 g, yet their reproductive biology is remarkably constant. The reported gestation and lactation lengths are unusually homogeneous. Ovulation is induced by copulation and a postpartum estrus is present. Testes are abdominal outside of the reproductive season. All species are polytoccous but some variation in litter size and the developmental state at birth exists. Delineating the differences between hares and rabbits should be especially useful as the two groups form a natural control for the effects of diet, habitat, and phylogeny on the production of altricial versus precocial young because both are polytocous, both have similar diets and similar phylogenetic ancestry, and they can be found sympatrically.

## Order Lagomorpha, Family Ochotonidae, Genus *Ochotona*

Pikas (*Ochotona* 100-400 g) are terrestrial, often gregarious herbivores in Asia and North America. Ovulation is triggered by coitus (8787, 12007b). The deciduate, hemochorial, discoid placenta has a mesometrial, superficial implantation into a duplex uterus (4463, 8053) and is probably eaten after birth (11655). Estrus occurs (6820, 9808, 10089) after the birth of blind and naked or slightly furred (4193, 5146, 8783, 8784, 8786, 9175, 11657) neonates whose eyes open in 8-10 d (8053, 8783, 8798, 9808, 9949, 11039, 11655). Females have 6-12 mammae (1474, 2550, 8784, 9175) and a cloaca (2771). Testes descend during the breeding season (8053, 9808). Details of anatomy (2772, 5804, 9808) and milk composition (8786) are available.

| | | |
|---|---|---|
| **NEONATAL MASS** | | |
| *Ochotona alpina* | 6-8 g | 8798 |
| | 9.6 g; n1 | 4193 |
| *Ochotona curzoniae* | 11.2 g; sd1.2; n36 | 12007a |
| *Ochotona pallasi* | 6-8 g | 8798 |
| *Ochotona princeps* | 8.5 g; r4.1-12.5; n34 | 9808 |
| | 9.0 g; r8.0-10.23; n5 | 6884 |
| | m: 11.5 g | 11657 |
| | 12.4 g; n71 | 3875, 11655, 11656 |
| | f: 12.7 g | 11657 |
| *Ochotona pusilla* | 6-8 g | 8798 |
| *Ochotona roylei* | 6-8 g | 8798 |
| *Ochotona rufescens* | 11.4 g; sd1.9; r8.5-12.25; n51 | 8783, 8784, 8786, 8788 |
| **NEONATAL SIZE** | | |
| *Ochotona curzoniae* | HB: 57.6 mm; sd4.3; n36 | 12007a |
| **WEANING MASS** | | |
| *Ochotona princeps* | 52.24 g; n28, 28 d | 9808 |
| | 75 g; sd13.2; r50.9-103.2; n37,; 6th wk | 11655, 11656 |
| | f: 121.5 g; 59 d | 11657 |
| | m: 132 g; 59 d | 11657 |
| *Ochotona rufescens* | 74.9 g; sd15.5; n51 | 8788 |
| | 82 g; 21 d | 8783 |
| **LITTER SIZE** | | |
| *Ochotona alpina* | ~1 wean | 1238 |
| | 2.2 yg | 9070 |
| | 2.4 emb, yg; r2-3; n5 f | 4193 |
| | 3-8 | 1033 |
| | mode 3; r1-5 | 5717 |
| | 3 yg; n1 lt | 8053 |
| | 3 emb; n1 f | 10516 |
| | 4-6 | 10517 |
| | 4.8 ps; r2-6 | 5519 |
| *Ochotona collaris* | 2-6 | 2088 |
| | 3 emb; n1 f | 6689 |
| | 4 emb; n1 f | 6689 |
| | 4 emb; n1 f | 12049 |
| *Ochotona curzoniae* | 2-5 yg | 7382 |
| | 4.8 yg; ±1.8; r1-8; n17 lt | 12007b |
| *Ochotona daurica* | 3-5 | 8053 |
| | 5.6 emb; r2-7; n5 f | 10516 |
| | 6.1-7.7; r1-11; n >1 population | 10090 |

| | | |
|---|---|---|
| *Ochotona macrotis* | 1-4 yg | 7382 |
| | 4.08; r2-6 | 10090 |
| | 5 yg; r3-7 | 12104 |
| *Ochotona pallasi* | 1 yg; n1 f | 8053 |
| | 6.5-7.5; r1-12; n >1 population | 10090 |
| | 8 emb; n1 f | 10516 |
| *Ochotona princeps* | 1.62 yg; n26 lt | 11657 |
| | 1.75; r1-3 | 10092 |
| | 1.83 wean; ±0.08; mode 2; n35 lt | 7301, 7302 |
| | 2-3 | 2359 |
| | 2.17; r1-3; n6 lt | 10089 |
| | 2.33 emb; ±0.07; mode 2-3; r1-4; n80 f | 7301, 7302 |
| | 2.64 cl, ca; mode 2,3; r2-4; n175 lt | 7301, 7302 |
| | 2.67 emb; r2-3; n3 f | 279 |
| | 2.7 emb; r2-3; n3 f | 279 |
| | 2.74 yg; mode 3; r1-4; n26 f | 3875, 11655, 11656 |
| | 2.78; mode 2-3; r1-5; n18 lt | 10089 |
| | 2.9 yg; se0.2; r1-6; n21 | 6820 |
| | mode 3-4 | 2360, 6884 |
| | mode 3-5 yg | 9807 |
| | 3 emb; n1 f | 5025 |
| | 3 emb; n1 f | 6560 |
| | 3 yg; n1 lt | 2360 |
| | 3 yg; n1 lt | 2550a |
| | 3 yg; n1 | 11039 |
| | 3.0 yg; r2-4; n2 lt, 1 f | 10089 |
| | 3.0; r2-4; n10 lt | 5804, 6820, 10089 |
| | 3.13 emb; n53 f | 10088a, 10089 |
| | 3.4; r2-5; n50 lt | 6820 |
| | 3.6 ps; se0.2; n33 f | 12049 |
| | 4 emb; n1 f | 6553 |
| | 4 emb; n1 f | 2550 |
| | 5 emb; n1 f | 9949 |
| *Ochotona pusilla* | r3-13 yg; 8.1-8.8; 1st lt; 9.1-10.6; 2nd, 3rd lt; 7.3-7.4; last lt | 1033 |
| | 4-12 | 8053 |
| | 6 yg n1 f | 8798 |
| | 7-13 | 8053 |
| | 9.7 emb; r6-12; n3 f | 4645 |
| | max 12 | 10090 |
| *Ochotona roylei* | 3 emb; r1-5 | 7382 |
| | 3.0; r1-5 | 8784 |
| *Ochotona rufescens* | 4.82 wean; n85 lt | 3553 |
| | 5.2 emb; se0.36; r4-10; n23 f | 9175 |
| | 6; r4-9 | 8783, 8784, 8786, 8788 |
| | 6.12 yg; r4-10; n85 lt, 24 f | 8785 |
| | 6.59; n182 lt | 10090 |
| *Ochotona rutila* | 4.2 | 8798 |
| **SEXUAL MATURITY** | | |
| *Ochotona alpina* | f/m: 25-30 d | 5717 |
| | f/m: not 1st summer | 10516 |
| *Ochotona collaris* | f: 1st birth: ~1 yr | 6689 |
| *Ochotona daurica* | f: 1st birth: 1st yr | 10090 |
| *Ochotona macrotis* | f/m: 1st mate: 1 yr | 12104 |
| | f/m: 1st repro: 1 yr | 10516 |
| *Ochotona pallasi* | f/m: 1st mate: 1st yr | 8798 |
| | f/m: 25-30 d | 7300 |
| *Ochotona princeps* | m: sperm: yearling | 9808 |
| | f/m: spring of yr after birth | 9949 |
| *Ochotona pusilla* | f: ~4-5 wk | 9949 |
| | m: yearling | 8798 |
| | f/m: 25-30 d | 8798 |

| | | |
|---|---|---|
| *Ochotona roylei* | f/m: 21-23 d | 3553 |
| *Ochotona rufescens* | f: 1st preg: 5-8 wk | 8784 |
| | f/m: 65 d | 8785 |
| | f/m: 60 d | 10090 |
| *Ochotona rutila* | f/m: 1st mate: yearling | 10517 |
| **GESTATION** | | |
| *Ochotona alpina* | est ~4 wk | 8798 |
| | 30 d | 8798 |
| *Ochotona curzoniae* | 22.5 d; ±0.9; r21-24; n17 lt | 12007b |
| *Ochotona pallasi* | 25 d | 7300, 7302 |
| *Ochotona princeps* | 30 d | 11656 |
| | 30 d | 10092 |
| | 30 d; n2 | 2359, 9807, 9808 |
| | 30 d 19 h; n8 | 8798 |
| *Ochotona pusilla* | 20-24 d | 8783, 8784, 8785, 8786, 8788 |
| *Ochotona rufescens* | 25-26.5 d | 8798 |
| **LACTATION** | | |
| *Ochotona alpina* | den emergence: 15 d | 8798 |
| | solid food: 18-20 d | 8798 |
| | wean: 20-22 d | 8798 |
| *Ochotona pallasi* | den emergence: 15 d | 8798 |
| | solid food: 18-20 d | 3875, 11655, 11656 |
| | wean: 20-22 d | 11039 |
| *Ochotona princeps* | solid food: 7 d | 11656 |
| | solid food: 12 d | 9808 |
| | wean: mode 35 d; min 18 | 2360 |
| | wean: ~3-4 wk | 11655 |
| | wean: <4 wk | 7302 |
| | wean: end 6th wk | 2359 |
| | wean: 3-4 wk | 11657 |
| | wean: 21-29 d | 8798 |
| | wean: 3-5 wk | 8798 |
| *Ochotona pusilla* | den emergence: 15 d | 8798 |
| | solid food: 18-20 d | 9949 |
| | wean: 20-22 d | 8798 |
| | wean: max 3 wk | 8798 |
| *Ochotona roylei* | den emergence: 15 d | 8798 |
| | solid food: 18-20 d | 8783, 8784 |
| | wean: 20-22 d | 9175 |
| *Ochotona rufescens* | solid food: 8 d | 8783, 8784, 8788 |
| | wean: 2-3 wk | 9808 |
| | wean: 21 d; min 9-10 | 10092 |
| **INTERLITTER INTERVAL** | | |
| *Ochotona princeps* | ~1 mo | 5519 |
| | ~1 mo | 1033 |
| **LITTERS/YEAR** | | |
| *Ochotona alpina* | est 1 | 9070 |
| | 2 | 2088 |
| | 2 | 5717 |
| | 2 | 6601a |
| | 2/season | 8798 |
| | 2 | 10090 |
| | 2-3/summer | 8053 |
| | 2-3 | 10517 |
| *Ochotona collaris* | mode 1; r1-2 | 10516 |
| *Ochotona daurica* | 1-3 | 8798 |
| | 2 | 10516 |
| *Ochotona pallasi* | 2-3 | 2088 |
| | 3 | 7300, 7301, 7302 |
| *Ochotona princeps* | mode 1; r1-2 | 6820 |
| | 2 | 2359 |
| | 2 | 2772 |
| | 2 | 10089, 10092 |

| | | |
|---|---|---|
| | max 2 | 9808 |
| | 3 | 1033 |
| | 3.0; r2-5 | 8053 |
| *Ochotona pusilla* | 2 | 8798 |
| | 2 | 9949 |
| | 2+; r2-3 | 4645 |
| | 3-4 | 10090 |
| | 3-5 adult; 1-3 juveniles | 9175 |
| *Ochotona roylei* | 1-2 | 8785 |
| *Ochotona rufescens* | est 2 | 8786 |
| | 2.8-5.2 | 8053, 10090 |
| | max 4 | 4193 |
| *Ochotona rutila* | 2-3 | 9070 |
| **SEASONALITY** | | |
| *Ochotona alpina* | estrus: end May-early June; Sojr-Charchan, Mongolia | 10516 |
| | preg f: June-early July; Japan | 5717 |
| | repro: mid Apr-?; Yakutia, USSR | 2088 |
| | repro: May-mid Aug; Mongolia | 10516 |
| | repro: ?-July; USSR | 12104 |
| *Ochotona collaris* | preg f: May-June; n2; AK | 6689 |
| | preg f: June; n1; AK | 6689 |
| | preg f: June; n1; YUK, Camada | 12049 |
| | birth: May-June; n BC, Canada, AK | 179 |
| | lact f: June; n1; AK | 6689 |
| | lact f: June-July; n2; YUK, Canada | 12049 |
| *Ochotona curzoniae* | birth: est May-Aug; Nepal | 7382 |
| *Ochotona daurica* | preg f: May-July; Mongolia | 7300 |
| *Ochotona macrotis* | repro: Apr-Aug; Tersky-Alatau, USSR | 9808 |
| *Ochotona pallasi* | birth: end May-July; Coney, Gobi, Mongolia | 9808 |
| *Ochotona princeps* | m fertile: May-June; Rocky Mt, ALB, Canada | 2359, 2360 |
| | testes: large Mar-May, small Oct, Dec; CA | 7302 |
| | estrus: early Mar-?; CA | 10092 |
| | mate: mid Apr-June; CO | 2772 |
| | mate: May-July; sw ALB, Canada | 7300, 7302 |
| | mate: end May-June; lab? | 5804, 6820 |
| | ovulate: June-July; UT, CO | 12049 |
| | conceive: May-early June; sw ALB, Canada | 9808 |
| | preg f: end Apr-July; CO | 2088 |
| | preg f: June; n1; YUK, Canada | 5146 |
| | preg f: ?-July; CA | 1474 |
| | birth: May-June; BC, Canada | 10092 |
| | birth: May-June; w US | 7302 |
| | birth: May-Aug; w US | 9949 |
| | birth: mid June-early Aug; lab? | 9175 |
| | birth: Aug; n1; captive | 2550a |
| | wean: end June-early Aug; sw ALB, Canada | 8786 |
| *Ochotona pusilla* | repro: Apr-June/Aug; Irtysh R, USSR | 8053 |
| *Ochotona roylei* | preg f: May, Aug; Nepal | 7382 |
| *Ochotona rufescens* | birth: 2nd lt June-July; Pakistan | 9175 |
| | repro: Apr-Sept; Afghanistan | 8786 |
| *Ochotona rutila* | birth: early June; USSR | 8053 |

## Order Lagomorpha, Family Leporidae

Rabbits and hares (*Bunolagus* 1-1.5 kg; *Caprolagus*, *Lepus* 1.3-7 kg; *Nesolagus* 365-390 mm; *Oryctolagus* 0.8-3 kg; *Pentalagus* 430-510 mm; *Poelagus* 2-3 kg; *Pronolagus* 1-3 kg; *Romerolagus* 340-530 g; *Sylvilagus* 250-3280 g) are terrestrial herbivores from the New World, Eurasia, and Africa. *Lepus* and *Sylvilagus* females are often larger than males (*Lepus* 1518, 3344, 3346, 3348, 4710, 5210, 6419, 6452, 8500, 8501, 8730, 11398; cf 265, 9080; *Sylvilagus* 1720, 5210, 6614, 10236), while each sex has been reported larger in *Oryctolagus* (1213, 4054, 8491, 10386).

Ovulation is induced by coitus (617, 1213, 1307, 1530, 1531, 1654, 4536, 5944, 6904, 6914, 8521). Superfetation may be caused by sperm storage (6913, 6914). The discoid, deciduate, labyrinthine, hemochorial or hemoendothelial placenta has a mesometrial, superficial implantation into a duplex uterus (231, 1005, 1214, 2959, 3011, 6934, 7563, 7567, 8053, 10310, 10487, 11966) and may be eaten after birth (4099, 8053). A postpartum estrus is common (1004, 1213, 1214, 1307, 1653, 1654, 1722, 2326, 3344, 3396, 4099, 4499, 5239, 5624, 5760, 5865, 6832, 7561, 7692, 8053, 9805, 11398), but in hares a prepartum estrus also occurs which may lead to superfetation (1003, 5883, 6909).

Leporids exhibit a range of neonatal development. The extremes are the furred *Lepus* neonates with open eyes (143, 502, 680, 1110, 1474, 2088, 2139, 2439, 2458, 3571, 4006, 4467, 4645, 5146, 5760, 6441, 7433, 8053, 8501, 8798, 9080, 9175, 9687, 10154, 10158, 11058, 11253, 11363, 11365) and the naked *Oryctolagus* neonates whose eyes open at 10 d (3571, 4645, 7433, 8053, 8798). Other genera are blind at birth, but furred (*Poelagus* 5760; *Pronolagus* 10158; *Romerolagus* eyes open 5-7 d, 1676, 2404, 2819, 2820; *Sylvilagus* eyes open 2-10 d, 893, 1290, 1474, 2891, 5146, 7561, 8798, 10222, 10583, 10731, 10908; cf 5099).

Females have 6 mammae (*Caprolagus* 3759; *Pronolagus* 9945), 6-10 mammae (*Lepus* 331, 1474, 2548, 2550, 3571, 4745, 6335, 7433, 8053, 9175, 9945; *Oryctolagus* 3571, 7433, 8053), or 8-10 mammae (*Sylvilagus* 1474, 1720, 2548, 2550, 4745). Males have seminal vesicles, a proprostate, paraprostate, prostate, and bulbourethral glands (*Oryctolagus* 1611, 4940) or inguinal and coagulating glands (*Sylvilagus* 2959).

Details of anatomy (1005, 1723, 4981), endocrinology (*Lepus* 1528, 1529, 1530, 1532, 1533, 1534, 1535, 1535a, 1535b, 2326, 2327, 2328, 2329, 5621, 6452, 6453, 10316, 10317, 10318; *Sylvilagus* 1535, 2964, 9524, 10384), copulation (9350a), reproductive cyclicity (9350), birth (1653), milk composition (277, 511, 6424), suckling (1306, 6908), and light (*Lepus* 2326; *Sylvilagus* 964) are available.

**NEONATAL MASS**

| | | |
|---|---|---|
| *Bunolagus monticularis* | 40-50 g | 2821c |
| *Lepus americanus* | 52 g; r48-58; n3 | 8005a |
| | 65.9 g; n1 emb | 1110 |
| | 67 g; sd13.2; n128 | 9805 |
| | 68.7 g; n3 | 3899 |
| | 81.7 g; r64-96; n22 | 2458 |
| *Lepus californicus* | 96-114 g | 4099 |
| | 110 g | 4467 |
| *Lepus capensis* | 80-130 g | 3350 |
| | 97 g; n21 | 3349 |
| | 100-130 g | 6101 |
| | 100-165 g | 3344, 3346 |
| | 107 g; r65-115; n54, 1-2 d | 8513 |
| | 113.2 g; ±15.6; r60-180; n283 | 6909 |
| | 130.2 g; r100-170; n11 | 4562 |
| | 150 g | 11058 |
| *Lepus saxatilis* | 98 g; n8 | 3349 |
| *Lepus timidus* | 70-130 g | 8798 |
| | 80-140 g | 4645 |
| | 80.7-132.7 g; across litter size | 8323 |
| | 84-140 g | 8053 |
| | 87.3 g; r71-106; n9 term emb | 3348 |
| | 104.8 g; n1, 2 d | 11414 |
| *Lepus townsendii* | 87 g; r84-90; n2 yg | 680 |
| | 89.25 g; r84-93; n4 term emb | 680 |
| | 105.33 g; r101-110; n3, 1 d | 680 |
| *Oryctolagus cuniculus* | 30-40 g | 8798 |
| | f: 35.23 g; se1.140; n40 | 1175 |
| | m: 36.99 g; se0.833; n55 | 1175 |
| | 40-45 g | 3571 |
| | 42.0-44.5 g | 1213 |
| | 45.8 g; sd1.62; n32 emb, 7 lt, 31 d gestation | 9270 |
| | 46.95 g; sd8.42; n126 emb, 20 lt, 30 d gestation | 10763 |
| *Romerolagus diazi* | 24.1 g; r22.6-25.0; n5 | 1677 |
| | f: 25-27 g | 6953b |
| | m: 32 g | 6953b |

| | | |
|---|---|---|
| *Sylvilagus aquaticus* | 53.5 g; r48.5-56.9; n6, 1 lt | 4920 |
| | 55.7 g; n1 | 10583 |
| | 61.4 g; n18 | 10222 |
| *Sylvilagus audubonii* | 32.8 g; n9 stillborn | 5142 |
| | 36 g; r33-40; n3 | 2550b |
| *Sylvilagus bachmani* | 24.3-30.2 g; term emb | 7561 |
| *Sylvilagus floridanus* | 24.8-31.2 g; 25.4 d gestation | 9251 |
| | 29.54 g; r23.3-33.0; n5 | 893 |
| | 30 g | 8798 |
| | 30.3-38.0 g; n6 term emb | 2891 |
| | 37.8 g; n22 yg, 1 d | 1137 |
| | 42.2 g; n6 | 4742 |
| **NEONATAL SIZE** | | |
| *Lepus americanus* | 105.3 mm; r97.0-113.4; n8 | 2458 |
| | CR: 110.7 mm | 1110 |
| | HB: 153.7 mm; n3 | 3899 |
| *Lepus capensis* | 130-170 mm | 11058 |
| *Oryctolagus cuniculus* | f: 117.70 mm; se1.181; n40; nose-tail | 1175 |
| | m: 119.07 mm; se0.964; n55; nose-tail | 1175 |
| *Romerolagus diazi* | TL: 78.3 mm; r74-83; n3 term emb | 1677 |
| | TL: 93.8 mm; r83-106; n5 | 1677 |
| *Sylvilagus aquaticus* | TL: 130 mm; n4 | 4920 |
| *Sylvilagus audubonii* | TL: 90 mm; n1 "very young" | 5142 |
| | 117 mm; n9 stillborn | 5142 |
| *Sylvilagus floridanus* | CR: 87 mm | 1137 |
| | 90-98 mm; n6 term emb | 2891 |
| | HB: 121-134 mm; 25.4 d gestation | 9251 |
| *Sylvilagus transitionalis* | CR?: 73.8 mm | 10731 |
| **WEANING MASS** | | |
| *Lepus alleni* | 3.0 kg; sd0.34; n13 | 9616 |
| *Lepus americanus* | ~400 g; 30 d | 8005a |
| | 486 g; sd84.4; n46; 28 d | 9805 |
| *Lepus californicus* | 1.47 kg; 12-13 wk | 10246 |
| | 2.3 kg; sd0.64; n9 | 9616 |
| *Lepus capensis* | 200 g; solid food | 3346 |
| | mode ~1 kg weaned | 3346 |
| | 808-1022 g | 6909 |
| *Lepus townsendii* | 737 g; ~4 wk | 10246 |
| *Oryctolagus cuniculus* | 198 g; ±23 | 11887 |
| | 200-250 g; 1st capture | 11887 |
| *Romerolagus diazi* | ~80 g; est 20 d | 5007 |
| | f: 80-130 g; 3 wk | 6953b |
| | m: 95-128 g; 3 wk | 6953b |
| *Sylvilagus floridanus* | ~112 g; leave nest | 893 |
| **LITTER SIZE** | | |
| *Bunolagus monticularis* | 1 yg | 2821c |
| *Caprolagus hispidus* | 2-5 | 3759 |
| *Lepus alleni* | 1.94 emb; mode 1,2; r1-5; n124 f | 11365 |
| | 3 emb; n1 f | 11363 |
| | 6 emb; n1 f | 5410 |
| *Lepus americanus* | 1 emb; n1 f | 2548 |
| | r1.64-3.05; across yr & season | 6132 |
| | 2 emb; n1 f | 9968 |
| | 2 ps; n2 | 10839 |
| | 2-4 cl | 7872 |
| | r2.2-4.9; 1st lt smaller | 5624 |
| | 2.27 emb; 1st lt: r2.96-3.26; later lt | 11895 |
| | 2.4 yg; mode 2; r1-4; n36 | 143 |
| | 2.5 emb; r1-4; n2 f | 10839 |
| | 2.68 yg; mode 2; r1-5; n71 f | 1111 |
| | 2.7 emb; mode 4; r1-4; n7 f | 2663 |
| | 2.73 emb, yg; r1-5; n26 lt | 6706 |
| | 2.82 emb; mode 3; r1-7; n266 f | 143 |
| | 2.88; r1-5; n81 lt | 9805 |

| | | |
|---|---|---|
| | 2.90 emb; r1-5; n140 f; r2.54-3.58; n7 yr | 4037 |
| | 2.90-5.67 across 4 mo; 2.6-6.5 by mo; n83 lt, 5 yr | 2326, 2328 |
| | 2.96 yg; mode 3; r1-6; n80 lt | 9806 |
| | 3.00; r2-4; 1st lt; 4.75; r2-7; later lt | 2672 |
| | 3.00; 1st lt; 4.84; later lt | 7243 |
| | 3 emb; n1 f | 4745 |
| | 3.0 yg; r1-4; n6 lt | 4006 |
| | 3.0; r1-4; n7 | 2663 |
| | 3.0 emb; r2-4; n11 f | 12049 |
| | 3.1 yg; n16 | 2664 |
| | 3.35; se0.12; n245 lt | 3510 |
| | 3.4 emb; 95%ci 0.3; n100 f | 1229 |
| | 3.44 emb; r1-4; n149 f | 2139 |
| | 3.56; r2-5; 1st lt; 5.91; r1-9; later lt | 2672 |
| | 3.7 cl; n146 tracts | 2664 |
| | 3.8 yg; n5 lt | 9579a |
| | 3.82 emb; mode 4; r1-7; n82 f | 9321 |
| | 3.83 yg; r2-6; n6 | 9579a |
| | 3.9 emb; n137 f | 2664 |
| | 3.9 emb; 95%ci 0.2; n129 yearlings | 1229 |
| | 4 emb; n1 f | 4006 |
| | 4.5 emb; r3-6; n2 f | 2545 |
| | 4.6 emb; 95%ci 0.4; n78 f | 1229 |
| | 4.7 emb; r1-8; n3 f | 10597 |
| | 5 emb; n1 f | 7809 |
| | 5 emb; n1 f | 8125 |
| | 5 emb; n1 f | 10598 |
| | 5 emb ea; n2 f | 4081a |
| | 5 emb ea; n3 f | 9579a |
| | 5.4 emb; r3-8; n5 f | 2550 |
| | 6 emb ea; n2 f | 5254a |
| | 7 emb; n1 f | 6553 |
| | 7 emb; n1 f | 10218 |
| *Lepus californicus* | 1.5-2.6 emb; across mo | 4332 |
| | 1.79 yg | 4467 |
| | mode 2-4; max 6 | 502 |
| | 2.24 emb; mode 1-2; r1-6; n70 f | 11365 |
| | 2.26 emb; mode 1-2; r1-8; n114 f | 5254a |
| | 2.5 yg | 6335 |
| | 2.5 emb; r2-3; n2 f | 522 |
| | 2.6 emb; se0.19; r1-5; n32 f | 1322 |
| | 2.8 emb; r2-4; n6 f | 2550 |
| | 3 emb; n1 f | 523a |
| | 3 emb; n1 f | 11228a |
| | 3.8 cl; r1-6 | 4099 |
| | 4 yg; r1-8 | 1871a |
| | 4.9 emb; ±1.4; r2-7; n22 f | 3197 |
| | 5.23 emb; r3-8; n39 f | 8126 |
| *Lepus capensis* | 1.0-2.8; across mo; r1-5 | 3344 |
| | 1-3 | 6452 |
| | 1-5 | 9125 |
| | 1-5 | 8798 |
| | 1-6 | 7433 |
| | 1 emb; n1 f | 2439 |
| | 1 emb; n1 f | 4925 |
| | 1 emb ea; n4 f | 6293 |
| | 1.17 emb | 6517 |
| | 1.2 emb; r1-2; n4 f | 331 |
| | 1.3 | 10318 |
| | 1.5 | 3349 |
| | 1.5-3.3 yg; across seasons | 5944 |
| | 1.6-1.8; 1st lt; 4.23-4.27; 2nd, 3rd lt | 8370 |
| | 1.65; se0.14; post partum mating; 2.24; se0.06; n>4000; prepartum mating | 1530, 1531 |

| | | |
|---|---|---|
| | 1.69; r1-3; n13 lt | 1452 |
| | 1.7-3.7 emb; across seasons | 4712, 5944 |
| | 1.7; r1-2; n3 lt | 8965 |
| | 1.77; mode 2; r1-4; n31 lt | 10487 |
| | 1.8; young f | 226 |
| | 1.9 emb; r1-3; n13 f | 10154 |
| | 1.9 emb; r1-6; n44 f | 11058 |
| | 1.9 emb; mode 1 50%; r1-4; n211 f | 9080 |
| | 2-4 | 5277 |
| | mode 2-4; r1-7 | 3571 |
| | 2-5; across seasons | 5944 |
| | 2 emb; n1 f | 6612 |
| | 2 emb; r1-4 | 8359 |
| | 2.1 | 2576 |
| | 2.1; mode 2; r1-3; n8 lt | 4562 |
| | 2.21 emb without resorption | 4712 |
| | 2.21 large emb; mode 1-3; r1-5; r1.0-2.8; across mo; n311 lt | 3346 |
| | 2.27; adult f | 226 |
| | 2.3; n155 lt | 7446 |
| | 2.3 | 8814 |
| | 2.3 emb; r2-3; n3 f | 8134 |
| | 2.34 yg; r1-5; n154 lt | 6909 |
| | 2.46 | 225 |
| | 2.5 | 8476 |
| | 2.55 ps; mode 3; r1-7; r1.0-3.6; across mo; n320 lt | 3346 |
| | 2.6; r1-4; n10 lt, 2 f | 8782 |
| | 2.64 | 6131 |
| | 2.66-3.40 yg; n4 groups | 3572 |
| | 2.68; r1.4-3.3; across mo | 1307 |
| | 2.69 emb; r1-4; n144 lt | 4709 |
| | 2.69; mode 2-3; r1-5; n960 lt | 4712 |
| | 2.76 | 4712 |
| | 2.8 cl; mode 3; r1-9; r1.0-3.8; across mo; n407 lt | 3346 |
| | 2.84; sd1.23; n248 lt | 3540 |
| | 3-5; rare 7 | 8053 |
| | mode 3-5; r3-7 | 4645 |
| | 3-6 | 3778 |
| | 3.25; r1-6; n63 lt | 9497 |
| | 3.75 emb; r3-4; n4 f | 10516 |
| | 4 yg; n1 lt | 1004 |
| | 5 emb ea; n4 f | 9774 |
| | max 6-7 | 6909 |
| *Lepus nigricollis* | 1.33; r1-2; n3 | 10616 |
| | 1.5 emb; r1-2; n2 f | 7382 |
| | 1.84 emb; mode 1; r1-4; n63 f | 8730 |
| | 2 | 9175 |
| | mode 2 | 8500 |
| | mode 2; r1-2 | 8501 |
| | 2.0 cl; r1-7; n60 f | 8730 |
| *Lepus oiostolus* | 2 emb; n1 f | 179 |
| *Lepus saxatilis* | 1-2 | 11253 |
| | 1-2 yg | 4354a |
| | 1; r1-2 | 8965 |
| | 1 emb ea; n2 f | 4925 |
| | 1.6 | 3349 |
| | 1.6 emb; r1-3; n86 f | 10154, 10158 |
| | 2 | 9945 |
| | mode 2 yg; r1-2 | 1149 |
| *Lepus timidus* | 1-7 | 7433 |
| | 1.59 emb; r1-3; n278 f | 4709 |
| | 1.74 emb; se0.05 | 4710a |
| | 1.88 emb; mode 1; r1-5; n72 f | 3348 |
| | 1.98 emb; se0.069 | 4710a |
| | 2.02 emb; se0.045 | 4710a |

| | | |
|---|---|---|
| | 2.06 yg; n16 fall lt | 7214 |
| | 2.10 implantation sites; mode 2; n78 | 3348 |
| | 2.3 emb; r1-5; n14 late-term f | 3348 |
| | 2.32 emb; se0.063 | 4710a |
| | 2.39 cl; mode 2; r1-5; n75 | 3348 |
| | 3 | 6441 |
| | 3; 1st lt; 5; r3-5; 2nd lt | 4645 |
| | 3; 1st lt; 5; 2nd lt; 3; 3rd lt | 8053 |
| | 3.11 yg; n84 spring lt | 7214 |
| | 3.3 viable emb; 95%ci 0.5; r2.5-4.5; across yr | 4548 |
| | 3.55; se0.2; r2-7; n40 lt | 8323 |
| | 3.72; n29 1st lt | 10876 |
| | 3.8; r3-5; n9 | 8233 |
| | 4-8 | 1474 |
| | 4.0 emb; n2 f | 6710 |
| | 4.33; n88 lt (without 1st lt) | 10876 |
| | 4.63; n8 lt | 10886 |
| | mode 6-8 emb | 5029a |
| | 6.3 emb, ps, yg; sd1.34; r3-8; n10 | 265 |
| | 6.5 cl; sd1.7; max 11; n15 f | 8233 |
| | 7 yg; n1 nest | 11414 |
| | max 8 yg | 5146 |
| *Lepus townsendii* | 3-6 | 1474 |
| | 3 emb; n1 f | 8126 |
| | 3 emb; n1 f | 6166 |
| | 3.3 emb; r1-5; n3 f | 5405b, |
| | 3.56 emb, 5.9 cl; n9 f | 5865 |
| | 3.6 yg; ±0.6; r1-5; n15 f | 5865 |
| | 4.5; r4-5 | 502 |
| | 4.5 emb; r4-5; n2 f | 1649a |
| | 4.6 yg | 5239 |
| | 4.9 emb | 5239 |
| | 5 emb, ps; r1-11; n25 f | 680 |
| | 5.55 emb; r4-8; n9 f | 5254a |
| | 5.67 emb; r5-6; n3 f | 261a |
| | 5.75 cl; ±0.94; r3-8; n12 f | 5865 |
| | 5.8 cl | 5239 |
| | 7 emb; n1 f | 3703 |
| *Oryctolagus cuniculus* | 1-14 | 8798 |
| | 2.0-5.2; across season and yr | 8228 |
| | mode 3-6 yg; n33 f; 4.5; n>100 lt | 7691 |
| | 3.3-6.1 emb; n13 sites | 5081 |
| | 4-7 | 11620 |
| | 4.16 yg; se0.19; r1-6; n56 lt | 1175 |
| | 4.3 emb; n233 f | 11887 |
| | 4.36 emb; r3.39-5.70; across months | 8491 |
| | 4.7 emb; se0.14; n101 f | 291, 293 |
| | 4.72; mode 5; r1-9; n191 lt | 7260 |
| | 4.8 yg; sd1.5; n22 lt | 7689 |
| | r4.85-5.22 emb; n4 locations | 8229a |
| | 4.88 emb; 5.81 cl; n1819 f | 1214 |
| | 4.9-5.6 cl | 1213 |
| | 4.923 ps; ±0.067 | 196 |
| | 5.0 yg; sd1.4; n61 lt | 7689 |
| | 5.0-5.33 emb; r3-7; n3 locations | 2898 |
| | 5.47 all emb; se0.28; 5.38 live emb; se0.10; n17 f; r3.0-5.49; across age | 8229 |
| | 5.54 | 752 |
| | 5.5 yg; n9 lt, 3 f | 7693 |
| | 5.676 cl; ±0.062 | 196 |
| | 5.77 cl; sd1.57; mode 5; r3-15; n656 f | 7070 |
| | 5.83 cl; sd1.76; mode 5; r3-15; n745 f | 11478 |
| | 5.95 emb; mode 5,7; r3-10; n57 f | 6516 |
| | 6 cl; 4-6 emb | 1178 |
| | 6-7.8; across 5 mo | 9918 |

| | | |
|---|---|---|
| | 6.32 yg; n22 | 7701 |
| | 6.69 emb; mode 4; r1-7; n71 f | 9270 |
| | 6.80 yg; se0.53; n20 lt | 60 |
| | 6.92 live emb; sd2.81; se0.17; r1-14; n270 f | 10763 |
| | 7.07 cl; ±1.49; 6.50 emb | 10044 |
| | 7.29; r5-9 | 10386 |
| | 9.7 cl; se0.2 | 10875 |
| | 11.6 cl; se0.2 | 10875 |
| *Poelagus marjorita* | 1 emb; n1 f | 4495 |
| | 1-2 | 5760 |
| *Pronolagus crassicaudatus* | 1-2 | 4180 |
| *Pronolagus randensis* | 1.1 emb; r1-2; n8 f | 10158 |
| *Pronolagus rupestris* | 2 yg; n1 nest | 10158 |
| *Romerolagus diazi* | 1-3 emb | 9231 |
| | 1.5; r1-2; n2 lt | 6478 |
| | 2.0 yg; n2 lt, n1 f | 2819, 2820 |
| | 2.1; r1-5 | 6569 |
| | 2.3 yg; n18 lt, 4 f | 5007 |
| | 2.33 yg; r2-3; n3 lt | 1676 |
| | 2.33 emb, ps; n12 f | 1677 |
| | 2.5 yg; n4 lt | 6953b |
| | ~3 yg; r1-4 | 6387 |
| *Sylvilagus aquaticus* | 1-6 yg | 10583 |
| | 2.0 yg; r1-3; n5 | 8214 |
| | 2-3; r1-5 | 1474 |
| | 2.6 emb, yg; r1-4; n5 lt | 6614 |
| | 2.8 large emb; mode 3; r1-4; n14 f | 10861 |
| | 2.83 emb; mode 2-3; r1-5; n29 f | 5099 |
| | 2.91; r1-6; n224 lt | 4740 |
| | 3.0 all emb; n24 f | 10861 |
| | 3.08; mode 3; r1-6; n38 lt | 10222 |
| | 3.3 ps; r3-4; n3 f | 6298 |
| | 3.4 ps; r3-4; n7 f | 10861 |
| | 3.7 emb; r3-5 | 10583 |
| | 3.7 cl; r2-6; n46 f | 10861 |
| | 4.8 yg; r4-6; n4 lt, 3-6 d | 4920 |
| | 6.7 cl; n5 f | 10753 |
| *Sylvilagus audubonii* | 1 emb; 1 f | 2550 |
| | 1-5 yg | 5146 |
| | 2 emb ea; n2 f | 502 |
| | 2.5; r1-4; n2 lt | 3705 |
| | 2.6 emb; r1-3; n10 f; 3.30 cl | 1727 |
| | 2.7 ps; n8 f | 10470 |
| | 2.9 emb; r2-4; mode 3; n56 f | 10236 |
| | 3 emb; n1 f | 5409, 5410 |
| | 3.0 yg; r1-5; n3 lt | 2550b |
| | 3 emb, yg; r1-5; n6 f | 5142 |
| | r3-6 emb; n≥5 f | 523a |
| | 3.3 emb; r2-6; n10 f | 10470 |
| | 3.6 emb; r2-6; n19 f | 8126 |
| | 4 emb; r3-5; n3 f | 5254a |
| | 4 ps; n1 f | 3703 |
| | 4.3 cl; r2-7; n18 f | 10470 |
| | 5 emb; n1 f | $5405b_1$ |
| | 6 emb; n1 f | 6166 |
| | 6 emb; n1 f | 10954 |
| | 6.4 emb; r5-7; n5 f | 261a |
| *Sylvilagus bachmani* | 2.67-3.27 emb, ps; n2 yr | 1722 |
| | 3.5 emb; r2-5; n11 f | 8126 |
| | 3.69 emb; r2-6; n26 f | 7561 |
| | 4 ps; r3-6; n14 f | 7561 |
| | 5 yg; n1 lt | 7561 |

| | | |
|---|---|---|
| *Sylvilagus brasiliensis* | 1.18 nestlings; mode 1; r1-2; n11 lt | 2817 |
| | 1.9 | 2816 |
| | 3-6 | 8798 |
| *Sylvilagus cunicularius* | 3 emb; n1 f | 5410 |
| | 5 emb; n1 f | 2341 |
| *Sylvilagus floridanus* | 2.6-6.6; 1st lt; r3.4-7.0 2nd lt across locations | 2010 |
| | 2.8-5.0 viable emb | 5807 |
| | 2.8 emb; n404 f | 10822 |
| | 2.9 emb, yg; r2-5; n9 lt | 6614 |
| | 2.9 emb; r2-4; n56 f | 10236 |
| | 3.0 emb; r1-4; n3 f | 3212 |
| | 3.0 emb; r2-3; n5 f | 522 |
| | 3.02 emb; mode 3; r1-6; n280 f | 1136 |
| | 3.1; r1-7; n36 | 10236 |
| | 3.13 cl; n107 lt | 8077 |
| | 3.14; n209 lt | 8340, 8341 |
| | 3.41 emb; r1-6; n29 f | 586 |
| | 3.5-5.6 yg; n3 samples | 3117 |
| | 3.5 emb; max 7; se0.0416; n611 lt | 4741, 4742 |
| | 3.52 emb; r2-5; n34 f | 11712 |
| | 3.57 emb; ±1.32; r1-6; n21 f | 965 |
| | r3.64-4.78 cl; r3.63-4.86 implantations; n3 yr | 2012 |
| | 3.64; sd0.23; n12 1st lt | 2014 |
| | 4 emb; n1 f | 4897 |
| | 4.0 emb; r3-5; n3 f; 4.66 cl | 1727 |
| | 4.0 emb, ps; n4 subadult | 7806 |
| | 4.06 cl; ±0.83; r3-5; n17 f | 965 |
| | 4.1-5.3 cl | 5807 |
| | 4.3 yg; n17 nests | 4276 |
| | 4.39 emb; ±0.28 | 10384 |
| | 4.4 emb; r1-8; n40 f | 9687 |
| | 4.4; mode 4; r3-9; n109 lt | 1386a |
| | 4.5 emb; r4-5; n2 f | 1290 |
| | 4.5 emb; r4-5; n2 f | 3703 |
| | 4.5 emb; r4-5; n2 f | 6298 |
| | 4.5; r4-5; n2 lt | 3703 |
| | 4.5 emb; r2-7; n22 f | 4276 |
| | 4.64 yg; r3-9; n11 lt | 586 |
| | 4.7 ps; r3-6; n5 f | 2891 |
| | 4.7 emb; r2-6; n21 f | 6515 |
| | 4.797 cl; n142 lt | 9382 |
| | 4.8 emb; 1st lt; 7.1 emb; 2nd lt | 5864 |
| | 4.8 yg; r1-7; mode 1-2; n35 lt, 7 f | 9877 |
| | 4.9 yg; n13 nests | 2891 |
| | 4.92; se0.13; mode 5; r2-9; n140 lt; 4.95 emb; se0.29; n20 f | 9250 |
| | 4.98 emb, yg; mode 4/5; r2-7; n44 f | 893 |
| | 5 emb; n1 f | 575 |
| | 5 yg ea; n2 nests | 169 |
| | 5.04; n27 lt | 10908 |
| | 5.06 emb; ±0.47 | 10384 |
| | 5.2; mode 5; r3-7; n38 lt | 1386a |
| | 5.22 emb; r4-7; n9 f | 169 |
| | 5.25 emb; ±0.42 | 10384 |
| | 5.31 emb; ±0.68; n~281 f | 6576 |
| | 5.34 emb; ±0.53 | 10384 |
| | 5.4 emb, yg; r4-7; n12 lt | 4499 |
| | 5.42 yg; r3-8; n26 lt | 892 |
| | 5.5 emb, ps; n124 f adult | 7806 |
| | 5.58 swellings; se0.14; 5.10 emb; se0.15; r2-8; n106 f | 10893 |
| | 5.86; n85 lt | 2014 |
| | 6 emb; n1 f | 2548 |
| | 6 emb; n1 f | 6166 |
| | 6 emb; r2-10; n82 f | 6371 |
| | 6.4 emb; r3-10; n8 f | 5405b, |
| | 6.5 emb; r3-8; n13 f | 2891 |

| | | |
|---|---|---|
| | 6.8 *emb*; n24 lt | 5003a |
| | 7 *emb*; n1 f | 2228a |
| | 8 emb; n1 f | 261a |
| | 9 yg; n1 nest | 585 |
| | 10 emb; n1 f | 6371 |
| | 12 *emb*; n1 f | 5003a |
| | 12 yg; n1 nest | 5802 |
| *Sylvilagus idahoensis* | 5.89 ps, emb; r4-8; n9 f | 3277 |
| | 5.93 *emb*; r4-8; n14 f | 5254a |
| | max 6 | 5146 |
| | 6 emb *ea*; n3 f | 8126 |
| | 6.2-6.3 ova; r5-8; n10 f | 3277 |
| *Sylvilagus nuttallii* | 2 yg | 2088 |
| | 3 emb; n1 f | 502 |
| | 3.5; r3-4; n2 lt | 10954 |
| | 4.6 emb; se0.2; r2-6; 4.3 live *emb*; se0.1; r1-6; 31 f | 8717 |
| | 4.7 emb; r4-5; n3 f | 2550 |
| | 5.0 *emb*; r4-6; n2 f | 1119 |
| | 5.0 cl; *se*0.2; r3-8; n31 f | 8717 |
| | 5.9 *emb*; r4-8; n14 f | 4034 |
| | 6 *emb* ea; n2 f | 4034 |
| | 6.1 *emb*; r4-8; n8 f | 8126 |
| | 6.62 emb; r5-8; n15 f | 5254a |
| | 7.7 cl; r7-8; n3 f | 4034 |
| | max 8 | 5146 |
| *Sylvilagus palustris* | 2.82 ps; se0.093; n121 f | 4921 |
| | mode 3-5 | 10866 |
| | 3.09 cl; se0.054; n132 f | 4921 |
| | 4.4 *emb*; n4 f | 976 |
| *Sylvilagus transitionalis* | 5.0; n3 lt, 1 f | 10731 |
| | 5.2 emb; n5 lt | 5003a |
| | 5.2 emb; r3-8; n19 f | 1719 |
| **SEXUAL MATURITY** | | |
| *Lepus americanus* | f: 1st mate: yr of their birth | 11228 |
| | f: *est* 90-100 d; 1060 g probably conceived 2 mo; 400 other juv f did not breed | 5622 |
| | f: 7-8 mo | 8798 |
| | m: 5-7 mo | 8798 |
| | f/m: 1.2-2.3 yr | 5624 |
| *Lepus californicus* | f: 7-8 mo | 8798 |
| | m: 5-7 mo | 8798 |
| *Lepus capensis* | f: rapid growth of ovaries: 3-3.5 kg | 4981 |
| | f: 1st cl: 4 mo; mode 6-7 mo | 1307 |
| | f: 1st conceive: 5 mo | 6452 |
| | f: 7-8 mo | 8798 |
| | m: epididymal sperm: 3.2 mo | 6452 |
| | m: 4 mo; @2.0-2.5 kg | 3346 |
| | m: 5-7 mo | 8798 |
| | f/m: 1st repro: 5-6 mo | 8359 |
| | f/m: 1st repro: 6 mo; born *June*; 9-12 mo; born Jan-Apr | 6909 |
| | f/m: 6-9 mo; fully grown: 1.25-1.5 yr | 3571 |
| | f/m: 6.5 mo | 1452 |
| | f/m: 8 mo | 5760 |
| *Lepus nigricollis* | f: 6 mo | 8501 |
| *Lepus timidus* | f: 7-8 mo | 8798 |
| | m: 5-7 | 8798 |
| | f/m: 8-9 mo | 3571 |
| *Lepus townsendii* | f/m: 1st mate: ~1 yr | 5239 |
| *Oryctolagus cuniculus* | f: 1st mate: 6 mo | 11887 |
| | f: 1st birth: 4 mo; n1 | 7701 |
| | f: 120-150 d | 8798 |
| | f: 5 mo | 11478 |
| | m: spermatogenesis: 7 mo | 293 |
| | m: 9 mo | 11478 |

| | | |
|---|---|---|
| | f/m: 1st mate: 4 mo | 1213 |
| | f/m: 5-8 mo | 8053 |
| | f/m: 6 mo | 4645 |
| | f/m: 6-8 mo | 3571 |
| *Romerolagus diazi* | f: 1st mate: 5-8 mo; 410-650 g | 5007 |
| | f: 403 g | 6569 |
| | m: testes descend: ~5 mo; n5 | 5007 |
| | m: 1st mate: 6.5 mo; 440 g | 5007 |
| | m: 543 g | 6569 |
| *Sylvilagus aquaticus* | m: sperm: 3-6 mo | 2013 |
| | f/m: 23-30 wk | 10222 |
| *Sylvilagus audubonii* | f: 1st mate: summer of birth | 10236 |
| | f: 1st conceive: est ~80 d | 10470 |
| | m: epididymal sperm: 107 d | 10470 |
| *Sylvilagus brasiliensis* | f/m: 125 d | 8798 |
| *Sylvilagus floridanus* | f: 1st mate: 2-5 mo | 8077 |
| | f: 1st mate: 1st summer; n1 | 4624 |
| | f: 1st mate: summer of birth | 2038 |
| | f: 1st conceive: 85, 90 d | 1655 |
| | f: 1st conceive: ~100 d | 4742 |
| | f: 1st preg: 1st summer | 2891 |
| | f: 2.5 mo | 7806 |
| | juv f, but not juv m, breed | 5722 |
| | m: epididymal sperm: 80 d | 8077 |
| | f/m: 125 d | 8798 |
| *Sylvilagus idahoensis* | m: spermatogenesis: 6 mo | 3277 |
| **CYCLE LENGTH** | | |
| *Lepus capensis* | 7 d | 1530 |
| *Lepus timidus* | estrus: 10-24 h | 8053 |
| *Oryctolagus cuniculus* | 7 d; mode 4.5-8; r4.5-41.5; n67 cycle, 34 f | 7260, 7690, 7692 |
| *Romerolagus diazi* | 13 d | 2404 |
| *Sylvilagus aquaticus* | 12 d | 6832 |
| **GESTATION** | | |
| *Lepus americanus* | 34 d | 1110 |
| | 35 d | 5624 |
| | 35.5 d; ibi | 11895 |
| | 36 d | 4006 |
| | 36.56 d; r35.85-37.23; 18 lt | 9805 |
| | 36.6 d; r34-38; n8 f | 2664 |
| | 37 d | 2458 |
| | 37 d | 2326 |
| | 37.2 d; r36-40; n37 f | 9806 |
| | 37.9 d; 19 lt | 9805 |
| | 39.375 d; n8 | 9805 |
| *Lepus californicus* | 40 d; field data | 4099 |
| | 42 d | 8798 |
| | 43 d; r41-47; n8 f | 4467 |
| *Lepus capensis* | 39.5 d; r38-41 | 10487 |
| | 40.2 d; se0.1; n66 | 6909 |
| | 41.1 d; se0.2; n43 | 6909 |
| | 41-42 d; n1 | 1005 |
| | 41.5 d; r41-42 | 1530 |
| | 42 d | 9125 |
| | 42 d | 3344 |
| | ~1.5 mo | 4645 |
| | ~6 wk; r40-44 d | 1307 |
| | mode 42,43 d; r42-69; n6 | 4562 |
| | 44 d; artificial insemination | 10318 |
| | 44-46 d | 10316 |
| *Lepus nigricollis* | ~1 mo | 8501 |
| | 42-44 d | 9175 |
| *Lepus saxatilis* | 5-6 wk | 4354a |

| | | |
|---|---|---|
| *Lepus timidus* | 44-52 d | 3571 |
| | 44-53 d | 7433 |
| | ~45 d | 10294 |
| | est 46 d | 265 |
| | 50 d; ppe to birth | 11398 |
| | 50 d; r48-51 | 8053 |
| | ~50 d | 3348 |
| | ~50 d; 1st cl to 1st yg seen | 8233 |
| | 51 d; se~0.2; n40 | 8323 |
| | 60 d | 6441 |
| *Lepus townsendii* | 1 mo | 6445 |
| | 36-43 d | 6445 |
| | 42-43 d; field data | 5865 |
| *Oryctolagus cuniculus* | implant: 7 d | 2162 |
| | implant: 7 d | 1214 |
| | 28-30 d | 1214 |
| | 28-31 d; max 37 | 3571 |
| | min 29 d | 1175 |
| | 30 d | 10763 |
| | 30-31 d | 7510 |
| | 30-32 d | 7690 |
| | 30-37 d | 7433 |
| | 30.27 d; r28-33; n191 lt, 91 f | 7260 |
| | 33 d | 196 |
| *Poelagus marjonita* | ~5 wk | 5760 |
| *Romerolagus diazi* | 1 mo | 9231 |
| | 38 d; n1 | 1676 |
| | 38-40 d; n1 | 2819, 2820 |
| | 39 d ea; n3 | 1677 |
| | 39.8 d; r39-41; n20 | 6953a, 6953b |
| | 40 d ea; n3 | 2404 |
| *Sylvilagus aquaticus* | 35-39 d; n18 lt, 9 f; assuming immediate ppe | 10222 |
| | mode 37 d 75%; r36-38; n8 lt | 4920 |
| | 39-40 d; n3 | 5099 |
| *Sylvilagus audubonii* | 26-30 d | 5146 |
| | ≤28 d; n3 | 2550b |
| *Sylvilagus bachmani* | ~27 d; n1 | 7561 |
| | 27 d; ±3 | 1722 |
| *Sylvilagus brasiliensis* | 26-30 d | 8798 |
| *Sylvilagus floridanus* | 25-28 d | 9251 |
| | 26-28 d; interval between peaks of mating | 3116 |
| | 26.5 d; n1 | 4623 |
| | 26.5 d; r25-28; n2 | 5003a |
| | 26.67; mode 26; r26-28; n6 | 6831 |
| | 29.3 d | 4499 |
| | >29-<36 d | 1386a |
| *Sylvilagus idahoensis* | 39 d; field data | 3277 |
| *Sylvilagus nuttallii* | 28-30 d | 2088 |
| *Sylvilagus transitionalis* | 28 d; n1 | 5003a |
| **LACTATION** | | |
| *Lepus americanus* | solid food: immediately | 2088 |
| | solid food: 7-9 d | 8798 |
| | solid food: 10 d | 5210 |
| | solid food: 11 d | 4006 |
| | solid food: 10-12 d | 9805 |
| | wean: 14-21 d | 8798 |
| | wean: max 3 wk | 2139 |
| | wean: 25-28 d | 9805 |
| | wean: 30 d | 8005a |
| | independent: ~6 wk | 1871a |
| *Lepus californicus* | solid food: 7-9 d | 8798 |
| | wean: 14-21 d | 8798 |
| | wean: 2-4 wk | 1871a |

| | | |
|---|---|---|
| *Lepus capensis* | solid food: 7 d | 6424 |
| | solid food: 7-9 d | 8798 |
| | solid food: 2 wk | 4645 |
| | wean: 14-21 d | 8798 |
| | wean: 1 mo | 5760 |
| | independent: ~1 mo | 10158 |
| | wean: 3 wk | 6908 |
| | wean: 3 wk; max 4 wk | 3571 |
| | wean: mode ~4 wk | 1306 |
| | wean: 8 wk | 8053 |
| | wean: ~2 mo | 4645 |
| *Lepus timidus* | den emergence: 2-4 d | 8053 |
| | solid food: 7-9 d | 8798 |
| | solid food: 8-9 d | 4645 |
| | wean: 14-21 d | 8798 |
| | wean: 3 wk | 3571 |
| *Lepus townsendii* | solid food: 15 d | 1871a |
| *Oryctolagus cuniculus* | den emergence: 21 d | 8798 |
| | solid food: 21 d | 8798 |
| | wean: 22-25 d | 8798 |
| | independent: 25-28 d | 8798 |
| | wean: 4 wk | 3571 |
| | wean: 31 d | 7894 |
| *Romerolagus diazi* | solid food: 15-16 d | 2819, 2820 |
| | wean: 3 wk | 6953b |
| | independent: 25-30 d | 2819, 2820 |
| *Sylvilagus audubonii* | leave nest: ≤2 wk | 1871a |
| | wean: 3-4 wk | 1871a |
| | wean: est 4 wk | 2550b |
| *Sylvilagus aquaticus* | leave nest: 10-17 d; still nursed | 10222 |
| *Sylvilagus bachmani* | nurse: 2-3 wk | 7561 |
| *Sylvilagus brasiliensis* | den emergence: 14 d | 8798 |
| | solid food: 14-18 d | 8798 |
| | independent: 25-28 d | 8798 |
| *Sylvilagus floridanus* | den emergence: 14 d | 8798 |
| | solid food: by 2 wk | 10908 |
| | solid food: 14-16 d | 2891 |
| | solid food: 14-18 d | 8798 |
| | wean: 14.7 d; r14-15; n3 | 5003a |
| | wean: ~15 d | 443a |
| | nurse: 20-25 d | 2891 |
| | independent: 25-28 d | 8798 |
| *Sylvilagus nuttallii* | independent: 4-5 wk | 1871a |
| *Sylvilagus transitionalis* | wean: 16 d; n1 | 5003a |
| | leave nest: 16 d | 10731 |
| **INTERLITTER INTERVAL** | | |
| *Lepus americanus* | 34.4 d | 7243 |
| | 35.5 d | 11895 |
| | 37 d | 6132 |
| | 39 d; r38-40; n2 | 3899 |
| | 38.4 d | 2672 |
| *Lepus capensis* | 25-42 d | 11058 |
| | 28-56 d | 9125 |
| | 30.25 d; r12-38; n4 | 1452 |
| | 38 d; r35-39 | 1530 |
| | 40.2 d; mode 38 (24%); r26-59; n108 | 6909 |
| | ~48 d | 1307 |
| *Lepus timidus* | 50.1 d; sd3.5; r46-54; n8 | 307 |
| *Oryctolagus cuniculus* | 29 d; min ibi | 1175 |
| *Sylvilagus bachmani* | 29.5 d; n2 | 1722 |
| *Sylvilagus floridanus* | 1 mo | 4742 |
| | 30.5 d | 1386a |
| *Sylvilagus idahoensis* | 39 d | 3277 |

| LITTERS/YEAR | | |
|---|---|---|
| *Caprolagus hispidus* | 2-3 | 3759 |
| *Lepus americanus* | 1-4 | 2326 |
| | 1-5 | 5623 |
| | min 2 | 2550 |
| | 2; CO; >2; UT | 2672 |
| | 2.3 | 4037 |
| | 2.60; r1-3 | 11895 |
| | 2.75 | 9321 |
| | 2.75-3 | 9806 |
| | 2.9-3.5 | 2664 |
| | 3 | 7872 |
| | 3.27 | 7243 |
| | 3.75, 3.92; n2 yr; ~4 central WI; ~3 n WI | 6132 |
| | max 4 | 9805 |
| | ~4 | 5624 |
| *Lepus californicus* | 3.6; r3-5 | 4099 |
| *Lepus capensis* | 1.77 | 225 |
| | 2/summer | 8053 |
| | mode 3,4; r1-6 | 6909 |
| | 2; central Russia; 4 Caucasus | 5944 |
| | 2-3 | 6452 |
| | 2-4 | 3572 |
| | 2.2; 3.0; 2.95; r1-4; 3 locations | 3540 |
| | 2.46/season | 226 |
| | 3 | 9497 |
| | 3 | 9125 |
| | 3 | 8370 |
| | 3-4 | 5944 |
| | 3-4 | 8476 |
| | 3-4; max 5 | 3571 |
| | 4.29 | 2576 |
| | 4.8; field data | 3344 |
| | 7.73 | 3349 |
| | max 5 | 11058 |
| *Lepus nigricollis* | 8-9 | 8730 |
| *Lepus saxatilis* | min 2 | 11253 |
| | 8.7 | 3349 |
| *Lepus timidus* | 1 | 265 |
| | 1 | 4548 |
| | 1 | 8233 |
| | est 1 | 5146 |
| | 2-3 | 6441 |
| | 2-3 | 307 |
| | 2.6 | 3348 |
| | 3 | 4711 |
| *Lepus townsendii* | 2; max 3-4 | 5865 |
| | est 3.29; max 4 | 5239 |
| *Oryctolagus cuniculus* | 1-3 | 10386 |
| | 2.8/7 mo | 11887 |
| | 3-5; r3-6 | 3571 |
| | 4 | 1213 |
| | ~4 | 1178 |
| | 4-5 | 4645 |
| | 4-5; max 7 | 8053 |
| | 5-6 | 7433 |
| | 5-7 | 8798 |
| *Sylvilagus aquaticus* | 3.09; r2-5; n34 lt, 11 f | 10222 |
| *Sylvilagus bachmani* | 3-4 | 5146 |
| *Sylvilagus brasiliensis* | 4-5 | 8798 |
| | 4.7; enclosure | 2816 |
| *Sylvilagus floridanus* | 1-4/season; n11 f | 4499 |
| | mode 2; max 4 | 10908 |
| | 2.9 | 9250 |

| | | |
|---|---|---|
| | 3-4 | 1386a |
| | 3-4 | 2891 |
| | 4.7 | 10822 |
| | 4.81 | 965 |
| | 7-8 | 2014 |
| | max 7 | 3117 |
| | max 7 | 4742 |
| | 7; n1 | 6832 |
| *Sylvilagus idahoensis* | 2 | 3277 |
| *Sylvilagus nuttallii* | 2-3 | 2088 |
| | min 3/season | 2550 |
| | 4-5 | 8717 |
| **SEASONALITY** | | |
| *Caprolagus hispidus* | mate: Jan-Mar, perhaps to June; Assam, India | 3759 |
| | juveniles found: June-July; Assam to Uttar Pradesh, India | 3759 |
| *Lepus alleni* | mate: Dec-Sept; NM, n Mexico | 1474 |
| | preg f: Jan-Oct; AZ | 11365 |
| *Lepus americanus* | spermatogenesis: Feb-Aug; MT | 1111 |
| | testes: large Apr-July, small Aug; Rochester, ALB, Canada | 2326 |
| | testes: large May-July, small Nov-early Feb; Mautoulin I, ONT, Canada | 7872 |
| | mate: Mar-?; ME | 9805 |
| | mate: Mar-Aug; NEW, Canada | 2664 |
| | mate: mid Mar-?; Canada | 2139 |
| | mate: Apr-?; UT | 2672 |
| | mate: Apr/May-?; CO | 2672 |
| | mate: Dec; NEW, Canada | 2663 |
| | conceive: Mar-July; central ALB, Canada | 5624 |
| | conceive: Apr-?; ALB, Canada | 7243 |
| | conceive: Apr-July; MI | 1111 |
| | conceive: Apr-July; Mautoulin I, Lake Huron, ONT, Canada | 7872 |
| | preg f: Feb-July; ID | 3510 |
| | preg f: Mar-Aug; MN | 143 |
| | preg f: Apr; Rochester, ALB, Canada | 2326 |
| | preg f: Apr-June; NS, Canada | 11895 |
| | preg f: Apr-July; ALB, Canada | 9321 |
| | preg f: Apr-Sept; Canada | 2139 |
| | preg f: May-June; YUK, Canada | 12049 |
| | preg f: June-July; n BC, Canada | 10597 |
| | preg f: Nov; n1; NH | 9968 |
| | preg f: end Nov-Jan; NEW, Canada | 2663 |
| | birth: Apr-Aug, peak June-July; WY | 1871a |
| | birth: Apr-Aug; WI | 5210 |
| | birth: Apr-Aug; ME | 9805, 9806 |
| | birth: Apr-Oct; ONT, Canada | 6706 |
| | birth: May-June; NM | 502 |
| | birth: May-July; AK | 8005a |
| | birth: May-Sept; MI | 1111 |
| | repro: summer; 9-21 wk; YUK, Canada | 6014 |
| *Lepus californicus* | mate: Dec-Sept; s N Am | 1474 |
| | preg f: Jan-Apr; UT | 5254a |
| | preg f: Jan-Aug, peak Apr; n94; KS | 1322 |
| | preg f: Jan-Aug, peak Feb-Apr; UT | 4099 |
| | preg f: Jan-Sept; s AZ | 11365 |
| | preg f: Feb-early Mar; se CO | 3086 |
| | preg f: Feb-Mar; n2; CA | 11228a |
| | preg f: Apr, June; n2; Coahuila, Mexico | 522 |
| | preg f: Apr, June, Dec-Feb; CA | 8126 |
| | preg f: June; n1; NE | 5405b, |
| | preg f: July; n1; Durango, Mexico | 523a |
| | preg f: yr rd, except Oct; sw TX | 4332 |
| | preg f: yr rd, peak Jan-Apr; CA | 6335 |
| | birth: yr rd; w US | 5146 |
| | lact f: Jan, Mar-Apr; few data; Coahuila, Mexico | 522 |

| | | |
|---|---|---|
| | lact f: Apr-Sept; NM | 502 |
| | lact f: July; n2; Durango, Mexico | 523a |
| | repro: mid Feb-mid June; ID | 3197 |
| *Lepus capensis* | spermatogenesis: reactivated end Nov; Netherlands | 1307 |
| | spermatogenesis: Dec-Aug; s Kazakhstan | 8370 |
| | testes: large Feb-Aug, small Sept-Oct; Germany | 7446 |
| | testes: small Sept-Dec; Norfolk, UK | 6453 |
| | testosterone: low July-Feb; Norfolk, UK | 6453 |
| | estrus: Jan-Aug; Germany | 7433 |
| | mate: Jan-June; Biserni Otak I, Yugoslavia | 11058 |
| | mate: Jan-Aug; Leeds, UK | 4922 |
| | mate: Jan-Aug; Germany | 3571 |
| | mate: Jan-Sept; Norfolk, UK | 6452 |
| | mate: Feb & earlier; Moscow, Russia | 8053 |
| | mate: Mar/Apr-?; n Eur USSR | 4645 |
| | mate: Jan-?; s Eur USSR | 4645 |
| | mate: July/Aug-Apr/May; Argentina | 2576 |
| | mate: Aug-Jan; Patagonia, Argentina | 226 |
| | mate: peak Oct, none Feb-July; Snowy Plain, Australia | 3778 |
| | mate: Dec-Sept; captive | 1534 |
| | mate: end Dec-?; Netherlands | 1307 |
| | mate: est yr rd; NEW, Canada | 3346 |
| | ovulation: Jan-Aug; captive | 1533 |
| | conceive: Mar-mid Sept; Germany | 9125 |
| | preg f: Jan; n4; Iran | 6293 |
| | preg f: Jan-Aug; ONT, Canada | 9080 |
| | preg f: Jan-Sept; Scotland, UK | 4709 |
| | preg f: 95% Feb-June, 10% Aug, none Dec; s ONT, Canada | 9080 |
| | preg f: >50% Feb-July, <40% Aug-Oct; no data Nov-Jan; Germany | 7446 |
| | preg f: Feb, June, Nov; few data; Botswana | 10154 |
| | preg f: Feb-Aug/Sept; Germany | 10420 |
| | preg f: mid Mar-Aug; Bisemi Otak I, Yugoslavia | 11058 |
| | preg f: May; n1; Kenya | 4925 |
| | preg f: Aug; n1; El Vergel, Malleco Province, Chile | 4054 |
| | preg f: yr rd, 90% Aug-Feb; New Zealand | 3344, 3345, 3346 |
| | preg f: yr rd, peak Aug-Oct; Riverine District, Australia | 3778 |
| | preg f: yr rd; n12 yr; Scotland, UK | 4712 |
| | preg f: Jan-Sept; n1 yr; Poland | 4712 |
| | birth: Jan-June; Karakun, USSR | 9497 |
| | birth: end Jan-Nov; captive | 6909 |
| | birth: Feb-Aug; captive | 10487 |
| | birth: Feb-Sept/Oct; s Sweden | 3540 |
| | birth: Feb-Oct; Germany | 3571 |
| | birth: Mar-July; s Moravia, Czechoslovakia | 4981 |
| | birth: Mar-Aug; Germany | 7433 |
| | birth: Mar-Aug; zoo | 1452 |
| | birth: Mar-Sept; Moscow, Russia | 8053 |
| | birth: Apr-May, summer; no data fall, winter; Turkestan | 9774 |
| | birth: Oct-Apr; Angola | 11697 |
| | birth: yr rd, peak Oct-Feb; S Africa | 10158 |
| | yg found: Apr-early May; Baluchistan | 9175 |
| | lact f: Feb; n1; El Vergel, Malleco Province, Chile | 4054 |
| | lact f: Feb, July, Nov-Dec; n8; Transvaal, S Africa | 8965 |
| | lact f: Aug-Sept; n2; Iran | 6293 |
| | lact f: < 10% Dec-Jan; Germany | 7446 |
| | repro: Jan-mid June; Karakumy desert, Turkmenia | 9499 |
| | repro: Feb-Sept; Byelorussia | 3572 |
| | repro: yr rd, peak spring-summer; Caucasus, Russia | 5944 |
| | repro: mid winter-fall; USSR | 5944 |
| | repro: Jan-Sept; captive | 6909 |
| *Lepus nigricollis* | testes: large Mar-Sept; India | 8730 |
| | preg f: Apr; n2; Nepal | 7382 |
| | preg f: yr rd, peak July-Sept; India | 8730 |

| | | |
|---|---|---|
| | birth: spring-summer; Pakistan | 9175 |
| | birth: yr rd; Sri Lanka | 8501 |
| | lact f: Sept; n1; Nepal | 7382 |
| | repro: not seasonal; Sri Lanka | 8500 |
| *Lepus saxatilis* | preg f: Jan, Mar-Apr, Aug, Nov-Dec; n13; Transvaal, S Africa | 8965 |
| | preg f: July; n2; Kenya | 4925 |
| | preg f: yr rd; Botswana | 10154 |
| | birth: Apr, Oct-Nov; Namibia | 9945 |
| | birth: almost yr rd; Transvaal, S Africa | 8965 |
| | birth: yr rd, peak Sept-Feb; S Africa | 10158 |
| | lact f: Feb, May, Oct; n8; Transvaal, S Africa | 8965 |
| | repro: yr rd | 1149 |
| *Lepus timidus* | estrus: mid Feb-July; Europe | 8053 |
| | estrus: Mar-Aug; Alps, Europe | 7433 |
| | mate: Jan-June; Scotland, UK | 4710 |
| | mate: early-mid Feb; USSR | 4645 |
| | mate: Mar-June; Germany | 3571 |
| | conceive: Feb-Aug; e Scotland, UK | 4709 |
| | conceive: Mar-?; Sweden | 307 |
| | conceive: Apr; w AK | 265 |
| | conceive: peak Apr; sw NEW, Canada | 4548 |
| | preg f: Jan-Sept, peak Mar-June; ne Scotland, UK | 4710a |
| | preg f: Mar-June; ne Scotland, UK | 3348 |
| | birth: Mar-Aug; ne Scotland, UK | 4710, 4711 |
| | birth: Mar-Sept; Europe | 8053 |
| | birth: Apr/May-July/Aug; Germany | 3571 |
| | birth: Apr-Aug; Alps, Europe | 7433 |
| | birth: May-early July; captive | 10294 |
| | birth: May-end Aug/Sept; Norway | 11398 |
| | birth: end May-early June; w AK | 265 |
| | birth: peak end May-mid June; sw NEW, Canada | 4548 |
| | birth: June; NWT, Canada | 8233 |
| | birth: June-July; N America | 1474 |
| *Lepus townsendii* | testes: large Feb-June; ND | 5239 |
| | mate: Feb-Mar; IA | 5865 |
| | preg f: end Feb-July; WY | 9228a |
| | preg f: Mar-July; UT | 5254a |
| | preg f: May; n3; SD | 261a |
| | preg f: May-mid July; CO | 680 |
| | preg f: June; n1; ND | 3703 |
| | preg f: July; n1; se MT | 6166 |
| | preg f: July-Aug; n3; NE | 5405b$_1$ |
| | preg f: Aug; n2; CO | 1649a |
| | birth: May-Aug; NM | 502 |
| | lact f: July; n1; se MT | 6166 |
| | repro: Feb-early July perhaps longer; IA | 5865 |
| *Oryctolagus cuniculus* | sperm: yr rd; sw Australia | 11620 |
| | spermatogenesis: yr rd; n Cambridgeshire, England, UK | 1178 |
| | testes: large Mar-Aug; s Sweden | 291 |
| | testes: large Apr-May, low Oct-Nov; n Cambridgeshire, England, UK | 1178 |
| | testes: large May, small Oct; s Sweden | 293 |
| | testes: large Sept, small Jan; New Zealand | 11478 |
| | testes: large spring, small summer; France | 9228 |
| | m accessory glands: peak May-June; Sweden | 2216 |
| | ovaries: large spring, small summer-fall; France | 9228 |
| | uterine mass: peak May-June, low Dec; s Sweden | 293 |
| | estrus: Feb-July; Wales, UK | 1213 |
| | mate: Jan-June; w Wales, UK | 8491 |
| | mate: Feb-June; Netherlands | 11422 |
| | mate: Feb-Aug; Germany | 7433 |
| | mate: Feb/Mar-Sept; Germany | 3571 |
| | mate: early Mar-Apr; sw Australia | 5746 |
| | mate: Mar/Apr-Oct; WA, Australia | 5745 |

| | | |
|---|---|---|
| | mate: mid Apr-mid Aug; s Sweden | 2217 |
| | mate: June-Nov; New Zealand | 11478 |
| | mate: July-May; sub Antarctic | 9918 |
| | mate: Aug-Mar; Macquarie I, sub Antarctic | 10044 |
| | mate: Nov-May, peak Mar-Apr; Sierra Morena, Spain | 10226 |
| | mate: yr rd; NSW, Australia | 11887 |
| | conceive: mid Mar/June-?; San Juan I, WA | 10386 |
| | preg f: Mar-Aug; captive | 1173 |
| | preg f: Mar/Apr-Aug; s Sweden | 291 |
| | preg f: mid June-early Nov; New Zealand | 7070 |
| | preg f: none Oct-Dec; n Cambridgeshire, England, UK | 1178 |
| | preg f: peak spring; France | 9228 |
| | preg f: yr rd, peak June-Oct; New Zealand | 11476, 11478 |
| | preg f: yr rd, high Aug-Nov, peak Oct; New Zealand | 7070 |
| | birth: Feb-July, peak Apr-June; Oxfordshire, s England, UK | 2084 |
| | birth: Mar-July, peak May; Netherlands | 11422 |
| | birth: Mar-July; s Sweden | 293 |
| | birth: Mar-Oct; Germany | 3571 |
| | birth: Mar-Oct; Germany | 7433 |
| | birth: peak May; Scotland, UK | 4619 |
| | birth: mid June-?; NSW, Australia | 8228 |
| | repro: June-Dec; captive, Australia | 7701 |
| *Poelagus marjorita* | mate: est yr rd; Angola, Uganda | 5760 |
| *Pronolagus randensis* | birth: est yr rd; S Africa | 10158 |
| *Romerolagus diazi* | mate: Mar-May; zoo | 2404 |
| | mate: Mar-July; Mexico | 9231 |
| | mate: Dec-July, peak Jan-Apr; Mexico | 6569 |
| | mate: yr rd; captive | 5007 |
| | preg f: May-June; Mexico | 6387 |
| | birth: May; n2; zoo | 6478 |
| | birth: peak spring; captive | 5007 |
| | lact f: Feb-Dec; Mexico | 1677 |
| | repro: Mar-June; Mexico | 9231 |
| *Sylvilagus aquaticus* | anestrus: Nov-Jan; MD | 10861 |
| | mate: Jan-Sept; mid-s US | 1474 |
| | mate: Feb; n AL | 4740 |
| | preg f: Feb-Sept; captive | 10222 |
| | preg f: Feb-?; MS | 8214 |
| *Sylvilagus audubonii* | sperm: yr rd, low Sept-Nov; AZ | 10470 |
| | testes: low Sept-Oct; AZ | 10236 |
| | mate: fall-spring; central CA | 3280 |
| | preg f: Jan-Aug; AZ | 10236 |
| | preg f: Feb-June, Dec; CA | 8126 |
| | preg f: Apr, July, Dec; few data; Mexico | 522, 523a |
| | preg f: May-July; n5; SD | 261a |
| | preg f: June; n1; Sinaloa, Mexico | 5409 |
| | preg f: July; n1; se MT | 6166 |
| | preg f: July; n2; NM | 502 |
| | preg f: July; n1; NE | 5405b, |
| | birth: yr rd; sw US | 1474 |
| | lact f: Apr, June-July; few data; Mexico | 522, 523a |
| | lact f: May; n1; NM | 502 |
| *Sylvilagus bachmani* | sperm: Oct-July, peak Jan-May; CA | 7561 |
| | mate: Jan-June; w US | 5146 |
| | mate: Jan-June; w US | 1474 |
| | preg f: Jan, Mar-June; CA | 8126 |
| | preg f: mid Feb-mid Aug; OR | 1722 |
| | preg f: Dec-June; CA | 7561 |
| | birth: Jan-May; w US | 1474 |
| | birth: est Aug; n1; OR | 11276 |
| *Sylvilagus brasiliensis* | repro: yr rd, low Mar-Apr; Venezuela | 2816 |
| *Sylvilagus floridanus* | sperm: yr rd, except Oct; OK | 935 |
| | spermatogenesis: Jan-July; VA | 9524 |
| | testes: large Feb-June/July, small July-Jan; piedmont GA | 8340 |

| | | |
|---|---|---|
| | testes: large Feb-July; IL | 9687 |
| | testes: large Feb-Aug; OK | 935 |
| | testes: large Mar-Aug, small Sept-Jan; coastal GA | 8340 |
| | testes: large Apr; VA | 9524 |
| | testes: large May, quiescent Sept-Dec; WI | 2964 |
| | testes: large end Dec-Oct; NY | 4276 |
| | epididymal sperm: Feb-Aug; IL | 2891 |
| | mate: Jan-Aug; OR | 10893 |
| | mate: Jan-Aug; MI | 10908 |
| | mate: end Jan-Aug; WI | 5210 |
| | mate: Feb-Aug, peak mid Feb, early July; se MO | 3116 |
| | mate: mid Feb-mid July, peak Apr-May; St Clements I, MD | 965 |
| | mate: mid Feb-Aug, if warm winter Dec-Oct; AL | 4742 |
| | mate: mid Feb-mid Sept; OK | 935 |
| | mate: Mar-?; IL | 9687 |
| | mate: Apr-Aug; OH | 10384 |
| | mate: yr rd; Venezuela | 8077 |
| | conceive: Jan-Sept; GA | 8342 |
| | conceive: Jan-Sept; GA | 8342 |
| | preg f: Jan; n1; IL | 6575 |
| | preg f: Jan-Feb; n1; MI | 169 |
| | preg f: Jan-Sept, peak June; KY | 1386a |
| | preg f: Feb-Aug, MO | 2014 |
| | preg f: Feb-Sept; NY | 4276 |
| | preg f: Feb-Sept; piedmont, GA | 8340 |
| | preg f: Feb-Oct; coastal, GA | 8340 |
| | preg f: Feb-Oct, peak Mar-June; GA | 8340 |
| | preg f: end Feb; captive, IA | 5864 |
| | preg f: end Feb-Aug; IL | 2891 |
| | preg f: Mar-Apr; FL | 7498a |
| | preg f: Mar-June; mountain, GA | 8340 |
| | preg f: Mar, Sept; OK | 935 |
| | preg f: peak Apr-May; VA | 6515 |
| | preg f: Mar-June; few data; Coahuila, Mexico | 522 |
| | preg f: Mar-Aug; MI | 4499 |
| | preg f: Apr-Sept; n24; CT | 5003a |
| | preg f: May-July; n8; NE | $5405b_1$ |
| | preg f: June; n1; se MT | 6166 |
| | preg f: June; n1; SD | 261a |
| | preg f: June-July; n2; ND | 3703 |
| | preg f: Nov; n1; WI | 6371 |
| | preg f: Dec-Jan; n2; PA | 1160 |
| | preg f: yr rd, low June-July; TX | 1136 |
| | birth: ?-early Sept; IL | 7290 |
| | birth: Feb-July; MD | 965 |
| | birth: end Feb-Sept, peak Apr-June; OH | 443a |
| | birth: Mar-Sept; PA | 893 |
| | birth: Mar-?; captive | 9250 |
| | lact f: Mar-June; n4; FL | 7498a |
| | lact f: Apr; few data; Coahuila, Mexico | 522 |
| | lact f: Apr-Sept; PA | 892 |
| | lact f: May, July-Aug; NE | $5405b_1$ |
| | yg found: ?-July; MI | 169 |
| | repro: Jan-Sept; AL | 586 |
| | repro: end Jan-early Oct; NC | 586 |
| *Sylvilagus idahoensis* | spermatogenesis: Jan-July, none Aug-Dec; no data Apr-Aug; ID | 3277 |
| | mate: spring-early summer; w US | 5146 |
| | conceive: end Feb-Apr; ID | 3277 |
| | preg f: end Feb-Mar; UT | 5254a |
| | preg f: end Mar-May; ID | 4034 |
| | birth: June-July; nw US | 1474 |

| | | |
|---|---|---|
| *Sylvilagus nuttallii* | testes: large Jan-July; OR | 8717 |
| | mate: end Feb-early July; OR | 8717 |
| | preg f: end Feb-July; OR | 8717 |
| | preg f: Apr-July; ne CA | 8126 |
| | preg f: Aug; n1; NM | 502 |
| | birth: Apr-July; BC, Canada | 2088 |
| *Sylvilagus palustris* | mate: Feb-Sept; se US | 1474 |
| | preg f: Feb; GA | 10866 |
| | preg f: yr rd; s FL | 4921 |
| | lact f: Nov; GA | 10866 |
| *Sylvilagus transitionalis* | preg f: Apr-June; n5; CT | 5003a |