Supporting Information

Carlo and Stevens 10.1073/pnas.1221398110




Fig. S1. Examples of histological material used for the cell counting in this study. Panel of images shows superior parietal lobule (SPL) area in four species (mouse, rat, cat, and monkey, from top to bottom) at low and high magnification (2× and 100× with oil). (*Left*) Low-magnification images; triangles indicate the region in which columns were outlined. (*Right*) High-magnification images; arrows indicate neurons—identified by the presence of a nucleolus—and asterisks mark glial cells, which are much smaller and have darkly stained punctuation. (Scale bars: *Left*, 500 μm; *Right*, 25 μm.)

Table S1. Species characteristics

Species	Common name	Specimen ID	Sex	Age, y	Cut section thickness, μm	Brain regions
Macaca mulatta	Rhesus macaque	M203	M	11.2	40	M1, S1, P, T, F
M. mulatta	Rhesus macaque	M104	M	6.5	40	M1, P
M. mulatta	Rhesus macaque	M105	M	2.8	40	S1, T
Felis catus	Cat	C105	Unknown	Unknown	40	M1, S1, P, T, F
Rattus sp.	Rat	R105	M	Adult	40	M1, S1, P, T
Mus sp.	Mouse	MS305	F	Adult	40	M1, S1, P, T
Mus sp.	Mouse	MS205	F	Adult	40	M1, S1, P, T

F, frontal; M1, primary motor cortex; P, parietal; S1, primary somatosensory cortex; T, temporal.