Comparative analysis of the *Eulemur* vocal repertoire: a dynamic time warping approach.

Authors: Marco Gamba[a], Olivier Friard[a], Isidoro Riondato[a], Roberta Righini[a], Camilla Colombo[a], Longondraza Miaretsoa[b], Valeria Torti[a], Bakri Nadhurou[a], Cristina Giacoma[a].

[a] Department of Life Sciences and Systems Biology, University of Torino, Italy.
[b] Faculté de Sciences, University of Mahajanga, Madagascar.

Corresponding author: Marco Gamba, Department of Life Sciences and Systems Biology, University of Torino, Via Accademia Albertina 13, 10123 Torino, Italy.
Tel. +390116704560
Fax. +390116704508
Email: marco.gamba@unito.it




Fig. 8 Spectrogram of a Long Grunt of *Eulemur macaco* (1a) with temporal parameters overlaid (Ptmin = percentage of signal duration before minimum F0; Ptmax = percentage of signal duration before maximum F0). Formant analysis is shown on the left on a LPC-smoothed frequency spectrum (1b, F1 = average first formant; F2 = average second formant; F3 = average third formant) and fundamental frequency (F0) analysis at the bottom of the figure (1c, StartF0 = F0 at the beginning of the signal; MinF0 = minimum F0; MaxF0 = maximum F0; EndF0 = F0 at the end of the signal)