Supplementary Material. Tables S1, S2, and S3 presenting the numerical data used for the analyses in this paper

Exploring the mammalian sensory space: co-operations and trade-offs among senses
Sirpa Nummela, Henry Pihlström, Kai Puolamäki, Mikael Fortelius, Simo Hemilä, Tom Reuter
Journal of Comparative Physiology A

Corresponding author: Sirpa Nummela, Department of Biosciences, University of Helsinki. Email: sirpa.nummela@helsinki.fi

Table S1 Log-transformed values of nasal-caudal (or nasal-temporal) diameter of spread out retina (RD, in mm), axial eye diameter (ED, in mm), square root of olfactory epithelium (OE) area (in mm^2), square root of cribriform plate (CP) area (in mm^2), basilar membrane length (BML, in mm), cochlear size (CS, in mm), and square root of tympanic membrane (TM) area (in mm^2). For species described in several studies, the values are averaged. Epithelium areas represent the sum of both right and left sides of the nose cavity (see Pihlström et al. 2005). Note that species in brackets are not included in Fig. 1, or in the regression line calculations. Data on CS from Spoor et al. (2002). Other data from: (1) Lauruschkus (1942); (2) Allison and Turner Warwick (1949); (3) Negus (1958); (4) von Békésy (1960); (5) Stone (1965); (6) Gurtovoi (1966); (7) Mulvaney and Heist (1970); (8) Kolb (1971); (9) Adams (1972); (10) Bhatnagar and Kallen (1975); (11) Hughes (1975), (12) Kolb (1975); (13) Güntherschultze (1979); (14) Söllner and Kraft (1980); (15) Gross et al. (1982); (16) Ritland (1982); (17) Bruns et al. (1983-84); (18) Henderson (1985); (19) Pye (1985-86); (20) Sigmund and Sedláček (1985); (21) West (1985); (22) Pettigrew et al. (1988); (23) Stone and Halasz (1989), (24) Fay (1992); (25), Mass and Supin (1992); (26) Peichl (1992); (27) Müller et al. (1993); (28) Gonzalez-Soriano et al. (1997); (29) Peichl and Moutairou (1998); (30) Ahnelt and Kolb (2000); (31) Ketten (2000); (32) Mass and Supin (2000); (33) Mass and Supin (2003); (34) Howland et al. (2004); (35) Rowe et al. (2005); (36) Manoussaki et al., (2008); (37) Heffner and Heffner (2010); (38) Pettigrew et al. (2010). For data sources for ED, CP, and TM, see Supplementary Material Table S2

Order Species	log RD	log ED	RD ref.	log $(OE)^{0.5}$	log $(CP)^{0.5}$	OE ref.	log BML	log CS	log $(TM)^{0.5}$	BML ref.
Didelphimorphia										
Didelphis marsupialis	1.164	0.940	30							
Monodelphis domestica				1.280	0.669	35	0.806	-	0.359	27
Notoryctemorphia										
Notoryctes typhlops							-	0.176	0.293	

Diprotodontia										
Dendrolagus dorianus	1.401	1.259	11							
Macropus rufus	1.653	1.491	11							
Proboscidea										
Elephas maximus							1.778	1.021	1.329	4, 21
Loxodonta africana	1.710	1.544	23,38							
Erinaceomorpha										
Erinaceus europaeus							-	0.380	0.618	
Erinaceus concolor				1.646	1.028	20				
Hemiechinus auritus				1.607	0.900	9				
Soricomorpha										
Crocidura russula							0.690	-	0.236	21
Neomys fodiens				0.987	0.541	14				
Sorex araneus				1.149	0.548	6,14	0.597	-	0.142	21
Talpa europaea							-	0.176	0.489	
Chiroptera										
(*Rhinolophus ferrumequinum*							1.207	-	0.176)	17, 19
Rhinolophus rouxii	0.380	0.255	22							
Macroderma gigas	1.055	0.845	22							
Megaderma lyra	0.746	0.623	22							
Taphozous georgianus	0.775	0.568	22							
Artibeus jamaicensis				1.183	0.619	10				
Artibeus cinereus	0.760	0.643	22							
Nyctophilus gouldi	0.428	0.279	22							
(*Pipistrellus pipistrellus*							-	0.255	0.129)	
Plecotus auritus				0.773	0.385	8				

Myotis blythii				0.918	0.423	6				
Myotis myotis				1.137	0.457	8				
(*Nyctalus noctula*				1.123	0.435	8	1.073	-	0.266)	19
Carnivora										
Canis lupus	1.544	1.371	26							
Canis familiaris				2.072	1.381	1	-	0.681	0.900	
Callorhinus ursinus	1.785	1.617	25							
(*Zalophus californianus*							1.735	-	0.619)	31
Odobenus rosmarus							-	0.892	0.972	
Pagophilus groenlandicus	1.802	1.623	33							
Halichoerus grypus							-	0.851	0.941	
Phoca vitulina							-	0.886	0.947	
Mustela nivalis							-	0.342	0.489	
Mustela putorius	1.072	0.929	18							
Lutra lutra							-	0.519	0.692	
Enhydra lutris	1.250	1.079	32							
Felis catus	1.543	1.342	5	1.723	0.976	1,3	1.352	0.663	0.799	24
Puma concolor							-	0.690	0.900	
Panthera leo							-	0.875	1.019	
Perissodactyla										
Equus caballus							-	0.826	0.892	
Artiodactyla										
Sus scrofa				2.230	1.644	13	-	0.653	0.897	
Capreolus capreolus				1.977	1.385	12				
Bos taurus							1.580	-	0.918	31
Capra hircus	1.627	1.415	28							

Ovis aries							-	0.716	0.840	
Rodentia										
Sciurus vulgaris							-	0.447	0.716	
Dipodomys merriami							0.991	-	0.666	19
Cricetomys gambianus	1.021	0.826	29							
Myodes glareolus				1.102	0.480	6				
Mesocricetus auratus							0.863	-	0.423	37
Peromyscus maniculatus				1.111	0.560	9				
Meriones unguiculatus							1.083	-	0.613	31
Rattus norvegicus				1.415	0.662	15	1.000	0.362	0.479	21, 31
Mus musculus				1.068	0.450	15	0.845	0.114	0.253	21
Chinchilla lanigera							1.267	-	0.902	21
Cavia porcellus							1.267	-	0.694	21
Lagomorpha										
Oryctolagus cuniculus				1.460	0.916	2,3,7,9	1.185	0.447	0.725	21
Primates										
Nycticebus coucang							-	0.447	0.679	
Saimiri sciureus							1.188	-	0.699	36
Callithrix jacchus							-	0.505	0.597	
Macaca mulatta							1.398	0.681	0.753	24
Papio cynocephalus							-	0.633	0.827	
Homo sapiens				1.527	1.060	3	1.535	0.820	0.904	4

Table S2 Body mass (in kg), axial eye diameter ("Eye", in mm), cribriform plate area ("Nose", in mm^2), tympanic membrane area ("Ear", in mm^2), and the typical habitats and diets of 119 mammalian species. Code = species code (cf. Figs. 3, 5 and 6, and Table 1). Habitats: Ae = aerial; Aq = aquatic; Ar = arboreal; F = fossorial; T = terrestrial. Diets: C = carnivore; P = piscivore; I = insectivore; H = herbivore; Om = omnivore; Other. Residuals ("Res") for eye, nose, and ear shown. Data on habitat, diet, and body mass from Nowak (1999); other data from: (1) Steindorff (1947); (2) Werner (1960); (3) Henson (1961); (4) Suthers and Wallis (1970); (5) Hughes (1975); (6) Ritland (1982); (7) Pettigrew et al. (1988); (8) Plassmann and Brändle (1992); (9) Rosowski (1992); (10) Nummela (1995); (11) Mason (2001); (12) Supin et al. (2001); (13) Howland et al. (2004); (14) Lange et al. (2004); (15) Pihlström et al. (2005); (16) Kirk (2006); (17) Nummela et al. (2004); (18) Nummela and Sánchez-Villagra (2006); (19) This study

Order *Species*	Code	Hab	Diet	Body mass	Eye	Nose	Ear	Res eye	Res nose	Res ear	Eye source	Nose source	Ear source
Monotremata													
Tachyglossus aculeatus	Ta	T	I	4.750	8.400	98.00	11.50	-0.177	-0.012	-0.178	5,13	15	9,11
Didelphimorphia													
Caluromys philander	cp	Ar	Om	0.350	8.700	32.95	4.080	0.114	0.006	-0.239	6	19	18
Didelphis marsupialis	Dm	T	Om	1.500	9.340	104.0	7.180	-0.009	0.113	-0.208	6	19	11
Marmosa robinsoni	mR	Ar	I	0.085	6.600	26.73	5.380	0.143	0.098	-0.089	6	19	18
Dasyuromorphia													
Dasyurus viverrinus	dv	T	C	1.075	10.60	55.10	15.13	0.081	0.008	-0.025	6	19	18
Diprotodontia													
Phalanger orientalis	pO	Ar	H	2.948	11.40	61.94	15.20	0.006	-0.066	-0.087	6	19	18
Trichosurus vulpecula	Tv	Ar	H	2.395	13.70	80.80	29.80	0.107	0.013	0.072	6	19	18
Macropus eugenii	me	T	H	4.250	23.70	109.2	27.00	0.285	0.022	0.014	6	19	18
Macropus rufus	mr	T	H	55.00	31.30	196.0	43.00	0.135	-0.101	-0.046	1	19	2
Tubulidentata													
Orycteropus afer	Oa	T	I	70.00	20.50	1760	93.30	-0.075	0.352	0.107	6	15	11

Afrosoricida													
Hemicentetes semispinosus	Hs	T	I	0.180	1.930	33.76	6.200	-0.470	0.076	-0.106	6	19	11
Setifer setosus	Ss	T	I	0.225	3.350	56.40	12.60	-0.254	0.166	0.034	6	19	11
Tenrec ecaudatus	te	T	I	2.000	4.380	83.36	10.37	-0.369	0.037	-0.146	6	19	11
Proboscidea													
Elephas maximus	Em	T	H	4060	33.47	4840	454.0	-0.291	0.174	0.194	1,6,13	19	10
Loxodonta africana	La	T	H	4350	45.30	10200	855.0	-0.167	0.329	0.327	6,13	19	10
Cingulata													
Tolypeutes matacus	Tm	T	I	1.295	6.500	180.4	12.00	-0.151	0.247	-0.087	6	19	11
Zaedyus pichiy	Zp	T	I	1.500	6.200	193.0	32.40	-0.187	0.247	0.119	6	15	19
Pilosa													
Cyclopes didactylus	Cy	Ar	I	0.266	10.20	28.20	8.760	0.212	-0.001	-0.055	6	19	19
Myrmecophaga tridactyla	My	T	I	28.50	10.70	900.0	58.10	-0.262	0.294	0.061	6	15	11
Erinaceomorpha													
Erinaceus europaeus	Ee	T	I	0.750	6.500	94.28	17.22	-0.094	0.160	0.027	1	15	11
Hemiechinus auritus	Ha	T	I	0.160	6.400	63.20	22.94	0.063	0.224	0.185	6	19	10
Soricomorpha													
Suncus murinus	Su	T	I	0.003	1.950	28.73	3.920	-0.032	0.441	0.053	6	19	11
Blarina brevicauda	Bb	T	I	0.023	1.140	18.15	4.330	-0.481	0.142	-0.054	6,13	19	11
Talpa europaea	tE	F	I	0.093	1.400	28.52	9.490	-0.539	0.104	0.028	6	15	10,11
Scalopus aquaticus	Sa	F	I	0.090	0.900	27.46	10.82	-0.728	0.099	0.059	6	15	3,11
Chiroptera													
Desmodus rotundus	Dr	Ae	Other	0.033	2.800	14.90	4.670	-0.129	0.064	-0.060	6	15	19
Glossophaga soricina	Gs	Ae	H	0.011	2.400	7.780	2.400	-0.080	0.031	-0.136	6	15	3
Carollia perspicillata	Cp	Ae	H	0.030	2.620	13.32	2.370	-0.148	0.049	-0.202	4	15	11
Artibeus jamaicensis	Aj	Ae	H	0.042	4.010	17.30	3.500	0.002	0.073	-0.138	6,13	15	19

Molossus molossus	Mo	Ae	I	0.014	2.200	6.600	2.340	-0.143	-0.029	-0.156	6	15	19
Plecotus auritus	pl	Ae	I	0.010	2.400	5.900	4.750	-0.069	-0.020	0.019	13	15	11
Myotis myotis	mm	Ae	I	0.033	3.100	8.180	5.290	-0.085	-0.066	-0.033	4	15	10
Nyctalus noctula	Nn	Ae	I	0.028	1.7	7.4	3.39	-0.328	-0.072	-0.120	4	19	19
Pholidota													
Manis javanica	Mj	Ar	I	6.000	7.400	268.0	29.30	-0.257	0.183	0.010	6	15	19
Carnivora													
Urocyon cinereoargenteus	Uc	T	C	4.750	16.50	207.1	34.34	0.116	0.150	0.059	6,13	15	19
Vulpes vulpes	Vv	T	C	9.000	20.55	511.0	49.00	0.145	0.284	0.096	6	15	11
Canis latrans	Cl	T	C	13.50	18.66	570.0	89.50	0.059	0.268	0.202	13	15	19
Canis lupus	cL	T	C	49.00	22.80	640.0	86.10	0.009	0.167	0.112	1,6,13	15	10
Ursus arctos	Ua	T	Om	200.0	18.00	1004	65.80	-0.242	0.127	-0.035	6	15	19
Ursus maritimus	Um	T	C	475.0	20.50	1634	74.00	-0.277	0.148	-0.064	1	19	2
Ailurus fulgens	Af	T	H	4.500	13.00	169.0	29.80	0.018	0.111	0.032	6	15	19
Callorhinus ursinus	Cu	Aq	P	158.0	41.40	506.0	24.20	0.145	0.001	-0.238	13	15	11
Eumetopias jubatus	Ej	Aq	P	695.0	31.25	500.9	25.90	-0.134	-0.146	-0.316	1,13	19	11
Zalophus californianus	Zc	Aq	P	225.0	37.00	547.0	17.31	0.058	-0.016	-0.333	6	15	11
Odobenus rosmarus	Or	Aq	Other	1050	26.50	291.0	88.00	-0.250	-0.304	-0.077	1,12,13	19	2
Erignathus barbatus	eB	Aq	P	280.0	33.30	310.0	90.80	-0.011	-0.161	0.013	13	19	11
Pagophilus groenlandicus	Pg	Aq	P	127.5	35.65	254.1	99.00	0.103	-0.127	0.082	12,13	19	2
Halichoerus grypus	Hg	Aq	P	207.5	33.00	386.0	76.00	0.017	-0.084	-0.007	1	15	10
Pusa hispida	pH	Aq	P	104.5	34.70	159.0	71.06	0.112	-0.210	0.022	13	15	10
Phoca vitulina	PV	Aq	P	110.0	35.25	302.0	78.50	0.114	-0.075	0.041	1	19	2,10
Cystophora cristata	Cc	Aq	P	272.0	55.00	508.0	98.40	0.210	-0.051	0.032	13	15	11
Bassariscus astutus	Ba	T	Om	1.080	12.50	132.9	24.10	0.152	0.198	0.076	6	15	19
Procyon lotor	Pl	T	Om	7.000	12.60	248.0	35.50	-0.043	0.151	0.042	13	19	19

Meles meles	ME	T	Om	13.00	9.500	360.0	50.40	-0.230	0.172	0.079	1,6	15	11,19
Gulo gulo	Gg	T	C	19.50	14.00	350.0	54.00	-0.105	0.126	0.069	1	15	19
Mustela nivalis	Mn	T	C	0.138	5.400	20.60	9.490	0.005	-0.005	0.003	6	15	10
Mustela putorius	Mp	T	C	0.958	7.500	175.0	19.98	-0.057	0.270	0.043	13	19	2
Lutra lutra	Ll	Aq	P	8.550	13.28	110.0	24.20	-0.041	-0.045	-0.054	13	15	19
Enhydra lutris	El	Aq	Other	30.00	12.35	271.9	20.61	-0.205	0.029	-0.168	13	15	19
Ichneumia albicauda	Ia	T	C	3.500	13.65	212.0	30.80	0.066	0.185	0.055	6,13	15	19
Felis catus	fC	T	C	3.900	22.10	117.1	39.70	0.264	0.045	0.103	7,13	19	8,9,10
Lynx lynx	Fl	T	C	23.00	26.82	194.0	60.70	0.160	-0.018	0.084	13	15	19
Acinonyx jubatus	aJ	T	C	46.50	36.70	156.0	70.10	0.222	-0.135	0.070	6	15	19
Puma concolor	Fc	T	C	69.50	32.00	347.0	63.20	0.120	0	0.023	1	15	19
Prionailurus viverrinus	Fv	T	C	10.85	24.20	60.40	47.75	0.195	-0.198	0.079	6	19	19
Panthera uncia	Pu	T	C	50.00	31.40	248.0	77.40	0.146	-0.041	0.087	6	15	19
Panthera leo	PL	T	C	185.0	37.90	624.0	109.0	0.089	0.031	0.079	1,6,13	15	10
Perissodactyla													
Equus caballus	Ec	T	H	530.0	41.67	785.0	60.84	0.020	-0.022	-0.114	1,6,13	19	2,10
Equus burchelli	Eb	T	H	280.0	41.90	1408	56.25	0.089	0.168	-0.090	6,13	15	19
Diceros bicornis	Db	T	H	1100	24.90	2120	46.24	-0.282	0.123	-0.219	1,6,13	15	10,17
Artiodactyla													
Camelus bactrianus	Cb	T	H	495.0	40.00	655.0	138.1	0.009	-0.054	0.068	6	19	19
Lama guanicoe	Lg	T	H	110.0	36.00	433.9	79.60	0.122	0.003	0.044	13	15	19
Hippopotamus amphibius	hA	Aq	H	2750	29.20	1940	111.5	-0.309	0.014	-0.086	13	15	10,17
Sus scrofa	SS	T	Om	195.0	24.80	1940	62.10	-0.100	0.273	-0.046	13	15	10
Giraffa camelopardalis	Gc	T	H	800.0	44.40	801.0	64.30	0.003	-0.057	-0.128	1,13	19	17
Capreolus capreolus	CC	T	H	32.50	24.35	588.1	18.84	0.082	0.189	-0.192	1,13	15	11
Rangifer tarandus	Rt	T	H	189.0	30.00	1450	36.70	-0.014	0.212	-0.159	13	15	19

Odocoileus virginianus	Ov	T	H	125.0	24.50	1170	35.64	-0.059	0.206	-0.139	6,13	15	19
Sylvicapra grimmia	Sg	T	H	18.50	26.00	418.2	40.10	0.169	0.170	0.007	6	15	19
Antilope cervicapra	Ac	T	H	37.50	28.00	490.2	61.50	0.127	0.135	0.056	6	15	19
Eudorcas thomsonii	Gt	T	H	23.00	25.00	364.0	64.50	0.129	0.118	0.097	13	15	19
Rupicapra rupicapra	Rr	T	H	37.00	28.90	456.0	36.60	0.142	0.121	-0.056	13	15	19
Ovis dalli	Od	T	H	62.70	31.60	573.0	48.16	0.125	0.119	-0.030	6	15	19
Rodentia													
Sciurus carolinensis	Sc	Ar	H	0.540	10.30	55.50	24.30	0.141	0.077	0.121	6,13	19	11
Marmota marmota	Mm	T	H	5.250	14.66	77.26	51.80	0.054	-0.074	0.142	13	19	19
Pteromys volans	pv	Ar	H	0.130	10.80	14.98	24.60	0.312	-0.068	0.214	6	15	19
Glis glis	gg	Ar	H	0.125	7.300	14.60	14.75	0.146	-0.070	0.105	6	15	19
Castor fiber	cF	Aq	H	18.50	11.28	139.0	37.30	-0.193	-0.070	-0.008	6	15	11
Ondatra zibethicus	Oz	Aq	H	1.249	10.33	24.60	18.32	0.054	-0.182	0.007	13	15	10
Cricetus cricetus	cc	T	H	0.510	4.990	26.10	12.60	-0.168	-0.082	-0.018	13	19	11
Peromyscus maniculatus	Pm	T	H	0.017	4.900	13.18	4.040	0.184	0.102	-0.050	6	15	19
Rattus rattus	RR	Ar	Om	0.200	6.310	21.44	9.550	0.033	-0.033	-0.019	6	19	10
Rattus norvegicus	RN	T	Om	0.300	5.940	21.07	9.000	-0.036	-0.076	-0.057	7,13	15	8,10,14
Mus musculus	mu	T	Om	0.021	4.340	7.950	3.200	0.109	-0.028	-0.114	7,13	15	9,11
Georychus capensis	gC	F	H	0.181	2.900	15.05	11.70	-0.294	-0.100	0.032	6	19	11
Hystrix cristata	Hc	T	H	20.00	11.24	322.0	34.20	-0.203	0.105	-0.032	6,13	19	11
Erethizon dorsatum	Ed	T	H	10.75	11.05	152.0	36.00	-0.145	0.003	0.018	6,13	15	11
Chinchilla lanigera	CL	T	H	0.650	10.10	23.30	63.70	0.113	-0.130	0.319	6	15	9,10
Cavia porcellus	cP	T	H	1.000	10.00	22.75	24.40	0.063	-0.177	0.083	6	15	10
Lagomorpha													
Oryctolagus cuniculus	Oc	T	H	1.800	18.10	68.10	28.20	0.259	0.003	0.078	13	15	10,11
Scandentia													

Tupaia minor	tm	T	I	0.200	7.500	29.05	6.760	0.108	0.034	-0.094	6	19	11
Primates													
Eulemur macaco	EM	Ar	H	2.500	15.30	77.97	18.50	0.151	0.001	-0.034	16	15	19
Varecia variegata	vV	Ar	H	3.850	18.50	59.00	20.00	0.188	-0.102	-0.045	16	19	19
Nycticebus coucang	Nc	Ar	Om	1.188	15.53	35.80	22.75	0.236	-0.096	0.058	6,16	19	10
Perodicticus potto	po	Ar	Om	1.225	12.55	41.50	25.60	0.140	-0.067	0.081	6,16	15	10
Galago senegalensis	gS	A	I	0.198	12.90	14.82	19.00	0.345	-0.112	0.131	6	19	10
Otolemur crassicaudatus	oC	Ar	I	1.300	16.30	48.30	19.60	0.248	-0.039	0.020	16	19	11
Saimiri sciureus	ss	Ar	H	0.925	14.95	3.460	25.00	0.246	-0.579	0.094	6,7	19	10
Callithrix jacchus	Cj	Ar	Om	0.342	11.60	4.160	15.60	0.241	-0.441	0.054	6,13,16	19	11
Saguinus midas	sM	Ar	Om	0.563	12.20	1.930	18.23	0.210	-0.657	0.056	16	15	19
Aotus trivirgatus	at	Ar	H	0.800	20.30	12.90	26.00	0.394	-0.279	0.111	6	19	10
Macaca mulatta	mM	Ar	H	8.000	18.53	11.00	32.00	0.111	-0.538	0.011	7,13,16	19	2
Papio cynocephalus	PC	T	Om	17.50	19.80	30.90	45.00	0.057	-0.391	0.036	13	19	10
Cercopithecus diana	cD	Ar	H	4.550	19.50	20.60	30.90	0.193	-0.347	0.039	16	15	19
Cercopithecus mitis	cM	Ar	H	9.000	18.00	16.10	37.20	0.086	-0.467	0.036	13	15	10
Pongo pygmaeus	pp	Ar	H	60.00	22.60	75.00	37.95	-0.016	-0.319	-0.079	6	15	11
Gorilla gorilla	gG	T	H	172.5	22.50	135.0	97.00	-0.130	-0.294	0.058	13	15	11
Pan troglodytes	pt	T	H	48.00	20.40	96.80	91.60	-0.037	-0.241	0.127	13,16	19	10
Homo sapiens	hs	T	Om	68.50	24.10	132.0	64.30	-0.002	-0.209	0.027	7	19	8,9,10

Table S3 Log-transformed values of infraorbital foramen (IOF) area (mm^2), eye diameter (ED) squared (mm^2), and cribriform plate (CP) area (mm^2). Original IOF data from Muchlinski (2010), ED and CP area data from Steindorff (1947), Ritland (1982), Pettigrew et al. (1988), Howland et al. (2004), Pihlström et al. (2005), Kirk (2006), and present study. See also Supplementary Material Table S2

Species	log IOF area	log ED sq	log CP area
Caluromys philander	0.196	1.88	1.52
Didelphis marsupialis	0.555	1.94	2.02
Monodelphis domestica	0.057	-	1.34
Notoryctes typhlops	0.114	0.008	-
Dasyurus viverrinus	0.400	2.06	1.74
Vombatus ursinus	1.061	-	2.48
Phalanger orientalis	0.348	2.12	1.80
Trichosurus vulpecula	0.380	2.28	1.90
Petauroides volans	0.083	1.90	1.56
Bettongia penicillata	0.360	2.28	-
Wallabia bicolor	0.747	2.12	2.18
Hemicentetes semispinosus	-0.056	0.58	1.52
Setifer setosus	0.573	1.06	1.76
Tenrec ecaudatus	0.641	1.28	1.92
Choloepus didactylus	0.507	-	2.48
Cyclopes didactylus	-0.187	2.02	1.46
Myrmecophaga tridactyla	0.711	2.06	2.96
Erinaceus europaeus	0.491	1.62	1.98
Hemiechinus auritus	0.303	1.62	1.80
Suncus murinus	0.121	0.58	1.46
Blarina brevicauda	-0.114	0.12	1.26
Neomys fodiens	-0.004	-	1.08

Sorex araneus	-0.509	-	1.10
Talpa europaea	0.053	0.30	1.46
Scalopus aquaticus	-0.004	-0.10	1.44
Desmodus rotundus	-0.310	0.90	1.18
Glossophaga soricina	-0.620	0.76	0.90
Urocyon cinereoargentatus	0.834	2.44	2.32
Vulpes vulpes	0.970	2.62	2.70
Canis lupus	1.420	2.72	2.80
Lycaon pictus	1.370	-	2.78
Ursus arctos	1.390	2.52	3.00
Bassariscus astutus	0.924	2.20	2.12
Procyon lotor	1.070	2.20	2.40
Neovison vison	0.945	-	1.84
Enhydra lutris	1.630	2.18	2.44
Ichneumia albicauda	0.764	2.28	2.32
Puma concolor	1.440	3.02	2.54
Panthera leo	2.420	3.16	2.80
Tapirus terrestris	2.020	2.64	3.16
Lama guanicoe	1.550	3.12	2.64
Hippopotamus amphibius	2.590	2.94	3.28
Giraffa camelopardalis	2.150	3.30	2.90
Antilope cervicapra	1.030	2.90	2.70
Rupicapra rupicapra	0.921	2.92	2.66
Aplodontia rufa	1.028	1.40	1.42
Sciurus carolinensis	0.314	2.02	1.74
Glis glis	0.380	1.72	1.16

Castor fiber	1.160	2.10	2.14
Ondatra zibethicus	0.549	2.02	1.40
Rattus rattus	0.678	1.60	1.34
Oryctolagus cuniculus	0.465	2.52	1.84
Tupaia minor	-0.167	1.76	1.46
Cynocephalus volans	0.053	2.30	1.44
Cheirogaleus major	0.173	2.20	1.24
Eulemur macaco	0.356	2.36	1.90
Varecia variegata	0.292	2.54	1.78
Propithecus diadema	1.000	2.62	1.92
Propithecus verreauxi	0.262	2.50	1.68
Nycticebus coucang	0.065	2.38	1.56
Perodicticus potto	0.143	2.20	1.62
Galago senegalensis	-0.456	2.22	1.18
Otolemur crassicaudatus	0.193	2.42	1.68
Saimiri sciureus	-0.180	2.34	0.54
Callithrix jacchus	-0.292	2.12	0.62
Saguinus midas	-0.071	2.18	0.28
Aotus trivirgatus	0.258	2.62	1.12
Macaca mulatta	0.207	2.54	1.04
Cercopithecus diana	0.575	2.58	1.32
Cercopithecus mitis	0.217	2.52	1.20
Pongo pygmaeus	1.066	2.70	1.88
Gorilla gorilla	0.984	2.70	2.14
Pan troglodytes	0.917	2.62	1.98
Homo sapiens	0.893	2.76	2.12

References

Adams DR (1972) Olfactory and non-olfactory epithelia in the nasal cavity of the mouse, *Peromyscus*. Am J Anat 133:37-50

Ahnelt PK, Kolb H (2000) The mammalian photoreceptor mosaic–adaptive design. Progr Retinal Eye Res 19:711-777

Allison AC, Turner Warwick RT (1949) Quantitative observations on the olfactory system of the rabbit. Brain 72:186-197

Békésy G von (1960) Experiments in hearing. McGraw-Hill, New York, pp 1-745

Bhatnagar KP, Kallen FC (1975) Quantitative observations on the nasal epithelia and olfactory innervation in bats. Acta Anat 91:272-282

Bruns V, Fiedler J, Kraus H-J (1983-1984) Structural diversity of the inner ear of bats. Myotis 21-22:52-61

Fay RR (1992) Structure and function in sound discrimination among vertebrates. In: Webster DB, Fay RR, Popper AN (eds) The evolutionary biology of hearing, Springer-Verlag, New York, pp 229-263

Gonzalez-Soriano J, Mayayo-Vicente S, Martinez-Sainz P, Contreras-Rodriguez J, Rodriguez-Veiga E (1997) A quantitative study of ganglion cells in the goat retina. Anat Histol Embryol 26:39-44

Gross EA, Swenberg JA, Fields S, Popp JA (1982) Comparative morphometry of the nasal cavity in rats and mice. J Anat 135:83-88

Güntherschulze J (1979) Studien zur Kenntnis der Regio olfactoria von Wild- und Hausschwein (*Sus scrofa scrofa* L. 1768 und *Sus scrofa f. domestica*). Zool Anz 202:256-279

Gurtovoi NN (1966) Ecological-morphological differences in the structure of the nasal cavity in the representatives of the orders Insectivora, Chiroptera and Rodentia. Zool Zh 45:1536-1551 [In Russian with English summary]

Heffner RS, Heffner HE (2010) Explaining high-frequency hearing. Anat Rec 293:2080-2082

in secondarily aquatic vertebrates. University of California Press, Berkeley, pp. 29-33.

Henderson Z (1985) Distribution of ganglion cells in the retina of adult pigmented ferret. Brain Res 358:221-228

Henson OW Jr (1961) Some morphological and functional aspects of certain structures of the middle ear in bats and insectivores. Univ Kansas Science Bull 42:151-255

Howland HC, Merola S, Basarab JR (2004) The allometry and scaling of the size of vertebrate eyes. Vision Res 44:2043-2065

Hughes A (1975) A comparison of retinal ganglion cell topography in the plains and tree kangaroo. J Physiol 244:61P-63P

Ketten DR (2000) Cetacean ears. In: Au WWL, Popper AN, Fay RR (eds) Hearing by whales and dolphins. Springer, New York and Berlin, pp 43-108

Kirk EC (2006) Effects of activity pattern on eye size and orbital aperture size in primates. J Hum Evol 51:159-170

Kolb A (1971) Licht- und elektronenmikroskopische Untersuchungen der Nasenhöhle und des Riechepithels einiger Fledermausarten. Z Säugertierk 36:202-213

Kolb A (1975) Lichtmikroskopische Untersuchungen am Riechepithel des Rehes (*Capreolus capreolus*). Anat Anz 137:417-428

Lange S, Stalleicken J, Burda H (2004) Functional morphology of the ear in fossorial rodents, *Microtus arvalis* and *Arvicola terrestris*. J Morphol 262:770-779

Lauruschkus G (1942) Über Riechfeldgrösse und Riechfeldkoefficient bei einigen Hunderassen und der Katze. Arch Tierheilk 77:473-497

Manoussaki D, Chadwick RS, Ketten DR, Arruda J, Dimitriadis EK, O´Malley JT (2008) The influence of cochlear shape on low-frequency hearing. Proc Natl Acad Sci USA 105:6162-6166

Mason MJ (2001) Middle ear structures in fossorial mammals: a comparison with non-fossorial species. J Zool 255:467-486

Mass AM, Supin AY (1992) Peak density, size and regional distribution of ganglion cells in the retina of the fur seal *Callorhinus ursinus*. Brain Behav Evol 39:69-76

Mass AM, Supin AY (2000) Ganglion cells density and retinal resolution in the sea otter, *Enhydra lutris*. Brain Behav Evol 55:111-119

Mass AM, Supin AY (2003) Retinal topography of the harp seal *Pagophilus groenlandicus.* Brain Behav Evol 62:212-222
Muchlinski MN (2010) A comparative analysis of vibrissa count and infraorbital foramen area in primates and other mammals. J Hum Evol 58:447-473

Müller M, Wess F-P, Bruns V (1993) Cochlear place-frequency map in the marsupial *Monodelphis domestica*. Hearing Res 67:198-202

Mulvaney BD, Heist HE (1970) Mapping of rabbit olfactory cells. J Anat 107:19-30

Negus V (1958) The comparative anatomy and physiology of the nose and paranasal sinuses. E. & S. Livingstone Ltd, Edinburgh and London, pp 1-402

Nowak RM (1999) Walker´s mammals of the world. 1-2 VI ed. The Johns Hopkins University Press, Baltimore, pp 1-2015

Nummela S (1995) Scaling of the mammalian middle ear. Hearing Res 85:18-30

Nummela S, Thewissen JGM, Bajpai S, Hussain ST, Kumar K (2004) Eocene evolution of whale hearing. Nature 430:776-778

Nummela S, Sánchez-Villagra MR (2006) Scaling of the marsupial middle ear and its functional significance. J Zool 270:256-267

Peichl L (1992) Topography of ganglion cells in the dog and wolf retina. J Comp Neurol 324:603-620

Peichl L, Moutairou K (1998) Absence of short-wavelength sensitive cones in the retinae of seals (Carnivora) and African giant rats (Rodentia). Eur J Neurosci 10:2586-2594

Pettigrew JD, Dreher B, Hopkins CS, McCall MJ, Brown M (1988) Peak density and distribution of ganglion cells in the retinae of microchiropteran bats: implications for visual acuity. Brain Behav Evol 32:39-56

Pettigrew JD, Bhagwandin A, Haagensen M, Manger PR (2010) Visual acuity and heterogeneities of retinal ganglion cell densities and the tapetum lucidum of the African elephant (*Loxodonta africana*). Brain Behav Evol 75:251-261

Pihlström H, Fortelius M, Hemilä S, Forsman R, Reuter T (2005) Scaling of mammalian ethmoid bones can predict olfactory organ size and performance. Proc R Soc B 272:957-962

Plassmann W, Brändle K (1992) A functional model of the auditory system in mammals and its evolutionary implications. In: Webster WB, Fay RR, Popper AN (eds) The evolutionary biology of hearing. Springer, New York and Berlin, pp 637-653

Pye A (1985-1986) Analysis of some cochlear components from surface preparations in bats. Myotis 23-24:51-55

Ritland SM (1982) The allometry of the vertebrate eye. PhD Thesis, The University of Chicago

Rosowski JJ (1992) Hearing in transitional mammals: predictions from the middle-ear anatomy and hearing capabilities. In: Webster WB, Fay RR, Popper AN (eds) The evolutionary biology of hearing. Springer, New York and Berlin, pp 615-631

Rowe TB, Eiting TP, Macrini TE, Ketcham RA (2005) Organization of the olfactory and respiratory skeleton in the nose of the gray short-tailed opossum *Monodelphis domestica*. J Mammal Evol 12:303-336

Sigmund L, Sedláček F (1985) Morphometry of the olfactory organ and olfactory thresholds of some fatty acids in *Sorex araneus*. Acta Zool Fenn 173:249-251

Söllner B, Kraft R (1980) Anatomie und Histologie der Nasenhöhle der Europeischen Wasserspitzmaus, *Neomys fodiens* (Pennant 1771), und anderer mitteleuropäischer Soriciden. Spixiana 3:251-272

Spoor F, Bajpai S, Hussain ST, Kumar K, Thewissen JGM (2002) Vestibular evidence for the evolution of aquatic behaviour in early cetaceans. Nature 417:163-166

Steindorff K (1947) Deskriptive Anatomie des Auges der Wirbeltiere und des Menschen. Tabulae Biol 22:166-297

Stone J (1965) A quantitative analysis of the distribution of ganglion cells in the cat's retina. J Comp Neurol 124:337-352

Stone J, Halasz P (1989) Topography of the retina in the elephant *Loxodonta africana*. Brain Behav Evol 34:84-95

Supin AY, Popov VV, Mass AM (2001) The sensory physiology of aquatic mammals. Kluwer Academic Publishers, Boston

Suthers RA, Wallis NE (1970) Optics of the eyes of echolocation bats. Vision Res 10:1165-1173

Werner CF (1960) Das Gehörorgan der Wirbeltiere und des Menschen. VEB Georg Thieme, Leipzig

West CD (1985) The relationship of the spiral turns of the cochlea and the length of the basilar membrane to the range of audible frequencies in ground dwelling mammals. J Acoust Soc Am 77:1091-1101