

10 Tables

Table 1. List of abbreviations used for brain structures and key to tables containing data on these structures.Size is given in mm³ if not indicated otherwise.

N = number of species.

Abbr.	structure name	<u>T a b l e n u m b e r s</u>				
		N	Size	Index	Aver. Index	%
LIN	linear brain measures (mm)	46	2	3	4	
BrW	brain weight (mg)	53	5,6	6	7	6,7
NET	net brain volume	50	8	6	7	8,9
VENT	ventricles	50	8	-	-	8,9
REST	rests	50	8	-	-	8,9
OBL	medulla oblongata	50	10	11	12	
OBL in	species with OBL subdivision	30	16	17	18	
TR	compl. sensorius trigeminalis	30	13	14	15	
TSO	ncl. tractus solitarii	30	13	14	15	
FUN	funicular nuclei	30	13	14	15	
FGR	ncl. fascicularis gracilis	30	13	14	15	
FCM	ncl. fasc. cuneatus medialis	30	13	14	15	
FCE	ncl. fasc. cuneatus externus	30	13	14	15	
RLN	relay nuclei	30				
REL	ncl. reticularis lateralis	30	16	17	18	
INO	complexus olivaris inferior	30	16	17	18	
PRP	ncl. prepositus hypoglossi	30	16	17	18	
COE	ncl. locus coeruleus	30	16	17	18	
MOT	motor nuclei	30	19	20	21	
V	ncl. mot. nervi trigemini	30	19	20	21	
VII	ncl. nervi facialis	30	19	20	21	
X	ncl. dorsalis mot. nervi vagi	30	19	20	21	
XII	ncl. nervi hypoglossi	30	19	20	21	
VC	vestibular complex	30	22	23	24	
VM	ncl. vestibularis medialis	30	22	23	24	
VI	ncl. vestibularis inferior	30	22	23	24	
VL	ncl. vestibularis lateralis	30	22	23	24	
VS	ncl. vestibularis superior	30	22	23	24	
AUD	auditory (acoustic) nuclei	30	25	26	27	
CON	cochlear nuclei	30	25	26	27	
DCO	ncl. cochlearis dorsalis	30	25	26	27	
VCO	ncl. cochlearis ventralis	30	25	26	27	
OLS	complexus olivaris superior	30	25	26	27	

Table 1. (cont.)

Abbr.	structure name	Table numbers			
		N	Vol.	Index	Aver. Index
MES	mesencephalon	50	10	11	12
MES in	species with MES subdivision	10	28	29	30
MTG	tegmentum mesencephali	10	28	29	30
NLL	ncl. lemnisci lateralis	10	28	29	30
MTC	tectum mesencephali	10	28	29	30
SUC	colliculi superiores	10	28	29	30
INC	colliculi inferiores	10	28	29	30
SCB	corpus subcommissuralis	23	40	41	42
CER	cerebellum	50	10	11	12
CER in	species with CER ncl. measurements	10	31	32	33
VPO	ventral pons	2	31	32	33
TCN	total cerebellar nuclei	10	31	32	33
MCN	ncl. medialis cerebelli	10	31	32	33
ICN	ncl. interpositus cerebelli	10	31	32	33
LCN	ncl. lateralis cerebelli	10	31	32	33
DIE	diencephalon	50	10	11	12
DIE in	species with DIE subdivision	10	34	35	36
ETH	epithalamus	10	34	35	36
HAM	ncl. habenularis medialis	38	40	41	42
EPI	epiphysis	38	40	41	42
THA	thalamus	10	34	35	36
CGM	corpus geniculatum mediale	10	37	38	39
CGL	corpus geniculatum laterale	10	37	38	39
GLD	dorsal part of CGL	10	37	38	39
GLV	ventral part of CGL	10	37	38	39
STH	subthalamus	10	34	35	36
PALL	pallidum	10	37	38	39
LUY	corpus luyi	10	37	38	39
HTH	hypothalamus	10	34	35	36
CAI	capsula interna	10	34	35	36
TRO	tractus opticus	10	34	35	36
TEL	telencephalon	50	10	11	12
BOL	olfactory bulbs	50	43	44	45
PAL	paleocortex	50	43	44	45
STR	striatum	50	43	44	45
SEP	septum	50	43	44	45
AMY	amygdala	50	43	44	45
HIP	hippocampus	50	43	44	45
SCH	schizocortex	50	43	44	45
NEO	neocortex	50	43	44	45

Table 1. (cont.)

Abbr.	structure name	<u>Table numbers</u>			
		N	Vol.	Index	Aver. Index
MOB	main olfactory bulb	50	46	47	48
L 1+2	ext. fiber + glomerular layers	43	46	47	48
L 3	external plexiform layer	43	46	47	48
L 4-6	mitral, int. plex., and gran. layers	43	46	47	48
PvZ	periventricular zone	43	46	47	48
AOB	accessory olfactory bulb	50	49	50	51
L1+2	vomerolnasal fiber and glom. layers	41	49	50	51
L3-5	ext. plex., mitral and int. plex. layers	41	49	50	51
L6	granular layer	41	49	50	51
PAL	paleocortex	50	43	44	45
RB	regio retrobulbaris	50	52	53	54
PRPI	regio prepiriformis	50	52	53	54
TOL	tuberculum olfactorium	50	52	53	54
OLC	olfactory cortices	50	52	53	54
TRL	tractus olfactorius lateralis	50	52	53	54
COA	commissura anterior	50	52	53	54
SIN	substantia innominata	50	52	53	54
SEP	septum	50	43	44	45
SET	ncl. triangularis septi	23	40	41	42
SFB	corpus subfornicale	23	40	41	42
AMY	amygdala, amygdaloid complex	50	43	44	45
MAM	medial amygdaloid division	50	55	56	57
NTO	ncl. tract. olf. lateralis	38	55	56	57
LAM	lateral amygdaloid division	50	55	56	57
MCB	basal ncl., magnocell. part	38	55	56	57
HIP	hippocampus	50	43	44	45
HIP in	species with HIP subdivision	10	58	59	60
HR	hipp. retrocommissuralis	10	58	59	60
HA	hipp. anterior (HS + HP)	10	58	59	60
HS	hipp. supracommissuralis	10	58	59	60
HP	hipp. precommissuralis	10	58	59	60
NEO	neocortex	50	43	44	45
NW	neocortical white matter	50	61	62	63
NG	neocortical grey matter	50	61	62	63
NG 1	neoc. grey, molecular layer	50	61	62	63
NG 2-6	neoc. grey, cell dense layers	50	61	62	63
ASG	area striata grey	10	64	64	64

Table 1. (cont.)

Abbr.	structure name	<u>Table numbers</u>			
		N	Vol.	Index	Aver. Index
Bauchot's data of diencephalic components (converted)					
ETH	epithalamus	20	65	66	67
THA	thalamus	20	65	66	67
STH	subthalamus	20	65	66	67
HTH	hypothalamus	20	65	66	67
RTH	ncl. reticularis	20	65	66	67
PTE	regio pretectalis	20	65	66	67
Bauchot's data of thalamic structures (converted)					
ANT	ncl. anterior thalami	20	68	69	70
MLT	ncl. mediales thalami	20	68	69	70
MNT	ncl. mediani thalami	20	68	69	70
DOT	ncl. dorsales thalami	20	68	69	70
VET	ncl. ventrales thalami	20	68	69	70
CGM	corpus geniculatum mediale	20	68	69	70
CGL	corpus geniculatum laterale	20	68	69	70
COM	area measures of commissures (mm ²)	43	71	72	73
Summarized size characteristics of all structures					74

Table 2. Linear measures of brains in mm

For proportion indices in species see Table 3

For average proportion indices in supraspecific taxonomic units see Table 4

	n	BL	HemL	BW	BH	n	CCL
Tenrecidae							
<i>Tenrec ecaudatus</i>	3	26.6	11.7	16.2	10.6	1	2.01
<i>Setifer setosus</i>	6	20.6	9.4	16.2	9.3	1	1.82
<i>Hemicentetes semispin.</i>	5	14.7	7.8	12.7	8.0	1	1.64
<i>Echinops telfairi</i>	6	16.4	7.9	11.7	7.1	1	1.24
<i>Geogale aurita</i>	1	8.5	4.4	7.0	4.0	1	0.33
<i>Oryzorictes talpoides</i>	1	12.6	7.3	11.6	6.9	1	2.68
<i>Microgale dobsoni</i>	2	14.6	7.7	11.0	7.0	1	3.13
<i>M. talazaci</i>	7	16.3	8.8	11.4	7.9	1	3.93
<i>Limnogale mergulus</i>	2	16.7	9.5	14.4	7.9	1	4.29
<i>Potamogale velox</i>	8	28.7	17.6	19.2	13.5	1	9.60
<i>Micropotam. ruwenzorii</i>	4	18.0	11.1	13.8	8.6	1	4.03
Chrysochloridae							
<i>Chrysochloris asiatica</i>	1	12.1	7.8	13.5	9.0	1	.-
<i>C. stuhlmanni</i>	2	12.5	8.6	14.7	9.0	1	2.86
Solenodontidae							
<i>Solenodon paradoxus</i>	2	34.5	16.9	21.0	12.8	1	4.30
Erinaceidae							
<i>Echinosorex gymnurus</i>	3	35.2	19.9	23.7	15.5	1	7.53
<i>Hylomys suillus</i>	1	18.6	10.4	13.1	8.5	1	3.73
<i>Atelerix algirus</i>	4	29.9	16.4	19.4	11.8	1	3.49
<i>Erinaceus europaeus</i>	2	26.4	14.3	20.5	11.3	1	2.54
<i>Hemiechinus auritus</i>	7	23.9	12.8	17.5	10.7	1	3.61
Talpidae							
<i>Desmana moschata</i>	3	24.9	19.3	23.2	11.5	1	9.93
<i>Galemys pyrenaicus</i>	2	17.7	14.1	15.2	10.4	1	7.48
<i>Talpa europaea</i>	2	17.6	11.2	15.3	7.3	1	5.31
<i>T. micrura</i>	1	16.5	10.8	13.4	6.5	1	.-
<i>Parascalops breweri</i>	6	17.2	10.7	13.8	7.4	1	4.19
<i>Scalopus aquaticus</i>	1	20.0	12.1	18.2	7.4	1	5.04

Table 2 (cont.)

	n	BL	HemL	BW	BH	n	CCL
Soricidae							
<i>Sorex alpinus</i>	-	-.	-.	-.	-.	1	2.48
<i>S. araneus</i>	2	9.7	6.7	9.2	4.8	1	2.39
<i>S. cinereus</i>	29	8.8	5.9	7.8	4.7	1	2.15
<i>S. fumeus</i>	2	10.3	6.9	8.7	5.4	1	2.86
<i>S. minutus</i>	-	-.	-.	-.	-.	1	1.85
<i>Microsorex hoyi</i>	7	7.8	4.9	6.2	3.8	1	1.91
<i>Neomys anomalus</i>	1	9.8	7.0	9.6	6.0	1	2.39
<i>N. fodiens</i>	4	11.1	7.4	10.3	6.0	1	2.59
<i>Blarina brevicauda</i>	40	11.8	7.5	11.6	6.3	1	3.18
<i>Cryptotis parva</i>	6	9.9	6.0	9.8	5.2	1	2.26
<i>Anourosorex squamipes</i>	7	11.2	6.8	11.9	6.3	1	1.87
<i>Crocidura attenuata</i>	1	10.0	5.6	8.8	4.4	1	-.
<i>C. flavescens</i>	26	13.8	7.3	11.1	5.6	4	1.89
<i>C. hildegardeae</i>	4	10.5	5.4	8.3	4.5	1	-.
<i>C. russula</i>	21	9.9	5.5	8.8	4.5	1	1.44
<i>C. suaveolens</i>	1	9.4	5.3	8.5	4.3	1	-.
<i>Suncus etruscus</i>	6	6.7	3.8	5.8	2.8	1	0.97
<i>S. murinus</i>	13	14.4	7.1	10.8	5.1	1	2.24
<i>Scutisorex somereni</i>	4	14.8	8.1	11.9	7.0	1	2.16
<i>Sylvisorex granti</i>	6	9.2	5.2	7.9	4.5	1	1.45
<i>S. megalura</i>	11	10.2	5.3	7.9	4.6	1	1.68
<i>Ruwenzorisorex suncoides</i>	1	12.2	7.0	10.6	5.4	1	2.68
<i>Myosorex babaulti</i>	1	11.6	7.1	10.5	5.8	1	2.75

Number of individuals = n

Abbreviations:

BH	brain height
BL	brain length
BW	brain width
CCL	corpus callosum length
HemL	hemisphere length

Table 3. Proportion indices (= linear measures of brains and corpus callosum given in Table 2 related to the cube roots of the standard brain volumes)

For linear measures in species see Table 2

For average proportion indices in supraspecific taxonomic units see Table 4

	BrV	CR	<u>BL</u> CR	<u>HemL</u> CR	<u>BW</u> CR	<u>BH</u> CR	<u>CCL</u> CR
Tenrecidae							
<i>Tenrec ecaudatus</i>	2498	13.57	1.96	0.86	1.19	0.78	0.15
<i>Setifer setosus</i>	1463	11.35	1.81	0.83	1.43	0.82	0.16
<i>Hemicentetes semispin.</i>	810	9.32	1.58	0.84	1.36	0.86	0.18
<i>Echinops telfairi</i>	601	8.44	1.94	0.94	1.39	0.84	0.15
<i>Geogale aurita</i>	125	5.00	1.70	0.88	1.40	0.80	0.07
<i>Oryzorictes talpoides</i>	560	8.24	1.53	0.89	1.41	0.84	0.33
<i>Microgale dobsoni</i>	538	8.13	1.80	0.95	1.35	0.86	0.38
<i>M. talazaci</i>	739	9.04	1.80	0.97	1.26	0.87	0.43
<i>Limnogale mergulus</i>	1110	10.35	1.61	0.92	1.39	0.76	0.41
<i>Potamogale velox</i>	4008	15.88	1.81	1.11	1.21	0.85	0.60
<i>Micropotam. ruwenzorii</i>	1095	10.31	1.74	1.08	1.34	0.83	0.39
Chrysochloridae							
<i>Chrysochloris asiatica</i>	676	8.77	1.38	0.89	1.54	1.03	-. -
<i>C. stuhlmanni</i>	710	8.92	1.40	0.96	1.65	1.01	0.32
Solenodontidae							
<i>Solenodon paradoxus</i>	4559	16.58	2.08	1.02	1.27	0.77	0.26
Erinaceidae							
<i>Echinorex gymnurus</i>	5873	18.04	1.95	1.10	1.31	0.86	0.42
<i>Hylomys suillus</i>	1158	10.50	1.77	0.99	1.25	0.81	0.36
<i>Atelerix algirus</i>	3151	14.66	2.04	1.12	1.32	0.80	0.24
<i>Erinaceus europaeus</i>	3250	14.81	1.78	0.97	1.38	0.76	0.17
<i>Hemiechinus auritus</i>	1815	12.20	1.96	1.05	1.43	0.88	0.30
Talpidae							
<i>Desmana moschata</i>	3861	15.69	1.59	1.23	1.48	0.73	0.63
<i>Galemys pyrenaicus</i>	1283	10.87	1.63	1.30	1.40	0.96	0.69
<i>Talpa europaea</i>	988	9.96	1.77	1.12	1.54	0.73	0.53
<i>T. micrura</i>	788	9.24	1.79	1.17	1.44	0.70	-. -
<i>Parascalops breweri</i>	849	9.47	1.82	1.13	1.45	0.78	0.44
<i>Scalopus aquaticus</i>	1264	10.81	1.85	1.12	1.68	0.68	0.47

Table 3 (cont.)

	BrV	CR	<u>BL</u> CR	<u>HemL</u> CR	<u>BW</u> CR	<u>BH</u> CR	<u>CCL</u> CR
Soricidae							
<i>Sorex alpinus</i>	253	6.32	--	--	--	--	0.39
<i>S. araneus</i>	208	5.93	1.63	1.13	1.55	0.81	0.40
<i>S. cinereus</i>	162	5.45	1.62	1.07	1.43	0.87	0.39
<i>S. fumeus</i>	233	6.15	1.67	1.11	1.41	0.87	0.47
<i>S. minutus</i>	111	4.81	--	--	--	--	0.38
<i>Microsorex hoyi</i>	93	4.53	1.73	1.09	1.38	0.84	0.42
<i>Neomys anomalus</i>	272	6.48	1.51	1.08	1.48	0.93	0.37
<i>N. fodiens</i>	317	6.82	1.62	1.08	1.51	0.87	0.38
<i>Blarina brevicauda</i>	379	7.24	1.62	1.03	1.60	0.87	0.44
<i>Cryptotis parva</i>	236	6.18	1.61	0.98	1.58	0.84	0.37
<i>Anourosorex squamipes</i>	375	7.21	1.55	0.95	1.65	0.87	0.26
<i>Crocidura attenuata</i>	202	5.86	1.71	0.96	1.50	0.75	--
<i>C. flavescens</i>	407	7.41	1.86	0.99	1.49	0.75	0.26
<i>C. hildegardeae</i>	206	5.90	1.78	0.92	1.41	0.76	--
<i>C. russula</i>	190	5.75	1.72	0.96	1.52	0.78	0.25
<i>C. suaveolens</i>	183	5.68	1.65	0.94	1.49	0.76	--
<i>Suncus etruscus</i>	60	3.91	1.71	0.97	1.49	0.71	0.25
<i>S. murinus</i>	372	7.19	2.00	0.99	1.50	0.71	0.31
<i>Scutisorex somereni</i>	618	8.52	1.74	0.95	1.40	0.82	0.25
<i>Sylvisorex granti</i>	159	5.42	1.70	0.96	1.45	0.82	0.37
<i>S. megalura</i>	181	5.66	1.80	0.93	1.39	0.82	0.30
<i>Ruwenzorisorex suncoides</i>	346	7.02	1.74	1.00	1.51	0.77	0.38
<i>Myosorex babaulti</i>	347	7.03	1.65	1.01	1.49	0.83	0.39

Abbreviations:

BH	brain height
BL	brain length
BrV	standard brain volume = $\frac{\text{BrW}}{1.036}$
BW	brain width
CCL	corpus callosum length
CR	cube root of standard BrV
HemL	hemisphere length

Table 4. Average proportion indices (= linear measures of brains and corpus callosum related to the cube roots of the standard brain volumes)

For linear measures in species see Table 2
For proportion indices in species see Table 3

	N	<u>BL</u> CR	<u>HemL</u> CR	<u>BW</u> CR	<u>BH</u> CR	<u>CCL</u> CR
Insectivora (AvIS)	46	1.73	1.01	1.44	0.82	0.35
Tenrecidae	10	1.76	0.94	1.33	0.83	0.30
Tenrecinae*	4	1.82	0.87	1.34	0.83	0.16
Geogalinae*	1	1.70	0.88	1.40	0.80	0.07
Oryzorictinae*	4	1.69	0.93	1.35	0.83	0.39
<i>Microgale</i>	2	1.80	0.96	1.31	0.87	0.41
terr. Oryzorictinae	3	1.71	0.94	1.34	0.86	0.38
<i>Limnogale</i>	1	1.61	0.92	1.39	0.76	0.41
Potamogalinae*	2	1.78	1.10	1.28	0.84	0.50
Chrysochloridae*	2	1.39	0.93	1.60	1.02	0.32
Solenodontidae*	1	2.08	1.02	1.27	0.77	0.26
Erinaceidae	5	1.90	1.05	1.34	0.82	0.30
Echinosoricinae*	2	1.86	1.05	1.28	0.84	0.39
Erinaceinae*	3	1.93	1.05	1.38	0.81	0.24
Talpidae	6	1.74	1.18	1.50	0.76	0.55
Desmaninae*	2	1.61	1.27	1.44	0.85	0.66
Talpinae*	4	1.81	1.14	1.53	0.72	0.48
<i>Talpa</i>	2	1.78	1.15	1.49	0.72	0.53
Soricidae	21	1.70	1.00	1.49	0.81	0.35
Soricinae*	9	1.62	1.06	1.51	0.86	0.39
<i>Sorex</i>	3	1.64	1.10	1.46	0.85	0.41
terr. Soricinae	7	1.63	1.05	1.51	0.85	0.39
<i>Neomys</i>	2	1.57	1.08	1.50	0.90	0.37
Crocidurinae*	12	1.76	0.97	1.47	0.77	0.30
<i>Crocidura</i>	5	1.74	0.95	1.48	0.76	0.25
<i>Suncus</i>	2	1.86	0.98	1.50	0.71	0.28
<i>Sylvisorex</i>	2	1.75	0.95	1.42	0.82	0.28
AvIF of the 12 families and subfamilies marked by *		1.75	1.02	1.40	0.83	0.34

Number of species = N

Abbreviations:

BH	brain height	CCL	corpus callosum length
BL	brain length	CR	cube root of standard BrV
BW	brain width	HemL	hemisphere length

Table 5. Average body weights (BoW) and brain weights (BrW) of 53 species of Insectivora. n = number of individuals; CV = coefficient of variation.

	BoW in g	CV in %	BrW in mg	CV in %	n BoW	n BrW	Source of data
Tenrecidae							
Tenrecinae							
<i>Tenrec ecaudatus</i>	845	14.6	2633	13.3	9	9	8,40
	885	1.6	2490	10.4	2	4	8,10a,40
	852		2588		11	13	
<i>Setifer setosus</i>	248	27.7	1516	11.6	16	16	40
	223	28.4			12	0	40
	237		1516		28	16	
<i>Hemicentetes semispin.</i>	117	12.7	839	3.9	7	7	40
	113	24.4			2	0	40
	116		839		9	7	
<i>Echinops telfairi</i>	87.5	25.9	623	10.2	6	6	S
Geogalinae							
<i>Geogale aurita</i>	10		130				b)
Oryzorictinae							
<i>Oryzorictes talpoides</i>	44.2		580		1	1	40
<i>Microgale dobsoni</i>	31.9	9.6	557	1.0	3	3	S
<i>M. talazaci</i>	48.2	16.0	783	8.0	13	13	S
	48.2	21.8	693	3.0	4	3	S
	48.2		766		17	16	
<i>Limnogale mergulus</i>	92		1150				40
Potamogalinae							
<i>Potamogale velox</i>	618	12.9	4152	6.9	13	13	S
<i>Micropotamog. lamottei</i>	64.2		800				57
<i>M. ruwenzorii</i>	98.5	14.9	1134	9.9	4	4	S
	95.0	28.8			4	0	30
	96.8		1134		8	4	
Chrysochloridae							
<i>Chrysochloris asiatica</i>	49		700		1	1	S
<i>C. stuhlmanni</i>	39.9	15.5	741	7.7	9	9	S
	45.3	15.2			5	0	S
	41.8		741		14	9	
mm+ff median	40.2		736				a)
mm	46.0		780		9	5	
ff	34.4		692		5	4	

Table 5 (cont.)

	BoW in g	CV in %	BrW in mg	CV in %	n BoW	n BrW	Source of data
Solenodontidae							
<i>Solenodon paradoxus</i>	674	3.7	4795	5.0	4	4	42,S
	663		4580	0.3	1	2	S
	672		4723		5	6	
Erinaceidae							
Echinosoricinae							
<i>Echinosorex gymnurus</i>	823	12.5	6208	5.7	2	2	S
			5960		0	2	S
	823		6084		2	4	
<i>Hylomys suillus</i>	57		1200				c)
Erinaceinae							
<i>Atelerix algirus</i>	736	14.4	3264	4.8	7	7	S
<i>Erinaceus europaeus</i>	839	15.5	3378	6.4	21	21	2,3,4,8,13, 23,32,59,S
	860		3300		16	3	5,7,8,15,S
	849		3367		35	22	
<i>Hemiechinus auritus</i>	235	26.2	1880	8.3	7	7	S
Talpidae							
Desmaninae							
<i>Desmana moschata</i>	443	10.9	4000	2.5	3	3	40
<i>Galemys pyrenaicus</i>	61.1	4.6	1328	3.8	3	3	26,40
	59.6		1330		44	2	26,40
	59.7		1329		47	5	
Talpinae							
<i>Talpa europaea</i>	82.5	11.4	1016	6.9	11	11	5,6,11, 32,S
	78.0		1055		1	3	
	82.1		1024		12	14	
<i>T. micrura</i>	41.4		816		1	1	S
<i>Parascalops breweri</i>	53.8	5.7	880	4.1	6	6	S
<i>Scalopus aquaticus</i>	115		1310		1	1	S
Soricidae							
Soricinae							
<i>Sorex alpinus</i>	10.3	24.7	259	5.2	4	4	S
	11.1		265	4.1	41	4	50,S
	11.0		262		45	8	
<i>S. araneus</i>	8.5	18.4	205	7.7	40	40	5,6,9,40,S
	10.3		218		717	249	12,15,17,19, 20,22,24,27, 29,31,35,37, 38,39,47,S
	10.2		216		757	289	

Table 5 (cont.)

	BoW in g	CV in %	BrW in mg	CV in %	n BoW	n BrW	Source of data
Soricinae (cont.)							
<i>Sorex cinereus</i>	4.20	25.0	174	7.5	29	29	S,B
	3.9				28	0	16,18,25,34,B
	4.05		174		57	29	
	5.16	22.8	168	10.3	11	11	S,B (definite adults)
<i>S. fumeus</i>	8.4	9.9	241	3.5	5	2	S,B
<i>S. minutus</i>	4.11	22.8	120	13.9	11	11	S
	4.42		115		338	113	1,12,17,21,28,
	4.41		115		349	124	29,31,40,41, 43,44,47
<i>S. palustris</i>	14.6		278		1	1	B
<i>Microsorex hoyi</i>	2.63	7.5	96	12.0	7	7	S,B
<i>Neomys anomalus</i>	9.42	2.4	282	3.5	5	5	S
	11.8				76	0	17,47,48,53
	11.6		282		81	5	
<i>N. fodiens</i>	15.0	13.2	327	5.1	17	17	S
	15.4		333		254	4	15,17,29,31,
	15.3		328		271	21	36,41,43,47, 48,49,53,S
<i>Blarina brevicauda</i>	18.2	16.4	397	4.8	42	42	S,B
	17.4		348		68	68	14 (preserved)
	19.3				16	0	7,16,34,B
	17.9		367		126	110	
	19.7	18.9	393	2.9	7	7	S,B (definite adults)
<i>Cryptotis parva</i>	9.5	8.3	244	1.8	5	5	S
	10.3		248		4	1	S
	9.9		245		9	6	
<i>Anourosorex squamipes</i>	20.0	9.4	389	5.7	7	7	S
	20.5	17.2			2	0	S
	20.1		389		9	7	
Crocidurinae							
<i>Crocidura attenuata</i>	9.3		209		1	1	S
<i>C. flavescens</i>	31.6	23.5	426	9.0	52	52	S
	34.3		394		58	7	51,56,S
	33.0		422		110	59	
	29.3		414				a)
mm + ff median	29.3		414				
mm	32.7		432		63	43	
ff	25.8		396		22	16	
<i>C. giffardi</i>	82.0		545		1	1	10

Table 5 (cont.)

	BoW in g	CV in %	BrW in mg	CV in %	n BoW	n BrW	Source of data
Crocidurinae (cont.)							
<i>C. hildegardae</i>	9.1	17.5	213	10.5	9	9	S
	10.5				41	0	40,51,S
	10.2		213		50	9	
<i>C. jacksoni</i>	10.3		250		1	1	S
	13.25				4	0	S
	12.7		250		5	1	
<i>C. leucodon</i>	13.5		190		1	1	S
<i>C. russula</i>	11.4	12.5	195	6.8	24	24	58,S
	11.0		201		61	12	22,54,55,56,
	11.1		197		85	36	58,F,S
<i>C. suaveolens</i>	10.3		190		1	1	F,S
<i>Suncus etruscus</i>	2.07	10.1	62	3.7	6	6	F,S
	1.85				194	0	45,46,52,56
	1.86		62		200	6	
<i>S. murinus</i>	34.1	31.9	383	8.7	26	26	33,S
	37.6	26.4	449		6	1	33,S
	34.8		385		32	27	
mm + ff median	33.8		383				a)
mm	41.9		400		18	15	
ff	25.7		366		14	12	
<i>Scutisorex somereni</i>	58.9	16.3	640	2.5	4	4	S
	63.7				70	0	51
	63.4		640		74	4	
<i>Sylvisorex granti</i>	3.88	5.5	165	3.1	6	6	S
	3.90				25	0	51,H
	3.90		165		31	6	
<i>S. megalura</i>	5.55	9.9	188	7.1	13	13	S
	5.47				40	0	40,51,56,S
	5.49		188		53	13	
<i>Ruwenzorisorex suncoides</i>	18.2		370		1	1	S
	18.5		358		1	1	S d)
<i>Myosorex babaulti</i>	17.0		360		1	1	S
	17.0				12	0	51
	17.0		360		13	1	

Notes to Table 5:

In general there are 3 lines for each species. The first line includes all the pairs of body and brain weights we are aware of, the second line single values (either body weights or brain weights), and the third line the combined data from the two preceding lines. If there are more than three lines, they include data of preserved material (as e.g. in *Blarina*), or data from definite adults (if appropriate).

Table 5 (cont.)

- a) Median values between males and females are given in the case of statistically significant sex differences (99% for BoW, $P < .01$; 95% for BrW, $P < .05$).
- b) From *Geogale aurita* only secondary inferred data, taken from Rehkämper et al. (1986), are available (see text).
- c) From *Hylomys suillus* we have only one brain of a juvenile animal (BoW 31.4 g, BrW 957 mg). Its brain was used to measure the brain composition. Based on BoW data of Miller (1942), Rudd (1965), Harrison (1966), Lim and Heyneman (1968) and Medway (1969) adults have a BoW of about 57 g. Based on cranial capacities ($n = 7$) the expected brain weight is 1200 mg.
- d) In the second line of *Ruwenzorisorex suncoides* a specimen of uncertain determination is listed, which in Fons et al. (1984) was designated as *Sylvisorex (lunaris?)*. According to its BrW it may have been a *Ruwenzorisorex suncoides*.

Key to the source of body and brain weight data:

1: Snell (1892); 2: Weber (1896); 3: Dubois (1897); 4: Flatau and Jacobsohn (1899); 5: Ziehen (1899a); 6: Welcker und Brandt (1903); 7: Hrdlicka (1905); 8: Warncke (1908); 9: Lapicque (1912); 10: Waterlot (1912); 10a: Brodmann (1913); 11: Slifer (1924); 12: Middleton (1931); 13: Bonin (1937); 14: Crile and Quiring (1940); 15: Cantuel (1943); 16: Morrison and Pearson (1946); 17: Borowski and Dehnel (1953); 18: Pruitt (1954); 19: Siivonen (1954); 20: Pucek (1955); 21: Cabon (1956); 22: Niethammer (1956); 23: Haug (1958); 24: Schubarth (1958); 25: Morrison et al. (1959); 26: Stephan and Bauchot (1959); 27: Bielak and Pucek (1960); 28: Hawkins et al. (1960); 29: Niethammer (1960); 30: Rahm (1960, 1961); 31: Hawkins and Jewell (1962); 32: Portmann (1962); 33: Bauchot and Stephan (1964); 34: Buckner (1964); 35: Pucek, Z. (1964); 36: Gebczynska and Gebczynski (1965); 37: Gebczynski (1965); 38: Pucek, M. (1965); 39: Pucek, Z. (1965); 40: Bauchot and Stephan (1966); 41: Mangold-Wirz (1966); 42: Boller (1969); 43: Gebczynski (1971a); 44: Gebczynski (1971b); 45: Fons (1974); 46: Fons et Saint Girons (1975); 47: Gebczynski (1977); 48: Niethammer (1977); 49: Niethammer (1978); 50: Spitzenberger (1978); 51: Dieterlen und Heim de Balsac (1979); 52: Fons (1979); 53: Spitzenberger (1980); 54: Genoud (1981); 55: Genoud and Vogel (1981); 56: Vogel et al. (1981); 57: Stephan and Kuhn (1982); 58: Schwerdtfeger (1984); 59: Hofer (pers. comm.).

F: collection Roger Fons; H: collection Rainer Hutterer; S: collection Heinz Stephan in cooperation with Georg Baron (B), Kunwar Bhatnagar, Roland Bauchot, Petra Friesleber, Kabongo Ka Mubalamata, Linjanja Lulyo, Otmar Lütt, José Regidor, Gerd Rehkämper, Wolfbernard Spatz, Michael and Roland Stephan, Gottfried Vauk, Songsakdi Yenbutra, and others.

Table 6. Standards of body and brain weights, percentages of brain weight relative to body weight, encephalization indices (EIs) derived from the standards and size indices (SIs) of NET brain volumes. EIs and SIs are related to the average of the four Tenrecinae (AvTn = 100)

For average encephalization indices in supraspecific taxonomic units see					
For NET brain volumes see Table 8				Table 7	
	<u>Standards</u>		<u>BrW</u>	BrW	NET
	BoW	BrW	BoW	EI	SI
slope				.66	.66
y-intercept				1.544	1.506
Tenrecidae					
<i>Tenrec ecaudatus</i>	852	2588	0.30	86	85
<i>Setifer setosus</i>	237	1516	0.64	117	119
<i>Hemicentetes semispinosus</i>	116	839	0.72	104	104
<i>Echinops telfairi</i>	87.5	623	0.71	93	93
<i>Geogale aurita</i>	10.0	130	1.30	81	83
<i>Oryzorictes talpoides</i>	44.2	580	1.31	136	138
<i>Microgale dobsoni</i>	31.9	557	1.75	162	162
<i>M. talazaci</i>	48.2	766	1.59	170	174
<i>Limnogale mergulus</i>	92.0	1150	1.25	166	165
<i>Potamogale velox</i>	618	4152	0.67	171	174
<i>Micropotamogale lamottei</i>	64.2	800	1.25	147	149
<i>M. ruwenzorii</i>	96.8	1134	1.17	158	162
Chrysochloridae					
<i>Chrysochloris asiatica</i>	49.0	700	1.43	153	157
<i>C. stuhlmanni</i>	40.2	736	1.83	184	188
Solenodontidae					
<i>Solenodon paradoxus</i>	672	4723	0.70	184	184
Erinaceidae					
<i>Echinosorex gymnurus</i>	823	6084	0.74	207	209
<i>Hylomys suillus</i>	57.0	1200	2.11	238	241
<i>Atelerix algirus</i>	736	3264	0.44	120	117
<i>Erinaceus europaeus</i>	849	3367	0.40	112	112
<i>Hemiechinus auritus</i>	235	1880	0.80	146	144
Talpidae					
<i>Desmana moschata</i>	443	4000	0.90	205	202
<i>Galemys pyrenaicus</i>	59.7	1329	2.23	255	258
<i>Talpa europaea</i>	82.1	1024	1.25	160	163
<i>T. micrura</i>	41.4	816	1.97	200	205
<i>Parascalops breweri</i>	53.8	880	1.64	181	188
<i>Scalopus aquaticus</i>	115	1310	1.14	163	170

Table 6 (cont.)

	<u>Standards</u>		<u>BrW</u>	<u>BrW</u>	<u>NET</u>
	BoW	BrW	BoW	EI	SI
Soricidae					
<i>Sorex alpinus</i>	11.0	262	2.38	154	160
<i>S. araneus</i>	10.2	216	2.12	133	137
<i>S. cinereus</i>	5.16	168	3.26	163	169
<i>S. fumeus</i>	8.40	241	2.87	169	175
<i>S. minutus</i>	4.41	115	2.61	123	128
<i>S. palustris</i>	14.6	278	1.90	135	-
<i>Microsorex hoyi</i>	2.63	96	3.65	145	150
<i>Neomys anomalus</i>	11.6	282	2.43	160	166
<i>N. fodiens</i>	15.3	328	2.14	155	159
<i>Blarina brevicauda</i>	19.7	393	1.99	157	162
<i>Cryptotis parva</i>	9.90	245	2.47	154	160
<i>Anourosorex squamipes</i>	20.1	389	1.94	153	158
<i>Crocidura attenuata</i>	9.30	209	2.25	137	142
<i>C. flavescens</i>	29.3	414	1.41	127	130
<i>C. giffardi</i>	82.0	545	0.66	85	-
<i>C. hildegardeae</i>	10.2	213	2.09	131	136
<i>C. jacksoni</i>	12.7	250	1.97	133	136
<i>C. leucodon</i>	13.5	190	1.41	97	-
<i>C. russula</i>	11.1	197	1.77	115	118
<i>C. suaveolens</i>	10.3	190	1.84	116	120
<i>Suncus etruscus</i>	1.86	62	3.33	118	121
<i>S. murinus</i>	33.8	383	1.13	107	110
<i>Scutisorex somereni</i>	63.4	640	1.01	118	122
<i>Sylvisorex granti</i>	3.90	165	4.23	192	199
<i>S. megalura</i>	5.49	188	3.42	175	181
<i>Ruwenzorisorex suncoides</i>	18.2	370	2.03	156	161
<i>Myosorex babaulti</i>	17.0	360	2.12	159	164

Abbreviations and notes:

- BoW body weight in g
 BrW brain weight in mg
 EI BrW encephalization indices
 SI NET size indices of the NET brain volume

Table 7. Average percentages of brain weight related to body weight, and averages of the encephalization indices (EIs) of BrW and size indices (SIs) of the NET brain volumes in supraspecific taxonomic units

For encephalization indices in species see Table 6

	N BrW	N NET	BrW BoW	BrW EI	NET SI
slope				.66	.66
y-intercept				1.544	1.506
Insectivora (AvIS)	53	50	1.71	148	154
Tenrecidae	12	12	1.06	133	134
Tenrecinae*	4	4	0.59	100	100
Geogalinae*	1	1	1.30	81	83
Oryzorictinae*	4	4	1.47	158	160
<i>Microgale</i>	2	2	1.67	166	168
terr.Oryzorictinae	3	3	1.55	156	158
<i>Limnogale</i>	1	1	1.25	166	165
Potamogalinae*	3	3	1.03	159	162
<i>Micropotamogale</i>	2	2	1.21	153	155
Chrysochloridae*	2	2	1.63	169	172
Solenodontidae*	1	1	0.70	184	184
Erinaceidae	5	5	0.90	165	165
Echinosoricinae*	2	2	1.42	222	225
Erinaceinae*	3	3	0.55	126	124
Talpidae	6	6	1.52	194	198
Desmaninae*	2	2	1.56	230	230
Talpinae*	4	4	1.50	176	181
<i>Talpa</i>	2	2	1.61	180	184
Soricidae	27	24	2.24	140	148
Soricinae*	12	11	2.48	151	157
<i>Sorex</i>	6	5	2.52	146	154
terr. Soricinae	10	9	2.52	149	155
<i>Neomys</i>	2	2	2.29	157	162
Crocidurinae*	15	13	2.05	137	142
<i>Crocidura</i>	8	6	1.68	118	130
<i>Suncus</i>	2	2	2.23	112	115
<i>Sylvisorex</i>	2	2	3.83	183	190
AvIF of the 12 families and subfamilies marked by *			1.36	158	160

Number of species = N
Abbreviations as in Table 6

Table 8. Volumes of net brain (NET), ventricles (VENT) and rests (REST) and their percentages relative to standard brain volumes (= 100%)

For standard brain volumes (BrV) see Table 3

For average percentages in supraspecific taxonomic units see Table 9

	n	NET		VENT		REST	
		mm ³	%	mm ³	%	mm ³	%
Tenrecidae							
<i>Tenrec ecaudatus</i>	3	2331	93.3	65.7	2.6	101.6	4.1
<i>Setifer setosus</i>	3	1404	96.0	22.0	1.5	36.8	2.5
<i>Hemicentetes semispin.</i>	2	770	95.0	16.9	2.1	23.3	2.9
<i>Echinops telfairi</i>	2	569	94.6	8.4	1.4	24.0	4.0
<i>Geogale aurita</i>	2	122	97.4	1.2	0.9	2.1	1.7
<i>Oryzorictes talpoides</i>	1	538	96.0	4.8	0.9	17.3	3.1
<i>Microgale dobsoni</i>	1	511	95.0	5.3	1.0	21.8	4.0
<i>M. talazaci</i>	2	718	97.2	5.6	0.7	15.5	2.1
<i>Limnogale mergulus</i>	1	1046	94.3	12.7	1.1	51.0	4.6
<i>Potamogale velox</i>	5	3874	96.7	18.4	0.4	115.4	2.9
<i>Micropotamogale lam.</i>	3	743	96.2	11.9	1.6	17.2	2.2
<i>M. ruwenzorii</i>	4	1064	97.2	8.4	0.8	22.7	2.0
Chrysochloridae							
<i>Chrysochloris asiatica</i>	1	657	97.2	8.1	1.2	10.7	1.6
<i>C. stuhlmanni</i>	2	690	97.1	7.0	1.0	13.7	1.9
Solenodontidae							
<i>Solenodon paradoxus</i>	2	4331	95.0	57.1	1.3	171.1	3.7
Erinaceidae							
<i>Echinosorex gymnurus</i>	3	5638	96.0	121.4	2.1	112.9	1.9
<i>Hylomys suillus</i>	1	1116	96.3	23.2	2.0	19.4	1.7
<i>Atelerix algirus</i>	3	2916	92.6	123.4	3.9	111.1	3.5
<i>Erinaceus europaeus</i>	5	3065	94.3	105.8	3.3	79.2	2.4
<i>Hemiechinus auritus</i>	3	1692	93.3	81.7	4.5	40.7	2.2
Talpidae							
<i>Desmana moschata</i>	1	3620	93.8	137.6	3.5	103.7	2.7
<i>Galemys pyrenaicus</i>	2	1229	95.8	27.4	2.2	25.9	2.0
<i>Talpa europaea</i>	2	957	96.8	10.9	1.1	20.7	2.1
<i>T. micrura</i>	2	768	97.5	9.8	1.3	9.7	1.2
<i>Parascalops breweri</i>	3	835	98.3	10.3	1.2	4.5	0.5
<i>Scalopus aquaticus</i>	1	1246	98.5	8.6	0.7	10.0	0.8

Table 8 (cont.)

	n	NET		VENT		REST	
		mm ³	%	mm ³	%	mm ³	%
Soricidae							
<i>Sorex alpinus</i>	3	250	98.7	1.2	0.5	2.2	0.8
<i>S. araneus</i>	5	203	97.5	1.3	0.6	4.0	1.9
<i>S. cinereus</i>	3	160	98.5	0.8	0.5	1.7	1.0
<i>S. fumeus</i>	2	229	98.3	1.8	0.8	2.1	0.9
<i>S. minutus</i>	4	109	98.2	0.4	0.4	1.5	1.4
<i>Microsorex hoyi</i>	3	91	98.4	0.5	0.6	1.0	1.0
<i>Neomys anomalus</i>	3	268	98.5	1.5	0.5	2.6	1.0
<i>N. fodiens</i>	2	308	97.4	1.7	0.5	6.7	2.1
<i>Blarina brevicauda</i>	3	372	98.1	2.7	0.7	4.7	1.2
<i>Cryptotis parva</i>	3	233	98.4	1.5	0.6	2.3	1.0
<i>Anourosorex squamipes</i>	4	368	98.0	2.9	0.8	4.5	1.2
<i>Crocidura attenuata</i>	1	198	98.3	1.4	0.7	2.1	1.0
<i>C. flavescens</i>	3	387	96.9	2.7	0.7	9.6	2.4
<i>C. hildegardeae</i>	4	202	98.2	1.1	0.5	2.6	1.3
<i>C. jacksoni</i>	1	234	97.0	1.2	0.5	6.0	2.5
<i>C. russula</i>	5	186	97.7	1.5	0.8	2.9	1.5
<i>C. suaveolens</i>	3	180	98.0	1.6	0.9	2.2	1.1
<i>Suncus etruscus</i>	3	58	97.8	0.5	0.8	0.9	1.4
<i>S. murinus</i>	3	359	97.0	3.1	0.9	7.8	2.1
<i>Scutisorex somereni</i>	3	604	97.8	4.6	0.8	8.9	1.4
<i>Sylvisorex granti</i>	3	157	98.4	0.9	0.5	1.7	1.1
<i>S. megalura</i>	3	178	98.3	1.1	0.6	2.0	1.1
<i>Ruwenzorisorex sunc.</i>	1	351	98.3	2.1	0.6	3.9	1.1
<i>Myosorex babaulti</i>	1	342	98.3	1.5	0.5	4.2	1.2

Number of individuals = n

Abbreviations:

- NET net brain volume
- REST rests including meninges, nerves, hypophysis, ect.
- VENT ventricle volumes

Table 9. Average percentages of NET brain, ventricles and rests relative to standard brain volumes (BrV = 100%)

For volumes and percentages in species see Table 8

	N	NET	VENT	REST
Insectivora (AvIS)	50	96.9	1.2	1.9
Tenrecidae	12	95.7	1.3	3.0
Tenrecinae*	4	94.7	1.9	3.4
Geogalinae*	1	97.4	0.9	1.7
Oryzorictinae*	4	95.6	0.9	3.5
<i>Microgale</i>	2	96.1	0.9	3.0
terr. Oryzorictinae	3	96.1	0.8	3.1
<i>Limnogale</i>	1	94.3	1.1	4.6
Potamogalinae*	3	96.7	0.9	2.4
<i>Micropotamogale</i>	2	96.7	1.2	2.1
Chrysochloridae*	2	97.1	1.1	1.8
Solenodontidae*	1	95.0	1.3	3.7
Erinaceidae	5	94.5	3.1	2.4
Echinosoricinae*	2	96.2	2.0	1.8
Erinaceinae*	3	93.4	3.9	2.7
Talpidae	6	96.8	1.6	1.6
Desmaninae*	2	94.8	2.8	2.4
Talpinae*	4	97.8	1.1	1.1
<i>Talpa</i>	2	97.1	1.2	1.7
Soricidae	24	98.0	0.6	1.4
Soricinae*	11	98.2	0.6	1.2
<i>Sorex</i>	5	98.2	0.6	1.2
terr. Soricinae	9	98.2	0.6	1.2
<i>Neomys</i>	2	98.0	0.5	1.5
Crocidurinae*	13	97.8	0.7	1.5
<i>Crocidura</i>	6	97.7	0.7	1.6
<i>Suncus</i>	2	97.4	0.8	1.8
<i>Sylvisorex</i>	2	98.3	0.6	1.1
AvIF of the 12 families and subfamilies marked by *		96.2	1.5	2.3

Number of species = N

Abbreviations as in Table 8

Table 10. Volumes of the five fundamental brain parts

For size indices in species see Table 11
For average size indices in supraspecific taxonomic units see Table 12

	n	OBL	MES	CER	DIE	TEL
Tenrecidae						
<i>Tenrec ecaudatus</i>	3	315.5	133.3	311.5	153.7	1417
<i>Setifer setosus</i>	3	180.6	76.6	146.8	93.1	907
<i>Hemicentetes semispinosus</i>	2	117.0	45.9	120.5	47.7	439
<i>Echinops telfairi</i>	2	90.3	44.0	58.2	40.3	336
<i>Geogale aurita</i>	2	34.0	12.7	12.5	7.9	55
<i>Oryzorictes talpoides</i>	1	61.0	26.6	81.3	36.9	332
<i>Microgale dobsoni</i>	1	58.7	37.6	57.0	32.6	325
<i>M. talazaci</i>	2	84.0	45.5	94.9	45.4	449
<i>Limnogale mergulus</i>	1	203.8	67.4	176.0	81.0	518
<i>Potamogale velox</i>	5	722.8	259.9	622.7	300.3	1968
<i>Micropotamogale lamottei</i>	3	134.9	58.6	82.7	62.8	404
<i>M. ruwenzorii</i>	4	198.0	72.5	149.3	88.8	555
Chrysochloridae						
<i>Chrysochloris asiatica</i>	1	58.5	43.2	72.6	52.8	430
<i>C. stuhlmanni</i>	2	68.0	41.7	74.3	51.6	454
Solenodontidae						
<i>Solenodon paradoxus</i>	2	471.6	208.7	761.2	272.3	2617
Erinaceidae						
<i>Echinosorex gymnurus</i>	3	622.4	360.4	801.0	395.1	3459
<i>Hylomys suillus</i>	1	125.4	83.9	157.2	85.9	663
<i>Atelerix algirus</i>	3	343.0	156.4	364.7	251.0	1801
<i>Erinaceus europaeus</i>	5	335.0	151.3	395.8	242.4	1941
<i>Hemiechinus auritus</i>	3	185.2	109.0	196.1	140.2	1062
Talpidae						
<i>Desmana moschata</i>	1	402.4	218.7	514.5	343.8	2140
<i>Galemys pyrenaicus</i>	2	147.1	62.1	183.6	104.5	732
<i>Talpa europaea</i>	2	95.7	50.0	174.2	80.7	556
<i>T. micrura</i>	2	77.1	45.0	116.0	60.8	469
<i>Parascalops breweri</i>	3	74.7	35.6	121.0	66.1	537
<i>Scalopus aquaticus</i>	1	118.5	57.7	197.7	91.7	780

Table 10 (cont.)

	n	OBL	MES	CER	DIE	TEL
Soricidae						
<i>Sorex alpinus</i>	3	31.2	12.5	31.5	17.0	157
<i>S. araneus</i>	5	29.1	11.5	22.0	14.8	126
<i>S. cinereus</i>	3	17.2	8.3	16.5	10.9	107
<i>S. fumeus</i>	2	22.6	11.1	23.4	15.8	156
<i>S. minutus</i>	4	15.0	6.4	12.1	7.7	68
<i>Microsorex hoyi</i>	3	10.9	4.8	8.7	6.9	60
<i>Neomys anomalus</i>	3	34.2	16.0	32.4	21.5	164
<i>N. fodiens</i>	2	45.3	19.2	41.0	24.1	179
<i>Blarina brevicauda</i>	3	43.0	18.8	42.3	26.3	242
<i>Cryptotis parva</i>	3	30.1	12.7	24.2	17.1	149
<i>Anourosorex squamipes</i>	4	44.1	18.8	39.7	25.4	240
<i>Crocidura attenuata</i>	1	32.4	15.0	24.3	13.0	114
<i>C. flavescens</i>	3	59.9	26.7	51.9	25.1	224
<i>C. hildegardae</i>	4	31.4	14.9	23.0	14.0	118
<i>C. jacksoni</i>	1	42.3	17.9	28.5	16.3	129
<i>C. russula</i>	5	28.7	13.7	21.0	13.9	109
<i>C. suaveolens</i>	3	27.0	13.7	21.9	12.7	105
<i>Suncus etruscus</i>	3	11.1	4.7	5.9	4.4	32
<i>S. murinus</i>	3	54.4	24.6	42.8	24.1	213
<i>Scutisorex somereni</i>	3	82.9	36.6	83.9	40.7	360
<i>Sylvisorex granti</i>	3	21.5	11.4	18.7	10.1	95
<i>S. megalura</i>	3	26.8	14.2	24.9	12.3	100
<i>Ruwenzorisorex suncooides</i>	1	54.9	25.0	52.7	26.2	192
<i>Myosorex babaulti</i>	1	46.2	19.0	42.4	23.6	211

Number of individuals = n

Abbreviations and notes:

- CER cerebellum; for volumes of components see Table 31
 DIE diencephalon; for volumes of components see Tables 34, 37 and 40
 MES mesencephalon; for volumes of components see Table 28
 OBL medulla oblongata; for volumes of components see Tables 13, 16, 19, 22,
 TEL telencephalon; for volumes of components see Table 43 and 25

Table 11. Size indices of the five fundamental brain parts related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 10
For average size indices in supraspecific taxonomic units see Table 12

	n	OBL	MES	CER	DIE	TEL
slope		.61	.58	.68	.65	.68
y-intercept		.776	.481	.552	.350	1.242
Tenrecidae						
<i>Tenrec ecaudatus</i>	3	86	88	89	85	83
<i>Setifer setosus</i>	3	108	106	100	119	126
<i>Hemicentetes semispin.</i>	2	108	96	133	97	99
<i>Echinops telfairi</i>	2	99	109	78	98	92
<i>Geogale aurita</i>	2	140	110	73	79	66
<i>Oryzorictes talpoides</i>	1	101	98	173	141	145
<i>Microgale dobsoni</i>	1	119	167	152	153	177
<i>M. talazaci</i>	2	132	159	191	163	184
<i>Limnogale mergulus</i>	1	216	162	228	192	137
<i>Potamogale velox</i>	5	240	207	221	206	143
<i>Micropotamogale lam.</i>	3	178	173	137	188	137
<i>M. ruwenzorii</i>	4	204	169	187	203	142
Chrysochloridae						
<i>Chrysochloris asiatica</i>	1	91	149	144	188	175
<i>C. stuhlmanni</i>	2	120	162	169	209	211
Solenodontidae						
<i>Solenodon paradoxus</i>	2	149	158	255	177	179
Erinaceidae						
<i>Echinorex gymnurus</i>	3	174	243	234	225	206
<i>Hylomys suillus</i>	1	178	266	282	277	243
<i>Atelerix algerus</i>	3	102	112	115	154	116
<i>Erinaceus europaeus</i>	5	92	100	113	135	113
<i>Hemiechinus auritus</i>	3	111	152	134	180	148
Talpidae						
<i>Desmana moschata</i>	1	164	211	229	292	195
<i>Galemys pyrenaicus</i>	2	203	192	319	327	260
<i>Talpa europaea</i>	2	109	128	244	205	159
<i>T. micrura</i>	2	133	171	259	242	214
<i>Parascalops breweri</i>	3	110	117	226	221	205
<i>Scalopus aquaticus</i>	1	110	122	220	188	177

Table 11 (cont.)

	n	OBL	MES	CER	DIE	TEL
Soricidae						
<i>Sorex alpinus</i>	3	121	103	173	160	176
<i>S. araneus</i>	5	118	99	127	146	149
<i>S. cinereus</i>	3	106	106	151	167	200
<i>S. fumeus</i>	2	104	107	154	177	210
<i>S. minutus</i>	4	102	90	124	131	142
<i>Microsorex hoyi</i>	3	101	91	127	165	178
<i>Neomys anomalus</i>	3	128	128	171	195	177
<i>N. fodiens</i>	2	144	131	180	183	160
<i>Blarina brevicauda</i>	3	117	110	156	169	182
<i>Cryptotis parva</i>	3	124	111	143	172	179
<i>Anourosorex squamipes</i>	4	119	109	145	161	179
<i>Crocidura attenuata</i>	1	139	136	150	136	143
<i>C. flavescens</i>	3	128	125	146	125	129
<i>C. hildegardae</i>	4	128	128	133	139	140
<i>C. jacksoni</i>	1	150	135	142	139	131
<i>C. russula</i>	5	111	112	114	129	121
<i>C. suaveolens</i>	3	109	117	126	124	123
<i>Suncus etruscus</i>	3	127	109	109	132	121
<i>S. murinus</i>	3	106	105	110	109	111
<i>Scutisorex somereni</i>	3	110	109	140	122	123
<i>Sylvisorex granti</i>	3	157	171	208	186	216
<i>S. megalura</i>	3	159	174	220	182	180
<i>Ruwenzorisorex suncoides</i>	1	157	154	205	177	153
<i>Myosorex babaulti</i>	1	137	121	173	167	176

Number of individuals = n

Abbreviations and notes:

- CER cerebellum; for size indices of components see Table 32
 DIE diencephalon; for size indices of components see Tables 35, 38 and 41
 MES mesencephalon; for size indices of components see Table 29
 OBL medulla oblongata; for size indices of components see Tables 14, 17, 20, 23,
 TEL telencephalon; for size indices of components see Table 44 and 26

Table 12. Average size indices of the five fundamental brain parts related to the average of the four Tenrecinae (AvTn = 100)

	N	OBL	MES	CER	DIE	TEL
slope		.61	.58	.68	.65	.68
y-intercept		.776	.481	.552	.350	1.242
Insectivora (AvIS)	50	132	136	167	169	159
Tenrecidae	12	144	137	147	144	128
Tenrecinae*	4	100	100	100	100	100
Geogalinae*	1	140	110	73	79	66
Oryzorictinae*	4	142	146	186	162	161
<i>Microgale</i>	2	126	163	171	158	180
terr. Oryzorictinae	3	117	141	172	152	169
<i>Limnogale</i>	1	216	162	228	192	137
Potamogalinae*	3	207	183	182	199	140
<i>Micropotamogale</i>	2	191	171	162	195	139
Chrysochloridae*	2	105	156	157	198	193
Solenodontidae*	1	149	158	255	177	179
Erinaceidae	5	131	175	176	194	165
Echinosoricinae*	2	176	254	258	251	225
Erinaceinae*	3	102	121	121	156	126
Talpidae	6	138	157	249	246	202
Desmaninae*	2	184	201	274	310	227
Talpinae*	4	116	134	237	214	189
<i>Talpa</i>	2	121	150	251	223	186
Soricidae	24	125	120	151	154	158
Soricinae*	11	117	108	150	166	176
<i>Sorex</i>	5	110	101	146	156	175
terr. Soricinae	9	112	103	144	161	177
<i>Neomys</i>	2	136	129	176	189	169
Crocidurinae*	13	132	130	152	144	144
<i>Crocidura</i>	6	127	125	135	132	131
<i>Suncus</i>	2	117	107	109	120	116
<i>Sylvisorex</i>	2	158	173	214	184	198
AvIF of the 12 families and subfamilies marked by *		139	150	179	180	161

Number of species = N

Abbreviations and notes:

- CER cerebellum; for av. size indices of components see Table 33
DIE diencephalon; for av. size indices of components see Tables 36, 39 and 42
MES mesencephalon; for av. size indices of components see Table 30
OBL medulla oblongata; for av. size indices of components see Tables 15, 18, 21, 24
TEL telencephalon; for av. size indices of components see Table 45 and 27

Table 13. Volumes of the sensory trigeminal complex and of the tractus solitarii and funicular nuclei of the medulla oblongata

For size indices in species see Table 14

For average size indices in supraspecific taxonomic units see Table 15

	n	TR	TSO	FUN	FGR	FCM	FCE
Tenrecidae							
<i>Tenrec ecaudatus</i>	1	49.3	5.37	5.56	1.351	2.325	1.879
<i>Setifer setosus</i>	1	27.5	2.37	3.02	0.770	1.511	0.737
<i>Hemicentetes semispin.</i>	1	17.6	2.20	2.40	0.599	0.990	0.809
<i>Echinops telfairi</i>	1	12.7	1.71	2.58	0.707	0.986	0.883
<i>Microgale talazaci</i>	1	13.8	1.73	1.70	0.444	0.752	0.506
<i>Potamogale velox</i>	1	151.4	9.70	8.25	2.813	2.820	2.616
<i>Micropotamogale ruwenz.</i>	1	47.6	3.63	2.70	0.989	1.293	0.422
Chrysochloridae							
<i>Chrysochloris stuhlmanni</i>	1	14.4	1.21	1.59	0.466	0.705	0.414
Solenodontidae							
<i>Solenodon paradoxus</i>	1	69.2	9.89	11.03	3.579	5.373	2.073
Erinaceidae							
<i>Echinosorex gymnurus</i>	1	90.5	9.53	10.66	3.417	3.094	4.153
<i>Hylomys suillus</i>	1	14.5	2.18	2.27	0.484	0.633	1.149
<i>Atelerix algirus</i>	1	39.9	6.18	9.23	2.079	3.073	4.075
<i>Erinaceus europaeus</i>	1	44.8	7.18	10.49	1.837	3.385	5.266
<i>Hemiechinus auritus</i>	1	23.9	3.63	4.46	0.941	1.485	2.029
Talpidae							
<i>Desmana moschata</i>	1	97.0	9.00	9.12	2.559	3.932	2.632
<i>Galemys pyrenaicus</i>	1	32.3	2.97	3.23	0.680	1.738	0.812
<i>Talpa europaea</i>	1	13.3	1.88	4.07	0.626	2.468	0.973
Soricidae							
<i>Sorex araneus</i>	1	6.0	0.70	0.31	0.095	0.132	0.087
<i>S. cinereus</i>	1	4.2	0.42	0.22	0.067	0.102	0.051
<i>Microsorex hoyi</i>	1	2.2	0.30	0.18	0.059	0.078	0.044
<i>Neomys fodiens</i>	1	9.7	1.24	0.66	0.212	0.254	0.195
<i>Blarina brevicauda</i>	1	8.1	1.26	0.70	0.216	0.260	0.227
<i>Crocidura flavescens</i>	1	10.8	1.07	0.89	0.384	0.408	0.094
<i>C. hildegardeae</i>	1	6.8	0.70	0.40	0.154	0.183	0.066
<i>C. russula</i>	1	5.3	0.53	0.32	0.151	0.126	0.039
<i>Suncus etruscus</i>	1	2.3	0.22	0.17	0.059	0.070	0.037
<i>S. murinus</i>	1	9.8	1.10	0.84	0.247	0.302	0.287
<i>Scutisorex somereni</i>	1	15.5	1.53	1.54	0.452	0.642	0.448
<i>Sylvisorex megalura</i>	1	4.9	0.61	0.51	0.168	0.173	0.165
<i>Ruwenzorisorex sunc.</i>	1	12.0	1.12	0.82	0.267	0.350	0.205

Number of individuals = n

Abbreviations as in Table 15

Table 14. Size indices of the sensory trigeminal complex and of the tractus solitarii and funicular nuclei of the medulla oblongata (AvTn = 100)

For volumes in species see Table 13

For average size indices in supraspecific taxonomic units see Table 15

	n	TR	TSO	FUN	FGR	FCM	FCE
slope		.59	.58	.62	.62	.62	.61
y-intercept		-.001	-.926	-.927	-1.515	-1.300	-1.400
Tenrecidae							
<i>Tenrec ecaudatus</i>	1	92	90	72	67	71	77
<i>Setifer setosus</i>	1	110	84	86	85	102	66
<i>Hemicentetes semispinosus</i>	1	107	118	106	103	104	112
<i>Echinops telfairi</i>	1	91	108	136	145	123	145
<i>Microgale talazaci</i>	1	140	154	130	131	136	120
<i>Potamogale velox</i>	1	342	197	130	171	105	130
<i>Micropotamogale ruwenzorii</i>	1	322	216	134	190	151	65
Chrysochloridae							
<i>Chrysochloris stuhlmanni</i>	1	163	120	136	154	142	109
Solenodontidae							
<i>Solenodon paradoxus</i>	1	149	191	165	207	189	98
Erinaceidae							
<i>Echisosorex gymnurus</i>	1	173	164	140	174	96	174
<i>Hylomys suillus</i>	1	133	176	156	129	103	245
<i>Atelerix algirus</i>	1	81	113	130	114	102	183
<i>Erinaceus europaeus</i>	1	84	121	135	92	103	216
<i>Hemiechinus auritus</i>	1	96	129	128	104	100	182
Talpidae							
<i>Desmana moschata</i>	1	267	221	176	192	179	161
<i>Galemys pyrenaicus</i>	1	290	234	216	176	275	168
<i>Talpa europaea</i>	1	99	123	224	133	320	166
Soricidae							
<i>Sorex araneus</i>	1	153	153	63	74	63	53
<i>S. cinereus</i>	1	161	138	67	79	74	47
<i>Microsorex hoyi</i>	1	122	145	84	106	86	61
<i>Neomys fodiens</i>	1	194	215	103	128	93	93
<i>Blarina brevicauda</i>	1	140	189	94	111	82	93
<i>Crociodura flavescens</i>	1	148	127	92	155	100	30
<i>C. hildegardeae</i>	1	175	155	82	121	88	40
<i>C. russula</i>	1	128	111	60	111	57	23
<i>Suncus etruscus</i>	1	162	127	96	132	95	64
<i>S. murinus</i>	1	123	121	80	91	68	84
<i>Scutisorex somereni</i>	1	134	116	99	113	98	90
<i>Sylvisorex megalura</i>	1	179	191	149	191	120	147
<i>Ruwenzorisorex suncooides</i>	1	217	175	115	145	116	88

Number of individuals = n

Abbreviations as in Table 15

Table 15. Average size indices of the sensory trigeminal complex and of the tractus solitarii and funicular nuclei related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 13

For size indices in species see Table 14

	N	TR	TSO	FUN	FGR	FCM	FCE
slope		.59	.58	.62	.62	.62	.61
y-intercept		-.001	-.926	-.927	-1.515	-1.300	-1.400
Insectivora (AvIS)	30	159	151	119	131	118	111
Tenrecidae	7	172	138	113	127	113	102
Tenrecinae*	4	100	100	100	100	100	100
Oryzorictinae*	1	140	154	130	131	136	120
Potamogalinae*	2	332	206	132	181	128	98
Chrysochloridae*	1	163	120	136	154	142	109
Solenodontidae*	1	149	191	165	207	189	98
Erinaceidae	5	113	141	138	123	101	200
Echinosoricinae*	2	153	170	148	152	100	209
Erinaceinae*	3	87	121	131	103	102	194
Talpidae	3	219	193	205	167	258	165
Desmaninae*	2	279	228	196	184	227	165
Talpinae*	1	99	123	224	133	320	166
Soricidae	13	157	151	91	120	88	70
Soricinae*	5	154	168	82	100	79	69
<i>Sorex</i>	2	157	145	65	76	68	50
terr. Soricinae	4	144	156	77	93	76	63
<i>Neomys</i>	1	194	215	103	128	93	93
Crociodurinae*	8	158	140	97	132	93	71
<i>Crociodura</i>	3	150	131	78	129	82	31
<i>Suncus</i>	2	143	124	88	112	82	74
AvIF of the 11 families and subfamilies marked by *		165	156	140	143	147	127

Number of species = N

Abbreviations:

FCE	ncl. fasc. cuneatus externus
FCM	ncl. fasc. cuneatus medialis
FGR	nucleus fascicularis gracilis
FUN	funicular nuclei
TR	compl. sensorius trigeminalis
TSO	nucleus tractus solitarii

Table 16. Volumes of some relay nuclei of the medulla oblongata

For size indices in species see Table 17

For average size indices in supraspecific taxonomic units see Table 18

	OBL	n	REL	INO	PRP	COE
Tenrecidae						
<i>Tenrec ecaudatus</i>	315.5	1	2.083	2.989	1.664	0.317
<i>Setifer setosus</i>	180.6	1	0.717	1.019	0.802	0.184
<i>Hemicentetes semispinosus</i>	117.0	1	0.799	1.009	0.826	0.172
<i>Echinops telfairi</i>	90.3	1	0.596	0.640	0.599	0.153
<i>Microgale talazaci</i>	84.0	1	0.616	0.894	0.603	0.105
<i>Potamogale velox</i>	722.8	1	4.799	5.567	3.089	0.651
<i>Micropotamogale ruwenz.</i>	198.0	1	1.327	1.377	1.088	0.299
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	68.0	1	0.433	1.007	0.523	0.074
Solenodontidae						
<i>Solenodon paradoxus</i>	471.6	1	4.181	4.933	2.894	0.566
Erinaceidae						
<i>Echinorex gymnurus</i>	622.4	1	4.947	7.576	3.669	0.594
<i>Hylomys suillus</i>	125.4	1	1.381	1.452	1.007	0.172
<i>Atelerix algirus</i>	343.0	1	3.590	3.401	1.923	—
<i>Erinaceus europaeus</i>	335.0	1	3.130	5.299	2.175	0.621
<i>Hemiechinus auritus</i>	185.2	1	1.813	1.959	0.987	0.324
Talpidae						
<i>Desmana moschata</i>	402.4	1	2.916	4.894	2.837	0.436
<i>Galemys pyrenaicus</i>	147.1	1	0.889	1.807	0.775	0.103
<i>Talpa europaea</i>	95.7	1	0.994	1.714	0.570	0.128
Soricidae						
<i>Sorex araneus</i>	29.1	1	0.240	0.342	0.171	0.035
<i>S. cinereus</i>	17.2	1	0.218	0.224	0.169	0.014
<i>Microsorex hoyi</i>	10.9	1	0.084	0.120	0.095	0.019
<i>Neomys fodiens</i>	45.3	1	0.242	0.486	0.351	0.039
<i>Blarina brevicauda</i>	43.0	1	0.307	0.558	0.379	0.036
<i>Crocidura flavescens</i>	59.9	1	0.554	0.458	0.422	0.041
<i>C. hildegardeae</i>	31.4	1	0.291	0.303	0.239	0.051
<i>C. russula</i>	28.7	1	0.165	0.285	0.213	0.032
<i>Suncus etruscus</i>	11.1	1	0.092	0.131	0.090	0.008
<i>S. murinus</i>	54.4	1	0.414	0.612	0.330	0.043
<i>Scutisorex somereni</i>	82.9	1	0.630	0.980	0.659	0.085
<i>Sylvisorex megalura</i>	26.8	1	0.268	0.268	0.207	0.020
<i>Ruwenzorisorex suncoides</i>	54.9	1	0.454	0.467	0.384	0.064

Number of individuals = n

Abbreviations:

COE nucleus locus coeruleus
INO complexus olivaris inferior
OBL medulla oblongata

PRP nucleus prepositus hypoglossi
REL nucleus reticularis lateralis

Table 17. Size indices of some relay nuclei of the medulla oblongata related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 16

For average size indices in supraspecific taxonomic units see Table 18

	OBL	n	REL	INO	PRP	COE
slope	.61		.58	.61	.52	.64
y-intercept	.776		-1.379	-1.340	-1.250	-2.176
Tenrecidae						
<i>Tenrec ecaudatus</i>	86	1	100	107	89	63
<i>Setifer setosus</i>	108	1	72	79	83	83
<i>Hemicentetes semispinosus</i>	108	1	121	121	124	123
<i>Echinops telfairi</i>	99	1	107	92	104	131
<i>Microgale talazaci</i>	132	1	156	184	143	132
<i>Potamogale velox</i>	240	1	276	242	194	160
<i>Micropotamogale ruwenz.</i>	204	1	224	185	179	240
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	120	1	122	231	136	104
Solenodontidae						
<i>Solenodon paradoxus</i>	149	1	229	203	174	132
Erinaceidae						
<i>Echinosorex gymnurus</i>	174	1	241	276	199	121
<i>Hylomys suillus</i>	178	1	317	270	219	194
<i>Atelerix algirus</i>	102	1	187	133	110	--
<i>Erinaceus europaeus</i>	92	1	150	189	116	124
<i>Hemiechinus auritus</i>	111	1	183	153	103	148
Talpidae						
<i>Desmana moschata</i>	164	1	204	260	212	132
<i>Galemys pyrenaicus</i>	203	1	199	326	164	113
<i>Talpa europaea</i>	109	1	185	255	102	114
Soricidae						
<i>Sorex araneus</i>	118	1	149	181	91	119
<i>S. cinereus</i>	106	1	201	180	128	72
<i>Microsorex hoyi</i>	101	1	115	146	102	154
<i>Neomys fodiens</i>	144	1	119	201	151	102
<i>Blarina brevicauda</i>	117	1	130	198	143	80
<i>Crocidura flavescens</i>	128	1	187	128	130	71
<i>C. hildegardeae</i>	128	1	183	163	128	177
<i>C. russula</i>	111	1	98	144	108	104
<i>Suncus etruscus</i>	127	1	154	196	116	79
<i>S. murinus</i>	106	1	129	156	94	67
<i>Scutisorex somereni</i>	110	1	136	171	135	89
<i>Sylvisorex megalura</i>	159	1	239	207	152	101
<i>Ruwenzorisorex suncoides</i>	157	1	202	174	151	150

Number of individuals = n

Abbreviations as in Tables 16 and 18

Table 18. Average size indices of some relay nuclei of the medulla oblongata related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 16
For size indices in species see Table 17

	N	OBL	REL	INO	PRP	COE
slope		.61	.58	.61	.52	.64
y-intercept		.776	-1.379	-1.340	-1.250	-2.176
Insectivora (AvIS)	30+)	133	170	185	136	120
Tenrecidae	7	140	151	144	131	133
Tenrecinae*	4	100	100	100	100	100
Oryzorictinae*	1	132	156	184	143	132
Potamogalinae*	2	222	250	213	187	200
Chrysochloridae*	1	120	122	231	136	104
Solenodontidae*	1	149	229	203	174	132
Erinaceidae	5 +)	131	216	204	149	147
Echinosoricinae*	2	176	279	273	209	158
Erinaceinae*	3 +)	102	173	158	110	136
Talpidae	3	159	196	280	159	120
Desmaninae*	2	184	201	293	188	123
Talpinae*	1	109	185	255	102	114
Soricidae	13	124	157	173	125	105
Soricinae*	5	117	143	181	123	105
<i>Sorex</i>	2	112	175	181	109	96
terr. Soricinae	4	111	149	176	116	106
<i>Neomys</i>	1	144	119	201	151	102
Crocidurinae*	8	128	166	167	127	105
<i>Crocidura</i>	3	122	156	145	122	117
<i>Suncus</i>	2	117	141	176	105	73
AvIF of the 11 families and subfamilies marked by *		140	182	205	145	128

Number of species = N +) for COE = N – 1
Abbreviations:

- COE nucleus locus coeruleus
- INO complexus olivaris inferior
- OBL medulla oblongata
- PRP nucleus prepositus hypoglossi
- REL nucleus reticularis lateralis

Table 19. Volumes of motor nuclei of the medulla oblongata

For size indices in species see Table 20

For average size indices in supraspecific taxonomic units see Table 21

	n	MOT	V	VII	X	XII
Tenrecidae						
<i>Tenrec ecaudatus</i>	1	7.88	1.890	3.585	0.829	1.572
<i>Setifer setosus</i>	1	4.14	0.832	2.099	0.372	0.841
<i>Hemicentetes semispinosus</i>	1	2.57	0.427	1.342	0.278	0.521
<i>Echinops telfairi</i>	1	2.66	0.385	1.444	0.281	0.549
<i>Microgale talazaci</i>	1	2.00	0.391	1.109	0.208	0.295
<i>Potamogale velox</i>	1	15.81	2.255	10.839	1.117	1.598
<i>Micropotamogale ruwenzorii</i>	1	3.77	0.783	2.154	0.355	0.482
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	1	1.53	0.328	0.691	0.190	0.322
Solenodontidae						
<i>Solenodon paradoxus</i>	1	9.48	2.378	4.325	0.990	1.785
Erinaceidae						
<i>Echinosorex gymnurus</i>	1	16.68	3.378	8.810	1.460	3.036
<i>Hylomys suillus</i>	1	3.41	0.601	1.814	0.391	0.601
<i>Atelerix algirus</i>	1	9.65	1.553	4.560	0.920	2.612
<i>Erinaceus europaeus</i>	1	11.14	1.693	5.366	1.293	2.785
<i>Hemiechinus auritus</i>	1	5.30	0.818	2.468	0.486	1.530
Talpidae						
<i>Desmana moschata</i>	1	8.45	1.295	5.003	0.799	1.355
<i>Galemys pyrenaicus</i>	1	2.86	0.528	1.496	0.271	0.564
<i>Talpa europaea</i>	1	2.01	0.385	1.023	0.284	0.323
Soricidae						
<i>Sorex araneus</i>	1	0.76	0.143	0.344	0.098	0.171
<i>S. cinereus</i>	1	0.35	0.049	0.175	0.039	0.087
<i>Microsorex hoyi</i>	1	0.28	0.038	0.124	0.044	0.071
<i>Neomys fodiens</i>	1	1.03	0.181	0.500	0.116	0.234
<i>Blarina brevicauda</i>	1	1.04	0.169	0.418	0.216	0.236
<i>Crocidura flavescens</i>	1	1.11	0.186	0.549	0.160	0.216
<i>C. hildegardeae</i>	1	0.61	0.108	0.303	0.089	0.108
<i>C. russula</i>	1	0.58	0.070	0.329	0.075	0.107
<i>Suncus etruscus</i>	1	0.24	0.030	0.130	0.028	0.048
<i>S. murinus</i>	1	1.49	0.217	0.798	0.152	0.322
<i>Scutisorex somereni</i>	1	1.71	0.241	0.875	0.262	0.334
<i>Sylvisorex megalura</i>	1	0.54	0.068	0.298	0.075	0.103
<i>Ruwenzorisorex suncoides</i>	1	1.12	0.136	0.614	0.157	0.216

Number of individuals = n

Abbreviations:

MOT sum of measured motor nuclei

V ncl. motorius nervi trigemini

VII ncl. nervi facialis

X

ncl. dorsalis motorius nervi vagi

XII

ncl. nervi hypoglossi

Table 20. Size indices of motor nuclei of the medulla oblongata related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 19

For average size indices in supraspecific taxonomic units see Table 21

	n	MOT	V	VII	X	XII
slope		.62	.67	.62	.62	.59
y-intercept		-.853	-1.705	-1.146	-1.844	-1.476
Tenrecidae						
<i>Tenrec ecaudatus</i>	1	86	104	76	88	88
<i>Setifer setosus</i>	1	100	108	99	88	100
<i>Hemicentetes semispinosus</i>	1	96	90	99	102	94
<i>Echinops telfairi</i>	1	118	98	126	123	117
<i>Microgale talazaci</i>	1	129	148	140	131	90
<i>Potamogale velox</i>	1	210	154	282	145	108
<i>Micropotamogale ruwenzorii</i>	1	158	185	177	146	97
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	1	110	140	98	134	109
Solenodontidae						
<i>Solenodon paradoxus</i>	1	119	154	107	122	115
Erinaceidae						
<i>Echinosorex gymnurus</i>	1	185	191	192	159	173
<i>Hylomys suillus</i>	1	198	203	207	223	166
<i>Atelerix algirus</i>	1	115	94	107	107	159
<i>Erinaceus europaeus</i>	1	121	94	115	138	156
<i>Hemiechinus auritus</i>	1	128	107	117	115	183
Talpidae						
<i>Desmana moschata</i>	1	138	111	160	128	111
<i>Galemys pyrenaicus</i>	1	161	173	166	150	151
<i>Talpa europaea</i>	1	93	102	93	129	72
Soricidae						
<i>Sorex araneus</i>	1	128	153	114	162	130
<i>S. cinereus</i>	1	90	83	88	99	98
<i>Microsorex hoyi</i>	1	108	101	95	168	119
<i>Neomys fodiens</i>	1	135	148	129	149	140
<i>Blarina brevicauda</i>	1	117	116	92	238	122
<i>Crocidura flavescens</i>	1	98	98	95	138	88
<i>C. hildegardeae</i>	1	104	117	102	148	83
<i>C. russula</i>	1	93	70	104	117	77
<i>Suncus etruscus</i>	1	115	99	124	134	100
<i>S. murinus</i>	1	120	104	126	120	121
<i>Scutisorex somereni</i>	1	93	76	93	140	86
<i>Sylvisorex megalura</i>	1	135	109	145	182	113
<i>Ruwenzorisorex suncoides</i>	1	132	99	142	181	117

Number of individuals = n

Abbreviations as in Tables 19 and 21

Table 21. Average size indices of motor nuclei of the medulla oblongata related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 19

For size indices in species see Table 20

	N	MOT	V	VII	X	XII
slope		.62	.67	.62	.62	.59
y-intercept		-.853	-1.705	-1.146	-1.844	-1.476
Insectivora (AvIS)	30	124	121	127	140	116
Tenrecidae	7	128	127	143	118	99
Tenrecinae*	4	100	100	100	100	100
Oryzorictinae*	1	129	148	140	131	90
Potamogalinae*	2	184	170	230	145	103
Chrysochloridae*	1	110	140	98	134	109
Solenodontidae*	1	119	154	107	122	115
Erinaceidae	5	149	138	148	148	167
Echinosoricinae*	2	192	197	200	191	169
Erinaceinae*	3	121	98	113	120	166
Talpidae	3	131	129	140	136	111
Desmaninae*	2	150	142	163	139	131
Talpinae*	1	93	102	93	129	72
Soricidae	13	113	106	111	152	107
Soricinae*	5	116	120	104	163	122
<i>Sorex</i>	2	109	118	101	131	114
terr. Soricinae	4	111	113	97	167	117
<i>Neomys</i>	1	135	148	129	149	140
Crocidurinae*	8	111	97	116	145	98
<i>Crocidura</i>	3	98	95	100	134	83
<i>Suncus</i>	2	117	102	125	127	110
AvIF of the 11 families and subfamilies marked by *		130	133	133	138	116

Number of species = N

Abbreviations:

- MOT sum. of measured motor nuclei
 V ncl. motorius nervi trigemini
 VII ncl. nervi facialis
 X ncl. dorsalis motorius nervi vagi
 XII ncl. nervi hypoglossi

Table 22. Volumes of the vestibular nuclei of the medulla oblongata

For size indices in species see Table 23

For average size indices in supraspecific taxonomic units see Table 24

	n	VC	VM	VI	VL	VS
Tenrecidae						
<i>Tenrec ecaudatus</i>	1	14.44	5.152	4.270	2.885	2.132
<i>Setifer setosus</i>	1	7.18	2.705	2.009	1.090	1.379
<i>Hemicentetes semispinosus</i>	1	5.69	2.125	1.823	0.995	0.748
<i>Echinops telfairi</i>	1	3.85	1.482	1.111	0.625	0.634
<i>Microgale talazaci</i>	1	3.44	1.519	0.883	0.681	0.361
<i>Potamogale velox</i>	1	25.00	8.039	9.248	3.451	4.266
<i>Micropotamogale ruwenzorii</i>	1	8.58	3.259	2.626	1.277	1.421
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	1	3.04	1.286	0.796	0.504	0.456
Solenodontidae						
<i>Solenodon paradoxus</i>	1	22.05	8.800	7.285	2.936	3.030
Erinaceidae						
<i>Echinosorex gymnurus</i>	1	33.49	14.390	8.991	7.053	3.055
<i>Hylomys suillus</i>	1	8.52	3.603	2.332	1.462	1.122
<i>Atelerix algirus</i>	1	15.56	6.417	4.774	2.399	1.972
<i>Erinaceus europaeus</i>	1	15.90	6.438	3.979	3.256	2.231
<i>Hemiechinus auritus</i>	1	8.59	3.422	2.273	1.716	1.180
Talpidae						
<i>Desmana moschata</i>	1	19.49	8.125	6.485	2.541	2.341
<i>Galemys pyrenaicus</i>	1	6.16	2.612	2.085	0.793	0.666
<i>Talpa europaea</i>	1	4.14	1.671	1.418	0.460	0.590
Soricidae						
<i>Sorex araneus</i>	1	1.29	0.539	0.266	0.264	0.219
<i>S. cinereus</i>	1	0.88	0.367	0.262	0.148	0.106
<i>Microsorex hoyi</i>	1	0.55	0.206	0.156	0.108	0.076
<i>Neomys fodiens</i>	1	2.32	0.964	0.728	0.356	0.270
<i>Blarina brevicauda</i>	1	2.22	0.997	0.668	0.224	0.326
<i>Crocidura flavescens</i>	1	2.15	0.924	0.554	0.422	0.246
<i>C. hildegardeae</i>	1	1.57	0.674	0.470	0.237	0.185
<i>C. russula</i>	1	1.19	0.544	0.346	0.171	0.131
<i>Suncus etruscus</i>	1	0.52	0.265	0.133	0.059	0.061
<i>S. murinus</i>	1	2.14	0.932	0.595	0.385	0.230
<i>Scutisorex somereni</i>	1	3.55	1.416	1.069	0.613	0.448
<i>Sylvisorex megalura</i>	1	1.40	0.644	0.325	0.273	0.158
<i>Ruwenzorisorex suncoides</i>	1	2.51	1.129	0.886	0.235	0.259

Number of individuals = n		Abbreviations:	
VC	vestibular complex	VM	nucleus vestibularis medialis
VI	nucleus vestibularis inferior	VS	nucleus vestibularis superior
VL	nucleus vestibularis lateralis		

Table 23. Size indices of the vestibular nuclei of the medulla oblongata related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 22

For average size indices in supraspecific taxonomic units see Table 24

	n	VC	VM	VI	VL	VS
slope		.53	.53	.55	.54	.51
y-intercept		-.393	-.822	-.966	-1.179	-1.149
Tenrecidae						
<i>Tenrec ecaudatus</i>	1	100	96	97	114	96
<i>Setifer setosus</i>	1	98	99	92	86	120
<i>Hemicentetes semispinosus</i>	1	113	114	123	115	93
<i>Echinops telfairi</i>	1	89	92	88	84	91
<i>Microgale talazaci</i>	1	109	129	97	127	70
<i>Potamogale velox</i>	1	205	177	249	162	227
<i>Micropotamogale ruwenzorii</i>	1	188	192	196	163	194
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	1	106	121	97	104	98
Solenodontidae						
<i>Solenodon paradoxus</i>	1	173	185	188	132	154
Erinaceidae						
<i>Echinosorex gymnurus</i>	1	236	272	207	284	140
<i>Hylomys suillus</i>	1	247	281	233	249	201
<i>Atelerix algirus</i>	1	116	129	117	103	96
<i>Erinaceus europaeus</i>	1	110	120	90	129	101
<i>Hemiechinus auritus</i>	1	118	126	104	136	103
Talpidae						
<i>Desmana moschata</i>	1	191	213	210	143	147
<i>Galemys pyrenaicus</i>	1	174	198	203	132	117
<i>Talpa europaea</i>	1	99	107	116	64	88
Soricidae						
<i>Sorex araneus</i>	1	93	104	69	114	94
<i>S. cinereus</i>	1	91	102	98	92	65
<i>Microsorex hoyi</i>	1	81	82	85	97	66
<i>Neomys fodiens</i>	1	135	151	150	123	95
<i>Blarina brevicauda</i>	1	113	136	120	68	100
<i>Crocidura flavescens</i>	1	89	102	80	103	62
<i>C. hildegardeae</i>	1	114	132	122	103	81
<i>C. russula</i>	1	82	101	85	70	54
<i>Suncus etruscus</i>	1	92	127	87	64	62
<i>S. murinus</i>	1	82	96	79	87	54
<i>Scutisorex somereni</i>	1	97	104	101	98	76
<i>Sylvisorex megalura</i>	1	140	173	118	164	93
<i>Ruwenzorisorex suncoides</i>	1	133	161	166	74	83

Number of individuals = n

Abbreviations as in Tables 22 and 24

Table 24. Average size indices of vestibular nuclei of the medulla oblongata related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 22

For size indices in species see Table 23

	N	VC	VM	VI	VL	VS
slope		.53	.53	.55	.54	.51
y-intercept		-.393	-.822	-.966	-1.179	-1.149
Insectivora (AvIS)	30	127	141	129	119	104
Tenrecidae	7	129	128	135	122	127
Tenrecinae*	4	100	100	100	100	100
Oryzorictinae*	1	109	129	97	127	70
Potamogalinae*	2	197	184	223	163	211
Chrysochloridae*	1	106	121	97	104	98
Solenodontidae*	1	173	185	188	132	154
Erinaceidae	5	165	186	150	180	128
Echinisoricinae*	2	241	276	220	266	171
Erinaceinae*	3	115	125	104	122	100
Talpidae	3	155	173	176	113	117
Desmaninae*	2	182	206	207	137	132
Talpinae*	1	99	107	116	64	88
Soricidae	13	103	121	105	97	76
Soricinae*	5	103	115	104	99	84
<i>Sorex</i>	2	92	103	83	103	80
terr. Soricinae	4	95	106	93	93	81
<i>Neomys</i>	1	135	151	150	123	95
Crocidurinae*	8	104	125	105	96	71
<i>Crocidura</i>	3	95	112	96	92	66
<i>Suncus</i>	2	87	111	83	75	58
AvIF of the 11 families and subfamilies marked by *		139	152	142	128	116

Number of species = N

Abbreviations:

VC	vestibular complex
VI	nucleus vestibularis inferior
VL	nucleus vestibularis lateralis
VM	nucleus vestibularis medialis
VS	nucleus vestibularis superior

Table 25. Volumes of the auditory nuclei of the medulla oblongata

For size indices in species see Table 26

For average size indices in supraspecific taxonomic units see Table 27

	n	AUD	CON	DCO	VCO	OLS
Tenrecidae						
<i>Tenrec ecaudatus</i>	1	14.14	8.43	4.23	4.20	5.71
<i>Setifer setosus</i>	1	9.59	5.54	2.00	3.54	4.05
<i>Hemicentetes semispinosus</i>	1	4.78	2.87	1.01	1.86	1.91
<i>Echinops telfairi</i>	1	6.81	4.39	1.63	2.76	2.42
<i>Microgale talazaci</i>	1	4.83	3.13	1.31	1.82	1.70
<i>Potamogale velox</i>	1	19.72	10.48	4.06	6.42	9.24
<i>Micropotamogale ruwenzorii</i>	1	5.36	2.94	1.33	1.61	2.42
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	1	3.09	1.71	0.77	0.94	1.38
Solenodontidae						
<i>Solenodon paradoxus</i>	1	19.32	11.70	4.65	7.05	7.62
Erinaceidae						
<i>Echinorex gymnurus</i>	1	35.20	22.16	11.82	10.34	13.04
<i>Hylomys suillus</i>	1	8.27	4.69	1.91	2.78	3.58
<i>Atelerix algirus</i>	1	14.93	8.56	3.55	5.01	6.37
<i>Erinaceus europaeus</i>	1	10.95	6.84	2.15	4.69	4.11
<i>Hemiechinus auritus</i>	1	11.01	5.95	2.07	3.88	5.06
Talpidae						
<i>Desmana moschata</i>	1	11.45	5.31	2.47	2.84	6.14
<i>Galemys pyrenaicus</i>	1	3.25	1.88	0.85	1.03	1.37
<i>Talpa europaea</i>	1	3.53	2.04	0.99	1.05	1.49
Soricidae						
<i>Sorex araneus</i>	1	0.90	0.61	0.30	0.31	0.29
<i>S. cinereus</i>	1	0.62	0.38	0.18	0.20	0.24
<i>Microsorex hoyi</i>	1	0.34	0.22	0.10	0.12	0.12
<i>Neomys fodiens</i>	1	1.22	0.72	0.33	0.39	0.50
<i>Blarina brevicauda</i>	1	1.42	0.82	0.37	0.45	0.60
<i>Crocidura flavescens</i>	1	3.11	1.92	1.01	0.91	1.19
<i>C. hildegardeae</i>	1	1.89	1.12	0.50	0.62	0.77
<i>C. russula</i>	1	1.47	0.90	0.45	0.45	0.57
<i>Suncus etruscus</i>	1	0.50	0.30	0.14	0.16	0.20
<i>S. murinus</i>	1	2.87	1.86	0.83	1.03	1.01
<i>Scutisorex somereni</i>	1	5.45	3.59	1.36	2.23	1.86
<i>Sylvisorex megalura</i>	1	1.74	1.07	0.52	0.55	0.67
<i>Ruwenzorisorex suncoides</i>	1	2.53	1.59	0.77	0.82	0.94

Number of individuals = n

AUD auditory (acoustic) nuclei

CON cochlear nuclei

DCO nucleus cochlearis dorsalis

Abbreviations:

OLS complexus olivaris superior

VCO nucleus cochlearis ventralis

Table 26. Size indices of the auditory nuclei related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 25

For average size indices in supraspecific taxonomic units see Table 27

	n	AUD	CON	DCO	VCO	OLS
slope		.51	.50	.50	.51	.54
y-intercept		-.267	-.461	-.867	-.700	-.742
Tenrecidae						
<i>Tenrec ecaudatus</i>	1	84	83	107	67	82
<i>Setifer setosus</i>	1	109	104	96	109	117
<i>Hemicentetes semispinosus</i>	1	78	77	69	82	81
<i>Echinops telfairi</i>	1	129	136	128	141	119
<i>Microgale talazaci</i>	1	124	130	139	126	116
<i>Potamogale velox</i>	1	138	122	120	121	159
<i>Micropotamogale ruwenzorii</i>	1	96	86	99	78	113
Chrysochloridae						
<i>Chrysochloris stuhlmanni</i>	1	87	78	90	71	104
Solenodontidae						
<i>Solenodon paradoxus</i>	1	129	130	132	128	125
Erinaceidae						
<i>Echinorex gymnurus</i>	1	212	223	303	169	192
<i>Hylomys suillus</i>	1	195	180	187	177	223
<i>Atelerix algirus</i>	1	95	91	96	87	99
<i>Erinaceus europaeus</i>	1	65	68	54	75	59
<i>Hemiechinus auritus</i>	1	126	112	100	120	147
Talpidae						
<i>Desmana moschata</i>	1	95	73	86	64	126
<i>Galemys pyrenaicus</i>	1	75	70	81	64	83
<i>Talpa europaea</i>	1	69	65	81	56	76
Soricidae						
<i>Sorex araneus</i>	1	51	55	70	47	46
<i>S. cinereus</i>	1	50	49	57	45	54
<i>Microsorex hoyi</i>	1	39	39	43	38	40
<i>Neomys fodiens</i>	1	56	53	62	49	64
<i>Blarina brevicauda</i>	1	57	53	61	50	66
<i>Crociodura flavescens</i>	1	103	103	137	82	106
<i>C. hildegardeae</i>	1	108	102	116	96	122
<i>C. russula</i>	1	80	78	100	66	86
<i>Suncus etruscus</i>	1	67	64	78	57	77
<i>S. murinus</i>	1	88	93	105	86	83
<i>Scutisorex somereni</i>	1	122	130	126	134	109
<i>Sylvisorex megalura</i>	1	135	132	162	116	148
<i>Ruwenzorisorex suncoides</i>	1	107	108	132	94	109

Number of individuals = n

Abbreviations as in Tables 25 and 27

Table 27. Average size indices of the auditory nuclei related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 25

For size indices in species see Table 26

	N	AUD	CON	DCO	VCO	OLS
slope		.51	.50	.50	.51	.54
y-intercept		-.267	-.461	-.867	-.700	-.742
Insectivora (AvIS)	30	99	96	107	90	104
Tenrecidae	7	108	105	108	103	112
Tenrecinae*	4	100	100	100	100	100
Oryzorictinae*	1	124	130	139	126	116
Potamogalinae*	2	117	104	110	100	136
Chrysochloridae*	1	87	78	90	71	104
Solenodontidae*	1	129	130	132	128	125
Erinaceidae	5	139	135	148	126	144
Echinosoricinae*	2	203	201	245	173	207
Erinaceinae*	3	95	90	83	94	102
Talpidae	3	80	69	83	61	95
Desmaninae*	2	85	72	84	64	105
Talpinae*	1	69	65	81	56	76
Soricidae	13	82	81	96	74	85
Soricinae*	5	51	50	59	46	54
<i>Sorex</i>	2	50	52	63	46	50
terr. Soricinae	4	49	49	58	45	51
<i>Neomys</i>	1	56	53	62	49	64
Crocidurinae*	8	101	101	120	91	105
<i>Crocidura</i>	3	97	94	118	81	105
<i>Suncus</i>	2	78	78	92	72	80
AvIF of the 11 families and subfamilies marked by *		106	102	113	95	112

Number of species = N

Abbreviations:

- AUD auditory (acoustic) nuclei
- CON cochlear nuclei
- DCO nucleus cochlearis dorsalis
- OLS complexus olivaris superior
- VCO nucleus cochlearis ventralis

Table 28. Volumes of mesencephalic components

For size indices in species see Table 29
For average size indices in supraspecific taxonomic units see Table 30
For volumes of subcommissural body (SCB) see Table 40

	n	MES	n	MTG	NLL	MTC	SUC	INC
Tenrecidae								
<i>Tenrec ecaudatus</i>	3	133.3	1	72.82	4.48	60.49	24.73	35.76
<i>Setifer setosus</i>	3	76.6	1	44.94	3.24	31.68	9.69	21.99
<i>Hemicentetes semisp.</i>	2	45.9	1	27.17	2.40	18.69	8.33	10.36
<i>Echinops telfairi</i>	2	44.0	1	19.72	2.23	24.29	8.50	15.79
Erinaceidae								
<i>Erinaceus europaeus</i>	5	151.3	1	92.55	3.72	58.74	29.57	29.17
Soricidae								
<i>Sorex araneus</i>	5	11.5	1	7.15	0.62	4.34	2.69	1.65
<i>S. minutus</i>	4	6.4	1	3.90	0.19	2.53	1.62	0.91
<i>Crocidura flavescens</i>	3	26.7	1	14.58	1.33	12.15	5.54	6.61
<i>C. russula</i>	5	13.7	1	7.78	0.65	5.89	3.04	2.85
<i>Suncus murinus</i>	3	24.6	1	13.98	1.01	10.61	5.09	5.52

Number of individuals = n

Abbreviations:

- INC colliculi inferiores
- MES mesencephalon
- MTC tectum mesencephali (colliculi superiores + inferiores)
- MTG tegmentum mesencephali
- NLL nucleus lemnisci lateralis
- SUC colliculi superiores

Table 29. Size indices of mesencephalic components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 28

For average size indices in supraspecific taxonomic units see Table 30

For size indices of subcommissural body (SCB) see Table 41

	n	MES	n	MTG	NLL	MTC	SUC	INC
slope		.58		.58	.58	.58	.58	.58
y-intercept		.481		.213	-.865	.141	-.287	-.061
Tenrecidae								
<i>Tenrec ecaudatus</i>	3	88	1	89	66	87	96	82
<i>Setifer setosus</i>	3	106	1	115	100	96	79	106
<i>Hemicentetes semispin.</i>	2	96	1	106	112	86	102	76
<i>Echinops telfairi</i>	2	109	1	90	122	131	123	136
Erinaceidae								
<i>Erinaceus europaeus</i>	5	100	1	113	54	85	115	67
Soricidae								
<i>Sorex araneus</i>	5	99	1	114	117	82	135	49
<i>S. minutus</i>	4	90	1	101	60	77	132	44
<i>Crocidura flavescens</i>	3	125	1	126	137	124	151	107
<i>C. russula</i>	5	112	1	118	118	105	146	81
<i>Suncus murinus</i>	3	105	1	111	96	100	128	82

Number of individuals = n

Abbreviations:

INC	colliculi inferiores
MES	mesencephalon
MTC	tectum mesencephali (colliculi superiores + inferiores)
MTG	tegmentum mesencephali
NLL	nucleus lemnisci lateralis
SUC	colliculi superiores

Table 30. Average size indices of mesencephalic components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 28							
For size indices in species see Table 29							
For average size indices of subcommissural body (SCB) see Table 42							
	N	MES	MTG	NLL	MTC	SUC	INC
slope		.58	.58	.58	.58	.58	.58
y-intercept		.481	.213	-.865	.141	-.287	-.061
Insectivora (AvIS)	10	103	108	98	97	121	83
Tenrecidae							
Tenrecinae*	4	100	100	100	100	100	100
Erinaceidae							
Erinaceinae*	1	100	113	54	85	115	67
Soricidae	5	106	114	106	98	138	73
Soricinae*	2	94	107	88	79	134	47
Crœcidurinae*	3	114	118	117	110	142	90
<i>Crocidura</i>	2	118	122	128	115	148	94
AvIF of the 4 subfamilies marked by *		102	110	90	94	123	76

Number of species = N

Abbreviations:

INC	colliculi inferiores
MES	mesencephalon
MTC	tectum mesencephali (colliculi superiores + inferiores)
MTG	tegmentum mesencephali
NLL	nucleus lemnisci lateralis
SUC	colliculi superiores

Table 31. Volumes of cerebellar nuclei and ventral pons

For size indices in species see Table 32

For average size indices in supraspecific taxonomic units see Table 33

	n	CER	n	VPO	TCN	MCN	ICN	LCN
Tenrecidae								
<i>Tenrec ecaudatus</i>	3	311.5	1	2.524	9.890	2.074	4.465	3.351
<i>Setifer setosus</i>	3	146.8	1	.-	4.331	1.239	1.845	1.247
<i>Hemicentetes semisp.</i>	2	120.5	1	.-	3.332	0.563	1.572	1.197
<i>Echinops telfairi</i>	2	58.2	1	0.494	1.720	0.390	0.870	0.460
Erinaceidae								
<i>Erinaceus europaeus</i>	5	395.8	1	.-	12.716	3.315	5.556	3.845
Soricidae								
<i>Sorex araneus</i>	5	22.0	1	.-	0.579	0.179	0.185	0.215
<i>S. minutus</i>	4	12.1	1	.-	0.359	0.095	0.128	0.136
<i>Crocidura flavescens</i>	3	51.9	1	.-	1.145	0.268	0.415	0.462
<i>C. russula</i>	5	21.0	1	.-	0.564	0.114	0.225	0.225
<i>Suncus murinus</i>	3	42.8	1	.-	1.077	0.238	0.389	0.450

Number of individuals = n

Abbreviations:

CER	cerebellum
ICN	nucleus interpositus cerebelli
LCN	nucleus lateralis cerebelli
MCN	nucleus medialis cerebelli
TCN	total cerebellar nuclei
VPO	ventral pons

Table 32. Size indices of cerebellar nuclei and ventral pons related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 31

For average size indices in supraspecific taxonomic units see Table 33

	n	CER	n	VPO	TCN	MCN	ICN	LCN
slope		.68		.68	.68	.68	.68	.68
y-intercept		.552		-1.608	-.979	-1.638	-1.314	-1.476
Tenrecidae								
<i>Tenrec ecaudatus</i>	3	89	1	104	96	92	94	102
<i>Setifer setosus</i>	3	100	1	--	100	131	92	91
<i>Hemicentetes semispin.</i>	2	133	1	--	125	97	128	141
<i>Echinops telfairi</i>	2	78	1	96	78	81	86	66
Erinaceidae								
<i>Erinaceus europaeus</i>	5	113	1	--	124	147	117	117
Soricidae								
<i>Sorex araneus</i>	5	127	1	--	114	160	79	133
<i>S. minutus</i>	4	124	1	--	125	150	96	148
<i>Crocidura flavescens</i>	3	146	1	--	110	117	86	139
<i>C. russula</i>	5	114	1	--	105	96	90	131
<i>Suncus murinus</i>	3	110	1	--	94	94	73	123

Number of individuals = n

Abbreviations:

- CER cerebellum
- ICN nucleus interpositus cerebelli
- LCN nucleus lateralis cerebelli
- MCN nucleus medialis cerebelli
- TCN total cerebellar nuclei
- VPO ventral pons

Table 33. Average size indices of cerebellar nuclei and ventral pons related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 31

For size indices in species see Table 32

	N	CER	TCN	MCN	ICN	LCN
slope		.68	.68	.68	.68	.68
y-intercept		.552	-.979	-1.638	-1.314	-1.476
Insectivora (AvIS)	10	113	107	117	94	119
Tenrecidae						
Tenrecinae*	4	100	100	100	100	100
Erinaceidae						
Erinaceinae*	1	113	124	147	117	117
Soricidae	5	124	110	123	85	135
Soricinae*	2	126	119	155	87	140
Crocidurinae*	3	123	103	102	83	131
<i>Crocidura</i>	2	130	107	107	88	135
AvIF of the 4 subfamilies marked by *		115	112	126	97	122

Number of species = N

Abbreviations:

CER	cerebellum
ICN	nucleus interpositus cerebelli
LCN	nucleus lateralis cerebelli
MCN	nucleus medialis cerebelli
TCN	total cerebellar nuclei

Table 34. Volumes of diencephalic components

For size indices in species see Table 35
For average size indices in supraspecific taxonomic units see Table 36
For volumes of subthalamic components and geniculate bodies see Table 37
For volumes of medial nuclei of habenula (HAM) see Table 40

	n	DIE	ETH	THA	STH	HTH	CAI	TRO
Tenrecidae								
<i>Tenrec ecaudatus</i>	2	153.7	2.890	66.67	12.29	54.94	14.03	2.903
<i>Setifer setosus</i>	1	93.1	1.557	39.89	5.58	36.82	8.24	1.029
<i>Hemicentetes semispin.</i>	1	47.7	1.261	20.66	3.60	18.59	3.05	0.551
<i>Echinops telfairi</i>	1	40.3	0.965	17.67	2.65	16.19	2.20	0.597
Erinaceidae								
<i>Erinaceus europaeus</i>	2	242.4	3.789	119.50	14.49	81.51	19.09	4.041
Soricidae								
<i>Sorex araneus</i>	2	14.8	0.269	7.38	1.11	5.26	0.62	0.146
<i>S. minutus</i>	1	7.7	0.150	3.65	0.66	2.86	0.27	0.099
<i>Crocidura flavescens</i>	2	25.1	0.564	12.50	1.52	8.40	1.88	0.217
<i>C. russula</i>	1	13.9	0.360	7.31	0.85	4.51	0.71	0.113
<i>Suncus murinus</i>	1	24.1	0.498	11.56	1.59	9.08	1.11	0.218

Number of individuals = n

Abbreviations:

- CAI capsula interna
- DIE diencephalon
- ETH epithalamus
- HTH hypothalamus
- STH subthalamus
- THA thalamus
- TRO tractus opticus

Table 35. Size indices of diencephalic components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 34

For average size indices in supraspecific taxonomic units see Table 36

For size indices of subthalamic components and geniculate bodies see Table 38

For size indices of medial nuclei of the habenula (HAM) see Table 41

	n	DIE	ETH	THA	STH	HTH	CAI	TRO
slope		.65	.63	.64	.71	.62	.78	.61
y-intercept		.350	-1.276	.010	-.948	.007	-1.083	-1.416
Tenrecidae								
<i>Tenrec ecaudatus</i>	2	85	78	87	91	82	88	123
<i>Setifer setosus</i>	1	119	94	118	102	122	140	95
<i>Hemicentetes semispin.</i>	1	97	119	96	109	96	91	79
<i>Echinops telfairi</i>	1	98	109	99	98	100	81	102
Erinaceidae								
<i>Erinaceus europaeus</i>	2	135	102	156	107	123	120	172
Soricidae								
<i>Sorex araneus</i>	2	146	117	163	189	123	123	92
<i>S. minutus</i>	1	131	111	138	204	112	102	104
<i>Crocidura flavescens</i>	2	125	127	141	123	102	163	72
<i>C. russula</i>	1	129	149	153	137	100	132	68
<i>Suncus murinus</i>	1	109	102	119	116	101	86	66

Number of individuals = n

Abbreviations:

CAI	capsula interna
DIE	diencephalon
ETH	epithalamus
HTH	hypothalamus
STH	subthalamus
THA	thalamus
TRO	tractus opticus

Table 36. Average size indices of diencephalic components related to the average of the four Tenrecidae (AvTn = 100)

For volumes in species see Table 34

For size indices in species see Table 35 39

For av. size indices of subthalamus components and geniculate bodies see Table

For average size indices of medial nuclei of the habenula (HAM) see Table 42

	N	DIE	ETH	THA	STH	HTH	CAI	TRO
slope		.65	.63	.64	.71	.62	.78	.61
y-intercept		.350	-1.276	.010	-.948	.007	-1.083	-1.416
Insectivora (AvIS)	10	117	111	127	127	106	113	97
Tenrecidae								
Tenrecinae*	4	100	100	100	100	100	100	100
Erinaceidae								
Erinaceinae*	1	135	102	156	107	123	120	172
Soricidae	5	128	121	143	154	108	121	80
Soricinae*	2	138	114	151	196	117	112	98
Crocidurinae*	3	121	126	138	125	101	127	69
<i>Crocidura</i>	2	127	138	147	130	101	147	70
AvIF of the 4 subfamilies marked by *		124	111	136	132	110	115	110

Number of species = N

Abbreviations:

- CAI capsula interna
- DIE diencephalon
- ETH epithalamus
- HTH hypothalamus
- STH subthalamus
- THA thalamus
- TRO tractus opticus

Table 37. Volumes of subthalamic components and geniculate bodies

For size indices in species see Table 38

For average size indices in supraspecific taxonomic units see Table 39

For volumes of other diencephalic components see Tables 65 and 68

	n	STH	PALL	LUY	CGM	CGL	GLD	GLV
Tenrecidae								
<i>Tenrec ecaudatus</i>	2	12.29	11.78	0.513	10.406	4.419	3.134	1.285
<i>Setifer setosus</i>	1	5.58	5.28	0.304	5.288	2.916	2.022	0.894
<i>Hemicentetes semisp.</i>	1	3.60	3.39	0.205	2.624	1.194	0.798	0.396
<i>Echinops telfairi</i>	1	2.65	2.41	0.239	2.124	1.126	0.736	0.390
Erinaceidae								
<i>Erinaceus europaeus</i>	2	14.49	13.56	0.927	13.984	10.752	7.956	2.796
Soricidae								
<i>Sorex araneus</i>	2	1.11	1.00	0.107	0.670	0.264	0.150	0.114
<i>S. minutus</i>	1	0.66	0.61	0.053	0.306	0.159	0.110	0.049
<i>Crocidura flavescens</i>	2	1.52	1.40	0.122	1.204	0.670	0.478	0.192
<i>C. russula</i>	1	0.85	0.76	0.087	0.656	0.239	0.150	0.089
<i>Suncus murinus</i>	1	1.59	1.48	0.109	1.458	0.492	0.336	0.156

Number of individuals = n

Abbreviations:

CGL	corpus geniculatum laterale
CGM	corpus geniculatum mediale
GLD	dorsal part of CGL
GLV	ventral part of CGL
LUY	nucleus subthalamicus Luysi
PALL	globus pallidus, pallidum
STH	subthalamus

Table 38. Size indices of subthalamic components and geniculate bodies related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 37

For average size indices in supraspecific taxonomic units see Table 39

For size indices of other diencephalic components see Tables 66 and 69

	n	STH	PALL	LUY	CGM	CGL	GLD	GLV
slope		.71	.71	.71	.72	.60	.60	.60
y-intercept		-.948	-.975	-2.162	-1.052	-1.080	-1.246	-1.579
Tenrecidae								
<i>Tenrec ecaudatus</i>	2	91	92	62	91	93	96	85
<i>Setifer setosus</i>	1	102	103	91	116	132	134	127
<i>Hemicentetes semispin.</i>	1	109	110	102	97	83	81	87
<i>Echinops telfairi</i>	1	98	95	145	96	93	89	101
Erinaceidae								
<i>Erinaceus europaeus</i>	2	107	107	112	123	226	245	185
Soricidae								
<i>Sorex araneus</i>	2	189	181	297	142	79	66	107
<i>S. minutus</i>	1	204	199	268	119	78	80	77
<i>Crocidura flavescens</i>	2	123	120	161	119	106	111	96
<i>C. russula</i>	1	137	131	229	131	68	62	80
<i>Suncus murinus</i>	1	116	115	130	130	72	72	72

Number of individuals = n

Abbreviations:

- CGL corpus geniculatum laterale
- CGM corpus geniculatum mediale
- GLD dorsal part of CGL
- GLV ventral part of CGL
- LUY nucleus subthalamicus Luysi
- PALL globus pallidus, pallidum
- STH subthalamus

Table 39. Average size indices of subthalamic components and geniculate bodies related to the average of the four Tenrecidae (AvTn = 100)

For volumes in species see Table 37

For size indices in species see Table 38

For average size indices of other diencephalic components see Tables 67 and 70

	N	STH	PALL	LUY	CGM	CGL	GLD	GLV
slope		.71	.71	.71	.72	.60	.60	.60
y-intercept		-.948	-.975	-2.162	-1.052	-1.080	-1.246	-1.579
Insectivora (AvIS)	10	127	125	160	116	103	104	102
Tenrecidae								
Tenrecinae*	4	100	100	100	100	100	100	100
Erinaceidae								
Erinaceinae*	1	107	107	112	123	226	245	185
Soricidae	5	154	149	217	128	81	78	86
Soricinae*	2	196	190	283	130	79	73	92
Crocicurinae*	3	125	122	173	127	82	82	83
<i>Crocicidura</i>	2	130	125	195	125	87	87	88
AvIF of the 4 subfamilies marked by *		132	130	167	120	122	125	115

Number of species = N

Abbreviations:

CGL	corpus geniculatum laterale
CGM	corpus geniculatum mediale
GLD	dorsal part of CGL
GLV	ventral part of CGL
LUY	nucleus subthalamicus Luysi
PALL	globus pallidus, pallidum
STH	subthalamus

Table 40. Volumes of some periventricular and septal structures

For size indices in species see Table 41

For average size indices in supraspecific taxonomic units see Table 42

	n	SET	n	SFB	n	HAM	n	EPI	n	SCB
Tenrecidae										
<i>Tenrec ecaudatus</i>	1	0.702	1	0.124	1	0.986	1	0.479	1	0.0112
<i>Setifer setosus</i>	1	0.329	1	0.060	1	0.426	1	0.445	1	0.0031
<i>Hemicentetes semis.</i>	1	0.344	1	0.033	1	0.432	1	0.184	1	0.0039
<i>Echinops telfairi</i>	1	0.338	1	0.029	1	0.365	1	0.243	1	0.0021
<i>Oryzorictes talpoides</i>	1	0.331	1	0.018	1	0.257	1	0.051	1	0.0007
<i>Microgale dobsoni</i>	1	0.278	1	0.008	1	0.241	1	0.035	1	0.0010
<i>M. talazaci</i>	1	0.367	1	0.019	1	0.370	1	0.087	1	0.0015
<i>Limnogale mergulus</i>	1	0.602	1	0.019	1	0.469	1	0.122	1	0.0020
<i>Potamogale velox</i>	1	1.108	1	0.090	1	1.462	1	0.003	1	0.0017
<i>Micropotamogale ruwenz.</i>	1	0.367	1	0.070	1	0.592	1	0.013	-	-.-
Chrysochloridae										
<i>Chrysochloris asiatica</i>	1	0.485	1	0.015	1	0.449	1	0.025	1	0.0081
<i>C. stuhlmanni</i>	6	0.394	6	0.019	6	0.379	6	0.096	6	(0.0)
Solenodontidae										
<i>Solenodon paradoxus</i>	1	0.884	1	0.084	1	2.109	1	1.289	1	0.0442
Erinaceidae										
<i>Echinosorex gymnurus</i>	1	1.664	1	0.110	1	2.400	1	0	-	-.-
<i>Hylomys suillus</i>	1	0.480	1	0.028	1	0.591	1	0.197	-	-.-
<i>Atelerix algirus</i>	1	1.199	1	0.063	1	0.948	1	0.303	1	0.0226
<i>Erinaceus europaeus</i>	2	0.885	2	0.059	2	1.177	2	0.424	2	0.0313
<i>Hemiechinus auritus</i>	1	0.419	1	0.080	1	0.747	1	0.121	-	-.-
Talpidae										
<i>Desmana moschata</i>	1	1.058	1	0.035	1	1.051	1	0.459	1	0.0278
<i>Galemys pyrenaicus</i>	1	0.446	5	0.038	1	0.424	1	0.071	1	0.0055
<i>Talpa europaea</i>	1	0.329	5	0.009	1	0.365	1	0.050	1	0.0102
<i>T. micrura</i>	1	0.386	1	0.018	1	0.470	1	0.058	-	-.-
<i>Parascalops breweri</i>	1	0.530	1	0.018	1	0.458	1	0.068	-	-.-
<i>Scalopus aquaticus</i>	1	0.504	1	0.019	1	0.603	1	0.095	1	0.0260

Table 40 (cont.)

	n	SET	n	SFB	n	HAM	n	EPI	n	SCB
Soricidae										
<i>Sorex araneus</i>	2	0.124	10	0.004	1	0.081	10	0.011	1	0.0003
<i>S. minutus</i>	1	0.074	1	0.003	1	0.062	1	0.004	1	0.0003
<i>Microsorex hoyi</i>	1	0.071	1	0.003	1	0.065	1	0.003	-	-, -
<i>Neomys fodiens</i>	1	0.269	1	0.007	1	0.210	1	0.028	1	0.0004
<i>Anourosorex squamipes</i>	1	0.215	1	0.019	1	0.181	1	0.025	-	-, -
<i>Crocidura flavescens</i>	1	0.261	1	0.018	1	0.191	1	0.037	1	0.0006
<i>C. russula</i>	2	0.126	2	0.010	1	0.141	5	0.029	1	0.0008
<i>Suncus etruscus</i>	1	0.052	1	0.004	1	0.070	1	0.007	-	-, -
<i>S. murinus</i>	1	0.203	1	0.015	1	0.170	1	0.064	1	0.0022
<i>Scutisorex somereni</i>	1	0.370	1	0.026	1	0.283	1	0.035	-	-, -
<i>Sylvisorex granti</i>	1	0.081	1	0.006	1	0.088	1	0.017	-	-, -
<i>S. megalura</i>	1	0.084	1	0.007	1	0.117	1	0.016	-	-, -
<i>Ruwenzorisorex sunc.</i>	1	0.204	1	0.011	1	0.266	1	0.020	-	-, -
<i>Myosorex babaulti</i>	1	0.148	1	0.009	1	0.157	1	0.003	-	-, -

Number of individuals = n

Abbreviations:

- EPI epiphysis cerebri, corpus pineale
- HAM nucleus medialis habenulae
- SCB corpus subcommissuralis
- SET nucleus triangularis septi
- SFB corpus subformicale

Table 41. Size indices of some periventricular and septal structures related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 40

For average size indices in supraspecific taxonomic units see Table 42

	n SET		n SFB		n HAM		n EPI		n SCB	
slope		.51		.44		.58		.58		.55
y-intercept		-1.570		-2.304		-1.637		-1.838		-3.652
Tenrecidae										
<i>Tenrec ecaudatus</i>	1	84	1	128	1	85	1	66	1	123
<i>Setifer setosus</i>	1	75	1	109	1	77	1	129	1	69
<i>Hemicentetes semispin.</i>	1	113	1	81	1	119	1	80	1	127
<i>Echinops telfairi</i>	1	128	1	82	1	118	1	125	1	81
<i>Oryzorictes talpoides</i>	1	178	1	68	1	124	1	39	1	39
<i>Microgale dobsoni</i>	1	177	1	33	1	140	1	33	1	66
<i>M. talazaci</i>	1	189	1	68	1	169	1	63	1	78
<i>Limnogale mergulus</i>	1	223	1	53	1	148	1	61	1	75
<i>Potamogale velox</i>	1	155	1	107	1	153	1	0.5	1	23
<i>Micropotamogale ruwenz.</i>	1	132	1	189	1	181	1	6	-	--
Chrysochloridae										
<i>Chrysochloris asiatica</i>	1	248	1	53	1	204	1	18	1	427
<i>C. stuhlmanni</i>	6	223	6	75	6	193	6	77	6	(0)
Solenodontidae										
<i>Solenodon paradoxus</i>	1	119	1	97	1	209	1	203	1	553
Erinaceidae										
<i>Echinosorex gymnurus</i>	1	202	1	115	1	212	1	0	-	--
<i>Hylomys suillus</i>	1	227	1	95	1	245	1	130	-	--
<i>Atelerix algirus</i>	1	154	1	70	1	89	1	45	1	269
<i>Erinaceus europaeus</i>	2	105	2	61	2	102	2	58	2	344
<i>Hemiechinus auritus</i>	1	96	1	146	1	136	1	35	-	--
Talpidae										
<i>Desmana moschata</i>	1	176	1	49	1	133	1	92	1	437
<i>Galemys pyrenaicus</i>	1	206	5	126	1	171	1	46	1	261
<i>Talpa europaea</i>	1	129	5	25	1	123	1	26	1	404
<i>T. micrura</i>	1	215	1	69	1	235	1	46	-	--
<i>Parascalops breweri</i>	1	258	1	64	1	197	1	46	-	--
<i>Scalopus aquaticus</i>	1	166	1	48	1	167	1	42	1	857

Table 41 (cont.)

	n	SET	n	SFB	n	HAM	n	EPI	n	SCB
Soricidae										
<i>Sorex araneus</i>	2	141	10	26	1	91	10	20	1	40
<i>S. minutus</i>	1	129	1	31	1	114	1	11	1	63
<i>Microsorex hoyi</i>	1	161	1	45	1	160	1	13	-	--
<i>Neomys fodiens</i>	1	249	1	43	1	187	1	39	1	41
<i>Anourosorex squamipes</i>	1	173	1	104	1	137	1	30	-	--
<i>Crocidura flavescens</i>	1	173	1	81	1	117	1	36	1	40
<i>C. russula</i>	2	137	2	72	1	151	5	50	1	100
<i>Suncus etruscus</i>	1	141	1	60	1	212	1	35	-	--
<i>S. murinus</i>	1	125	1	64	1	96	1	57	1	144
<i>Scutisorex somereni</i>	1	166	1	85	1	111	1	21	-	--
<i>Sylvisorex granti</i>	1	150	1	65	1	172	1	54	-	--
<i>S. megalura</i>	1	130	1	70	1	190	1	41	-	--
<i>Ruwenzorisorex sunc.</i>	1	172	1	63	1	214	1	26	-	--
<i>Myosorex babaulti</i>	1	129	1	51	1	131	1	4	-	--

Number of individuals = n

Abbreviations:

- EPI epiphysis cerebri, corpus pineale
HAM nucleus medialis habenulae
SCB corpus subcommissuralis
SET nucleus triangularis septi
SFB corpus subfornicale

Table 42. Average size indices of some periventricular and septal structures related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 40

For size indices in species see Table 41

	N	SET	SFB	HAM	EPI	N	SCB
slope		.51	.44	.58	.58		.55
y-intercept		-1.570	-2.304	-1.637	-1.838		-3.652
Insectivora (AvIS)	38	162	75	153	50	23	203
Tenrecidae	10	145	92	131	60	9	75
Tenrecinae*	4	100	100	100	100	4	100
Oryzorictinae*	4	192	56	145	49	4	64
<i>Microgale</i>	2	183	50	155	48	2	72
terr. Oryzorictinae	3	181	56	144	45	3	61
<i>Limnogale</i>	1	223	53	148	61	1	75
Potamogalinae*	2	144	148	167	3	1	23
Chrysochloridae*	2	235	64	198	48	1	427
Solenodontidae*	1	119	97	209	203	1	553
Erinaceidae	5	157	97	157	54	-	--
Echinosoricinae*	2	214	105	229	65	-	--
Erinaceinae*	3	118	92	109	46	2	306
Talpidae	6	192	63	171	50	4	490
Desmaninae*	2	191	87	152	69	2	349
Talpinae*	4	192	51	180	40	2	630
<i>Talpa</i>	2	172	47	179	36	1	404
Soricidae	14	155	61	149	31	6	72
Soricinae*	5	170	50	138	23	3	48
<i>Sorex</i>	2	135	28	103	15	2	52
Crocidurinae*	9	147	68	155	36	3	95
<i>Crocidura</i>	2	155	76	134	43	2	70
<i>Suncus</i>	2	133	62	154	46	1	144
<i>Sylvisorex</i>	2	140	67	181	48	-	--
AvIF of the families and subfamilies marked by *	(11)	166	83	162	62	(10)	260

Number of species = N

Abbreviations:

- EPI epiphysis cerebri, corpus pineale
HAM nucleus medialis habenulae
SCB corpus subcommissuralis
SET nucleus triangularis septi
SFB corpus subfornicale

Table 43. Volumes of telencephalic components

For size indices in species see Table 44

For average size indices in supraspecific taxonomic units see Table 45

	BOL	PAL	STR	SEP	AMY	HIP	SCH	NEO
Tenrecidae								
<i>Tenrec ecaudatus</i>	316.7	378.3	98.81	37.15	88.80	147.1	77.07	272.8
<i>Setifer setosus</i>	213.3	262.9	57.65	24.76	67.62	96.6	47.98	136.6
<i>Hemicentetes semisp.</i>	93.2	114.6	27.78	13.33	20.56	63.1	29.14	76.9
<i>Echinops telfairi</i>	71.8	94.5	24.20	11.54	22.51	46.7	13.18	51.7
<i>Geogale aurita</i>	11.7	13.3	3.81	2.04	4.12	10.9	2.02	7.2
<i>Oryzorictes talpoides</i>	47.0	67.2	32.53	10.05	20.75	64.3	23.47	66.5
<i>Microgale dobsoni</i>	53.5	76.2	22.95	9.52	16.89	66.7	16.26	62.6
<i>M. talazaci</i>	68.4	107.1	32.30	11.37	24.44	93.5	22.61	88.9
<i>Limnogale mergulus</i>	43.2	60.6	57.90	19.54	26.82	78.2	36.41	195.6
<i>Potamogale velox</i>	83.9	128.4	148.49	49.30	62.16	322.4	123.63	1050.0
<i>Micropotamogale lam.</i>	34.1	48.2	32.64	15.06	19.05	77.7	32.92	144.4
<i>M. ruwenzorii</i>	33.7	59.5	59.47	17.25	22.10	97.8	37.22	227.8
Chrysochloridae								
<i>Chrysochloris asiatica</i>	58.6	94.9	44.32	12.12	20.05	66.3	20.93	112.6
<i>C. stuhlmanni</i>	60.5	107.3	43.24	11.29	23.17	58.8	21.32	128.6
Solenodontidae								
<i>Solenodon paradox.</i>	469.8	561.6	257.51	66.17	135.36	299.7	158.14	668.6
Erinaceidae								
<i>Echinorex gymnur.</i>	427.4	682.8	209.44	76.75	170.14	609.1	198.79	1085.0
<i>Hylomys suillus</i>	85.8	128.0	38.38	17.50	37.43	127.0	51.47	177.8
<i>Atelerix algirus</i>	292.4	365.0	143.04	43.73	103.30	231.8	111.50	510.3
<i>Erinaceus europaeus</i>	335.6	398.6	147.95	51.97	108.67	237.9	134.80	525.1
<i>Hemiechinus auritus</i>	174.6	205.3	80.25	28.07	71.06	128.3	67.94	306.2
Talpidae								
<i>Desmana moschata</i>	141.6	229.5	221.87	42.62	90.80	267.3	163.27	983.4
<i>Galemys pyrenaicus</i>	39.2	70.0	74.38	20.11	36.43	108.6	44.82	338.7
<i>Talpa europaea</i>	60.2	85.9	60.44	15.07	26.41	87.1	39.76	181.2
<i>T. micrura</i>	53.2	78.6	48.08	13.30	22.69	67.4	28.93	157.2
<i>Parascalops breweri</i>	58.9	89.5	60.43	13.42	22.58	83.0	33.83	175.6
<i>Scalopus aquaticus</i>	86.6	139.0	76.18	17.34	30.87	109.9	67.52	252.9

Table 43 (cont.)

	BOL	PAL	STR	SEP	AMY	HIP	SCH	NEO
Soricidae								
<i>Sorex alpinus</i>	16.6	27.9	16.63	4.02	8.05	28.9	9.73	45.5
<i>S. araneus</i>	17.2	27.9	11.50	3.39	7.70	22.8	7.26	28.2
<i>S. cinereus</i>	11.7	22.2	11.77	2.63	4.89	19.5	5.45	28.8
<i>S. fumeus</i>	15.1	29.3	14.42	3.74	7.43	27.8	10.01	48.0
<i>S. minutus</i>	8.9	14.0	7.05	1.82	3.46	12.4	4.24	15.9
<i>Microsorex hoyi</i>	6.4	11.5	6.28	1.69	3.28	11.7	3.76	15.2
<i>Neomys anomalus</i>	13.1	23.5	18.73	4.29	8.18	25.2	12.00	59.0
<i>N. fodiens</i>	16.4	26.0	20.00	5.71	8.59	34.4	10.93	56.6
<i>Blarina brevicauda</i>	28.9	44.4	22.28	6.79	10.93	44.7	14.89	68.7
<i>Cryptotis parva</i>	20.5	30.0	12.88	4.11	9.10	26.2	9.64	36.2
<i>Anourosorex squamip.</i>	32.3	50.4	20.13	6.11	15.35	34.0	14.32	67.6
<i>Crociodura attenuata</i>	19.7	25.5	7.58	3.06	6.66	19.3	6.59	25.3
<i>C. flavescens</i>	35.4	48.2	16.93	6.01	12.63	38.3	13.25	53.1
<i>C. hildegardeae</i>	19.5	30.4	8.87	3.31	6.89	17.8	5.65	26.0
<i>C. jacksoni</i>	19.4	30.9	9.74	3.42	7.62	22.4	7.11	28.6
<i>C. russula</i>	17.1	24.0	8.64	3.22	6.65	17.9	5.85	25.3
<i>C. suaveolens</i>	16.3	23.3	8.06	2.89	5.94	17.2	6.24	24.6
<i>Suncus etruscus</i>	4.7	7.1	2.71	1.04	1.73	6.2	1.89	7.1
<i>S. murinus</i>	36.9	47.1	15.16	5.98	11.09	33.4	12.77	50.5
<i>Scutisorex somereni</i>	58.8	82.8	26.08	10.86	17.85	57.4	19.19	87.2
<i>Sylvisorex granti</i>	15.4	23.4	7.62	2.68	5.06	15.8	4.33	20.6
<i>S. megalura</i>	17.0	25.3	9.55	2.93	5.58	12.0	3.88	23.9
<i>Ruwenzorisorex sunc.</i>	17.6	29.0	17.43	6.02	10.25	39.5	13.02	59.5
<i>Myosorex babaulti</i>	34.6	52.3	18.62	5.58	12.08	33.8	8.27	45.3

Number of individuals as in Table 10

Abbreviations and notes:

- AMY amygdala; for volumes of components see Table 55
- BOL bulbi olfactorii (main + accessory); for volumes of components see Tables 46 and 49
- HIP hippocampus; for volumes of components see Table 58
- NEO neocortex; for volumes of components see Table 61
- PAL paleocortex; for volumes of components see Table 52
- SCH schizocortex
- SEP septum
- STR striatum

Table 44. Size indices of telencephalic components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 43

For average size indices in supraspecific taxonomic units see Table 45

	BOL	PAL	STR	SEP	AMY	HIP	SCH	NEO
slope	.64	.63	.67	.61	.60	.57	.64	.73
y-intercept	.682	.799	.089	-.129	.232	.581	.062	.346
Tenrecidae								
<i>Tenrec ecaudatus</i>	88	86	88	81	91	82	89	89
<i>Setifer setosus</i>	134	133	120	118	149	112	126	114
<i>Hemicentetes semispin.</i>	93	91	94	99	70	110	121	108
<i>Echinops telfairi</i>	85	90	99	101	90	96	65	89
<i>Geogale aurita</i>	56	50	66	67	61	77	40	61
<i>Oryzorictes talpoides</i>	86	98	209	134	125	195	180	189
<i>Microgale dobsoni</i>	121	137	184	155	124	243	154	225
<i>M. talazaci</i>	119	148	196	144	140	270	164	237
<i>Limnogale mergulus</i>	50	56	228	167	104	156	175	325
<i>Potamogale velox</i>	29	36	163	131	77	217	175	434
<i>Micropotamogale lam.</i>	49	56	164	160	92	190	199	312
<i>M. ruwenzorii</i>	38	53	226	143	83	189	173	365
Chrysochloridae								
<i>Chrysochloris asiatica</i>	101	130	266	152	114	189	150	296
<i>C. stuhlmanni</i>	118	166	297	159	148	188	174	391
Solenodontidae								
<i>Solenodon paradoxus</i>	151	148	268	168	160	192	213	260
Erinaceidae								
<i>Echinosorex gymnurus</i>	121	158	190	172	178	348	235	364
<i>Hylomys suillus</i>	134	159	208	200	194	333	336	419
<i>Atelerix algirus</i>	89	91	140	105	115	141	141	186
<i>Erinaceus europaeus</i>	93	90	131	114	111	134	156	172
<i>Hemiechinus auritus</i>	110	105	169	135	157	150	179	257
Talpidae								
<i>Desmana moschata</i>	60	78	305	139	137	218	287	519
<i>Galemys pyrenaicus</i>	60	85	391	223	184	277	284	772
<i>Talpa europaea</i>	75	85	257	138	110	185	205	327
<i>T. micrura</i>	102	120	323	185	142	212	231	468
<i>Parascalops breweri</i>	96	115	341	159	121	225	229	432
<i>Scalopus aquaticus</i>	86	111	258	129	105	193	281	357

Table 44 (cont.)

	BOL	PAL	STR	SEP	AMY	HIP	SCH	NEO
Soricidae								
<i>Sorex alpinus</i>	74	98	272	125	112	193	182	356
<i>S. araneus</i>	81	102	198	111	112	159	142	233
<i>S. cinereus</i>	85	125	319	130	107	201	165	391
<i>S. fumeus</i>	80	122	282	137	121	217	222	458
<i>S. minutus</i>	72	87	213	99	83	140	142	243
<i>Microsorex hoyi</i>	72	99	268	126	108	176	175	339
<i>Neomys anomalus</i>	57	80	295	129	110	164	217	445
<i>N. fodiens</i>	60	74	262	145	98	191	165	348
<i>Blarina brevicauda</i>	89	108	246	148	107	214	192	351
<i>Cryptotis parva</i>	98	112	226	137	135	186	193	306
<i>Anourosorex squam.</i>	98	121	220	132	149	161	182	341
<i>Crocidura attenuata</i>	98	99	139	106	102	142	137	224
<i>C. flavescens</i>	85	91	144	103	98	147	132	203
<i>C. hildegardae</i>	92	112	153	108	100	124	111	215
<i>C. jacksoni</i>	79	99	145	97	97	138	121	202
<i>C. russula</i>	76	84	140	100	92	119	109	197
<i>C. suaveolens</i>	76	85	138	94	86	120	122	202
<i>Suncus etruscus</i>	65	76	146	96	70	113	110	203
<i>S. murinus</i>	81	81	117	94	79	118	116	174
<i>Scutisorex somereni</i>	86	96	132	116	87	142	117	190
<i>Sylvisorex granti</i>	134	158	249	157	131	191	157	345
<i>S. megalura</i>	119	138	249	139	118	119	113	311
<i>Ruwenzorisorex sunc.</i>	57	74	203	138	105	198	176	323
<i>Myosorex babaulti</i>	117	139	227	133	129	176	117	258

Number of individuals as in Table 10

Abbreviations and notes:

- AMY amygdala; for size indices of components see Table 56
BOL bulbi olfactorii (main + accessory); for size indices of components see Tables
HIP hippocampus; for size indices of components see Table 59 47 and 50
NEO neocortex; for size indices of components see Table 62
PAL paleocortex; for size indices of components see Table 53
SCH schizocortex
SEP septum
STR striatum

Table 45. Average size indices of telencephalic components related to the average of the four Tenrecinae (AvTn = 100)

	BOL	PAL	STR	SEP	AMY	HIP	SCH	NEO
slope	.64	.63	.67	.61	.60	.57	.64	.73
y-intercept	.682	.799	.089	-.129	.232	.581	.062	.346
Insectivora (AvIS)	88	103	207	132	114	175	168	292
Tenrecidae	79	86	153	125	101	161	138	212
Tenrecinae*	100	100	100	100	100	100	100	100
Geogalinae*	56	50	66	67	61	77	40	61
Oryzoricinae*	94	110	204	150	123	216	168	244
<i>Microgale</i>	120	142	190	149	132	257	159	231
terr. Oryzoricinae	109	128	196	144	130	236	166	217
<i>Limnogale</i>	50	56	228	167	104	156	175	325
Potamogalinae*	39	48	184	145	84	199	182	370
<i>Micropotamogale</i>	44	54	195	151	88	190	186	338
Chrysochloridae*	110	148	281	156	131	189	162	344
Solenodontidae*	151	148	268	168	160	192	213	260
Erinaceidae	109	121	168	145	151	221	209	280
Echinosoricinae*	128	159	199	186	186	340	285	391
Erinaceinae*	97	95	147	118	128	142	159	205
Talpidae	80	99	313	162	133	218	253	479
Desmaninae*	60	82	348	181	161	247	285	645
Talpinae*	90	108	295	152	120	204	237	396
<i>Talpa</i>	88	102	290	161	126	199	218	397
Soricidae	85	103	208	121	106	160	151	286
Soricinae*	79	103	255	129	113	182	180	347
<i>Sorex</i>	78	107	257	120	107	182	171	336
terr. Soricinae	83	108	249	127	115	183	177	335
<i>Neomys</i>	58	77	279	137	104	177	191	397
Crocidurinae*	90	103	168	114	100	142	126	234
<i>Crocidura</i>	84	95	143	101	96	132	122	207
<i>Suncus</i>	73	79	131	95	74	116	113	188
<i>Sylvisorex</i>	126	148	249	148	124	155	135	328
AvIF of the 12 families and sub-families marked by *	91	104	210	139	122	186	178	300

Number of species as in Table 12

Abbreviations and notes:

- AMY amygdala; for av. size indices of components see Table 57
 BOL bulbi olfactorii (main + accessory); for av. size indices of components see Tables
 HIP hippocampus; for av. size indices of components see Table 60 48 and 51
 NEO neocortex; for av. size indices of components see Table 63
 PAL paleocortex; for av. size indices of components see Table 54
 SEP septum SCH schizocortex STR striatum

Table 46. Volumes of main olfactory bulbs and some of its layers

For size indices in species see Table 47

For average size indices in supraspecific taxonomic units see Table 48

	n	MOB	n	L1+2	L3	L4-6	PvZ
Tenrecidae							
<i>Tenrec ecaudatus</i>	2	315.9	1	130.15	57.36	102.49	25.93
<i>Setifer setosus</i>	2	212.8	1	73.13	51.41	81.99	6.23
<i>Hemicentetes semis.</i>	1	93.0	1	33.39	22.24	34.13	3.24
<i>Echinops telfairi</i>	2	71.6	1	27.06	17.14	26.12	1.26
<i>Geogale aurita</i>	1	11.6	1	3.90	2.93	4.62	0.13
<i>Oryzorictes talpoides</i>	2	46.9	1	11.58	13.83	20.65	0.80
<i>Microgale dobsoni</i>	1	53.4	1	17.88	14.69	20.20	0.59
<i>M. talazaci</i>	1	68.0	1	19.20	21.82	26.16	0.84
<i>Limnogale mergulus</i>	1	42.9	1	16.20	11.40	14.70	0.57
<i>Potamogale velox</i>	2	81.7	1	27.30	20.31	31.34	2.76
<i>Micropotamogale lam.</i>	1	34.1	1	11.45	9.53	12.79	0.29
<i>M. ruwenzorii</i>	4	33.7	1	10.22	8.87	14.44	0.21
Chrysochloridae							
<i>Chrysochloris asiatica</i>	1	58.5	1	19.96	13.63	23.26	1.62
<i>C. stuhlmanni</i>	2	60.3	1	19.42	14.70	24.36	1.80
Solenodontidae							
<i>Solenodon paradoxus</i>	2	469.6	1	224.95	84.14	133.09	27.37
Erinaceidae							
<i>Echinorex gymnurus</i>	1	424.2	1	148.10	83.88	160.71	31.53
<i>Hylomys suillus</i>	1	84.4	1	32.82	19.15	29.59	2.85
<i>Atelerix algirus</i>	2	290.2	1	121.53	49.28	97.03	22.39
<i>Erinaceus europaeus</i>	2	333.7	1	138.08	60.47	109.72	25.38
<i>Hemiechinus auritus</i>	3	173.3	1	62.93	37.34	60.60	12.42
Talpidae							
<i>Desmana moschata</i>	1	141.4	1	47.52	27.42	60.25	6.24
<i>Galemys pyrenaicus</i>	2	39.0	1	12.57	10.52	15.07	0.86
<i>Talpa europaea</i>	2	59.9	1	21.02	14.33	22.75	1.77
<i>T. micrura</i>	1	53.0	1	19.46	12.33	20.54	0.70
<i>Parascalops breweri</i>	1	58.8	1	20.32	14.03	23.24	1.25
<i>Scalopus aquaticus</i>	1	86.5	1	34.05	20.63	30.63	1.14

Table 46 (cont.)

	n	MOB	n	L1+2	L3	L4-6	PvZ
Soricidae							
<i>Sorex alpinus</i>	1	16.5	--	--	--	--	--
<i>S. araneus</i>	2	17.1	1	4.85	4.77	7.33	0.15
<i>S. cinereus</i>	1	11.6	--	--	--	--	--
<i>S. fumeus</i>	1	15.0	--	--	--	--	--
<i>S. minutus</i>	1	8.9	1	2.72	2.72	3.41	0.07
<i>Microsorex hoyi</i>	1	6.4	1	2.01	1.85	2.53	0.02
<i>Neomys anomalus</i>	1	13.0	1	3.92	3.40	5.40	0.25
<i>N. fodiens</i>	2	16.3	1	5.51	5.04	5.60	0.15
<i>Blarina brevicauda</i>	1	28.9	1	9.77	7.93	10.86	0.32
<i>Cryptotis parva</i>	1	20.5	1	5.90	5.90	8.56	0.14
<i>Anourosorex squamipes</i>	1	32.2	1	9.10	8.98	13.47	0.68
<i>Crociodura attenuata</i>	2	19.6	--	--	--	--	--
<i>C. flavescens</i>	1	35.2	1	11.72	9.85	13.01	0.60
<i>C. hildegardeae</i>	1	19.4	--	--	--	--	--
<i>C. jacksoni</i>	2	19.3	--	--	--	--	--
<i>C. russula</i>	1	17.0	1	5.89	4.84	6.16	0.14
<i>C. suaveolens</i>	1	16.2	--	--	--	--	--
<i>Suncus etruscus</i>	1	4.7	1	1.58	1.36	1.69	0.02
<i>S. murinus</i>	1	36.7	1	11.88	10.16	13.20	1.44
<i>Scutisorex somereni</i>	1	58.6	1	20.00	14.50	22.90	1.21
<i>Sylvisorex granti</i>	1	15.4	1	4.80	4.43	6.10	0.08
<i>S. megalura</i>	1	16.9	1	5.44	4.62	6.71	0.14
<i>Ruwenzorisorex sunc.</i>	1	17.5	1	5.77	4.45	7.06	0.26
<i>Myosorex babaulti</i>	1	34.5	1	10.63	9.26	14.45	0.16

Number of individuals = n

Abbreviations:

- L1+2 external fiber and glomerular layers
- L3 external plexiform layer
- L4-6 mitral, internal plexiform, and granular layers
- MOB main olfactory bulb
- PvZ periventricular zone

Table 47. Size indices of the main olfactory bulbs and of some of the MOB layers related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 46

For average size indices in supraspecific taxonomic units see Table 48

	n	MOB	n	L1+2	L3	L4-6	PvZ
slope		.64		.64	.64	.64	.64
y-intercept		.681		.248	.037	.241	-.721
Tenrecidae							
<i>Tenrec ecaudatus</i>	2	88	1	98	70	78	182
<i>Setifer setosus</i>	2	134	1	125	143	142	99
<i>Hemicentetes semispin.</i>	1	93	1	90	97	94	81
<i>Echinops telfairi</i>	2	85	1	87	90	86	38
<i>Geogale aurita</i>	1	55	1	50	62	61	16
<i>Oryzorictes talpoides</i>	2	86	1	58	112	105	37
<i>Microgale dobsoni</i>	1	121	1	110	147	126	34
<i>M. talazaci</i>	1	119	1	91	168	126	37
<i>Limnogale mergulus</i>	1	49	1	51	58	47	17
<i>Potamogale velox</i>	2	28	1	25	31	29	24
<i>Micropotamogale lam.</i>	1	49	1	45	61	51	10
<i>M. ruwenzorii</i>	4	38	1	31	44	44	6
Chrysochloridae							
<i>Chrysochloris asiatica</i>	1	101	1	93	104	111	70
<i>C. stuhlmanni</i>	2	118	1	103	127	132	89
Solenodontidae							
<i>Solenodon paradoxus</i>	2	152	1	197	120	118	223
Erinaceidae							
<i>Echinorex gymnurus</i>	1	120	1	114	105	126	226
<i>Hylomys suillus</i>	1	132	1	139	132	128	113
<i>Atelerix algirus</i>	2	88	1	100	66	81	172
<i>Erinaceus europaeus</i>	2	93	1	104	74	84	178
<i>Hemiechinus auritus</i>	3	110	1	108	104	106	198
Talpidae							
<i>Desmana moschata</i>	1	60	1	54	51	70	66
<i>Galemys pyrenaicus</i>	2	59	1	52	71	63	33
<i>Talpa europaea</i>	2	74	1	71	78	78	55
<i>T. micrura</i>	1	102	1	101	104	109	34
<i>Parascalops breweri</i>	1	96	1	90	101	104	51
<i>Scalopus aquaticus</i>	1	86	1	92	91	84	29

Table 47 (cont.)

	n	MOB	n	L1+2	L3	L4-6	PvZ
Soricidae							
<i>Sorex alpinus</i>	1	74	--	--	--	--	--
<i>S. araneus</i>	2	81	1	62	99	95	18
<i>S. cinereus</i>	1	85	--	--	--	--	--
<i>S. fumeus</i>	1	80	--	--	--	--	--
<i>S. minutus</i>	1	72	1	60	97	76	13
<i>Microsorex hoyi</i>	1	72	1	61	91	78	6
<i>Neomys anomalus</i>	1	56	1	46	65	65	28
<i>N. fodiens</i>	2	59	1	54	81	56	14
<i>Blarina brevicauda</i>	1	89	1	82	108	93	25
<i>Cryptotis parva</i>	1	99	1	77	125	113	17
<i>Anourosorex squamipes</i>	1	98	1	75	121	113	52
<i>Crociodura attenuata</i>	2	98	--	--	--	--	--
<i>C. flavescens</i>	1	84	1	76	104	86	36
<i>C. hildegardeae</i>	1	91	--	--	--	--	--
<i>C. jacksoni</i>	2	79	--	--	--	--	--
<i>C. russula</i>	1	76	1	71	95	76	16
<i>C. suaveolens</i>	1	76	--	--	--	--	--
<i>Suncus etruscus</i>	1	65	1	60	84	65	8
<i>S. murinus</i>	1	80	1	70	98	80	79
<i>Scutisorex somereni</i>	1	86	1	79	94	92	45
<i>Sylvisorex granti</i>	1	134	1	113	170	147	18
<i>S. megalura</i>	1	118	1	103	143	129	25
<i>Ruwenzorisorex sunc.</i>	1	57	1	51	64	63	21
<i>Myosorex babaulti</i>	1	117	1	98	139	135	14

Number of individuals = n

Abbreviations:

- L1+2 external fiber and glomerular layers
 L3 external plexiform layer
 L4-6 mitral, internal plexiform, and granular layers
 MOB main olfactory bulb
 PvZ periventricular zone

Table 48. Average size indices of the main olfactory bulbs and of some of the MOB layers related to the average of the four Tenrecinae (AvTn = 100)

	N	MOB	N	L1+2	L3	L4-6	PvZ
slope		.64		.64	.64	.64	.64
y-intercept		.681		.248	.037	.241	-.721
Insectivora (AvIS)	50	87	43	82	97	92	59
Tenrecidae	12	79	12	72	90	82	48
Tenrecinae*	4	100	4	100	100	100	100
Geogalinae*	1	55	1	50	62	61	16
Oryzorictinae*	4	94	4	77	121	101	31
<i>Microgale</i>	2	120	2	100	157	126	35
terr. Oryzorictinae	3	109	3	86	142	119	36
<i>Limnogale</i>	1	49	1	51	58	47	17
Potamogalinae*	3	38	3	34	45	42	13
<i>Micropotamogale</i>	2	44	2	38	52	48	8
Chrysochloridae*	2	110	2	98	115	121	80
Solenodontidae*	1	152	1	197	120	118	223
Erinaceidae	5	109	5	113	96	105	178
Echinosoricinae*	2	126	2	127	119	127	169
Erinaceinae*	3	97	3	104	81	90	183
Talpidae	6	80	6	77	83	85	45
Desmaninae*	2	60	2	53	61	67	50
Talpinae*	4	90	4	89	94	94	42
<i>Talpa</i>	2	88	2	86	91	93	45
Soricidae	24	85	17	73	105	92	26
Soricinae*	11	79	8	65	98	86	22
<i>Sorex</i>	5	78	2	61	98	85	16
terr. Soricinae	9	83	6	69	107	95	22
<i>Neomys</i>	2	58	2	50	73	60	21
Crocidurinae*	13	89	9	80	110	97	29
<i>Crocidura</i>	6	84	2	74	100	81	26
<i>Suncus</i>	2	73	2	65	91	72	44
<i>Sylvisorex</i>	2	126	2	108	156	138	22
AvIF of the 12 families and subfamilies marked by *		91		90	94	92	80

Number of species = N

Abbreviations:

- L1+2 external fiber and glomerular layers
- L3 external plexiform layer
- L4-6 mitral, internal plexiform, and granular layers
- MOB main olfactory bulbs
- PvZ periventricular zone

Table 49. Volumes of accessory olfactory bulbs and some of its layers

For size indices in species see Table 50

For average size indices in supraspecific taxonomic units see Table 51

	n	AOB	n	L1+2	L3-5	L6
Tenrecidae						
<i>Tenrec ecaudatus</i>	2	0.730	1	0.393	0.218	0.119
<i>Setifer setosus</i>	2	0.544	1	0.276	0.184	0.084
<i>Hemicentetes semispin.</i>	1	0.194	1	0.086	0.097	0.011
<i>Echinops telfairi</i>	2	0.255	1	0.114	0.107	0.034
<i>Geogale aurita</i>	1	0.086	1	0.023	0.045	0.018
<i>Oryzorictes talpoides</i>	2	0.120	1	0.037	0.076	0.007
<i>Microgale dobsoni</i>	1	0.177	1	0.059	0.099	0.019
<i>M. talazaci</i>	1	0.338	1	0.093	0.168	0.077
<i>Limnogale mergulus</i>	1	0.323	1	0.080	0.165	0.078
<i>Potamogale velox</i>	2	2.202	1	1.194	0.702	0.306
<i>Micropotamogale lam.</i>	1	0.067	1	0.026	0.035	0.006
<i>M. ruwenzorii</i>	4	0.015	--	--	--	--
Chrysochloridae						
<i>Chrysochloris asiatica</i>	1	0.132	1	0.037	0.078	0.017
<i>C. stuhlmanni</i>	2	0.202	1	0.066	0.108	0.028
Solenodontidae						
<i>Solenodon paradoxus</i>	2	0.222	1	0.052	0.149	0.021
Erinaceidae						
<i>Echinosorex gymnurus</i>	1	3.208	1	1.413	0.952	0.843
<i>Hylomys suillus</i>	1	1.347	1	0.511	0.490	0.346
<i>Atelerix algirus</i>	2	2.163	1	0.756	0.839	0.568
<i>Erinaceus europaeus</i>	2	1.922	1	0.627	0.922	0.373
<i>Hemiechinus auritus</i>	3	1.337	1	0.426	0.614	0.297
Talpidae						
<i>Desmana moschata</i>	1	0.218	1	0.096	0.092	0.030
<i>Galemys pyrenaicus</i>	2	0.206	1	0.050	0.128	0.028
<i>Talpa europaea</i>	2	0.300	1	0.090	0.141	0.069
<i>T. micrura</i>	1	0.119	1	0.045	0.046	0.028
<i>Parascalops breweri</i>	1	0.075	1	0.028	0.033	0.014
<i>Scalopus aquaticus</i>	1	0.108	1	0.041	0.050	0.017

Table 49 (cont.)

	n	AOB	n	L1+2	L3-5	L6
Soricidae						
<i>Sorex alpinus</i>	1	0.050	--	-.-	-.-	-.-
<i>S. araneus</i>	2	0.087	1	0.025	0.040	0.022
<i>S. cinereus</i>	1	0.022	--	-.-	-.-	-.-
<i>S. fumeus</i>	1	0.046	1	0.016	0.025	0.005
<i>S. minutus</i>	1	0.021	1	0.006	0.011	0.004
<i>Microsorex hoyi</i>	1	0.021	--	-.-	-.-	-.-
<i>Neomys anomalus</i>	1	0.079	1	0.031	0.038	0.010
<i>N. fodiens</i>	2	0.103	1	0.035	0.050	0.018
<i>Blarina brevicauda</i>	1	0.060	--	-.-	-.-	-.-
<i>Cryptotis parva</i>	1	0.019	1	0.006	0.009	0.004
<i>Anourosorex squamipes</i>	1	0.031	--	-.-	-.-	-.-
<i>Crociodura attenuata</i>	2	0.103	1	0.038	0.051	0.014
<i>C. flavescens</i>	1	0.208	1	0.080	0.089	0.039
<i>C. hildegardeae</i>	1	0.085	--	-.-	-.-	-.-
<i>C. jacksoni</i>	2	0.082	--	-.-	-.-	-.-
<i>C. russula</i>	1	0.045	1	0.013	0.025	0.007
<i>C. suaveolens</i>	1	0.078	--	-.-	-.-	-.-
<i>Suncus etruscus</i>	1	0.024	1	0.008	0.013	0.003
<i>S. murinus</i>	1	0.195	1	0.082	0.077	0.036
<i>Scutisorex somereni</i>	1	0.219	1	0.071	0.109	0.039
<i>Sylvisorex granti</i>	1	0.032	1	0.009	0.020	0.003
<i>S. megalura</i>	1	0.050	1	0.017	0.029	0.004
<i>Ruwenzorisorex sunc.</i>	1	0.079	1	0.030	0.044	0.005
<i>Myosorex babaulti</i>	1	0.088	1	0.031	0.047	0.010

Number of individuals = n

Abbreviations:

- AOB accessory olfactory bulb
L1+2 vomeronasal fiber and glomerular layers
L3-5 external plexiform, mitral, and internal plexiform layers
L6 internal granular layer

Table 50. Size indices of the accessory olfactory bulbs, and some of the AOB layers, related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 49

For average size indices in supraspecific taxonomic units see Table 51

	n	AOB	n	L1+2	L3-5	L6
slope		.64		.64	.64	.64
y-intercept		-1.903		-2.219	-2.318	-2.785
Tenrecidae						
<i>Tenrec ecaudatus</i>	2	78	1	87	60	97
<i>Setifer setosus</i>	2	131	1	138	116	154
<i>Hemicentetes semispinosus</i>	1	74	1	68	96	32
<i>Echinops telfairi</i>	2	116	1	107	128	117
<i>Geogale aurita</i>	1	157	1	85	215	250
<i>Oryzorictes talpoides</i>	2	85	1	54	139	38
<i>Microgale dobsoni</i>	1	154	1	106	225	126
<i>M. talazaci</i>	1	226	1	129	291	392
<i>Limnogale mergulus</i>	1	143	1	73	190	263
<i>Potamogale velox</i>	2	288	1	323	239	306
<i>Micropotamogale lamottei</i>	1	37	1	30	51	23
<i>M. ruwenzorii</i>	4	6	--	--	--	--
Chrysochloridae						
<i>Chrysochloris asiatica</i>	1	87	1	51	134	86
<i>C. stuhlmanni</i>	2	152	1	103	211	159
Solenodontidae						
<i>Solenodon paradoxus</i>	2	28	1	13	48	20
Erinaceidae						
<i>Echinosorex gymnurus</i>	1	349	1	319	270	700
<i>Hylomys suillus</i>	1	810	1	636	766	1587
<i>Atelerix algirus</i>	2	253	1	183	255	507
<i>Erinaceus europaeus</i>	2	205	1	139	256	303
<i>Hemiechinus auritus</i>	3	325	1	214	388	550
Talpidae						
<i>Desmana moschata</i>	1	35	1	32	39	37
<i>Galemys pyrenaicus</i>	2	120	1	60	194	124
<i>Talpa europaea</i>	2	143	1	88	175	251
<i>T. micrura</i>	1	88	1	69	89	155
<i>Parascalops breweri</i>	1	47	1	37	53	64
<i>Scalopus aquaticus</i>	1	41	1	32	50	49

Table 50 (cont.)

	n	AOB	n	L1+2	L3-5	L6
Soricidae						
<i>Sorex alpinus</i>	1	86	--	--	--	--
<i>S. araneus</i>	2	157	1	92	189	307
<i>S. cinereus</i>	1	62	--	--	--	--
<i>S. fumeus</i>	1	93	1	68	132	73
<i>S. minutus</i>	1	65	1	40	84	99
<i>Microsorex hoyi</i>	1	90	--	--	--	--
<i>Neomys anomalus</i>	1	132	1	106	165	132
<i>N. fodiens</i>	2	144	1	101	182	196
<i>Blarina brevicauda</i>	1	71	--	--	--	--
<i>Cryptotis parva</i>	1	35	1	23	45	49
<i>Anourosorex squamipes</i>	1	36	--	--	--	--
<i>Crociodura attenuata</i>	2	197	1	149	254	209
<i>C. flavescens</i>	1	191	1	152	214	270
<i>C. hildegardeae</i>	1	154	--	--	--	--
<i>C. jacksoni</i>	2	129	--	--	--	--
<i>C. russula</i>	1	77	1	45	112	97
<i>C. suaveolens</i>	1	140	--	--	--	--
<i>Suncus etruscus</i>	1	129	1	87	178	139
<i>S. murinus</i>	1	164	1	142	168	233
<i>Scutisorex somereni</i>	1	123	1	82	159	166
<i>Sylvisorex granti</i>	1	108	1	66	177	66
<i>S. megalura</i>	1	133	1	92	203	84
<i>Ruwenzorisorex suncoides</i>	1	98	1	77	142	49
<i>Myosorex babaulti</i>	1	115	1	84	158	103

Number of individuals = n

Abbreviations:

- AOB accessory olfactory bulb
L1+2 vomeronasal fiber and glomerular layers
L3-5 external plexiform, mitral, and internal plexiform layers
L6 granular layer

Table 51. Average size indices of the accessory olfactory bulbs and of some of the AOB layers related to the average of the four Tenrecinae (AvTn = 100)

	N	AOB	N	AOB	L1+2	L3-5	L6
slope		.64		.64	.64	.64	.64
y-intercept		-1.903		-1.903	-2.219	-2.318	-2.785
Insectivora (AvIS)	50	138	41	150	112	177	211
Tenrecidae	12	125	11	135	109	159	163
Tenrecinae*	4	100	4	100	100	100	100
Geogalinae*	1	157	1	157	85	215	250
Oryzorictinae*	4	152	4	152	91	211	205
<i>Microgale</i>	2	190	2	190	117	258	259
terr.Oryzorictinae	3	155	3	155	96	219	185
<i>Limnogale</i>	1	143	1	143	73	190	263
Potamogalinae*	3	111	2	163	177	145	164
<i>Micropotamogale</i>	2	22	1	37	30	51	23
Chrysochloridae*	2	120	2	120	77	173	123
Solenodontidae*	1	28	1	28	13	48	20
Erinaceidae	5	388	5	388	298	387	729
Echinosoricinae*	2	580	2	580	478	518	1143
Erinaceinae*	3	261	3	261	179	300	453
Talpidae	6	79	6	79	53	100	113
Desmaninae*	2	78	2	78	46	116	81
Talpinae*	4	80	4	80	57	92	130
<i>Talpa</i>	2	115	2	115	79	132	203
Soricidae	24	114	16	123	88	160	142
Soricinae*	11	88	6	104	72	133	143
<i>Sorex</i>	5	93	3	105	67	135	160
terr. Soricinae	9	77	4	88	56	113	132
<i>Neomys</i>	2	138	2	138	103	174	164
Crociodurinae*	13	135	10	134	98	176	142
<i>Crociodura</i>	6	148	3	155	115	193	192
<i>Suncus</i>	2	146	2	146	114	173	186
<i>Sylvisorex</i>	2	121	2	121	79	190	75
AvIF of the 12 families and subfamilies marked by *		158		163	123	186	246

Number of species = N

Abbreviations:

- AOB accessory olfactory bulb
L1+2 vomeronasal fiber and glomerular layers
L3-5 external plexiform, mitral, and internal plexiform layers
L6 granular layer

Table 52. Volumes of paleocortical components

For volumes of total paleocortex (PAL) see Table 43

For size indices in species see Table 53

For average size indices in supraspecific taxonomic units see Table 54

	n	RB	PRPI	TOL	OLC	TRL	COA	SIN
Tenrecidae								
<i>Tenrec ecaudatus</i>	1	50.44	227.1	51.37	328.9	15.248	24.242	9.881
<i>Setifer setosus</i>	1	39.47	146.5	45.78	231.7	12.157	10.908	8.080
<i>Hemicentetes semis.</i>	1	14.70	68.5	19.84	103.0	4.109	3.090	4.353
<i>Echinops telfairi</i>	1	12.38	58.3	15.20	85.9	2.870	2.443	3.264
<i>Geogale aurita</i>	2	1.85	7.9	2.14	11.8	0.568	0.404	0.487
<i>Oryzorictes talpoides</i>	1	9.77	37.2	14.18	61.2	2.094	1.257	2.657
<i>Microgale dobsoni</i>	1	10.12	47.8	12.96	70.8	2.387	1.364	1.570
<i>M. talazaci</i>	1	13.70	67.7	17.50	98.9	3.692	2.630	1.818
<i>Limnogale mergulus</i>	1	7.91	33.4	13.18	54.5	3.092	1.411	1.558
<i>Potamogale velox</i>	1	18.19	71.6	17.57	107.4	5.033	6.181	9.800
<i>Micropotamogale lam.</i>	1	7.65	27.1	8.69	43.5	2.428	0.922	1.407
<i>M. ruwenzorii</i>	1	7.46	33.6	11.62	52.7	1.725	1.847	3.207
Chrysochloridae								
<i>Chrysochloris asiatica</i>	1	14.15	52.3	20.71	87.1	3.047	2.387	2.269
<i>C. stuhlmanni</i>	1	14.88	59.7	24.45	99.0	3.332	2.539	2.356
Solenodontidae								
<i>Solenodon paradoxus</i>	1	72.65	316.2	116.90	505.8	19.529	21.167	15.183
Erinaceidae								
<i>Echinorex gymnurus</i>	3	95.41	384.0	114.40	593.8	30.474	39.034	19.447
<i>Hylomys suillus</i>	1	18.82	73.9	22.99	115.7	4.232	4.961	3.114
<i>Atelerix algirus</i>	1	53.86	220.8	54.38	329.0	13.017	13.859	9.133
<i>Erinaceus europaeus</i>	1	61.93	228.7	56.46	347.1	14.777	23.427	13.274
<i>Hemiechinus auritus</i>	1	21.53	127.4	35.41	184.4	6.812	7.718	6.394
Talpidae								
<i>Desmana moschata</i>	1	34.28	135.8	39.97	210.0	6.662	4.396	8.407
<i>Galemys pyrenaicus</i>	1	7.95	36.5	18.74	63.2	1.841	0.928	4.052
<i>Talpa europaea</i>	1	11.71	45.6	19.17	76.4	2.510	4.055	2.895
<i>T. micrura</i>	2	10.41	44.5	16.98	71.9	2.208	2.211	2.331
<i>Parascalops breweri</i>	3	12.78	49.5	19.80	82.0	2.571	1.753	3.125
<i>Scalopus aquaticus</i>	1	18.57	82.5	25.86	127.0	3.716	3.440	4.926

Table 52 (cont.)

	n	RB	PRPI	TOL	OLC	TRL	COA	SIN
Soricidae								
<i>Sorex alpinus</i>	3	4.37	15.4	6.28	26.1	0.464	0.648	0.715
<i>S. araneus</i>	1	3.96	15.9	5.98	25.8	0.676	0.530	0.826
<i>S. cinereus</i>	1	3.21	12.5	4.83	20.5	0.482	0.630	0.530
<i>S. fumeus</i>	1	4.36	17.0	6.03	27.4	0.523	0.735	0.660
<i>S. minutus</i>	1	1.75	7.2	3.75	12.7	0.342	0.363	0.578
<i>Microsorex hoyi</i>	1	1.87	5.8	2.86	10.5	0.293	0.245	0.461
<i>Neomys anomalus</i>	3	3.34	13.1	5.23	21.7	0.533	0.399	0.853
<i>N. fodiens</i>	1	4.35	12.4	6.04	22.8	0.712	0.842	1.663
<i>Blarina brevicauda</i>	1	6.88	23.5	9.61	40.0	1.442	1.492	1.470
<i>Cryptotis parva</i>	3	4.77	16.2	6.64	27.6	0.752	0.701	0.943
<i>Anourosorex squam.</i>	4	7.62	28.7	10.12	46.4	1.194	1.191	1.641
<i>Crociodura attenuata</i>	1	3.56	15.2	4.91	23.7	0.491	0.662	0.630
<i>C. flavescens</i>	1	8.14	27.6	7.21	42.9	1.682	1.906	1.653
<i>C. hildegardeae</i>	1	4.46	17.9	6.09	28.5	0.713	0.657	0.615
<i>C. jacksoni</i>	1	4.47	18.6	5.32	28.4	1.037	0.947	0.550
<i>C. russula</i>	1	3.64	13.7	4.67	22.0	0.678	0.597	0.747
<i>C. suaveolens</i>	3	3.55	13.7	4.41	21.6	0.531	0.465	0.655
<i>Suncus etruscus</i>	1	1.17	4.0	1.45	6.6	0.204	0.141	0.145
<i>S. murinus</i>	1	7.30	29.1	6.75	43.1	1.316	1.000	1.631
<i>Scutisorex somereni</i>	1	13.11	49.1	13.90	76.1	2.533	2.225	1.868
<i>Sylvisorex granti</i>	1	3.33	14.1	4.35	21.8	0.565	0.581	0.487
<i>S. megalura</i>	1	3.27	15.5	4.91	23.7	0.557	0.555	0.512
<i>Ruwenzorisorex sunc.</i>	1	4.12	17.8	5.05	27.0	0.709	0.625	0.632
<i>Myosorex babaulti</i>	1	7.30	30.0	11.00	48.3	1.435	1.368	1.213

Number of individuals = n

Abbreviations:

- COA complex of internal fibers passing through the anterior commissure
- OLC olfactory cortices (RB + PRPI + TOL)
- PRPI regio prepiriformis
- RB regio retrobulbaris (=nucleus olfactorius anterior)
- SIN substantia innominata
- TOL tuberculum olfactorium
- TRL tractus olfactorius lateralis

Table 53. Size indices of paleocortical components related to the average of the four Tenrecinae (AvTn = 100)

For size indices of total paleocortex (PAL) see Table 44

For volumes in species see Table 52

For average size indices in supraspecific taxonomic units see Table 54

	n	RB	PRPI	TOL	OLC	TRL	COA	SIN
slope		.63	.59	.53	.59	.67	.76	.64
y-intercept		-.064	.662	.242	.841	-.704	-.921	-.716
Tenrecidae								
<i>Tenrec ecaudatus</i>	1	83	92	82	89	84	120	68
<i>Setifer setosus</i>	1	146	127	145	133	158	143	127
<i>Hemicentetes semispin.</i>	1	85	90	92	90	86	69	108
<i>Echinops telfairi</i>	1	86	91	81	89	73	68	97
<i>Geogale aurita</i>	2	50	44	36	44	61	58	58
<i>Oryzorictes talpoides</i>	1	104	87	109	94	84	59	122
<i>Microgale dobsoni</i>	1	132	135	118	132	119	82	89
<i>M. talazaci</i>	1	138	150	128	145	139	115	79
<i>Limnogale mergulus</i>	1	53	51	69	55	76	38	45
<i>Potamogale velox</i>	1	37	35	33	35	34	39	83
<i>Micropotamogale lam.</i>	1	64	51	55	54	76	33	51
<i>M. ruwenzorii</i>	1	48	49	59	51	41	48	89
Chrysochloridae								
<i>Chrysochloris asiatica</i>	1	141	115	151	126	114	103	98
<i>C. stuhlmanni</i>	1	168	147	198	162	142	128	115
Solenodontidae								
<i>Solenodon paradoxus</i>	1	139	148	212	157	126	125	122
Erinaceidae								
<i>Echinorex gymnurus</i>	3	161	159	187	163	172	198	138
<i>Hylomys suillus</i>	1	171	148	155	154	143	191	122
<i>Atelerix algirus</i>	1	98	98	94	97	79	77	69
<i>Erinaceus europaeus</i>	1	102	93	91	94	82	116	92
<i>Hemiechinus auritus</i>	1	80	111	112	106	89	102	101
Talpidae								
<i>Desmana moschata</i>	1	85	81	91	83	57	36	88
<i>Galemys pyrenaicus</i>	1	70	71	123	82	60	35	154
<i>Talpa europaea</i>	1	84	74	106	82	66	119	90
<i>T. micrura</i>	2	116	108	135	115	92	109	112
<i>Parascalops breweri</i>	3	120	103	137	113	90	71	127
<i>Scalopus aquaticus</i>	1	108	109	120	111	78	78	123

Table 53 (cont.)

	n	RB	PRPI	TOL	OLC	TRL	COA	SIN
Soricidae								
<i>Sorex alpinus</i>	3	112	82	101	91	47	87	80
<i>S. araneus</i>	1	106	88	100	95	72	76	97
<i>S. cinereus</i>	1	132	103	116	112	81	151	96
<i>S. fumeus</i>	1	132	106	112	113	64	122	88
<i>S. minutus</i>	1	80	65	98	76	64	98	116
<i>Microsorex hoyi</i>	1	118	71	98	86	78	98	129
<i>Neomys anomalus</i>	3	83	67	82	74	52	52	92
<i>N. fodiens</i>	1	90	54	82	66	58	88	151
<i>Blarina brevicauda</i>	1	122	88	113	99	99	129	113
<i>Cryptotis parva</i>	3	130	91	113	103	82	102	113
<i>Anourosorex squam.</i>	4	133	106	118	114	81	101	125
<i>Crocidura attenuata</i>	1	101	89	86	92	56	101	79
<i>C. flavescens</i>	1	112	82	69	84	89	122	99
<i>C. hildegardeae</i>	1	120	99	102	104	76	94	72
<i>C. jacksoni</i>	1	105	90	79	91	96	114	56
<i>C. russula</i>	1	92	72	75	77	68	80	83
<i>C. suaveolens</i>	3	95	75	73	79	56	66	77
<i>Suncus etruscus</i>	1	92	60	60	66	68	73	51
<i>S. murinus</i>	1	92	79	60	78	63	57	89
<i>Scutisorex somereni</i>	1	111	93	88	95	79	79	68
<i>Sylvisorex granti</i>	1	164	138	121	141	115	172	106
<i>S. megalura</i>	1	129	124	114	125	90	127	90
<i>Ruwenzorisorex sunc.</i>	1	77	70	62	70	51	57	51
<i>Myosorex babaulti</i>	1	142	123	140	131	109	132	103

Number of individuals = n

Abbreviations:

COA	complex of internal fibers passing through the anterior commissure
OLC	olfactory cortices (RB + PRPI + TOL)
PRPI	regio prepiriformis
RB	regio retrobulbaris (=nucleus olfactorius anterior)
SIN	substantia innominata
TOL	tuberculum olfactorium
TRL	tractus olfactorius lateralis

Table 54. Average size indices of paleocortical components related to the average of the four Tenrecinae (AvTn = 100)

	N	RB	PRPI	TOL	OLC	TRL	COA	SIN
slope		.63	.59	.53	.59	.67	.76	.64
y-intercept		-.064	.662	.242	.841	-.704	-.921	-.716
Insectivora (AvIS)	50	107	94	104	98	84	95	96
Tenrecidae	12	86	83	84	84	86	73	85
Tenrecinae*	4	100	100	100	100	100	100	100
Geogalinae*	1	50	44	36	44	61	58	58
Oryzorictinae*	4	107	106	106	107	104	73	84
<i>Microgale</i>	2	135	142	123	139	129	99	84
terr. Oryzorictinae	3	125	124	119	124	114	85	97
<i>Limnogale</i>	1	53	51	69	55	76	38	45
Potamogalinae*	3	50	45	49	47	50	40	75
<i>Micropotamogale</i>	2	56	50	57	52	58	40	70
Chrysochloridae*	2	155	131	174	144	128	116	106
Solenodontidae*	1	139	148	212	157	126	125	122
Erinaceidae	5	122	122	128	123	113	137	104
Echinosoricinae*	2	166	154	171	158	157	195	130
Erinaeinae*	3	93	101	99	99	83	98	88
Talpidae	6	97	91	119	98	74	74	116
Desmaninae*	2	78	76	107	82	58	35	121
Talpinae*	4	107	98	125	105	82	94	113
<i>Talpa</i>	2	100	91	121	99	79	114	101
Soricidae	24	111	88	94	94	75	99	93
Soricinae*	11	113	84	103	93	71	100	109
<i>Sorex</i>	5	112	89	105	97	66	107	96
terr. Soricinae	9	118	89	108	99	74	107	107
<i>Neomys</i>	2	87	61	82	70	55	70	122
Crocidurinae*	13	110	92	87	95	78	98	79
<i>Crocidura</i>	6	104	85	81	88	73	96	78
<i>Suncus</i>	2	82	70	60	72	66	65	70
<i>Sylvisorex</i>	2	147	131	118	133	102	150	98
AvIF of the 12 families and subfamilies marked by *		106	98	114	103	92	94	99

Number of species = N

Abbreviations:

COA	complex of anterior commissure		
OLC	olfactory cortices (RB + PRPI + TOL)		
PRPI	regio prepiriformis	TOL	tuberculum olfactorium
RB	regio retrobulbaris (ncl. olf. ant.)		
SIN	substantia innominata	TRL	tractus olfactorius lateralis

Table 55. Volumes of amygdala and some amygdaloid components

For size indices in species see Table 56

For average size indices in supraspecific taxonomic units see Table 57

	AMY	n	MAM	LAM	n	NTO	MCB
Tenrecidae							
<i>Tenrec ecaudatus</i>	88.80	2	39.33	49.47	1	3.129	9.151
<i>Setifer setosus</i>	67.62	3	31.63	35.99	1	2.826	7.960
<i>Hemicentetes semis.</i>	20.56	2	9.52	11.04	1	1.185	2.221
<i>Echinops telfairi</i>	22.51	2	11.17	11.34	1	0.705	3.203
<i>Geogale aurita</i>	4.12	1	2.04	2.08	--	--	--
<i>Oryzorictes talpoides</i>	20.75	1	9.66	11.09	1	0.639	1.210
<i>Microgale dobsoni</i>	16.89	1	6.73	10.16	1	0.749	1.709
<i>M. talazaci</i>	24.44	1	11.72	12.72	1	0.724	1.647
<i>Limnogale mergulus</i>	26.82	2	11.04	15.78	1	0.310	2.530
<i>Potamogale velox</i>	62.16	1	28.65	33.51	1	0.455	5.323
<i>Micropotamogale lam.</i>	19.05	1	8.74	10.31	1	0.268	1.388
<i>M. ruwenzorii</i>	22.10	1	9.25	12.85	1	0.466	1.639
Chrysochloridae							
<i>Chrysochloris asiatica</i>	20.05	1	9.08	10.97	1	0.831	1.628
<i>C. stuhlmanni</i>	23.17	1	12.15	11.02	1	0.956	2.113
Solenodontidae							
<i>Solenodon paradoxus</i>	135.36	1	51.31	84.05	1	2.522	14.260
Erinaceidae							
<i>Echinosorex gymnurus</i>	170.14	3	69.57	100.57	--	--	--
<i>Hylomys suillus</i>	37.43	1	17.21	20.22	--	--	--
<i>Atelerix algirus</i>	103.30	1	46.82	56.48	1	3.134	7.650
<i>Erinaceus europaeus</i>	108.67	3	45.00	63.67	1	3.748	11.518
<i>Hemiechinus auritus</i>	71.06	1	29.55	41.51	1	2.767	4.878
Talpidae							
<i>Desmana moschata</i>	90.80	1	40.87	49.93	1	0.094	12.178
<i>Galemys pyrenaicus</i>	36.43	1	14.95	21.48	1	0.231	5.669
<i>Talpa europaea</i>	26.41	1	11.37	15.04	1	0.565	3.497
<i>T. micrura</i>	22.69	2	11.03	11.66	--	--	--
<i>Parascalops breweri</i>	22.58	3	11.14	11.44	--	--	--
<i>Scalopus aquaticus</i>	30.87	1	14.49	16.38	--	--	--

Table 55 (cont.)

	AMY	n	MAM	LAM	n	NTO	MCB
Soricidae							
<i>Sorex alpinus</i>	8.05	3	3.70	4.35	--	-.-	-.-
<i>S. araneus</i>	7.70	3	3.74	3.96	2	0.275	0.758
<i>S. cinereus</i>	4.89	1	2.38	2.51	1	0.244	0.541
<i>S. fumeus</i>	7.43	1	3.67	3.76	1	0.330	0.642
<i>S. minutus</i>	3.46	2	1.83	1.63	1	0.179	0.572
<i>Microsorex hoyi</i>	3.28	1	1.35	1.93	1	0.106	0.232
<i>Neomys anomalus</i>	8.18	3	3.94	4.24	--	-.-	-.-
<i>N. fodiens</i>	8.59	1	4.13	4.46	1	0.239	1.161
<i>Blarina brevicauda</i>	10.93	1	5.02	5.91	1	0.454	0.894
<i>Cryptotis parva</i>	9.10	3	4.55	4.55	--	-.-	-.-
<i>Anourosorex squamipes</i>	15.35	4	7.40	7.95	--	-.-	-.-
<i>Crocidura attenuata</i>	6.66	1	3.47	3.19	--	-.-	-.-
<i>C. flavescens</i>	12.63	2	6.20	6.43	1	0.415	1.203
<i>C. hildegardae</i>	6.89	1	3.53	3.36	1	0.336	0.592
<i>C. jacksoni</i>	7.62	1	3.50	4.12	1	0.237	0.539
<i>C. russula</i>	6.65	2	3.24	3.41	1	0.240	0.762
<i>C. suaveolens</i>	5.94	3	2.94	3.00	--	-.-	-.-
<i>Suncus etruscus</i>	1.73	1	0.92	0.81	1	0.084	0.118
<i>S. murinus</i>	11.09	2	5.43	5.66	1	0.442	0.793
<i>Scutisorex somereni</i>	17.85	1	8.02	9.83	1	0.790	1.664
<i>Sylvisorex granti</i>	5.06	1	2.42	2.64	1	0.320	0.541
<i>S. megalura</i>	5.58	1	2.90	2.68	1	0.268	0.454
<i>Ruwenzorisorex sunc.</i>	10.25	1	4.74	5.51	1	0.178	0.694
<i>Myosorex babaulti</i>	12.08	1	6.22	5.86	1	0.584	1.095

Number of individuals of the total amygdaloid complex (AMY) as in Table 10

Number of individuals = n

Abbreviations:

- AMY total amygdaloid complex, amygdala
 LAM lateral amygdaloid division, including basal, cortical, and lateral nuclei
 MAM medial amygdaloid division, including central and medial nuclei, and anterior amygdaloid area
 MCB magnocellular part of the basal nucleus
 NTO bed nucleus of the lateral olfactory tract

Table 56. Size indices of the amygdala and some amygdaloid components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 55

For average size indices in supraspecific taxonomic units see Table 57

	AMY	n	MAM	LAM	n	NTO	MCB
slope	.60		.60	.60		.60	.60
y-intercept	.232		-.098	-.043		-1.159	-.696
Tenrecidae							
<i>Tenrec ecaudatus</i>	91	2	86	95	1	79	79
<i>Setifer setosus</i>	149	3	149	149	1	153	149
<i>Hemicentetes semispin.</i>	70	2	69	70	1	99	64
<i>Echinops telfairi</i>	90	2	96	86	1	69	109
<i>Geogale aurita</i>	61	1	64	58	--	--	--
<i>Oryzorictes talpoides</i>	125	1	125	126	1	95	62
<i>Microgale dobsoni</i>	124	1	106	140	1	135	106
<i>M. talazaci</i>	140	1	144	137	1	102	80
<i>Limnogale mergulus</i>	104	2	92	116	1	30	83
<i>Potamogale velox</i>	77	1	76	78	1	14	56
<i>Micropotamogale lam.</i>	92	1	90	94	1	32	57
<i>M. ruwenzorii</i>	83	1	75	91	1	43	52
Chrysochloridae							
<i>Chrysochloris asiatica</i>	114	1	110	117	1	116	78
<i>C. stuhlmanni</i>	148	1	166	133	1	150	114
Solenodontidae							
<i>Solenodon paradoxus</i>	160	1	129	187	1	73	142
Erinaceidae							
<i>Echinorex gymnurus</i>	178	3	155	198	--	--	--
<i>Hylomys suillus</i>	194	1	191	197	--	--	--
<i>Atelerix algirus</i>	115	1	112	119	1	86	72
<i>Erinaceus europaeus</i>	111	3	99	123	1	95	100
<i>Hemiechinus auritus</i>	157	1	140	173	1	151	92
Talpidae							
<i>Desmana moschata</i>	137	1	132	142	1	4	156
<i>Galemys pyrenaicus</i>	184	1	161	204	1	29	242
<i>Talpa europaea</i>	110	1	101	118	1	58	123
<i>T. micrura</i>	142	2	148	138	--	--	--
<i>Parascalops breweri</i>	121	3	128	116	--	--	--
<i>Scalopus aquaticus</i>	105	1	105	105	--	--	--

Table 56 (cont.)

	AMY	n	MAM	LAM	n	NTO	MCB
Soricidae							
<i>Sorex alpinus</i>	112	3	110	114	--	--	--
<i>S. araneus</i>	112	3	116	109	2	98	93
<i>S. cinereus</i>	107	1	111	103	1	131	100
<i>S. fumeus</i>	121	1	128	116	1	133	89
<i>S. minutus</i>	83	2	94	74	1	106	117
<i>Microsorex hoyi</i>	108	1	94	119	1	86	64
<i>Neomys anomalus</i>	110	3	114	108	--	--	--
<i>N. fodiens</i>	98	1	101	96	1	67	112
<i>Blarina brevicauda</i>	107	1	105	109	1	109	74
<i>Cryptotis parva</i>	135	3	144	127	--	--	--
<i>Anourosorex squamipes</i>	149	4	153	145	--	--	--
<i>Crociodura attenuata</i>	102	1	114	92	--	--	--
<i>C. flavescens</i>	98	2	102	94	1	79	79
<i>C. hildegardeae</i>	100	1	110	92	1	120	73
<i>C. jacksoni</i>	97	1	95	99	1	74	58
<i>C. russula</i>	92	2	96	89	1	82	89
<i>C. suaveolens</i>	86	3	91	82	--	--	--
<i>Suncus etruscus</i>	70	1	79	62	1	83	40
<i>S. murinus</i>	79	2	82	76	1	77	48
<i>Scutisorex somereni</i>	87	1	83	90	1	94	69
<i>Sylvisorex granti</i>	131	1	134	129	1	204	119
<i>S. megalura</i>	118	1	131	106	1	139	81
<i>Ruwenzorisorex suncoides</i>	105	1	104	107	1	45	60
<i>Myosorex babaulti</i>	129	1	142	118	1	154	99

Number of individuals of the total amygdaloid complex (AMY) as in Table 10

Number of individuals = n

Abbreviations:

AMY total amygdaloid complex, amygdala

LAM lateral amygdaloid division, including basal, cortical, and lateral nuclei

MAM medial amygdaloid division, including central and medial nuclei, and anterior amygdaloid area

MCB magnocellular part of the basal nucleus

NTO bed nucleus of the lateral olfactory tract

Table 57. Average size indices of the amygdala and some amygdaloid components related to the average of the four Tenrecinae (AvTn = 100)

	N	AMY	MAM	LAM	N	NTO	MCB
slope		.60	.60	.60		.60	.60
y-intercept		.232	-.098	-.043		-1.159	-.696
Insectivora (AvIS)	50	114	114	115	38	92	92
Tenrecidae	12	101	98	103	11	77	82
Tenrecinae*	4	100	100	100	4	100	100
Geogalinae*	1	61	64	58	--	--	--
Oryzorictinae*	4	123	116	130	4	90	83
<i>Microgale</i>	2	132	125	139	2	119	93
terr. Oryzorictinae	3	130	125	135	3	111	83
<i>Limnogale</i>	1	104	92	116	1	30	83
Potamogalinae*	3	84	80	88	3	30	55
<i>Micropotamogale</i>	2	88	82	92	2	38	55
Chrysochloridae*	2	131	138	125	2	133	96
Solenodontidae*	1	160	129	187	1	73	142
Erinaceidae	5	151	139	162	3	110	88
Echinosoricinae*	2	186	173	198	--	--	--
Erinaceinae*	3	128	117	138	3	110	88
Talpidae	6	133	129	137	3	30	174
Desmaninae*	2	161	147	173	2	16	199
Talpinae*	4	120	121	119	1	58	123
<i>Talpa</i>	2	126	125	128	1	58	123
Soricidae	24	106	110	102	18	105	81
Soricinae*	11	113	116	111	7	104	93
<i>Sorex</i>	5	107	112	103	4	117	100
terr. Soricinae	9	115	118	113	6	111	90
<i>Neomys</i>	2	104	107	102	1	67	112
Crocidurinae*	13	100	105	95	11	105	74
<i>Crocidura</i>	6	96	101	91	4	89	75
<i>Suncus</i>	2	74	81	69	2	80	44
<i>Sylvisorex</i>	2	124	132	117	2	172	100
AvTn of the families and subfamilies marked by *	(12)	122	117	127	(10)	82	105

Number of species = N

Abbreviations:

AMY total amygdaloid complex
 LAM lateral amygdaloid division
 MAM medial amygdaloid division

MCB large-celled part of basal nucleus
 NTO bed nucleus of the lateral olfactory tract

Table 58. Volumes of hippocampal components

For size indices in species see Table 59
For average size indices in supraspecific taxonomic units see Table 60

	n	HIP	n	HR	HA	HS	HP
Tenrecidae							
<i>Tenrec ecaudatus</i>	3	147.1	1	136.3	10.82	2.086	8.73
<i>Setifer setosus</i>	3	96.6	1	89.9	6.69	1.041	5.65
<i>Hemicentetes semispin.</i>	2	63.1	1	58.3	4.84	0.644	4.20
<i>Echinops telfairi</i>	2	46.7	1	44.7	2.01	0.240	1.77
Erinaceidae							
<i>Erinaceus europaeus</i>	5	237.9	1	223.2	14.76	2.350	12.41
Soricidae							
<i>Sorex araneus</i>	5	22.8	1	21.2	1.51	0.244	1.27
<i>S. minutus</i>	4	12.4	1	11.7	0.68	0.120	0.56
<i>Crocidura flavescens</i>	3	38.3	1	35.8	2.58	0.515	2.06
<i>C. russula</i>	5	17.9	1	16.7	1.22	0.175	1.04
<i>Suncus murinus</i>	3	33.4	1	31.0	2.37	0.605	1.77

Number of individuals = n

Abbreviations:

- HA hippocampus anterior (hippocampus supra- + precommissuralis, HS + HP)
- HIP total hippocampus
- HP hippocampus precommissuralis
- HR hippocampus retrocommissuralis
- HS hippocampus supracommissuralis

Table 59. Size indices of hippocampal components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 58

For average size indices in supraspecific taxonomic units see Table 60

	n	HIP	n	HR	HA	HS	HP
slope		.57		.57	.57	.57	.57
y-intercept		.581		.552	-.599	-1.420	-.670
Tenrecidae							
<i>Tenrec ecaudatus</i>	3	82	1	82	92	117	87
<i>Setifer setosus</i>	3	112	1	112	118	121	117
<i>Hemicentetes semispin.</i>	2	110	1	109	128	113	131
<i>Echinops telfairi</i>	2	96	1	98	62	49	65
Erinaceidae							
<i>Erinaceus europaeus</i>	5	134	1	134	126	132	124
Soricidae							
<i>Sorex araneus</i>	5	159	1	159	160	171	158
<i>S. minutus</i>	4	140	1	141	116	135	112
<i>Crocidura flavescens</i>	3	147	1	146	149	198	141
<i>C. russula</i>	5	119	1	119	122	117	123
<i>Suncus murinus</i>	3	118	1	117	127	214	111

Number of individuals = n

Abbreviations:

HA	hippocampus anterior (hippocampus supra- + precommissuralis, HS + HP)
HIP	total hippocampus
HP	hippocampus precommissuralis
HR	hippocampus retrocommissuralis
HS	hippocampus supracommissuralis

Table 60. Average size indices of hippocampal components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 58
For size indices in species see Table 59

	N	HIP	HR	HA	HS	HP
slope		.57	.57	.57	.57	.57
y-intercept		.581	.552	-.599	-1.420	-.670
Insectivora (AvIS)	10	122	122	120	137	117
Tenrecidae						
Tenrecinae*	4	100	100	100	100	100
Erinaceidae						
Erinaceinae*	1	134	134	126	132	124
Soricidae	5	137	136	135	167	129
Soricinae*	2	150	150	138	153	135
Crocidurinae*	3	128	127	133	176	125
<i>Crocidura</i>	2	133	133	136	157	132
AvIF of the 4 subfamilies marked by *		128	128	124	140	121

Number of species = N

Abbreviations:

- HA hippocampus anterior (hippocampus supra- + precommissuralis, HS + HP)
- HIP total hippocampus
- HP hippocampus precommissuralis
- HR hippocampus retrocommissuralis
- HS hippocampus supracommissuralis

Table 61. Volumes of neocortical components

For size indices in species see Table 62

For average size indices in supraspecific taxonomic units see Table 63

	n	NEO	n	NW	NG	NG1	NG2-6
Tenrecidae							
<i>Tenrec ecaudatus</i>	3	272.8	1	35.76	237.0	73.32	163.7
<i>Setifer setosus</i>	3	136.6	1	17.73	118.9	38.18	80.7
<i>Hemicentetes semis.</i>	2	76.9	1	6.14	70.7	24.02	46.7
<i>Echinops telfairi</i>	2	51.7	1	5.25	46.5	15.33	31.1
<i>Geogale aurita</i>	2	7.2	1	0.53	6.7	1.81	4.9
<i>Oryzorictes talpoides</i>	1	66.5	1	5.91	60.6	18.63	41.9
<i>Microgale dobsoni</i>	1	62.6	1	5.59	57.0	16.79	40.2
<i>M. talazaci</i>	2	88.9	1	9.71	79.2	23.59	55.6
<i>Limnogale mergulus</i>	1	195.6	1	18.09	177.5	42.50	135.0
<i>Potamogale velox</i>	5	1050.0	1	140.89	909.1	141.57	767.6
<i>Micropotamogale lamottei</i>	3	144.4	1	14.36	130.0	30.25	99.8
<i>M. ruwenzorii</i>	4	227.8	1	23.33	204.4	41.12	163.3
Chrysochloridae							
<i>Chrysochloris asiatica</i>	1	112.6	1	5.80	106.8	29.05	77.8
<i>C. stuhlmanni</i>	2	128.3	1	7.61	120.7	31.25	89.4
Solenodontidae							
<i>Solenodon paradoxus</i>	2	668.6	1	68.55	600.1	162.66	437.4
Erinaceidae							
<i>Echinorex gymnurus</i>	3	1085.0	1	224.98	860.0	194.52	665.5
<i>Hylomys suillus</i>	1	177.8	1	18.12	159.6	44.82	114.8
<i>Atelerix algirus</i>	3	510.3	1	64.88	445.4	119.32	326.1
<i>Erinaceus europaeus</i>	5	525.1	1	68.70	456.4	129.79	326.6
<i>Hemiechinus auritus</i>	3	306.2	1	30.47	275.8	74.24	201.5
Talpidae							
<i>Desmana moschata</i>	1	983.4	1	129.55	853.8	157.34	696.5
<i>Galemys pyrenaicus</i>	2	338.7	1	33.73	305.0	53.73	251.2
<i>Talpa europaea</i>	2	181.2	1	14.13	167.1	38.01	129.1
<i>T. micrura</i>	2	157.2	1	10.49	146.7	33.06	113.6
<i>Parascalops breweri</i>	3	175.6	1	10.85	164.8	34.98	129.8
<i>Scalopus aquaticus</i>	1	252.9	1	20.26	232.6	58.99	173.6

Table 61 (cont.)

	n	NEO	n	NW	NG	NG1	NG2-6
Soricidae							
<i>Sorex alpinus</i>	3	45.5	1	2.34	43.2	10.42	32.8
<i>S. araneus</i>	5	28.2	3	2.55	25.6	5.56	20.0
<i>S. cinereus</i>	3	28.8	3	1.97	26.8	6.77	20.0
<i>S. fumeus</i>	2	48.0	2	2.80	45.2	10.45	34.8
<i>S. minutus</i>	4	15.9	1	1.19	14.7	3.20	11.5
<i>Microsorex hoyi</i>	3	15.2	3	0.85	14.4	4.05	10.3
<i>Neomys anomalus</i>	3	59.0	1	3.96	55.1	12.32	42.8
<i>N. fodiens</i>	2	56.6	2	5.19	51.4	11.10	40.3
<i>Blarina brevicauda</i>	3	68.7	3	5.17	63.5	15.33	48.2
<i>Cryptotis parva</i>	3	36.2	1	2.67	33.5	8.18	25.3
<i>Anourosorex squamipes</i>	4	67.6	1	3.32	64.3	17.71	46.6
<i>Crocidura attenuata</i>	1	25.3	1	1.90	23.4	6.63	16.8
<i>C. flavescens</i>	3	53.1	3	4.18	48.9	14.81	34.1
<i>C. hildegardeae</i>	4	26.0	1	1.92	24.1	7.00	17.1
<i>C. jacksoni</i>	1	28.6	1	2.16	26.4	6.82	19.6
<i>C. russula</i>	5	25.3	5	1.77	23.6	6.89	16.7
<i>C. suaveolens</i>	3	24.6	1	1.68	22.9	5.91	17.0
<i>Suncus etruscus</i>	3	7.1	3	0.48	6.6	2.11	4.5
<i>S. murinus</i>	3	50.5	3	4.63	45.9	13.62	32.3
<i>Scutisorex somereni</i>	3	87.2	1	6.81	80.3	21.79	58.6
<i>Sylvisorex granti</i>	3	20.6	1	1.42	19.2	5.25	14.0
<i>S. megalura</i>	3	23.9	1	1.33	22.6	6.04	16.6
<i>Ruwenzorisorex suncoides</i>	1	59.5	1	3.96	55.6	15.23	40.4
<i>Myosorex babaulti</i>	1	45.3	1	3.68	41.6	10.63	31.0

Number of individuals = n

Abbreviations:

- NEO total neocortex
- NG neocortical grey matter
- NG1 molecular layer of neocortical grey matter
- NG2-6 cell layers 2-6 of neocortical grey matter
- NW neocortical white matter

Table 62. Size indices of neocortical components related to the average of the four Tenrecinae (AvTn = 100)

For volumes in species see Table 61

For average size indices in supraspecific taxonomic units see Table 63

	n	NEO	n	NW	NG	NG1	NG2-6
slope		.73		.83	.72	.68	.73
y-intercept		.346		-.847	.318	-.077	.124
Tenrecidae							
<i>Tenrec ecaudatus</i>	3	89	1	93	88	89	89
<i>Setifer setosus</i>	3	114	1	133	112	111	112
<i>Hemicentetes semispin.</i>	2	108	1	84	111	113	109
<i>Echinops telfairi</i>	2	89	1	90	89	87	89
<i>Geogale aurita</i>	2	61	1	55	61	45	68
<i>Oryzorictes talpoides</i>	1	189	1	179	190	169	198
<i>Microgale dobsoni</i>	1	225	1	222	227	190	241
<i>M. talazaci</i>	2	237	1	274	234	202	247
<i>Limnogale mergulus</i>	1	325	1	298	329	234	374
<i>Potamogale velox</i>	5	434	1	478	428	214	529
<i>Micropotamogale lam.</i>	3	312	1	319	312	213	359
<i>M. ruwenzorii</i>	4	365	1	369	365	219	436
Chrysochloridae							
<i>Chrysochloris asiatica</i>	1	296	1	161	312	246	341
<i>C. stuhlmanni</i>	2	390	1	250	406	303	453
Solenodontidae							
<i>Solenodon paradoxus</i>	2	260	1	217	266	232	284
Erinaceidae							
<i>Echinorex gymnurus</i>	3	364	1	602	329	242	372
<i>Hylomys suillus</i>	1	419	1	444	418	342	451
<i>Atelerix algirus</i>	3	186	1	190	185	160	198
<i>Erinaceus europaeus</i>	5	172	1	179	171	158	179
<i>Hemiechinus auritus</i>	3	257	1	231	260	216	281
Talpidae							
<i>Desmana moschata</i>	1	519	1	579	510	298	612
<i>Galemys pyrenaicus</i>	2	772	1	796	772	398	954
<i>Talpa europaea</i>	2	327	1	256	336	227	389
<i>T. micrura</i>	2	468	1	335	483	314	564
<i>Parascalops breweri</i>	3	432	1	279	450	278	532
<i>Scalopus aquaticus</i>	1	357	1	278	367	280	409

Table 62 (cont.)

	n	NEO	n	NW	NG	NG1	NG2-6
Soricidae							
<i>Sorex alpinus</i>	3	356	1	225	369	244	428
<i>S. araneus</i>	5	233	3	261	231	137	276
<i>S. cinereus</i>	3	391	3	354	395	265	454
<i>S. fumeus</i>	2	458	2	336	470	293	553
<i>S. minutus</i>	4	243	1	244	244	139	294
<i>Microsorex hoyi</i>	3	339	3	267	344	250	383
<i>Neomys anomalus</i>	3	445	1	364	454	278	537
<i>N. fodiens</i>	2	348	2	379	347	207	413
<i>Blarina brevicauda</i>	3	351	3	306	357	241	411
<i>Cryptotis parva</i>	3	306	1	280	309	206	357
<i>Anourosorex squamipes</i>	4	341	1	193	356	275	392
<i>Crociodura attenuata</i>	1	224	1	210	226	174	248
<i>C. flavescens</i>	3	203	3	178	206	178	218
<i>C. hildegardeae</i>	4	215	1	196	218	172	236
<i>C. jacksoni</i>	1	202	1	184	204	145	231
<i>C. russula</i>	5	197	5	169	200	160	216
<i>C. suaveolens</i>	3	202	1	170	205	144	233
<i>Suncus etruscus</i>	3	203	3	202	203	165	214
<i>S. murinus</i>	3	174	3	175	175	148	186
<i>Scutisorex somereni</i>	3	190	1	153	195	155	213
<i>Sylvisorex granti</i>	3	345	1	322	347	249	389
<i>S. megalura</i>	3	311	1	227	319	226	359
<i>Ruwenzorisorex sunc.</i>	1	323	1	250	331	253	365
<i>Myosorex babaulti</i>	1	258	1	246	260	185	294

Number of individuals = n

Abbreviations:

- NEO total neocortex
- NG neocortical grey matter
- NG1 molecular layer of neocortical grey matter
- NG2-6 cell layers 2-6 of neocortical grey matter
- NW neocortical white matter

Table 63. Average size indices of neocortical components related to the average of the four Tenrecinae (AvTn = 100)

	N	NEO	NW	NG	NG1	NG2-6
slope		.73	.83	.72	.68	.73
y-intercept		.346	-.847	.318	-.077	.124
Insectivora (AvIS)	50	292	266	296	209	335
Tenrecidae	12	212	216	212	157	238
Tenrecinae*	4	100	100	100	100	100
Geogalinae*	1	61	55	61	45	68
Oryzorictinae*	4	244	243	245	199	265
<i>Microgale</i>	2	231	248	230	196	244
terr. Oryzorictinae	3	217	225	217	187	229
<i>Limnogale</i>	1	325	298	329	234	374
Potamogalinae*	3	370	389	368	215	441
<i>Micropotamogale</i>	2	338	344	339	216	398
Chrysochloridae*	2	343	205	359	274	397
Solenodontidae*	1	260	217	266	232	284
Erinaceidae	5	279	329	273	224	296
Echinosoricinae*	2	391	523	373	292	412
Erinaceinae*	3	205	200	205	178	219
Talpidae	6	479	421	486	299	577
Desmaninae*	2	645	688	641	348	783
Talpinae*	4	396	287	409	274	473
<i>Talpa</i>	2	397	296	410	270	476
Soricidae	24	286	246	290	204	329
Soricinae*	11	347	292	352	230	409
<i>Sorex</i>	5	336	284	342	216	401
terr. Soricinae	9	335	274	342	228	394
<i>Neomys</i>	2	397	372	400	243	475
Crocidae*	13	234	207	238	181	262
<i>Crocida</i>	6	207	185	210	162	230
<i>Suncus</i>	2	188	189	189	157	200
<i>Sylvisorex</i>	2	328	275	333	237	374
AvIF of the 12 families and subfamilies marked by *		300	284	301	214	343

Number of species = N

Abbreviations:

- NEO total neocortex
- NG neocortical grey matter
- NG1 molecular layer of neocortical grey matter
- NG2-6 cell layers 2-6 of neocortical grey matter
- NW neocortical white matter

Table 64. Volumes of neocortical grey (NG), area striata grey (ASG), percentage of ASG relative to NG, and size indices of ASG related to the average of the four Tenrecinae (AvTn = 100)

ASG was estimated from GLD volumes (taken from Table 37) according to the interrelationship found in Primates: $\log \text{GLD} = -0.8801 + 0.8476 \times \log \text{ASG}$

	NG mm ³	ASG mm ³	<u>ASG</u> NG	ASG Index
				.55 = slope -.054 = y-intercept
Tenrecidae				
Tenrecinae				
<i>Tenrec ecaudatus</i>	237.00	42.04	17.7	116
<i>Setifer setosus</i>	118.90	25.07	21.1	140
<i>Hemicentetes semispin.</i>	70.77	8.37	11.8	69
<i>Echinops telfairi</i>	46.46	7.61	16.4	74
Erinaceidae				
Erinaceinae				
<i>Erinaceus europaeus</i>	456.43	126.18	27.6	350
Soricidae				
Soricinae				
<i>Sorex araneus</i>	25.60	1.16	4.5	37
<i>S. minutus</i>	14.75	0.81	5.5	41
Crocidurinae				
<i>Crocidura flavescens</i>	48.87	4.57	9.4	81
<i>C. russula</i>	23.56	1.16	4.9	35
<i>Suncus murinus</i>	45.90	3.02	6.6	49
Average values	N			
Insectivora (AvIS)	10		12.6	99
Tenrecinae*	4		16.8	100
Erinaceinae*	1		27.6	350
Soricinae*	2		5.0	39
Crocidurinae*	3		7.0	55
AvIF of the 4 subfamilies marked by *			14.1	136

Number of species = N

Abbreviations and notes:

- ASG area striata grey matter
- GLD dorsal part of lateral geniculate body
- NG total neocortex grey matter

Table 65. Volumes of diencephalon components based on data given by Bauchot (1979b)

For size indices in species see Table 66

For average size indices in supraspecific taxonomic units see Table 67

	DIE	ETH	THA	STH	HTH	RTH	PTE
Tenrecidae							
<i>Tenrec ecaudatus</i>	153.7	3.175	60.86	38.27	43.52	6.20	1.704
<i>Setifer setosus</i>	93.1	2.256	37.55	20.40	28.45	3.52	0.933
<i>Hemicentetes semispin.</i>	47.7	1.043	20.01	10.44	13.63	2.07	0.516
<i>Echinops telfairi</i>	40.3	0.878	17.10	8.69	11.01	2.07	0.532
<i>Microgale talazaci</i>	45.4	1.023	19.98	10.91	10.91	2.05	0.517
<i>Limnogale mergulus</i>	81.0	1.150	34.76	20.87	19.52	4.06	0.672
<i>Potamogale velox</i>	300.3	3.036	143.70	69.98	64.19	16.71	2.703
Chrysochloridae							
<i>Chrysochloris stuhlmanii</i>	51.6	1.164	20.55	16.88	10.58	2.02	0.373
Solenodontidae							
<i>Solenodon paradoxus</i>	272.3	5.630	111.16	85.99	56.97	10.30	2.225
Erinaceidae							
<i>Atelerix algirus</i>	251.0	4.236	98.41	67.15	69.53	9.09	2.544
<i>Erinaceus europaeus</i>	242.4	4.940	92.41	57.84	76.36	8.12	2.756
Talpidae							
<i>Desmana moschata</i>	343.8	4.126	125.89	126.91	70.90	13.03	2.914
<i>Galemys pyrenaicus</i>	104.5	1.532	41.67	35.14	21.46	3.93	0.732
<i>Talpa europaea</i>	80.7	1.366	29.58	27.83	18.51	2.87	0.534
Soricidae							
<i>Sorex araneus</i>	14.8	0.304	5.80	3.88	4.12	0.57	0.125
<i>S. minutus</i>	7.7	0.145	3.04	1.92	2.21	0.31	0.068
<i>Neomys fodiens</i>	24.1	0.444	9.67	7.51	5.27	0.98	0.197
<i>Crociodura flavescens</i>	25.1	0.514	10.92	6.02	6.15	1.20	0.285
<i>C. russula</i>	13.9	0.297	5.47	3.59	3.75	0.59	0.167
<i>Suncus murinus</i>	24.1	0.440	9.88	5.53	6.85	1.09	0.269

Abbreviations:

DIE	diencephalon
ETH	epithalamus
HTH	hypothalamus
PTE	regio prepectalis
RTH	nucleus reticularis
STH	subthalamus
THA	thalamus

Table 66. Size indices of diencephalon components related to the average of the four Tenrecinae (AvTn = 100) based on data given by Bauchot (1979b)

For volumes in species see Table 65

For average size indices in supraspecific taxonomic units see Table 67

	DIE	ETH	THA	STH	HTH	RTH	PTE
slope	.65	.63	.64	.71	.62	.63	.62
y-intercept	.350	-1.256	-.013	-.438	-.121	-.970	-1.531
Tenrecidae							
<i>Tenrec ecaudatus</i>	85	82	84	87	88	83	88
<i>Setifer setosus</i>	119	130	117	115	127	105	107
<i>Hemicentetes semispin.</i>	97	94	98	98	95	97	92
<i>Echinops telfairi</i>	98	95	101	100	91	115	113
<i>Microgale talazaci</i>	163	161	172	191	130	166	159
<i>Limnogale mergulus</i>	192	120	198	231	156	220	138
<i>Potamogale velox</i>	206	96	242	200	158	272	171
Chrysochloridae							
<i>Chrysochloris stuhlmanii</i>	209	205	199	336	141	183	128
Solenodontidae							
<i>Solenodon paradoxus</i>	177	168	178	232	133	159	133
Erinaceidae							
<i>Atelerix algirus</i>	154	119	148	170	153	133	144
<i>Erinaceus europaeus</i>	135	127	127	132	154	108	143
Talpidae							
<i>Desmana moschata</i>	292	160	263	460	214	262	226
<i>Galemys pyrenaicus</i>	327	210	313	528	225	279	197
<i>Talpa europaea</i>	205	153	181	334	159	166	118
Soricidae							
<i>Sorex araneus</i>	146	127	135	205	129	122	101
<i>S. minutus</i>	131	103	121	184	116	112	92
<i>Neomys fodiens</i>	183	144	174	297	128	165	123
<i>Crocidura flavescens</i>	125	110	129	150	100	133	119
<i>C. russula</i>	129	118	121	178	111	120	128
<i>Suncus murinus</i>	109	86	107	124	102	111	103

Abbreviations:

DIE	diencephalon		
ETH	epithalamus	RTH	nucleus reticularis
HTH	hypothalamus	STH	subthalamus
PTE	regio pretectalis	THA	thalamus

Table 67. Average size indices of diencephalon components related to the average of the four Tenrecinae (AvTn = 100) based on data given by Bauchot (1979b)

For volumes in species see Table 65

For size indices in species see Table 66

	N	DIE	ETH	THA	STH	HTH	RTH	PTE
slope		.65	.63	.64	.71	.62	.63	.62
y-intercept		.350	-1.256	-.013	-.438	-.121	-.970	-1.531
Insectivora (AvIS)	20	164	130	160	218	136	156	131
Tenrecidae	7	137	111	145	146	121	151	124
Tenrecinae*	4	100	100	100	100	100	100	100
Oryzorictinae*	2	177	140	185	211	143	193	149
Potamogalinae*	1	206	96	242	200	158	272	171
Chrysochloridae*	1	209	205	199	336	141	183	128
Solenodontidae*	1	177	168	178	232	133	159	133
Erinaceidae								
Erinaceinae*	2	144	123	138	151	154	120	144
Talpidae	3	275	174	253	441	199	236	180
Desmaninae*	2	310	185	288	494	219	270	212
Talpinae*	1	205	153	181	334	159	166	118
Soricidae	6	137	115	131	190	115	127	111
Soricinae*	3	153	124	143	228	125	133	105
<i>Sorex</i>	2	138	115	128	194	123	117	96
<i>Neomys</i>	1	183	144	174	297	128	165	123
Crocidurinae*	3	121	105	119	151	104	121	117
<i>Crocidura</i>	2	127	114	125	164	106	127	123
<i>Suncus</i>	1	109	86	107	124	102	111	103
AvIF of the 10 families and sub-families marked by *		180	140	177	244	144	172	138

Number of species = N

Abbreviations:

DIE	diencephalon
ETH	epithalamus
HTH	hypothalamus
PTE	regio prepectalis
RTH	nucleus reticularis
STH	subthalamus
THA	thalamus

Table 68. Volumes of thalamus components based on data given by Bauchot (1979b)

For size indices in species see Table 69

For average size indices in supraspecific taxonomic units see Table 70

	ANT	MLT	MNT	DOT	VET	CGM	CGL
Tenrecidae							
<i>Tenrec ecaudatus</i>	2.48	6.02	6.12	4.36	27.28	10.03	4.58
<i>Setifer setosus</i>	2.14	4.29	4.35	2.69	15.71	5.77	2.60
<i>Hemicentetes semispin.</i>	1.31	2.01	2.18	1.38	8.56	3.15	1.42
<i>Echinops telfairi</i>	1.05	1.49	1.82	0.94	7.56	2.78	1.47
<i>Microgale talazaci</i>	2.31	2.27	2.59	2.06	5.77	3.77	1.22
<i>Limnogale mergulus</i>	1.61	3.89	4.26	3.70	7.62	12.47	1.21
<i>Potamogale velox</i>	6.68	17.78	10.71	21.67	31.30	51.26	4.32
Chrysochloridae							
<i>Chrysochloris stuhlmanii</i>	1.36	2.60	3.56	2.72	4.86	4.95	0.50
Solenodontidae							
<i>Solenodon paradoxus</i>	7.00	13.03	19.58	12.50	39.54	14.54	4.98
Erinaceidae							
<i>Atelerix algirus</i>	6.48	15.29	11.40	11.99	34.61	12.72	5.93
<i>Erinaceus europaeus</i>	4.61	12.70	10.02	10.44	34.39	12.64	7.62
Talpidae							
<i>Desmana moschata</i>	5.87	17.75	17.37	19.80	23.03	37.72	4.34
<i>Galemys pyrenaicus</i>	2.66	6.40	6.49	7.59	6.61	10.82	1.10
<i>Talpa europaea</i>	1.92	3.92	4.76	3.68	7.22	7.36	0.72
Soricidae							
<i>Sorex araneus</i>	0.44	0.73	0.77	0.55	1.84	1.20	0.26
<i>S. minutus</i>	0.20	0.33	0.41	0.22	1.04	0.68	0.16
<i>Neomys fodiens</i>	0.61	1.01	1.25	0.97	2.08	3.40	0.35
<i>Crocidura flavescens</i>	0.80	1.30	1.25	1.13	3.50	2.28	0.65
<i>C. russula</i>	0.33	0.64	0.60	0.48	1.84	1.20	0.38
<i>Suncus murinus</i>	0.56	1.10	0.95	0.89	3.49	2.28	0.62

Abbreviations:

ANT	nuclei anteriores thalami
CGL	corpus geniculatum laterale
CGM	corpus geniculatum mediale
DOT	nuclei dorsales thalami
MLT	nuclei mediales thalami
MNT	nuclei mediani thalami
VET	nuclei ventrales thalami

Table 69. Size indices of thalamus components related to the average of the four Tenrecinae (AvTn = 100) based on data given by Bauchot (1979b)

For volumes in species see Table 68

For average size indices in supraspecific taxonomic units see Table 70

	ANT	MLT	MNT	DOT	VET	CGM	CGL
slope	.58	.62	.64	.68	.64	.72	.60
y-intercept	-1.123	-.964	-.978	-1.281	-.377	-.996	-1.046
Tenrecidae							
<i>Tenrec ecaudatus</i>	66	84	77	85	87	77	89
<i>Setifer setosus</i>	119	133	125	125	113	112	109
<i>Hemicentetes semispin.</i>	110	97	99	104	97	102	91
<i>Echinops telfairi</i>	104	86	99	86	103	110	111
<i>Microgale talazaci</i>	324	189	206	282	115	229	133
<i>Limnogale mergulus</i>	155	217	224	326	100	476	89
<i>Potamogale velox</i>	213	304	167	524	122	497	102
Chrysochloridae							
<i>Chrysochloris stuhlmanni</i>	212	242	318	421	109	343	61
Solenodontidae							
<i>Solenodon paradoxus</i>	213	212	289	285	146	133	111
Erinaceidae							
<i>Atelerix algirus</i>	187	235	158	257	121	109	125
<i>Erinaceus europaeus</i>	122	179	127	203	109	98	148
Talpidae							
<i>Desmana moschata</i>	227	374	334	600	111	465	125
<i>Galemys pyrenaicus</i>	330	466	451	899	115	564	105
<i>Talpa europaea</i>	198	234	270	351	102	305	57
Soricidae							
<i>Sorex araneus</i>	153	160	166	218	99	224	71
<i>S. minutus</i>	112	120	149	153	96	231	74
<i>Neomys fodiens</i>	168	171	207	290	86	473	76
<i>Crocidura flavescens</i>	150	147	137	218	96	199	95
<i>C. russula</i>	107	132	122	179	94	211	100
<i>Suncus murinus</i>	97	114	95	155	87	179	83

Abbreviations:

ANT	nuclei anteriores thalami
CGL	corpus geniculatum laterale
CGM	corpus geniculatum mediale
DOT	nuclei dorsales thalami
MLT	nuclei mediales thalami
MNT	nuclei mediani thalami
VET	nuclei ventrales thalami

Table 70. Average size indices of thalamus components related to the average of the four Tenrecinae (AvTn = 100) based on data given by Bauchot (1979b)

For volumes in species see Table 68

For size indices in species see Table 69

	N	ANT	MLT	MNT	DOT	VET	CGM	CGL
slope		.58	.62	.64	.68	.64	.72	.60
y-intercept		-1.123	-.964	-.978	-1.281	-.377	-.996	-1.046
Insectivora (AvIS)	20	168	195	191	288	105	257	98
Tenrecidae	7	156	159	142	219	105	229	103
Tenrecinae*	4	100	100	100	100	100	100	100
Oryzorictinae*	2	239	203	215	304	108	353	111
Potamogalinae*	1	213	304	167	524	122	497	102
Chrysochloridae*	1	212	242	318	421	109	343	61
Solenodontidae*	1	213	212	289	285	146	133	111
Erinaceidae								
Erinaceinae*	2	155	207	143	230	115	103	137
Talpidae	3	252	358	351	617	109	445	95
Desmaninae*	2	279	420	392	749	113	515	115
Talpinae*	1	198	234	270	351	102	305	57
Soricidae	6	131	141	146	202	93	253	83
Soricinae*	3	144	150	174	220	94	309	74
<i>Sorex</i>	2	132	140	157	185	97	228	73
<i>Neomys</i>	1	168	171	207	290	86	473	76
Crocidurinae*	3	118	131	118	184	92	196	93
<i>Crocidura</i>	2	129	140	130	198	95	205	98
<i>Suncus</i>	1	97	114	95	155	87	179	83
AvIF of the 10 families and sub-families marked by *		187	220	219	337	110	285	96

Abbreviations:

ANT	nuclei anteriores thalami
CGL	corpus geniculatum laterale
CGM	corpus geniculatum mediale
DOT	nuclei dorsales thalami
MLT	nuclei mediales thalami
MNT	nuclei mediani thalami
VET	nuclei ventrales thalami

Table 71. Area measurements of commissures and chiasma opticum in mm²

For size indices in species see Table 72

For average size indices in supraspecific taxonomic units see Table 73

	n	CCA	COA	CHO
Tenrecidae				
<i>Tenrec ecaudatus</i>	1	0.859	1.025	0.209
<i>Setifer setosus</i>	1	0.691	0.734	0.092
<i>Hemicentetes semispin.</i>	1	0.542	0.412	0.046
<i>Echinops telfairi</i>	1	0.230	0.384	0.121
<i>Geogale aurita</i>	1	0.068	0.139	0.022
<i>Oryzorictes talpoides</i>	1	0.646	0.284	0.017
<i>Microgale dobsoni</i>	1	0.689	0.300	0.029
<i>M. talazaci</i>	1	0.890	0.358	0.043
<i>Limnogale mergulus</i>	1	1.649	0.238	0.051
<i>Potamogale velox</i>	1	6.241	0.392	0.129
<i>Micropotamog. ruwenzorii</i>	1	1.098	0.156	0.038
Chrysochloridae				
<i>Chrysochloris stuhlmanni</i>	1	0.754	0.362	0.011
Solenodontidae				
<i>Solenodon paradoxus</i>	1	2.730	1.175	0.094
Erinaceidae				
<i>Echinosorex gymnurus</i>	1	7.016	1.347	0.407
<i>Hylomys suillus</i>	1	1.441	0.278	0.308
<i>Atelerix algirus</i>	2	2.535	0.967	0.397
<i>Erinaceus europaeus</i>	1	1.480	0.554	0.235
<i>Hemiechinus auritus</i>	3	1.574	0.541	0.236
Talpidae				
<i>Desmana moschata</i>	1	5.059	0.466	0.071
<i>Galemys pyrenaicus</i>	1	2.055	0.183	0.042
<i>Talpa europaea</i>	1	1.134	0.377	0.010
<i>Parascalops breweri</i>	1	0.739	0.292	0.011
<i>Scalopus aquaticus</i>	1	1.859	0.424	-,-

Table 71 (cont.)

	n	CCA	COA	CHO
Soricidae				
<i>Sorex alpinus</i>	1	0.379	0.158	0.016
<i>S. araneus</i>	1	0.312	0.146	0.034
<i>S. cinereus</i>	1	0.312	0.118	0.014
<i>S. fumeus</i>	1	0.543	0.129	0.020
<i>S. minutus</i>	1	0.145	0.048	0.015
<i>Microsorex hoyi</i>	1	0.194	0.081	0.009
<i>Neomys anomalus</i>	1	0.338	0.083	0.019
<i>N. fodiens</i>	1	0.376	0.095	0.021
<i>Blarina brevicauda</i>	1	0.534	0.205	0.023
<i>Cryptotis parva</i>	1	0.336	0.139	-.-
<i>Anourosorex squamipes</i>	1	0.327	0.181	0.008
<i>Crociodura flavescens</i>	4	0.362	0.213	0.039
<i>C. russula</i>	1	0.242	0.158	0.018
<i>Suncus etruscus</i>	1	0.095	0.057	0.008
<i>S. murinus</i>	1	0.440	0.183	0.033
<i>Scutisorex somereni</i>	1	0.556	0.315	0.033
<i>Sylvisorex granti</i>	1	0.209	0.096	0.012
<i>S. megalura</i>	1	0.169	0.122	0.018
<i>Ruwenzorisorex suncoides</i>	1	0.494	0.170	0.020
<i>Myosorex babaulti</i>	1	0.622	0.294	0.030

Number of individuals = n

Abbreviations:

CCA corpus callosum area
 CHO chiasma opticum
 COA commissura anterior

Table 72. Size indices of commissural areas and chiasma opticum area related to the average of the four Tenrecinae (AvTn = 100)

For area measurements in species see Table 71

For average size indices in supraspecific taxonomic units see Table 73

	n	CCA	COA	CHO
slope		.52	.44	.37
y-intercept		-1.474	-1.253	-1.821
Tenrecidae				
<i>Tenrec ecaudatus</i>	1	77	94	114
<i>Setifer setosus</i>	1	120	119	81
<i>Hemicentetes semispin.</i>	1	136	91	52
<i>Echinops telfairi</i>	1	67	96	153
<i>Geogale aurita</i>	1	61	90	61
<i>Oryzorictes talpoides</i>	1	268	96	27
<i>Microgale dobsoni</i>	1	339	117	53
<i>M. talazaci</i>	1	353	116	67
<i>Limnogale mergulus</i>	1	468	58	63
<i>Potamogale velox</i>	1	658	42	79
<i>Micropotamog. ruwenzorii</i>	1	303	37	46
Chrysochloridae				
<i>Chrysochloris stuhlmanni</i>	1	329	128	19
Solenodontidae				
<i>Solenodon paradoxus</i>	1	275	120	56
Erinaceidae				
<i>Echinorex gymnurus</i>	1	637	126	225
<i>Hylomys suillus</i>	1	524	84	457
<i>Atelerix algirus</i>	2	244	95	229
<i>Erinaceus europaeus</i>	1	132	51	129
<i>Hemiechinus auritus</i>	3	274	88	207
Talpidae				
<i>Desmana moschata</i>	1	634	57	49
<i>Galemys pyrenaicus</i>	1	730	54	61
<i>Talpa europaea</i>	1	341	97	13
<i>Parascalops breweri</i>	1	277	91	16
<i>Scalopus aquaticus</i>	1	470	94	--

Table 72 (cont.)

	n	CCA	COA	CHO
Soricidae				
<i>Sorex alpinus</i>	1	324	99	44
<i>S. araneus</i>	1	278	94	94
<i>S. cinereus</i>	1	396	102	51
<i>S. fumeus</i>	1	535	91	59
<i>S. minutus</i>	1	200	45	56
<i>Microsorex hoyi</i>	1	349	94	43
<i>Neomys anomalus</i>	1	281	50	49
<i>N. fodiens</i>	1	271	51	51
<i>Blarina brevicauda</i>	1	338	99	51
<i>Cryptotis parva</i>	1	304	91	--
<i>Anourosorex squamipes</i>	1	205	87	17
<i>Crociodura flavescens</i>	4	186	86	75
<i>C. russula</i>	1	206	98	49
<i>Suncus etruscus</i>	1	204	77	43
<i>S. murinus</i>	1	210	70	59
<i>Scutisorex somereni</i>	1	191	91	46
<i>Sylvisorex granti</i>	1	307	95	50
<i>S. megalura</i>	1	208	103	62
<i>Ruwenzorisorex suncoides</i>	1	325	85	46
<i>Myosorex babaulti</i>	1	425	151	68

Number of individuals = n

Abbreviations:

CCA corpus callosum area
CHO chiasma opticum
COA commissura anterior

Table 73. Average size indices of commissural and chiasma opticum areas related to the average of the four Tenrecinae (AvTn = 100)

For area measurements in species see Table 71

For size indices in species see Table 72

	N	CCA	COA	CHO
slope		.52	.44	.37
y-intercept		-1.474	-1.253	-1.821
Insectivora (AvIS)	43 §)	313	88	80
Tenrecidae	11	259	87	72
Tenrecinae*	4	100	100	100
Geogalinae*	1	61	90	61
Oryzorictinae*	4	357	97	53
<i>Microgale</i>	2	346	117	60
terr. Oryzorictinae	3	320	110	49
<i>Limnogale</i>	1	468	58	63
Potamogalinae*	2	480	39	63
Chrysochloridae*	1	329	128	19
Solenodontidae*	1	275	120	56
Erinaceidae	5	362	89	249
Echinosoricinae*	2	581	105	341
Erinaceinae*	3	217	78	188
Talpidae	5 +)	490	79	49
Desmaninae*	2	682	56	55
Talpinae*	3 +)	363	94	15
Soricidae	20 +)	287	88	53
Soricinae*	11 +)	316	82	52
<i>Sorex</i>	5	347	86	61
terr. Soricinae	9 +)	325	89	52
<i>Neomys</i>	2	276	51	50
Crocidurinae*	9	251	95	55
<i>Crocidura</i>	2	196	92	62
<i>Suncus</i>	2	207	74	51
<i>Sylvisorex</i>	2	257	99	56
AvIF of the 12 families and subfamilies marked by *		334	90	91

Number of species = N +) for CHO = N - 1 §) for CHO = N - 2

Abbreviations:

CCA corpus callosum area
 CHO chiasma opticum
 COA commissura anterior

Table 74. Size characteristics of the various brain structures in the average Insectivora relative to the average of the four Tenrecinae (AvTn = 100)

	N	Min	Max	<u>Max</u> Min	AvIS	CV	NN	AvIF	CV
NET	50	83	258	3.1	154	23.6	12	160	27.6
OBL	50	86	240	2.8	132	25.9	12	139	24.9
MES	50	88	266	3.0	136	29.1	12	150	29.8
CER	50	73	319	4.4	167	32.6	12	179	36.7
DIE	50	79	327	4.1	169	29.3	12	180	34.7
TEL	50	66	260	3.9	159	25.6	12	161	30.0
OBL	50	86	240	2.8	132	25.9	12	139	24.9
OBL limited	30	86	240	2.8	133	28.0	11	140	27.8
TR	30	81	342	4.2	159	42.7	11	165	45.7
TSO	30	84	234	2.8	151	27.3	11	156	25.8
FUN									
FGR	30	67	207	3.1	131	29.3	11	143	25.2
FCM	30	57	320	5.6	118	49.1	11	147	49.4
FCE	30	23	245	10.7	111	51.4	11	127	38.1
RLN									
REL	30	72	317	4.4	170	33.4	11	182	27.8
INO	30	79	326	4.1	185	30.9	11	205	27.3
PRP	30	83	219	2.6	136	27.5	11	145	26.2
COE	29	63	240	3.8	120	33.7	11	128	23.2
MOT									
V	30	70	203	2.9	121	29.1	11	133	24.9
VII	30	76	282	3.7	127	34.3	11	133	34.4
X	30	88	238	2.7	140	24.5	11	138	17.3
XII	30	72	183	2.5	116	25.4	11	116	25.9
VC									
VM	30	82	281	3.4	141	36.2	11	152	35.7
VI	30	69	249	3.6	129	40.3	11	142	38.4
VL	30	64	284	4.4	119	41.2	11	128	41.1
VS	30	54	227	4.2	104	41.6	11	116	39.0
AUD									
DCO	30	43	303	7.0	107	46.4	11	113	44.2
VCO	30	38	177	4.7	90	41.1	11	95	38.8
OLS	30	40	223	5.6	104	39.5	11	112	34.5

Table 74 (cont.)

	N	Min	Max	<u>Max</u> Min	AvIS	CV	NN	AvIF	CV
MES	50	88	266	3.0	136	29.1	12	150	29.8
MES limited	10	88	125	1.4	103	10.6	4	102	8.3
MTG	10	89	126	1.4	108	11.0	4	110	7.1
NLL	10	54	137	2.5	98	29.4	4	90	29.7
MTC	10	77	131	1.7	97	18.6	4	94	15.1
SUC	10	79	151	1.9	121	19.0	4	123	15.4
INC	10	44	136	3.1	83	33.1	4	76	31.3
SCB	23	23	857	37.3	203	104.7	10	260	86.5
CER	50	73	319	4.4	167	32.6	12	179	36.7
CER limited	10	78	146	1.9	113	18.1	4	115	10.2
TCN	10	78	125	1.6	107	14.4	4	112	10.6
MCN	10	81	160	2.0	117	24.5	4	126	23.1
ICN	10	73	128	1.8	94	17.7	4	97	15.8
LCN	10	66	148	2.2	119	21.8	4	122	14.3
DIE	50	79	327	4.1	169	29.3	12	180	34.7
DIE limited	10	85	146	1.7	117	16.6	4	124	14.0
ETH	10	78	149	1.9	111	17.5	4	111	10.9
HAM	38	77	245	3.2	153	29.2	11	162	24.7
EPI	38	0	203	∞	50	82.5	11	62	85.6
THA	10	87	163	1.9	127	21.4	4	136	18.6
CGM	10	91	142	1.6	116	14.5	4	120	11.4
CGL	10	68	226	3.3	103	45.8	4	122	57.6
GLD	10	62	245	4.0	104	52.4	4	125	64.6
GLV	10	72	185	2.6	102	33.2	4	115	41.0
STH	10	91	204	2.2	127	30.3	4	132	33.3
PALL	10	92	199	2.2	125	29.0	4	130	31.8
LUY	10	62	297	4.8	160	49.6	4	167	50.1
HTH	10	82	123	1.5	106	12.7	4	110	10.5
CAI	10	81	163	2.0	113	24.3	4	115	10.1
TRO	10	66	172	2.6	97	32.8	4	110	40.0

Number of species = N

Number of subfamilies and families = NN

Abbreviations and notes:

AvIF average of families and subfamilies

AvIS average of species

CV coefficient of variation (standard deviation of the mean in percent of the mean)

Max maximum index in species

Min minimum index in species

Max/Min quotient of maximum through minimum division

Table 74 (cont.)

	N	Min	Max	<u>Max</u> Min	AvIS	CV	NN	AvIF	CV
TEL	50	66	260	3.9	159	25.6	12	161	30.0
TEL components									
BOL	50	29	151	5.2	88	30.1	12	91	33.9
PAL	50	36	166	4.6	103	29.7	12	104	33.6
STR	50	66	391	5.9	207	35.4	12	210	39.7
SEP	50	67	223	3.3	132	23.0	12	139	25.0
AMY	50	61	194	3.2	114	25.8	12	122	28.5
HIP	50	77	348	4.5	175	32.1	12	186	37.0
SCH	50	40	336	8.4	168	34.1	12	178	40.1
NEO	50	61	772	12.7	292	43.9	12	300	51.4
 MOB	50	28	152	5.4	87	30.1	12	91	34.3
L1+2	43	25	197	7.9	82	38.5	12	90	47.9
L3	43	31	170	5.5	97	32.8	12	94	27.8
L4-6	43	29	147	5.1	92	31.5	12	92	27.5
PvZ	43	6	226	37.7	59	104.2	12	80	91.5
 AOB	50	6	810	135	138	87.9	12	158	91.9
AOB limited	41	28	810	28.9	150	86.8	12	163	87.9
L1+2	41	13	636	48.9	112	96.1	12	123	99.1
L3-5	41	39	766	19.6	177	69.2	12	186	66.9
L6	41	20	1587	79.4	211	126.0	12	246	122.8
 PAL	50	36	166	4.6	103	29.7	12	104	33.6
PAL components									
RB	50	37	171	4.6	107	29.5	12	106	34.3
PRPI	50	35	159	4.5	94	31.9	12	98	35.3
TOL	50	33	212	6.4	104	35.8	12	114	44.4
TRL	50	34	172	5.1	84	35.3	12	92	35.5
COA	50	33	198	6.0	95	40.8	12	94	45.1
SIN	50	45	154	3.4	96	27.1	12	99	22.3
 SEP	50	67	223	3.3	132	23.0	12	139	25.0
SEP components									
SET	38	75	258	3.4	162	28.4	11	166	26.2
SFB	38	25	189	7.6	75	45.6	11	83	35.4

Table 74 (cont.)

	N	Min	Max	<u>Max</u> Min	AvIS	CV	NN	AvIF	CV
TEL (cont.)									
AMY	50	61	194	3.2	114	25.8	12	122	28.5
AMY components									
MAM	50	64	191	3.0	114	24.3	12	117	24.7
NTO	38	4	204	51.0	92	47.4	10	82	45.5
LAM	50	58	204	3.5	115	29.2	12	127	33.0
MCB	38	40	242	6.1	92	40.9	10	105	38.8
 HIP	50	77	348	4.5	175	32.1	12	186	37.0
HIP limited	10	82	159	1.9	122	19.2	4	128	16.3
 HR	10	82	159	1.9	122	19.3	4	128	16.3
HS	10	49	214	4.4	137	34.5	4	140	23.0
HP	10	65	158	2.4	117	22.5	4	121	12.3
 NEO	50	61	772	12.7	292	43.9	12	300	51.4
NEO components									
NW	50	55	796	14.5	266	50.9	12	284	62.2
NG	50	61	772	12.7	296	43.8	12	301	50.9
NG1	50	45	398	8.8	209	33.2	12	214	38.8
NG2-6	50	68	954	14.0	335	48.1	12	343	55.5
ASG	10	35	350	10.0	99	95.5	4	136	106.6

Bauchot's data of the diencephalic components (converted)

ETH	20	82	210	2.6	130	28.1	10	140	26.6
THA	20	84	313	3.7	160	37.4	10	177	32.1
STH	20	87	528	6.1	218	55.0	10	244	47.5
HTH	20	88	225	2.6	136	27.2	10	144	23.4
RTH	20	83	279	3.4	156	38.4	10	172	34.9
PTE	20	88	226	2.6	131	27.1	10	138	24.5

Bauchot's data of thalamic structures (converted)

ANT	20	66	330	5.0	168	42.3	10	187	30.1
MLT	20	84	466	5.5	195	50.2	10	220	41.6
MNT	20	77	451	5.9	191	51.1	10	219	43.7
DOT	20	85	899	10.6	288	69.3	10	337	56.0
VET	20	86	146	1.7	105	13.7	10	110	14.3
CGM	20	77	564	7.3	257	61.6	10	285	52.9
CGL	20	57	148	2.6	98	24.6	10	96	26.4

11 Atlas of the Brain of *Atelerix algirus*

The brains for this atlas were obtained through the cooperation with
 Dr. José Regidor
 from the Anatomical Institute of the University of Las Palmas, Gran Canaria,
 who made available three specimens of *Atelerix algirus* and all laboratory
 facilities of his Institute to allow for the stereotaxic procedures.

Zero coordinates

The zero planes, based on anatomical landmarks of the head, and intersecting each other at right angles, were defined as follows:

1. The horizontal plane (H 0) passes through the centers of the external auditory meatus and the lower margins of the upper jaws between I¹ and I².
2. The anterior-posterior plane (AP 0) lies perpendicular to the horizontal plane and also passes through the centers of the external auditory meatus.
3. The left-right zero plane (LR 0) is the median sagittal plane.

Reference coordinates

Under deep Nembutal anesthesia the hedgehogs (*Atelerix algirus*) were perfused intracardially with 250 ml of 0.9% NaCl followed by 500 ml of Bouin's fluid as fixative. Then the heads were fixed in a stereotaxic headholder for rats (David Kopf Co.). Five reference needles were inserted into the intracranial brain by aid of the stereotaxic instrument in order (1) to orientate the brain for sectioning in one of the three fundamental planes (transversal, sagittal, horizontal), (2) to identify the position of each section and (3) to estimate the shrinkage during the embedding, cutting and staining procedures. The needles have the following coordinates and functions:


1. A pair of needles was inserted perpendicular to the horizontal plane at A 5.0 in left-right positions of L 6.0 and R 6.0. The distance of these vertical needles from the median sagittal plane in the sections indicate the degree of shrinkage in the left-right direction.

2. A pair of needles was inserted parallel to the horizontal plane in a distance of H +10.0 at AP 0 and A 15.0. These needles serve to determine the average thickness of the transversal section when related to the fresh brain. In the atlas brain (4989) the number of sections between A 15.0 and AP 0 (that is in a distance of 15 000 μm) is 622 instead of the expected 750 20 μm sections. Thus one section corresponds to 24.12 μm in the fresh brain, a value which is used to calculate the position of each transversal section in the AP direction.

3. A horizontal needle was inserted in a left position of L 3.0 at H +7.5 parallel to sagittal and horizontal planes. This longitudinal needle is used to identify each point in the transversal section according to its H level. The distance from the median sagittal plane is an additional criterion for the shrinkage in the left-right direction.


Histological procedures

Following the stereotaxic procedures the brains were removed and kept for four days in the fixative. At the beginning of the washing procedure the brains were photographed. Washing (4-6 weeks) was performed in an ascending series of ethanols. After being passed through methylbenzoate and benzene the brains were embedded in paraffine, orientated for sectioning by aid of the needles and cut in the transversal plan (brains 4989, 4990), sagittal plane (left hemisphere of 4988) and horizontal plane (right hemisphere of brain 4988). The serial sections were stained for cells with gallocyanine. Sagittal and horizontal sections were additionally used to determine the structural borders given in the atlas (transversal sections only).



A 12.5

A 12.5




AOB accessory olfactory bulb
MOB main olfactory bulb
PRPI prepiriform region

RB retrobulbar region
TRL lateral olfactory tract



A 10




COA	anterior commissure	PRPI	prepiriform region
HP	hipp. precommissuralis	SEP	telencephalic septum
NEO	neocortex	STR	striatum
NW	neocortical white matter	TOL	olfactory tubercle
N V	trigeminal nerve	TRL	lateral olfactory tract



A 9

A 9




COA anterior commissure
DIAG diagonal band of Broca
HP hipp. precommissuralis
NEO neocortex
NW neocortical white matter
N II optic nerve

N V trigeminal nerve
PRPI prepiriform region
SEP telencephalic septum
STR striatum
TOL olfactory tubercle
TRL lateral olfactory tract



A 8


A 8




CC	corpus callosum	POL	lateral preoptic area
COA	anterior commissure	POM	medial preoptic area
DIAG	diagonal band of Broca	PRPI	prepiriform region
HS	hipp. supracommissuralis	SEP	telencephalic septum
NEO	neocortex	SIN	substantia innominata
NW	neocortical white matter	STR	striatum
N II	optic nerve	TOL	olfactory tubercle
N V	trigeminal nerve	TRL	lateral olfactory tract




A 7



CC	corpus callosum	PALL	pallidum, globus pallidus
DIAG	diagonal band of Broca	POL	lateral preoptic area
F	fimbria-fornix complex	POM	medial preoptic area
HS	hipp. supracommissuralis	PRPI	prepiriform region
MAM	medial amygdaloid division	SEP	telencephalic septum
NEO	neocortex	SES	stria terminalis nuclei
NTO	nucleus of the lateral	SFB	subfornical body
	olfactory tract	SIN	substantia innominata
NW	neocortical white matter	SON	supraoptic nucleus
N II	optic nerve	STR	striatum
N V	trigeminal nerve		




A 6





A 5

A 5




AD	ncl. anterodorsalis thalami
CAI	capsula interna
CMT	ncl. centromedialis thalami
GLD	dorsal nucleus of CGL
GLV	ventral nucleus of CGL
HR	hipp. retrocommissuralis
HTL	lateral hypothalamus
LAM	lateral amygdaloid division
LD	ncl. laterodorsalis thalami
MAM	medial amygdaloid division
MD	ncl. med. thalami dorsalis
NEO	neocortex

NW	neocortical white matter
N V	trigeminal nerve
PRPI	prepiriform region
PT	ncl. paratenialis thalami
RE	nucleus reuniens thalami
RTH	nuclei reticulares thalami
SCH	schizocortex
TRO	optic tract
VLT	ncl. ventrolateralis thalami
VMH	ncl. ventromed. hypothalami
ZI	zona incerta



A 4




CLT	ncl. centrolateralis thalami
CMT	ncl. centromedialis thalami
DMH	ncl. dorsomed. hypothalami
GLD	dorsal nucleus of CGL
GLV	ventral nucleus of CGL
HAL	ncl. habenularis lateralis
HAM	ncl. habenularis medialis
HR	hipp. retrocommissuralis
HTL	hypothalamus lateralis
IMD	ncl. intermediadors. thalami
INF	nucleus infundibularis
LAM	lateral amygdaloid division
LD	ncl. laterodorsalis thalami
MAM	medial amygdaloid division

MD	ncl. med. thalami dorsalis
NEO	neocortex
NW	neocortical white matter
N V	trigeminal nerve
PRPI	prepiriform region
RE	nucleus reuniens thalami
RH	ncl. rhomboides thalami
RTH	nuclei reticulares thalami
SCH	schizocortex
VLT	ncl. ventrolateralis thalami
VMH	ncl. ventromed. hypothalami
VMT	ncl. ventromed. thalami
ZI	zona incerta




A 3.5




CLT	ncl. centrolateralis thalami	LUY	nucleus subthalamicus Luysi
CMT	ncl. centromedialis thalami	MD	ncl. med. thalami dorsalis
GLD	dorsal nucleus of CGL	NEO	neocortex
GLV	ventral nucleus of CGL	NW	neocortical white matter
HAL	ncl. habenularis lateralis	N V	trigeminal nerve
HAM	ncl. habenularis medialis	PRPI	prepiriform region
HR	hipp. retrocommissuralis	RE	nucleus reuniens thalami
HTL	hypothalamus lateralis	RTH	nuclei reticulares thalami
HTP	area hypothalamica posterior	SCH	schizocortex
IMD	ncl. intermediodors. thalami	VLT	ncl. ventrolateralis thalami
LAM	lateral amygdaloid division	VMT	ncl. ventromed. thalami
LD	ncl. laterodorsalis thalami	ZI	zona incerta




A 3



AVT	area ventralis tegmenti	LP	ncl. lateralis post. thalami
CM	corpus mamillare	MD	ncl. med. thalami dorsalis
CMT	ncl. centromedialis thalami	NEO	neocortex
GLD	dorsal nucleus of CGL	NW	neocortical white matter
GLV	ventral nucleus of CGL	N V	trigeminal nerve
HAL	ncl. habenularis lateralis	PC	pedunculus cerebri
HAM	ncl. habenularis medialis	PRPI	prepiriform region
HR	hipp. retrocommissuralis	RE	nucleus reuniens thalami
HTL	hypothalamus lateralis	RTH	nuclei reticulares thalami
HTP	area hypothalamica posterior	SCH	schizocortex
HYP	hypophysis	SN	substantia nigra
IMD	ncl. intermediadors. thalami	VP	thalamus ventroposterior
LAM	lateral amygdaloid division	ZI	zona incerta



A 2




APT	area pretectalis
AVT	area ventralis tegmenti
CA 1	area 1 of hippocampus (HR)
CA 3	area 3 of hippocampus (HR)
CGM	corpus geniculatum mediale
EPI	epiphysis, pineal body
FD	fascia dentata of HR
GC	griseum centrale
HR	hipp. retrocommissuralis
HYP	hypophysis
LAM	lateral amygdaloid division

LP	ncl. lateralis post. thalami
NEO	neocortex
NW	neocortical white matter
N V	trigeminal nerve
PC	pedunculus cerebri
RUB	nucleus ruber
SCH	schizocortex
SN	substantia nigra
SUB	subiculum (hippocampi)
SUC	colliculus superior
THP	thalamus posterior



A 1




AVT	area ventralis tegmenti
BCI	brachium colliculi inferioris
FD	fascia dentata of HR
FRTM	formatio reticularis tegmenti mesencephali
GC	griseum centrale
HR	hipp. retrocommissuralis
HYP	hypophysis

NEO	neocortex
N V	trigeminal nerve
PC	pedunculus cerebri
RUB	nucleus ruber
SCH	schizocortex
SN	substantia nigra
SUC	colliculus superior




AP 0




CU	nucleus cuneiformis
FR	formatio reticularis
FRTM	formatio reticularis tegmenti mesencephali
GC	griseum centrale
HYP	hypophysis
INC	colliculus inferior

IP	nucleus interpeduncularis
NEO	neocortex
N V	trigeminal nerve
RUB	nucleus ruber
SCH	schizocortex
SUC	colliculus superior
III	nucleus oculomotorius



P 1

P 1




CU	nucleus cuneiformis
DR	nucleus dorsalis raphe
FR	formatio reticularis
GC	griseum centrale
INC	colliculus inferior

NLL	nuclei of lateral lemniscus
N V	trigeminal nerve
SUC	colliculus superior
VPO	ventral pons




P 2

P 2




CER	cerebellum
FR	formatio reticularis
GC	griseum centrale
INC	colliculus inferior
NLL	nuclei of lateral lemniscus

OLS	superior olive
TR	complexus sensorius nervi trigemini
VPO	ventral pons
VT	ncl. ventralis tegmenti




P 2.5

P 2.5




CER	cerebellum
COE	nucleus locus coeruleus
FR	formatio reticularis
GC	griseum centrale
INC	colliculus inferior
OLS	superior olive
RA	nucleus raphes

TR	complexus sensorius nervi trigemini
TRM	nucleus mesencephalicus nervi trigemini
VCO	ncl. cochlearis ventralis
V	nucleus motorius nervi trigemini



P 3

P 3




CER	cerebellum
FR	formatio reticularis
GC	griseum centrale
N VII	nervus facialis
OLS	superior olive
RA	nucleus raphes
TR	complexus sensorius nervi trigemini

TRM	nucleus mesencephalicus nervi trigemini
VCO	ncl. cochlearis ventralis
VS	ncl. vestibularis superior
V	nucleus motorius nervi trigemini



P 3.5




CER	cerebellum
FR	formatio reticularis
GNF	genu nervi facialis
ICN	nucleus interpositus
	cerebelli
LCN	nucleus lateralis cerebelli
MCN	nucleus medialis cerebelli
N VII	nervus facialis

RA	nucleus raphes
TR	complexus sensorius nervi
	trigemini
VCO	ncl. cochlearis ventralis
VL	ncl. vestibularis lateralis
VM	ncl. vestibularis medialis
VS	ncl. vestibularis superior



P 4

P 4




CER	cerebellum
DCO	ncl. cochlearis dorsalis
FR	formatio reticularis
GNF	genu nervi facialis
ICN	nucleus interpositus
	cerebelli
LCN	nucleus lateralis cerebelli
MCN	nucleus medialis cerebelli
PRP	ncl. prepositus hypoglossi

RP	ncl. reticularis paramediani
TR	complexus sensorius nervi trigemini
VCO	ncl. cochlearis ventralis
VL	ncl. vestibularis lateralis
VM	ncl. vestibularis medialis
VI	nucleus nervi abducentis
VII	nucleus nervi facialis




P 5

P 5




CER	cerebellum
FR	formatio reticularis
ICN	nucleus interpositus cerebelli
MCN	nucleus medialis cerebelli
PRP	ncl. prepositus hypoglossi

TR	complexus sensorius nervi trigemini
TSO	nucleus tractus solitarii
VI	ncl. vestibularis inferior
VM	ncl. vestibularis medialis




P 6

P 6




CER	cerebellum
FCE	ncl. fascicularis cuneatus externus
FR	formatio reticularis
INO	inferior olive
PRP	ncl. prepositus hypoglossi

TR	complexus sensorius nervi trigemini
TSO	nucleus tractus solitarii
VI	ncl. vestibularis inferior
VM	ncl. vestibularis medialis
X	ncl. dorsalis motorius nervi vagi




P 7

P 7




CER	cerebellum
FCE	ncl. fascicularis cuneatus externus
FCM	ncl. fascicularis cuneatus medialis
FR	formatio reticularis
INO	inferior olive

REL	ncl. reticularis lateralis
TR	complexus sensorius nervi trigemini
TSO	nucleus tractus solitarii
X	ncl. dorsalis motorius nervi vagi
XII	nucleus nervi hypoglossi




P 8

P 8




AP	area postrema
CER	cerebellum
FCM	ncl. fascicularis cuneatus medialis
FGR	ncl. fascicularis gracilis
FR	formatio reticularis
INO	inferior olive



REL	ncl. reticularis lateralis
TR	complexus sensorius nervi trigemini
TSO	nucleus tractus solitarii
X	ncl. dorsalis motorius nervi vagi
XII	nucleus nervi hypoglossi



P 9



P 10



FCM	ncl. fascicularis cuneatus medialis	RMO	nucleus reticularis medullae oblongatae
FGR	ncl. fascicularis gracilis	TR	complexus sensorius nervi
RC	nucleus residualis cornus ventralis		trigemini

12 Subject Index

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

Accessory olfactory bulb (AOB)	103, 106, 107, 106-109, 150,
and components	151, 297, 322 - 324, 463 - 467,
	A 506 - 507
accessory olfactory system	322 - 324
adaptive radiation	298 - 311
advanced characters	6
allocortex (ALLO)	110
allometric methods	20, 21
ammons horn	see cornu ammonis
amygdala (AMY)	100, 122 - 127, 123, 126, 150,
	151, 153, 154, 297, 453 - 457
ancestral characters	6, 21, 22
<i>Anourosorex squamipes</i>	
brain and body weights	256
summary of brain characteristics	285
anterior amygdaloid area (AA)	122, 123
anterior commissure (COA)	111, 115, 116, 120, 151, 152,
	468 - 472, 493 - 497, A 508 - 513
anterior hippocampal commissure	120
anterior hypothalamic area (HTA)	A 516 - 517
apomorphic characters	6
archicortex (ARCH)	110
area hypothalamica posterior (HTP)	A 522 - 525
area measurements on brain parts in	
Chrysochloridae	187
Crocidae	273
Desmaninae	230
Erinaceinae	203 - 205
Oryzictinae	170
Potamogalinae	173
Solenodontidae	194
Soricinae	258 - 260
Talpinae	236, 237
Tenrecinae	162
area postrema (AP)	A 550 - 551
area preoptica (PO)	120, see also lateral preoptic area,
	medial preoptic area
area pretectalis (APT)	A 526 - 527

- area striata (AS) 136, 137, **156, 486**
- area ventralis tegmenti (AVT) **A 524 - 529**
- Atelerix algirus*
 - brain and body weights 202
 - summary of brain characteristics 227
 - brain atlas 503 - 553
- auditory nuclei (AUD) 75 - 78, **140, 141, 142**, 144, **433 - 435**
- auditory system 330 - 333
- average indices in Insectivora
 - families and subfamilies (AvIF) 22, 159
 - species (AvIS) 22, 159
 - Tenrecinae (AvTn) 22, 159
- Basal amygdaloid nucleus (BA) 123**
- basal Insectivora 17, 21
- best fit lines 20 - 23
- Blarina brevicauda*
 - brain and body weights 256
 - macromorphology of brain **36, 39, 42, 47**
 - summary of brain characteristics 284
- Blarini 307
- Bouin's fluid 12, 13, 15
- brachium colliculi inferioris (BCI) **A 528 - 529**
- brain weight, encephalization in
 - Chrysochloridae 185, 186
 - Crocidae 270 - 272
 - Desmaninae 229
 - Echinosoricinae 198
 - Erinaceinae 201, 202
 - Geogalinae 167
 - Oryzorictinae 169
 - Potamogalinae 172
 - Solenodontidae 193
 - Soricinae 255 - 257
 - Talpinae 234, 235
 - Tenrecinae 161
- bulbus olfactorius, bulbi olfactorii see olfactory bulbs, main olfactory bulb
- bulbus olfactorius accessorius see accessory olfactory bulb
- Capsula interna (CAI) 89, 90, 146, 442 - 444, A 516 - 519**
- caudatum see striatum
- cell measurements in
 - Chrysochloridae 187
 - Crocidae 273
 - Desmaninae 230
 - Erinaceinae 203, 204
 - Soricinae 258, 259

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

cell measurements in	
Talpinae	236, 237
central amygdaloid nucleus (CEA)	123
cerebellar nuclei (TCN)	86 - 88, 145, 439 - 441
cerebellum (CER)	29, 30, 52, 83 , 84 - 88, 85, 138, 140, 141, 145, 294, 416 - 420, A 534 - 551
cerebral sulci	28
chiasma opticum (CHO)	493 - 497
Chrysochloridae	185 - 192
adaptive radiation	303
brain and body weights, encephalization	185, 186, 405, 410, 412
cell and area measurements	187
histology	189, 190
index profiles	294 - 297
linear brain measures	400 - 404
macromorphology of brains	31, 186, 187
sense organs	191
summary of brain characteristics	191, 192
volume measurements	187 - 189
<i>Chrysochloris asiatica</i>	
summary of brain characteristics	192
<i>Chrysochloris stuhlmanni</i>	
macromorphology of brain	35, 38, 41, 45
summary of brain characteristics	192
cingular cortex	110, 133
classification	298
claustror cortex	133
cochlear nuclei (CON)	433 - 435
coefficient of variation (CV)	see variability
colliculus inferior (INC)	81, 82, 436 - 438, A 530 - 537
colliculus superior (SUC)	81, 82, 436 - 438, A 526 - 533
commissura anterior	see anterior commissure
complexus olivaris inferior (INO)	57, 61, 65, 66, 140, 142, 143, 157, 295, 424 - 426, A 546 - 551
complexus olivaris superior (OLS)	78, 140, 142, 144, 433 - 435, A 534 - 539
complexus sensorius trigeminalis (TR),	56 - 58, 57, 61, 140, 142, 143, 157, 295, 421 - 423, A 534 - 553
complexus sensorius nervi trigemini	
congeneric species, congeneric groups	23
conservative characters, conservative groups	6, 21, 22
conversion factors	15, 16
cornu ammonis (CA1 - CA4)	110, 128, 129, 131, 154, 155
cornu ammonis inversum	110

- corpus callosum (CC)
 length
 area in midsagittal sections (CCA)
 corpus geniculatum laterale (CGL)
 120, **A 512 - 517**
 13, 27, **400 - 404**
 27, **493 - 497**
 93, 94, **96**, 97, 136, **146**, 148,
 296, **445 - 447**, **490 - 492**,
 see also dorsal nucleus of CGL,
 ventral nucleus of CGL
 corpus geniculatum mediale (CGM)
 93, 94, **96**, 97, **146**, 148, 157,
 296, **445 - 447**, **490 - 492**,
 A 526 - 527
 A 524 - 525
 see nucleus subthalamicus
 123
- corpus mamillare (CM)
 corpus subthalamicum, corpus Luysi
 cortical amygdaloid nucleus (CAM)
 cortical maps
 of *Erinaceus*
 of *Talpa*
 110 (allocortex), **216**
 246
- cranial capacity of
 Echinosoricinae 198
 Geogalinae 167
 Potamogalinae 172
 Soricinae 255 - 257
- Crocidura attenuata*
 brain and body weights 271
 summary of brain characteristics 287
- Crocidura flavescens*
 brain and body weights 271
 summary of brain characteristics 287
- Crocidura giffardi*
 brain and body weights 271
- Crocidura* group
 index profiles **294**, **297**
 summary of brain characteristics 286 - 290
- Crocidura hildegardeae*
 brain and body weights 271
 summary of brain characteristics 288
- Crocidura jacksoni*
 brain and body weights 271
 summary of brain characteristics 288
- Crocidura leucodon*
 brain and body weights 271
- Crocidura russula*
 brain and body weights 271
 summary of brain characteristics 286, 287
- Crocidura suaveolens*
 brain and body weights 271
 summary of brain characteristics 288

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

Crociturinae	270 - 281, 285 - 292
adaptive radiation	309, 310
brain and body weights, encephalization	270 - 272, 407, 408, 412
cell and area measurements	273
histology, histochemistry	278, 279
index profiles	294 - 297
linear brain measures	400 - 404
macromorphology of brains	272
sense organs, physiology	280, 281
summary of brain characteristics	285 - 292
volume measurements	274 - 277
<i>Cryptotis parva</i>	
brain and body weights	256
summary of brain characteristics	284, 285
cubic root of brain weight or volume	13, 14, 400 - 404
Derived characters	6
<i>Desmana moschata</i>	
summary of brain characteristics	251
Desmaninae	229 - 233, 250, 251
brain and body weights, encephalization	229, 406, 410
cell and area measurements	230
histology	232, 233
index profiles	294 - 297
linear brain measures	400 - 404
macromorphology of brains	229, 230
sense organs	233
summary of brain characteristics	250, 251
volume measurements	230 - 232
diagonal band of Broca (DIAG)	110 , 119, 121, 153, A 510 - 515
diencephalon (DIE) and components	52, 83 , 88 - 100, 138, 140 , 141 , 146, 147 - 149, 157, 294 , 416 - 420
Dilambdodonta	298
dorsal hypothalamic area (HTD)	A 516 - 517
dorsal nucleus of CGL (GLD)	445 - 447 , A 518 - 525
<i>Echinops telfairi</i>	
summary of brain characteristics	179
<i>Echinorex gymnurus</i>	
brain and body weights	198
cranial capacities	198
macromorphology	35, 38, 41, 46
summary of brain characteristics	225, 226

- Echinosoricinae 197 - 200, 225, 226
 - adaptive radiation 305, 306
 - brain and body weights, encephalization 198, **406, 410, 412**
 - histology 200
 - index profiles **294, 295, 297**
 - linear brain measures **400 - 404**
 - macromorphology of brains 198, 199
 - sense organs 200
 - summary of brain characteristics 225, 226
 - volume measurements 199
- ecoethological adaptation 312 - 317
- elephant shrews, Macroscelidea 158
- encephalization, encephalization indices **48, 50, 410 - 412**
- entorhinal cortex, entorhinal region (ENT) **110, 128, 131, 132, 155**
- epiphysis (EPI) 91, 92, **146, 147, 448 - 452, A 526 - 527**
- epithalamus (ETH) 88 - 92, **89, 146, 147, 442 - 444, 487 - 489**
- Erinaceidae 197 - 228
 - adaptive radiation 304 - 306
 - body and brain weights **406, 410, 412**
 - linear brain measures **400, 402, 404**
 - macromorphology of brains 31
- Erinaceinae 200 - 224, 226 - 228
 - adaptive radiation 304 - 306
 - brain and body weights, encephalization 201, 202, **406, 410, 412**
 - cell and area measurements 203 - 205
 - fiber connections 218 - 221
 - hibernation 223
 - histology, histochemistry 207 - 218
 - index profiles **294 - 297**
 - linear brain measures **400 - 404**
 - macromorphology of brains 202, 203
 - physiology, evoked potentials 221 - 222
 - sense organs 223, 224
 - summary of brain characteristics 225, 226
 - volume measurements 205 - 207
- Erinaceus europaeus*
 - macromorphology of brain **35, 38, 41, 46**
 - summary of brain characteristic 226, 227
- evolved characters 6
- Fascia dentata (FD) **110, 128, 129, 131, 154,**
- fiber connections in brains of
 - Crocidurinae 278, 279
 - Erinaceinae 218 - 221
 - Tenrecinae 165
- fimbria-fornix complex (SFF or F) **120, 152, A 514 - 515**
- fixation of brains (Bouin's fluid) 12, 13, 15

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

formatio reticularis (FR)	A 530 - 551
formatio reticularis tegmenti	
mesencephali (FRTM)	A 528 - 531
fossorial species	314, 315
functional systems	318 - 340
funicular nuclei (FUN)	59 - 63, 421 - 423
<i>Galemys pyrenaicus</i>	
macromorphology of brain	36, 39, 42, 46
summary of brain characteristics	251
general body characteristics of	
Chrysochloridae	185
Desmaninae	229
Echinosoricinae	197
Erinaceidae	197
Erinaceinae	200, 201
Geogalinae	166
Insectivora	158
Oryzorictinae	168
Potamogalinae	171, 172
Solenodontidae	192, 193
Soricidae	254
Talpidae	228
Talpinae	234
Tenrecidae	160
Tenrecinae	160, 161
genu nervi facialis (GNF)	A 540 - 543
<i>Geogale</i>	22
<i>Geogale aurita</i> , Geogalinae	166 - 168, 180
cranial capacity, encephalization,	
brain and body weights	167, 405, 412
index profiles	294, 297
linear brain measures	400 - 404
macromorphology of brain	30, 34, 37, 40, 44 , 167
summary of brain characteristics	180
volume measurements	168
globus pallidus, pallidum (PALL)	98, 146 , 147, 157, 445 - 447 , A 514 - 517
griseum centrale (GC)	A 526 - 539

- Height of the brain (BH)** 13, 26, **400 - 404**
Hemicentetes semispinosus
 summary of brain characteristics 178, 179
Hemiechinus auritus
 brain and body weights 201
 summary of brain characteristics 227
 hippocampus (HIP) **100, 110, 123, 126, 128,**
 127 - 131, **150, 151, 154, 155,**
 157, **297, 453 - 457**
 hippocampus anterior (HA) 127, **478 - 480**
 hippocampus precommissuralis (HP) **110, 127, 130, 154, 155,**
 478 - 480, A 508 - 511
 hippocampus retro- or postcommiss. (HR) 127 - 130, **154, 155, 478 - 480,**
 with SUB, CA 1 - CA 4 and FD **A 516 - 529**
 hippocampus supracommissuralis (HS) **110, 120, 127, 130, 154, 155,**
 478 - 480, A 514 - 517
 histochemistry in brains of
 Crocidae 278, 279
 Erinaceidae 207 - 214
 Talpidae 243, 244
 histology in brains of
 Chrysochloridae 189, 190
 Crocidae 278, 279
 Desmanidae 232, 233
 Echinosoricidae 200
 Erinaceidae 207 - 218
 Oryzictidae 171
 Potamogalinae 175, 176
 Solenodontidae 196
 Soricidae 264 - 268
 Talpidae 240 - 247
 Tenrecidae 165, 166
Hylomys suillus
 brain and body weights 198
 cranial capacity 198
 summary of brain characteristics 225, 226
 hypophysis (HYP) **A 524 - 531**
 hypothalamus (HTH) **89, 90, 99, 100, 146, 147,**
 442 - 444, 487 - 491

Index profiles 294 - 297
 indices for interspecific comparisons 20, 22
 indusium griseum 127
 inferior olive see complexus olivaris inferior
 Insectivora
 adaptive radiation 298 - 311
 general body characteristics 158
 linear brain measures **26, 404**
 insular cortex 133

bold faced: page numbers of Tables
bold-faced italic: page numbers of Figures
bold-faced italic, marked A: page numbers of Atlas Figures

internal capsule	see capsula interna
interstructural comparisons	23, 24
isoponderous species	19, 21
Lateral amygdala,	
lateral amygdaloid division (LAM)	123 , 124, 125, 127, 153 , 154, 473 - 477 , A 516 - 527
lateral amygdaloid nucleus (LA)	123
lateral geniculate body	see corpus geniculatum laterale
lateral hypothalamus (HTL)	A 516 - 525
lateral olfactory tract (TRL)	106 , 107 , 111 , 115, 116, 151 , 468 - 472 , A 506 - 513
lateral preoptic area (POL)	A 512 - 515
lateral reticular nucleus (REL)	see nucleus reticularis lateralis
length of brain (BL)	13, 25, 400 - 404
length of cerebral hemispheres (HemL)	13, 26, 400 - 404
length of corpus callosum (CCL)	13, 27, 400 - 404
limbic system	127, 335 - 338
<i>Limnogale mergulus</i>	
summary of brain characteristics	182
linear measures on brains	13, 25, 400 - 404
lobulus petrosus	29, 30
locus coeruleus	see nucleus locus coeruleus
Macromorphology of brains in	25 - 47 , 26
Chrysochloridae	31, 186, 187
Crocidae	272
Desmaninae	229, 230
Echinosoricinae	198, 199
Erinaceidae	31
Erinaceinae	202, 203
Geogalinae	30, 167
Oryzorictinae	30, 169
Potamogalinae	31, 172, 173
Solenodontidae	31, 193, 194
Soricidae	32
Soricinae	257, 258
Talpidae	31
Talpinae	235, 236
Tenrecinae	30, 161, 162
magnocellular part of basal amygdaloid nucleus (MCB)	124, 127, 153 , 154, 473 - 477

- main olfactory bulb (MOB) and components **103**, 104 - 106, **106**, **107**, **110**,
150, 157, 297, **458** - **462**,
A 506 - **507**
- main olfactory system 318 - 322
- man, *Homo* 20, 52, 101, 137 - 157
- material and methods 12 - 24
- medial amygdala,
 medial amygdaloid division (MAM) **123**, 124, 125, **153**, 154,
473 - **477**, **A 514** - **521**
- medial amygdaloid nucleus (MA) **123**
- medial geniculate body see corpus geniculatum mediale
- medial preoptic area (POM) **A 512** - **515**
- medulla oblongata (OBL) and components **52**, 53 - 79, **54**, **138**, 139, **140**,
142, 142 - 144, **294**, **295**,
416 - **420**, **421** - **435**
- mesencephalic tectum (MTC) 28, 79, 82, **436** - **438**
- mesencephalic tegmentum (MTG) 79, 82, **436** - **438**
- mesencephalon (MES) and components **53**, 79 - 84, **83**, **138**, **140**, **141**,
144, **145**, **294**, **416** - **420**
- methods of comparison 19
- Microgale dobsoni*
- summary of brain characteristics 181, 182
- Microgale talazaci*
- macromorphology of brain **34**, **37**, **40**, **44**
- summary of brain characteristics 181, 182
- Micropotamogale lamottei*
- summary of brain characteristics 184
- Micropotamogale ruwenzorii*
- summary of brain characteristics 184
- Microsorex hoyi*
- brain and body weights 256
- summary of brain characteristics 283
- motor nuclei (MOT) 68 - 71, **427** - **429**
- motor systems 334, 335
- mouse-elephant line 22
- Myosorex babaulti*
- brain and body weights 272
- summary of brain characteristics 291, 292
- Neocortex (NEO) and components **85**, **100**, **120**, **128**, 132 - 137,
134, **150**, 151, **156**, 157, **297**,
338 - 340, **453** - **457**, **481** - **485**,
A 508 - **531**
- neocortical grey matter (NG) 135, 136, **156**, **481** - **485**
- neocortical white matter (NW) 135, 136, **156**, **481** - **485**,
A 508 - **527**
- Neomyini 308

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

Neomys anomalus

brain and body weights 256

summary of brain characteristics 284

Neomys fodiens

brain and body weights 256

summary of brain characteristics 284

nervus facialis (N VII)

A 538 - 541

net brain volumes (NET)

50 - 52, ***410 - 415***

nuclei anterior thalami (ANT)

94, ***146, 148, 149, 157, 296, 490 - 492, A 516 - 517***

nuclei dorsales thalami (DOT)

95, ***96, 146, 148, 149, 157, 296, 490 - 492***

nuclei mediales thalami (MLT)

94, ***146, 148, 149, 157, 296, 490 - 492***

nuclei mediani thalami (MNT)

95, ***146, 148, 149, 296, 490 - 492***

nuclei of lateral lemniscus

see nucleus lemnisci lateralis

nuclei reticulares thalami (RTH)

89, ***146, 487 - 489, A 516 - 525***

nuclei ventrales thalami (VET)

95, ***96, 146, 148, 149, 296, 490 - 492***

nucleus anterodorsalis thalami (AD)

A 518 - 519

nucleus centrolateralis thalami (CLT)

A 520 - 523

nucleus centromedialis thalami (CMT)

A 518 - 525

nucleus cochlearis dorsalis (DCO)

61, 75 - 77, 140, 142, 144, 295, 433 - 435, A 542 - 543

nucleus cochlearis ventralis (VCO)

75 - 77, ***140, 142, 144, 295, 433 - 435, A 536 - 543***

nucleus cuneiformis (CU)

A 530 - 533

nucleus dorsalis motorius nervi vagi (mot X)

57, 68 - 70, ***140, 142, 144, 295, 427 - 429, A 546 - 551***

nucleus dorsalis raphe (DR)

A 532 - 533

nucleus dorsalis septi (SED)

120, 121

nucleus dorsomedialis hypothalami (DMH)

A 520 - 521

nucleus fascicularis gracilis (FGR)

57, 60, ***140, 142, 295, 427 - 429, A 550 - 553***

nucleus fascicularis cuneatus medialis (FCM)

57, 62, ***140, 142, 295, 427 - 429, A 548 - 553***

nucleus fascicularis cuneatus externus (FCE)

63, ***140, 142, 295, 427 - 429, A 546 - 549***

nucleus fimbrialis septi (SEF)

120

nucleus habenularis lateralis (HAL)

A 520 - 525

nucleus habenularis medialis (HAM)

91, 92, ***146, 147, 448 - 452, A 520 - 525***

nucleus infundibularis (INF)

A 520 - 521

- nucleus intermediodorsalis thalami (IMD) *A 520 - 525*
nucleus interpeduncularis (IP) *A 530 - 531*
nucleus interpositus cerebelli (ICN) 85 - 88, **145**, **439 - 441**,
A 540 - 545
nucleus lateralis cerebelli (LCN) 86 - 88, **145**, 157, **439 - 441**,
A 540 - 543
nucleus lateralis posterior thalami (LP) *A 524 - 527*
nucleus lateralis septi (SEL) *120*, 152
nucleus laterodorsalis thalami (LD) *A 518 - 523*
nucleus lemnisci lateralis (NLL) 81, 82, **436 - 438**, *A 532 - 535*
nucleus locus coeruleus (COE) 67, **140**, **142**, 143, **295**,
424 - 426, *A 536 - 537*
nucleus medialis cerebelli (MCN) 85 - 87, **145**, **439 - 441**, *A 540 - 545*
nucleus medialis thalami dorsalis (MD) *A 518 - 525*
nucleus medianus hypothalami (MHT) *120*,
nucleus mesencephalicus nervi trigemini (TRM) *A 536 - 539*, see also complexus
sensorius trigeminalis
nucleus motorius nervi trigemini (mot V) 68, **140**, **142**, 143, 157, **295**,
427 - 429, *A 536 - 539*
nucleus nervi abducentis (mot VI) *A 542 - 543*
nucleus nervi facialis (mot VII) 69, **140**, **142**, 144, 157, **295**,
427 - 429, *A 540 - 543*
nucleus nervi hypoglossi (mot XII) 70, 71, **140**, **142**, 144, **295**,
427 - 429, *A 548 - 551*
nucleus oculomotorius (III) *A 530 - 531*
nucleus of anterior commissure (SEA) *120*
nucleus of lateral olfactory tract (NTO) *123*, 124, 125, **153**, 154,
473 - 477, *A 514 - 515*
nucleus of stria terminalis (SES) *120*, 121, 153, *A 514 - 517*
nucleus olfactorius anterior see retrobulbar region
nucleus parhypothalamicus (PH) *A 516 - 517*
nucleus paratenialis thalami (PT) *A 516 - 519*
nucleus prepositus hypoglossi (PRP) 66, **140**, **142**, 143, **295**,
424 - 426, *A 542 - 547*
nucleus raphe (RA) *A 536 - 541*
nucleus residualis cornus ventralis (RC) *A 552 - 553*
nucleus reticularis dorsalis (RED) *57*
nucleus reticularis lateralis (REL) *57*, 64, 65, **140**, **142**, 143, **295**,
424 - 426, *A 548 - 551*
nucleus reticularis medullae oblongatae (RMO) *A 552 - 553*
nucleus reticularis paramediani (RP) *A 542 - 543*
nucleus reticularis ventralis (REV) *57*
nucleus reuniens thalami (RE) *A 518 - 525*
nucleus rhomboides thalami (RH) *A 520 - 521*
nucleus ruber (RUB) *A 526 - 531*
nucleus sensorius principalis nervi trigemini see complexus sens. trigemin.
nucleus spinalis nervi trigemini see complexus sens. trigemin.
nucleus subthalamicus Luysi (LUY) 98, **146**, 147, 157, **445 - 447**,
A 522 - 523

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

nucleus suprachiasmaticus (SC)	<i>A 516 - 517</i>
nucleus supraopticus (SON)	<i>A 514 - 515</i>
nucleus tractus solitarii (TSO)	<i>57, 58, 59, 140, 142, 143, 295, 422 - 424, A 544 - 551</i>
nucleus triangularis septi (SET)	<i>120, 121, 153, 448 - 452</i>
nucleus ventralis tegmenti (VT)	<i>A 534 - 535</i>
nucleus ventrolateralis thalami (VLT)	<i>A 518 - 523</i>
nucleus ventromedialis hypothalami (VMH)	<i>A 518 - 521</i>
nucleus ventromedialis thalami (VMT)	<i>A 520 - 523</i>
nucleus vestibularis inferior (VI)	<i>73, 140, 142, 295, 430 - 432, A 544 - 547</i>
nucleus vestibularis lateralis (VL)	<i>74, 140, 142, 295, 430 - 432, A 540 - 543</i>
nucleus vestibularis medialis (VM)	<i>72, 73, 140, 142, 295, 430 - 432, A 540 - 547</i>
nucleus vestibularis superior (VS)	<i>61, 74, 75, 140, 142, 144, 295, 430 - 432, A 538 - 541</i>
Olfactory bulbs (BOL)	<i>100, 102, 104, 453 - 457</i>
olfactory cortices (OLC)	<i>468 - 472</i>
olfactory tubercle (TOL)	<i>107, 110, 111, 114, 115, 151, 152, 468 - 472</i>
oliva inferior	see complexus olivaris inferior
oliva superior	see complexus olivaris superior
optic nerve (N II)	<i>A 510 - 515</i>
optic tract	see tractus opticus
<i>Oryzorictes talpoides</i>	
summary of brain characteristics	181
Oryzoricetinae	
adaptive radiation	301 - 302
brain and body weights, encephalization	169, <i>405, 410, 412</i>
area measurements	170
histology	171
index profiles	<i>294 - 297</i>
linear brain measures	<i>400 - 404</i>
macromorphology of brains	30, 31, 169
summary of brain characteristics	180 - 182
volume measurements	170, 171
Paleocortex (PAL)	<i>100, 109 - 117, 110, 112, 151, 152, 157, 297, 453 - 457</i>
pallidum	see globus pallidus

- Parascalops breweri*
 brain and body weights 235
 summary of brain characteristics 253
 parasubicular cortical region (PAR) **128**, 131
 pedunculus cerebri (PC) **A 524 - 529**
 percentage values 19, 20, 49, 50, **52**, **100**, **111**,
410 - 415
 periamygdaloid cortical region (PAM) **110**, **111**, 112
 periarhicortex **110**
 peripaleocortex **110**
 perirhinal cortical region (PRH) 128, 131, 132, 155
 periseptal region 120
 physiology in
 Erinaceinae 221, 222
 physiology, sense organs in
 Crocidae 280, 281
 Soricinae 268 - 270
 Talpinae 247 - 250
 pineal gland, pinealis see epiphysis
 plesiomorphic characters 6, 21, 22
 pons 53
 postcommissural hippocampus see hippocampus retrocomm.
Potamogale velox
 macromorphology of brain **34**, **37**, **40**, **45**
 summary of brain characteristics 183
 Potamogalinae 171 - 176, 183, 184
 adaptive radiation 302, 303
 brain and body weights, encephalization 172, **405**, **410**, **412**
 area measurements 173
 histology 175, 176
 index profiles **294 - 297**
 linear brain measures **400 - 404**
 macromorphology of brains 172, 173
 summary of brain characteristics 183, 184
 volume measurements 173 - 175
 precentral cortical region 133
 precommissural hippocampus see hippocampus precommiss.
 preparation of brains 12
 prepiriform cortical region (PRPI) **107**, **110**, **111**, 113, 114, **123**,
151, 152, **468 - 472**, **A 506 - 525**
 presubicular cortical region (PRS) **110**, **128**, 131, 132, 155
 pretectal region (PTE) 89, **146**, **487 - 489**
 Primates 19, 20, 137 - 157
 primitive characters 6, 21, 22
 proportion indices (PI) 14, 25 - 27
 psalterium ventrale 120
 putamen see striatum

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

Reference lines	21, 22
regio periamygaloidea	see periamygdaloid region
regio prepiriformis	see prepiriform cortical region
regio retrobulbaris	see retrobulbar region
regression methods	20, 21
relative brain size (RBrS)	20, 22
relay nuclei (RLN)	62 - 67, 424 - 426
requirements for comparison	19
rest, remnants	17, 413 - 415
retrobulbar region (RB)	106, 107, 110, 111 , 114, 115, 151 , 152, 468 - 472 , A 506 - 507
retrocommissural hippocampus	see hippocampus retrocomm.
retrosplenial cortex	110 , 133
<i>Ruwenzorisorex suncoides</i>	
brain and body weights	272
index profiles	294, 297
summary of brain characteristics	291
<i>Scalopus aquaticus</i>	
brain and body weights	235
macromorphology of brain	36, 39, 42, 47
summary of brain characteristics	253, 254
Scandentia	158
schizocortex (SCH)	100, 126, 128 , 131, 132, 150 , 151, 155, 157, 297, 453 - 457 , A 518 - 531
<i>Scutisorex somereni</i>	
brain and body weights	271
summary of brain characteristics	289, 290
semiaquatic species	315, 316
sense organs in	
Chrysochloridae	191
Crocidae	280, 281
Desmaninae	233
Echinosoricinae	200
Erinaceinae	223, 224
Soricinae	268 - 270
Talpinae	247 - 250
Tenrecinae	166
sensory trigeminal complex	see complexus sensorius trig.
septum (SEP) and components	100, 110 , 119 - 122, 120, 126 , 150 , 151 - 153, 153, 297 , 453 - 457, A 508 - 515
septum pellucidum	119

- Setifer setosus*
 summary of brain characteristics 178
 shape of brains 13
 size indices 51
- Solenodon paradoxus*
 macromorphology **35, 38, 41, 45**
 summary of brain characteristics 196
- Solenodontidae 192 - 196
 adaptive radiation 304
 brain and body weights, encephalization 193, **406, 410, 412**
 area measurements 194
 histology 196
 index profiles **294 - 297**
 linear brain measures **400 - 404**
 macromorphology of brains 31, 193, 194
 summary of brain characteristics 196
 volume measurements 194, 195
- somatosensory system 327 - 330
- Sorex* 23
- Sorex alpinus*
 brain and body weights 256
 summary of brain characteristics 282
- Sorex araneus*
 brain and body weights 256
 summary of brain characteristics 282
- Sorex cinereus*
 brain and body weights 256
 summary of brain characteristics 282, 283
- Sorex fumeus*
 brain and body weights 256
 summary of brain characteristics 283
- Sorex minutus*
 brain and body weights 256
 summary of brain characteristics 283
- Sorex palustris*
 brain and body weights 256
- Soricidae 254 - 292
 adaptive radiation 306 - 310
 body and brain weights **406 - 408, 411, 412**
 linear brain measures **401, 403, 404**
 macromorphology of brains 32
- Soricinae 255 - 270, 281 - 285
 adaptive radiation 307 - 309
 brain and body weights, encephalization 255 - 257, **406, 407, 411, 412**
 cell and area measurements 258 - 260
 histology 264 - 268
 index profiles **294 - 297**
 linear brain measures **400 - 404**
 macromorphology of brains 257, 258

bold faced: page numbers of Tables

bold-faced italic: page numbers of Figures

bold-faced italic, marked A: page numbers of Atlas Figures

Soricinae (cont.)	
physiology	268 - 270
sense organs	268 - 270
summary of brain characteristics	281 - 285
volume measurements	260 - 263
Soricini	307
standard values of brain and body	16
striatum (STR)	100 , 117 - 119, <i>118</i> , <i>120</i> , 150 , 151, 157, <i>297</i> , <i>453 - 457</i> , <i>A 508 - 517</i>
subcommissural body (SCB)	80 , 82, 84, 144, <i>145</i> , <i>448 - 452</i>
subfornical body (SFB)	119, 121, 122, 153 , <i>448 - 452</i> , <i>A 514 - 515</i>
subgenual cortex	<i>110</i> , 133
subiculum (SUB)	<i>128</i> , 129, 131, 154
substantia innominata (SIN)	111 , 117, 151 , 152, <i>468 - 472</i> , <i>A 514 - 515</i>
substantia nigra (SN)	<i>A 524 - 529</i>
subterranean species	314, 315
subthalamus (STH)	89 , 90, 98, 99, 146 , 147, <i>442 - 444</i> , <i>487 - 489</i>
sulcus semiannularis (SAN)	<i>123</i>
summary of brain characteristics in	
Chrysochloridae	191, 192
Crocidae	285 - 292
<i>Crocidae</i> group	286 - 290
Desmaninae	250, 251
Echinosoricinae	225, 226
Erinaceidae	224 - 228
Erinaceinae	226 - 228
Geogalinae	180
Oryzorictinae	180 - 182
Potamogalinae	183, 184
Solenodontidae	196
Soricinae	281 - 285
<i>Sylvisorex</i> group	290 - 292
Talpidae	250 - 254
Talpinae	252 - 254
Tenrecidae	176 - 184
Tenrecinae	177 - 180
<i>Suncus etruscus</i>	
brain and body weights	271
macromorphology of brain	36 , 39 , 42 , 47
summary of brain characteristics	289

- Suncus murinus*
 brain and body weights 271
 summary of brain characteristics 289
 superior olive see complexus olivaris superior
 supracommissural hippocampus see hippocampus supracomm.
 surface dwellers 312, 313
 surface measurements, surface dimensions 52
- Sylvisorex granti*
 brain and body weights 271
 summary of brain characteristics 290
- Sylvisorex* group
 index profiles **294, 297**
 summary of brain characteristics 290 - 292
- Sylvisorex megalura*
 brain and body weights 272
 summary of brain characteristics 290, 291
- Sylvisorex* spp.
 index profiles **294, 297**
- Talpa europaea*
 summary of brain characteristics 252, 253
- Talpa micrura*
 brain and body weights 234
 summary of brain characteristics 253
- Talpidae 228 - 254
 adaptive radiation 310, 311
 body and brain weights **406, 410, 412**
 linear brain measures **400, 402, 404**
 macromorphology of brains 31
- Talpinae 234 - 250, 252 - 254
 adaptive radiation 310, 311
 brain and body weights, encephalization 234, 235, **406, 410, 412**
 cell and area measurements 236 - 237
 histology, histochemistry 240 - 247
 index profiles **294 - 297**
 linear brain measures **400 - 404**
 macromorphology of brains 235, 236
 physiology, sense organs 247 - 250
 summary of brain characteristics 252 - 254
 volume measurements 238 - 240
- tectum see mesencephalic tectum
 tegmentum see mesencephalic tegmentum
 telencephalon (TEL) and components **52, 100 - 137, 101, 138, 139, 140, 141, 149 - 157, 150, 294, 297, 418 - 422**
- Tenrec ecaudatus*
 macromorphology of brain **34, 37, 40, 44**
 summary of brain characteristics 177, 178

bold faced: page numbers of Tables
bold-faced italic: page numbers of Figures
bold-faced italic, marked A: page numbers of Atlas Figures

Tenrecidae	160 - 184
adaptive radiation	299 - 303
body and brain weights	405, 410, 412
linear brain measures	400, 402, 404
Tenrecinae	22, 160 - 166, 177 - 180
adaptive radiation	304 - 306
brain and body weights, encephalization	161, 405, 410, 412
area measurements	162
histology, fiber connections	165, 166
index profiles	294 - 297
linear brain measures	400, 402, 404
macromorphology of brains	30, 161, 162
sense organs	166
summary of brain characteristics	177 - 180
volume measurements	162 - 165
tenrecine average (AvTn)	22
thalamus (THA)	88, 89 , 92 - 97, 146 , 147 - 149, 442 - 444, 487 - 489
thalamus posterior (THP)	A 526 - 527
thalamus ventroposterior (VP)	A 524 - 525
tractus olfactorius lateralis	see lateral olfactory tract
tractus opticus (TRO)	89, 90, 146, 442 - 444, A 516 - 519
tree shrews	18, 158
trigeminal nerve (N V)	A 508 - 533
T-tests	85
tuberculum olfactorium	see olfactory tubercle
<i>Tupaia glis</i>	18, 158
Variability, variation	17, 18, 25, 405 - 408
ventral nucleus of CGL (GLV)	445 - 447, A 518 - 525
ventral pons (VPO)	439, 440, A 532 - 535
ventricle size of brains	413 - 415
vestibular complex (VC)	71 - 75, 144, 430 - 432
vestibular system	333, 334
visual system	324 - 327
volume determination	14
volume measurements in	
Chrysochloridae	187 - 189
Crocidae	274 - 277
Desmaninae	230 - 232
Echinosoricinae	199
Erinaceinae	205 - 207
Geogalinae	168
Oryzictinae	170, 171

volume measurements in	
Potamogalinae	173 - 175
Solenodontidae	194, 195
Soricinae	260 - 263
Talpinae	238 - 240
Tenrecinae	162 - 165
vomeronasal organ, vomeronasal system	322 - 324
Weight of body (BoW)	12, 13, 405 - 411
weight of brain (BrW)	12, 13, 48, 405 - 411
weight of brain parts	238, 261, 275
width of the brain (BW)	13, 26, 400 - 404
width of cerebral hemispheres	13, 26, 400 - 404
winter survival	316, 317
Zalambdodonta	298
zona incerta (ZI)	A 518 - 525