

# Completeness of Inst-saturated Sets of Clauses with Equality

Alexander Maringele

[alexander.maringele@gmail.com](mailto:alexander.maringele@gmail.com)

December 6th, 2017






Harald Ganzinger and Konstantin Korovin.

Integrating Equational Reasoning into Instantiation-Based Theorem Proving.

In *18th CSL 2004. Proceedings*, volume 3210 of *LNCS*, pages 71–84, 2004.

# Instantiation-based first order theorem proving

## The big picture



$S_0 = S$ ,  $S_{i+1}$  is inferred from  $S_i$  by a sound calculus.

# Preliminaries I

- ▶ a clause  $C$  is a multiset of literals
- ▶ literals are (in)equations of first order terms
- ▶ a closure  $C \cdot \sigma$  is a pair of clause  $C$  and substitution  $\sigma$

# Preliminaries II

## ► orderings

$\succ_{gr}$       order on ground terms, literals, and clauses defined by  
                  a total, well-founded, and monotone extension of  
                  a total simplification ordering  $\succ'_{gr}$  on ground terms

$\succ_{\ell}$       an arbitrary total well-founded extension of  $\succ_{gr}$  such that  

$$L\sigma \succ_{gr} L'\sigma' \Rightarrow L \cdot \sigma \succ_{\ell} L' \cdot \sigma'$$

$\succ_{cl}$       an arbitrary total well-founded extension of  $\succ_{gr}$  such that  

$$C\tau \succ_{gr} D\rho \Rightarrow C \cdot \tau \succ_{cl} D \cdot \rho$$

$$(C\tau = D\rho \text{ and } C\theta = D) \Rightarrow C \cdot \tau \succ_{cl} D \cdot \rho$$

# Unit Paramodulation

## Definition

$$\frac{(\ell \approx r) \cdot \sigma \quad L[\ell'] \cdot \sigma'}{L[r]\theta \cdot \rho} \theta \qquad \frac{(s \not\approx t) \cdot \tau}{\square} \mu$$

where

- ▶  $\ell\sigma \succ_{gr} r\sigma$ ,  $\theta = \text{mgu}(\ell, s)$ ,  $\ell\sigma = \ell'\sigma' = \ell'\theta\rho$ ,  $\ell' \notin \mathcal{V}$
- ▶  $s\tau = t\tau$ ,  $\mu = \text{mgu}(s, t)$

## Remark

The set of literal closures  $\{ (f(x) \approx b) \cdot \{x \rightarrow a\}, a \approx b, f(b) \not\approx b \}$  is inconsistent, but the empty clause is not derivable if  $a \succ_{gr} b$ .

# UP-Redundancy

- ▶ We define the set

$$\text{irred}_R(\mathcal{L}) = \{ L \cdot \sigma \in \mathcal{L} \mid \sigma \text{ is irreducible w.r.t. } R \}$$

for a set of literal closures  $\mathcal{L}$  and a ground rewrite system  $R$ .

- ▶ Let  $\mathcal{L}_{L \cdot \sigma \succ_\ell} = \{ L' \cdot \sigma' \in \mathcal{L} \mid L \cdot \sigma \succ_\ell L' \cdot \sigma' \}$ .
- ▶ A literal closure  $L \cdot \sigma$  is UP-redundant in  $\mathcal{L}$  if

$$R \cup \text{irred}_R(\mathcal{L}_{L \cdot \sigma \succ_\ell}) \models L\sigma$$

for every ground rewrite system  $R$   
oriented by  $\succ_{gr}$  where  $\sigma$  is irreducible w.r.t.  $R$ .

- ▶  $\mathcal{R}_{UP}(\mathcal{L})$  denotes the set of all UP-redundant closures in  $\mathcal{L}$ .

# UP-Saturation I

A UP-saturation process is

a sequence  $\{\mathcal{L}_i\}_{i=0}^{\infty}$  of sets of literal closures

where  $\mathcal{L}_{i+1}$  is obtained from  $\mathcal{L}_i$

by adding a conclusion of an UP-inference with premises in  $\mathcal{L}_i$

or by removing a UP-redundant closure w.r.t.  $\mathcal{L}_i$ .

$$\mathcal{L}_{i+1} = \begin{cases} \mathcal{L}_i \cup \square & \text{if } \mathcal{L}_i \ni (s \not\approx t) \cdot \tau, s\tau = t\tau, \mu = \text{mgu}(s, t) \\ \mathcal{L}_i \setminus L \cdot \sigma & \text{if } R \cup \text{irred}_R(\mathcal{L}_i, L \cdot \sigma \succ_{\ell}) \models L\sigma \\ \mathcal{L}_i \cup L[r]\theta \cdot \rho & \text{if } \begin{cases} (\ell \approx r) \cdot \sigma \in \mathcal{L}_i, L[\ell'] \cdot \sigma' \in \mathcal{L}_i \\ \ell\sigma \succ_{gr} r\sigma, \theta = \text{mgu}(\ell, \ell'), \\ \ell' \notin \mathcal{V}, \ell\sigma = \ell'\sigma' = \ell'\theta\rho, \end{cases} \\ \mathcal{L}_i & \text{otherwise} \end{cases}$$



## UP-Saturation II

### Definition

Let  $\mathcal{L}^\infty$  be the set of persistent closures, i.e. the lower limit of the sequence. The process is UP-fair if for every UP-inference with premises in  $\mathcal{L}^\infty$  the conclusion is UP-redundant w.r.t.  $\mathcal{L}_j$  for some  $j$ . For a set of literals  $\mathcal{L}$  we define the saturated set of literal closures  $\mathcal{L}^{sat} = \mathcal{L}^\infty \setminus \mathcal{R}_{UP}(\mathcal{L}^\infty)$  for some UP-saturation process  $\{\mathcal{L}_i\}_{i=0}^\infty$  with  $\mathcal{L}_0 = \mathcal{L}$ .

### Lemma

*The set  $\mathcal{L}^{sat}$  is unique because for any two UP-fair saturation processes  $\{\mathcal{L}_i\}_{i=0}^\infty$  and  $\{\mathcal{L}'_i\}_{i=0}^\infty$  with  $\mathcal{L}_0 = \mathcal{L}'_0$  we have*

$$\mathcal{L}^\infty \setminus \mathcal{R}_{UP}(\mathcal{L}^\infty) = \mathcal{L}'^\infty \setminus \mathcal{R}_{UP}(\mathcal{L}'^\infty)$$

# Inst-Redundancy

Let  $S$  be a set of clauses.

- ▶ A ground closure  $C$  is called Inst-redundant if there exist ground instances  $C_1, \dots, C_k$  of  $S$  such that
  - ▶  $C \succ_{cl} C_i$  for all  $i$
  - ▶  $C_1, \dots, C_k \models C$
- ▶ A (possible non-ground) clause  $C$  is called Inst-redundant in  $S$  if each ground closure  $C \cdot \sigma$  is Inst-redundant in  $S$
- ▶  $R_{Inst}(S)$  denotes the set of all Inst-redundant clauses in  $S$ .

# Selection

Let  $S$  be a set of clauses  $S$ , let  $I_{\perp}$  be a model of  $S_{\perp}$ .

- ▶ A selection function  $\text{sel}$  maps clauses to literals such that

$$\text{sel}(C) \in C \qquad I_{\perp} \models \text{sel}(C)_{\perp}$$

- ▶ The set of  $S$ -relevant literal closures

$$\mathcal{L}_S = \left\{ L \cdot \sigma \mid \begin{array}{l} L \vee C \in S, L = \text{sel}(L \vee C) \\ (L \vee C) \cdot \sigma \text{ is not Inst-redundant in } S, \end{array} \right\}$$

- ▶  $\mathcal{L}_S^{\text{sat}}$  denotes the saturation process of  $\mathcal{L}_S$ .
- ▶ A set of clauses  $S$  is Inst-saturated w.r.t. a selection function, if  $\mathcal{L}_S^{\text{sat}}$  does not contain the empty clause.

# Completeness

## Theorem

*If a set of clauses  $S$  is Inst-saturated, and  $S \perp$  is satisfiable, then  $S$  is also satisfiable.*

## Proof.

1. Construct candidate model
2. Assumed counterexample fails

Conclude candidate is model



# Model Construction I

Let  $S$  be an Inst-saturated set of clauses.

- ▶  $S \perp$  is satisfiable
- ▶  $\Box \notin \mathcal{L}_S^{sat}$

Let  $L = L' \cdot \sigma \in \mathcal{L}_S^{sat}$ . We define by induction on  $\succ_\ell$

- ▶  $I_L = \bigcup_{L \succ_\ell M} \epsilon_M$        $\epsilon_M$  already defined for all  $M$  with  $L \succ_\ell M$
- ▶  $R_L = \{s \rightarrow t \mid s \approx t \in I_L, s \succ_{gr} t\}$
- ▶  $\epsilon_L = \begin{cases} \emptyset & \text{if } L'\sigma \text{ reducible by } R_L \\ \emptyset & \text{if } I_L \models L'\sigma \text{ or } I_L \models \overline{L'}\sigma \text{ (defined)} \\ \{L'\sigma\} & \text{if } L'\sigma \text{ is productive (irreducible, undefined)} \end{cases}$

## Model Construction II

- ▶  $R_S = \bigcup_{L \in \mathcal{L}_S^{\text{sat}}} R_L$   $R_S$  is convergent and interreduced
- ▶  $I_S = \bigcup_{L \in \mathcal{L}_S^{\text{sat}}} \epsilon_L$   $I_S$  is consistent,  
 $L\sigma \in L_S$  is irreducible by  $R_S$
- ▶ Let  $\mathcal{I}$  be an arbitrary total consistent extension of  $I_S$ .

### Lemma

*$\mathcal{I}$  is a model for all ground instances of clauses in  $S$ .*

# Assumed Counterexample I

Assume  $\mathcal{I}$  is not a model of  $S$ .

$$\text{Let } D = \min_{\succ_{cl}} \{ C' \cdot \sigma \mid C' \in S, \mathcal{I} \not\models C'\sigma \}$$

Then

- ▶  $D = D' \cdot \sigma$  is not Inst-redundant. Otherwise  $D_1, \dots, D_n \models D$ ,  $D \succ_{cl} D_i$  for all  $i$ , and  $\mathcal{I} \not\models D_j$  for one  $j$  contradicts minimality.
- ▶  $x\sigma$  irreducible by  $R_S$  for every variable  $x$  in  $D'$ . Otherwise let  $(\ell \rightarrow r)\tau \in R_L$  and  $x\sigma = x\sigma[l\tau]_p$  for some variable  $x$  in  $D'$ . We define substitution  $\sigma'$  with  $x\sigma' = x\sigma[r\tau]_p$  and  $y\sigma' = y\sigma$  for  $y \neq x$ .  $\mathcal{I} \not\models D'\sigma'$  and  $D \succ_{cl} D' \cdot \sigma'$  contradicts minimality.

## Assumed Counterexample II

Since  $D$  is not Inst-redundant in  $S$ , we have for some literal  $L$ , that  $D' = L \vee D''$ ,  $\text{sel}(D') = L$ ,  $L \cdot \sigma \in \mathcal{L}_S$ ,  $L\sigma$  is false in  $\mathcal{I}$

Assume  $L \cdot \sigma$  is UP-redundant in  $\mathcal{L}_S^{\text{sat}}$ .

By construction  $\sigma$  is irreducible by  $R_S$ . Then we have

$$R_S \cup \text{irred}_{R_S}(\{L' \cdot \sigma' \in \mathcal{L}_S^{\text{sat}} \mid L \cdot \sigma \succ_{\ell} L' \cdot \sigma'\}) \models L\sigma$$

Therefore there is  $L' \cdot \sigma' \in \text{irred}_{R_S}(\mathcal{L}_S^{\text{sat}})$  that is false in  $I$ .

$$M \cdot \tau = \min_{\succ_{\ell}} \{ L' \cdot \tau' \mid L' \cdot \sigma' \in \text{irred}_{R_S}(\mathcal{L}_S^{\text{sat}}), \mathcal{I} \not\models M' \tau' \}$$

is irreducible by  $R_S$ .

$$M \cdot \tau = \min_{\succ_{\ell}} \{ L' \cdot \tau' \mid L' \cdot \sigma' \in \text{irred}_{R_S}(\mathcal{L}_S^{\text{sat}}), \mathcal{I} \not\models M' \tau' \}$$

Assume  $M \cdot \tau$  is reducible by  $(\ell \rightarrow r) \in R_S$

and  $(\ell \rightarrow r)$  is produced by  $(\ell' \approx r') \cdot \rho \in \mathcal{L}_S^{\text{sat}}$



# Assumed Counterexample III

$\tau$  is irreducible by  $R_S$ , hence UP-inference is applicable:

$$\frac{(\ell' \approx r') \cdot \rho \quad M[\ell''] \cdot \tau}{M[r']\theta \cdot \mu} \text{ UP}$$

$$\ell' \rho = \ell'' \tau = \ell'' \theta \mu, \theta = \text{mgu}(\ell', \ell''), \mathcal{I} \not\models M[r']\theta \mu$$

# Assumed Counterexample IV

- ▶ If  $M[r']\theta \cdot \mu$  is not UP-redundant in  $\mathcal{L}_S^{sat}$  then  $M[r']\theta \cdot \mu \in \mathcal{L}_S^{sat}$ .  
 $M[r']\theta \cdot \mu \in \text{irred}_{R_S}(\mathcal{L}_S^{sat})$  because  $\mu$  is irreducible.
- ▶ If  $M[r']\theta \cdot \mu$  is UP-redundant in  $\mathcal{L}_S^{sat}$ .

From definition:

$$R_S \cup \text{irred}_{R_S}(\{M' \cdot \tau' \in \mathcal{L}_S^{sat} \mid M[r']\theta \cdot \mu \succ_\ell M' \tau'\}) \models M[r']\theta \mu$$

Hence there is  $M' \cdot \tau' \in \mathcal{L}_S^{sat}$ ,  $M \cdot \tau \succ_\ell M[r']\theta \cdot \mu \succ_\ell M' \cdot \tau'$   
 false in  $\mathcal{I}$ . contradicts minimality of  $M \cdot \tau$ .

# Assumed Counterexample V

We have that  $M \cdot \tau$

- ▶ is false in  $\mathcal{I}$
- ▶ is in  $\mathcal{L}_S^{sat}$
- ▶ is irreducible by  $R_S$
- ▶ is not productive.

Hence  $I_{M \cdot \tau} \models \overline{M} \tau$  with two possible cases:

1.  $M \cdot \tau$  is equation  $(s \approx t) \cdot \tau$
2.  $M \cdot \tau$  is inequation  $(s \not\approx t) \cdot \tau$

$$I_{M \cdot \tau} \models (s \not\approx t) \tau$$

$$I_{M \cdot \tau} \models (s \approx t) \tau$$

# Assumed Counterexample VI

1. Assume  $M \cdot \tau$  is equation  $(s \approx t) \cdot \tau$ :

- ▶  $I_{M \cdot \tau} \models (s \not\approx t)\tau$
- ▶ All literals in  $I_{M \cdot \tau}$  are irreducible by  $R_{M \cdot \tau}$
- ▶  $s\tau$  and  $t\tau$  are irreducible by  $R_{M \cdot \tau}$
- ▶  $R_{M \cdot \tau}$  is a convergent term rewrite system

Hence  $(s \not\approx t)\tau \in I_{M \cdot \tau}$  and produced to  $I_{M \cdot \tau}$  by a  $(s' \not\approx t') \cdot \tau'$ .  
Contradicts the minimality of  $M \cdot \tau$ .

2. Assume  $M \cdot \tau$  is inequation  $(s \not\approx t) \cdot \tau$ :

- ▶  $I_{M \cdot \tau} \models (s \approx t)\tau$
- ▶  $s\tau$  and  $t\tau$  are irreducible by  $R_{M \cdot \tau}$

Hence  $s\tau = t\tau$  and equality resolution is applicable.  
Contradicts that the empty clause is not in  $\mathcal{L}_S^{\text{sat}}$ .