Chapter 7 lecture script: Grouped ecological response curves

Draft for the course author's review. **Target: 150 minutes in total, with 116 minutes of slides on 31 slides (including the discussion, the three practice questions and one live demo of 18 minutes), 14 minutes of slack and two breaks of 10 minutes.** The breaks separate the three blocks of the lecture. The only code that is run live is the demo of slide 22, from the script `demo-pooling.R`.

This script follows the Chapter 7 theory from the introduction to the recipe, with a short introduction to the R practical. One data set runs through the lecture, the CO_2 uptake curves of 12 grass plants: observations in groups and the conditional and marginal distributions that describe them, the three ways of sharing information among groups, the marginal likelihood and its Laplace approximation, the variance components, group-level predictors and shrinkage, the uncertainty of a variance among groups and REML, the curves of observed and new groups, and the checking and comparison of grouped models. The slides carry keywords only; the narrative is here.

Slide 3 uses your diagram `ModelCycle.png` from the Chapter 1 lecture folder. Slides 5 (`co2-data.png`), 14 (`laplace.png`), 20 (`shrinkage.png`), 24 (`new-plants.png`) and 26 (`residuals-grouped.png`) use figures in `lecture-assets/`. All five are unchanged copies of Chapter 7 figures: `fig-co2_data`, `fig-laplace`, `fig-shrinkage`, `fig-new_plants` and `fig-residuals_grouped`. Their code is in `lecture-assets/figures.R`, which runs the chapter's chunks in the chapter's order with the chapter's seed (701), so the residual plot is the one on the chapter page. If the chapter figures change, run that script again (it takes about 13 seconds).

Timing and slide sequence

Times are measured from the start of the session and include the breaks.

Slide	Topic	Minutes	Elapsed time
1	Ecological Models and Data Analysis	1	0–1
2	Overview of the lecture	2	1–3
3	Where we are: the modelling cycle	1	3–4
4	Observations in groups	3	4–7
5	Response curves measured on the same groups	3	7–10
6	Conditional distributions	4	10–14
7	Marginal distributions	4	14–18
8	A two-level model with a random effect	4	18–22

Slide	Topic	Minutes	Elapsed time
9	Fixed effects and random effects	3	22–25
10	Practice 1: two seedlings in one quadrat	3	25–28
11	Three ways to share information	4	28–32
12	Discussion: which pooling answers which question?	3	32–35
	Break	10	35–45
13	The marginal likelihood	4	45–49
14	The Laplace approximation	3	49–52
15	Random effects in RTMB	4	52–56
16	Variance components and predicted effects	4	56–60
17	Group-level predictors	4	60–64
18	The model matrix	4	64–68
19	Two ways to encode groups	3	68–71
20	Shrinkage	4	71–75
21	Practice 2: wing length of chicks in nests	3	75–78
	Break	10	78–88
22	Live demo: from pooling to partial pooling	2 + 16	88–106
23	Uncertainty of variance components and REML	5	106–111
24	Curves for observed and new groups	4	111–115
25	Practice 3: sapling growth in forest stands	3	115–118
26	Checking a grouped model	4	118–122
27	AIC for grouped models	4	122–126
28	Comparing the candidates	3	126–129
29	A recipe for a grouped response curve	2	129–131
30	The Chapter 7 R practical	2	131–133
31	Summary and closing questions	3	133–136
	Slack for discussion	14	136–150
	Total	**150**	

Slides 116 minutes (blocks of 35, 33 and 48, the demo included), slack 14, breaks 20: 150 minutes. The break before the demo lets you open R and run section 0 of `demo-pooling.R`.

Slides 1 and 2 — Ecological Models and Data Analysis; Overview of the lecture

In this lecture, we learn how to fit response curves to observations that come in groups. Repeated measurements on the same plants, animals, nests or sites share the conditions of their group, so they are not independent of each other. A grouped model describes three things together: a curve that is common to all the groups, how that curve varies among groups, and how the observations vary around the curve of their own group. The variation among groups is described by a probability distribution, and the model is fitted by maximum likelihood once the deviations of the groups have been integrated out of the likelihood.

The session has three blocks, separated by two breaks. The first block explains why grouped observations are not independent, introduces conditional and marginal distributions, writes the model with a random effect and compares three ways of sharing information among groups. The second is about the fit: the marginal likelihood, the Laplace approximation, random effects in RTMB, the variance components, group-level predictors and shrinkage. The third starts with a live demo that goes from pooling to partial pooling, then turns to the uncertainty of a variance among groups and REML, the curves of observed and new groups, and the checking and comparison of grouped models. It ends with the recipe, the practical and the summary.

Delivery

Read the overview at the pace of the list, one keyword and its question at a time, and do not explain any term yet.

Slide 3 — Where we are: the modelling cycle

This lecture adds one entry to two boxes of the cycle. In the model box it is "groups (7)": the model now has two levels, a distribution for the groups and a distribution for the observations of each group. In the estimation box it is "marginalise (7)": the effects of the groups are not observed, so they are integrated out of the likelihood before it is maximised. The comparison box keeps the information criteria of Chapters 4 and 6, of which grouped models use only AIC, for reasons that slide 27 explains.

Delivery

One minute. Point to "groups (7)" and "marginalise (7)", and to the comparison box.

Slide 4 — Observations in groups

The **observational unit** is the entity on which one observation is made, such as a quadrat, a visit to a nest or a measurement on a plant. In Chapters 4 and 6 the observations were independent. Often, however, observations come in **groups**: several observations are made on the same individual, at the same site, in the same nest or in the same tank. The observations of one group share conditions that other groups do not share (the genotype and condition of an individual, the microhabitat of a site, the parents of a brood, the observer of a survey), so they tend to resemble each other more than observations of different groups do. The variation among groups is part of the process variation of Chapter 1: it would remain with perfect measurements.

Treating grouped observations as independent overstates the information in the data. Another observation of a group that we have already observed tells us less than an observation of a new group, because part of what it shows is the condition of its group, which we already know something about. A model that treats every observation as independent therefore calculates standard errors that are too small and intervals that are too narrow, and the evidence for an effect looks stronger than it is.

We have met this twice. Exercise 1 of the Chapter 4 practical found that the numbers of tadpoles surviving in the reed frog tanks varied among tanks much more than a Binomial model allows: the tadpoles of a tank share its conditions. And Chapter 6 used the number of tanks, 16, and not the number of tadpoles as the number of observations in AICc and BIC, for the same reason. A grouped model describes the dependence explicitly: each group has a curve of its own, and a distribution describes how the curves vary among groups. That is the takeaway.

Delivery

Ask students for one example of grouped observations from their own field before the example box appears (repeated counts at the same transect, several leaves of one tree, chicks of one nest).

Slide 5 — Response curves measured on the same groups

A common design for response curves measures every group at the same values of the predictor, so that each group gives a whole curve. The slide shows this in general; the example box gives the function. The chapter describes the curves with the saturating exponential shifted along x, $\mu(x) = a\big(1 - e^{-b(x-c)}\big)$, where $\mu(x)$ is the expected response at x, a the asymptote, b the rate at which the curve approaches it (it reaches 63% of the asymptote at $x = c + 1/b$), and c the value of x at which the expected response is zero. A grouped model lets one or more parameters of the curve vary among groups; we choose the one in which the groups differ most.

The data come from Potvin, Lechowicz and Tardif (1990), "The statistical analysis of ecophysiological response curves obtained from experiments involving repeated measures", *Ecology* 71: 1389–1400, and are built into R as `CO2`. They come from an experiment on the cold tolerance of the grass *Echinochloa crus-galli*. Six plants came from Quebec and six from Mississippi, and three plants of each origin were chilled overnight before the measurements. The CO_2 uptake of every plant (μmol/m²/s) was measured at the same seven ambient concentrations of CO_2, from 95 to 1000 μL/L, so the design has 2 origins × 2 treatments × 3 plants × 7 concentrations, 84 measurements in total. The observational unit is one measurement and the group is the plant. The help page of `CO2` gives the concentration in mL/L, but the units stored with the data set are μL/L, which is the same as parts per million; 1000 mL/L would be pure CO_2. Mention this only if a student reads the help page.

The asymptote is the uptake of a plant at high concentrations of CO_2, a measure of its capacity for photosynthesis. In the figure it is the property in which the plants differ most visibly, so it is the parameter that we let vary among plants. In the curve, a is in μmol/m²/s, b in L/μL and c in μL/L.

Figure walkthrough

The four panels are the combinations of origin (columns) and treatment (rows); in each panel the three plants of that combination are blue lines with points, distinguished by line type and symbol. Every curve rises steeply up to about 250 µL/L and then levels off. The plants differ mainly in the level that they reach: the non-chilled plants from Quebec near 40 µmol/m²/s, the chilled plants from Mississippi lowest, near 15 to 22. Within a panel the three plants still differ. Ask students what they would want to know from these data (does chilling lower the uptake, and does it do so for both origins?), and point out that the 84 points are not 84 independent observations: the seven points of a line belong to one plant.

Slide 6 — Conditional distributions

Chapter 2 introduced the joint distribution of two random variables, the marginal distribution of each and independence. The **conditional distribution** of X_1 for a given value x_2 of X_2 gives the probabilities of the values of X_1 among the outcomes in which $X_2 = x_2$. It is the joint probability divided by the marginal probability of the given value, which must be larger than zero. The bar reads "for a given value of", as in the likelihood notation of Chapter 4; what is given now is the value of another random variable.

Two variables are independent when the conditional distribution of X_1 is the same for every value of X_2, and equal to the marginal distribution: knowing X_2 does not change the probabilities of X_1. When they are not independent, the conditional distribution changes with the given value, which Practice 1 (slide 10) shows.

In the table of the two seeds of Chapter 2, with $P(X_1 = 1) = 0.6$ and $P(X_2 = 1) = 0.3$, the joint probability that neither seed germinates is 0.28 and the probability that the second fails is 0.70. So the probability that the first seed does not germinate, for a given failure of the second, is $0.28/0.70 = 0.4$, which equals the marginal probability $P(X_1 = 0) = 0.4$. These seeds are independent.

Delivery

Write the two-seed table on the board if students do not remember it, or show the Chapter 2 page. The only new idea is the division by the marginal probability of the given value.

Slide 7 — Marginal distributions

Multiplying both sides of the definition of slide 6 by $P(X_2 = x_2)$ writes the joint distribution as a conditional distribution times a marginal one. Adding these joint probabilities over all the values of X_2 gives the marginal distribution of X_1, as in the tables of Chapter 2. For continuous random variables the same relations hold with densities in place of probabilities and an integral in place of the sum: the second equation on the slide, where $f_{X|Y}(x \mid y)$ is the conditional density of X for a given value y of Y and $f_Y(y)$ the marginal density of Y. In plain English, the marginal density of X is the average of its conditional densities over all the values of Y, each weighted by the density of that value.

This is the form in which a grouped model is written: first the distribution of the value of a group, then the distribution of the observations for a given value. The marginal distribution of the observations follows by integration.

We have met a marginal distribution of this kind before. The Beta-Binomial distribution of Chapter 6 is the marginal distribution of a count that is Binomial for a given probability, when the probability varies among sampling units according to a Beta distribution: each reed frog tank had its own probability that a tadpole was killed, and the Beta-Binomial distribution describes the counts with those probabilities integrated out. For this combination of distributions the integral has a closed form, which is why we could write its probability mass directly. A **random effect** is the same idea with a Normal distribution for the value of each group, usually on a transformed scale (the logarithm of a positive parameter or the logit of a probability). For most such models the integral has no closed form, and slide 14 shows how it is approximated.

Transition

The next slide writes the CO_2 model in this form.

Slide 8 — A two-level model with a random effect

Let Y_{ij} be the i-th observation of group j and u_j the deviation of group j from the common curve. The deviation acts on one parameter of the curve, on a scale on which a Normal deviation is plausible: the logarithm of a positive parameter or the logit of a probability. The deviations follow a Normal distribution with mean zero and variance σ^2_{group}, and for a given deviation the observations follow a Normal distribution around the curve of their group, $\mu_j(x)$, with variance σ^2_{obs}. For the CO_2 data the parameter is the asymptote, and the curve is the one in the example box. The asymptote of group j is $a_j = a\,e^{u_j}$, that is, $\log(a_j) = \log(a) + u_j$, so every asymptote is positive and the deviations act by multiplication: a group with $u_j = 0.1$ has an asymptote about 10% above a, because $e^{0.1} \approx 1.105$.

The model has two levels. For a given value of u_j, the observations of group j are independent: each varies around the curve of its group with variance σ^2_{obs}. Marginally, that is, without knowing u_j, they are not independent, because they share u_j: an observation above the common curve makes it more probable that the group has a high asymptote, and so that its other observations are also above the common curve. This is the dependence of slide 4, now written as a model. Observations of different groups are independent, because the deviations of different groups are independent. That is the takeaway.

For the CO_2 data, $j = 1, \ldots, 12$ are the plants and $i = 1, \ldots, 7$ the measurements of a plant, x_{ij} is the concentration of measurement i of plant j, and the variance among groups is written σ^2_{plant}. The rate b and the location c are shared by all the plants. Any parameter of a curve can vary among groups in the same way; the chapter lets only the asymptote vary.

Delivery

Point at the bar in $Y_{ij} \mid u_j$ and connect it to slide 6: the observations are described for a given value of the group effect.

Slide 9 — Fixed effects and random effects

The parameters that are shared by all the groups are called **fixed effects**: here a, b and c, and the effects of the group-level predictors that slide 17 adds. The deviations u_j, which are described by a distribution, are the **random effects**, and a model with both is a **mixed-effects model** (or mixed model). Because it describes the data in levels, it is also called a **hierarchical** or **multilevel model**. The three names are used for the same kind of model in different fields.

"Random" says that the deviations are described by a distribution. It does not mean that the plants were allocated at random, nor that their differences have no cause: the plants differ for reasons (genotype, condition), and slide 17 explains part of the differences with predictors. The variances of the model, here σ^2_{plant} and σ^2_{obs}, are estimated together with the fixed effects; the random effects themselves are not estimated, as slides 13 and 16 explain.

Slide 10 — Practice 1: two seedlings in one quadrat

Short activity — allow about two minutes of work and one to discuss

Students work on paper. Expected answers, from the chapter's solution:

1. For a given quality the seedlings are independent, so their joint probability is the product of their probabilities. The quality was not recorded, so the products are added over the two qualities, each weighted by its probability; this is the discrete version of the integral of slide 7. $P(X_1 = 1, X_2 = 1) = 0.5 \times 0.8 \times 0.8 + 0.5 \times 0.2 \times 0.2 = 0.32 + 0.02 = 0.34$; in the same way $P(X_1 = 1, X_2 = 0) = P(X_1 = 0, X_2 = 1) = 0.08 + 0.08 = 0.16$ and $P(X_1 = 0, X_2 = 0) = 0.02 + 0.32 = 0.34$, which add up to one. Then $P(X_1 = 1) = 0.34 + 0.16 = 0.5$.
2. $P(X_2 = 1) = 0.5$ as well, so $P(X_1 = 1 \mid X_2 = 1) = 0.34/0.5 = 0.68$.
3. The seedlings are not independent: the survival of one makes it more probable that the quadrat is favourable, and so that the other survives too (0.68 against 0.5). For a given quality they are independent, by construction.

The third answer is the structure of a grouped model: independence for a given value of the group effect, and dependence when the group effect is integrated out. Students who calculate $P(X_1 = 1)P(X_2 = 1) = 0.25$ for the joint probability have assumed the independence that the question asks about.

Slide 11 — Three ways to share information

When a parameter of a curve may differ among groups, a model can treat it in three ways. With **complete pooling**, all the groups share one value of the parameter: the groups are ignored and the observations are treated as independent, as in Chapter 6. With **no pooling**, each group has its own value, an ordinary parameter estimated from the observations of that group, and nothing connects the values of different groups. With **partial pooling**, the values of the groups are drawn from a distribution whose variance is estimated, as in the model of slide 8: each group has its own value, the values of all the groups inform the distribution, and the distribution in turn informs the value of each group. That is the takeaway.

The three ways differ in what they assume and in the question that they answer. Complete pooling assumes that the groups do not differ. When they do, the observation variance absorbs the variation among groups, and the standard errors are too small because every observation is treated as independent. No pooling describes the measured groups, and only them: it says how each of these groups responds, but nothing about a group that was not measured, because it does not describe how groups vary. Partial pooling describes the measured groups and, through the distribution of the group values, groups in general, which is usually the question of an ecological study (plants of this species, nests of this bird, sites of this habitat). Its costs are an assumption about the shape of that distribution, which is Normal in this chapter, and an integral in the likelihood (slides 13 and 14).

For the CO_2 data, complete pooling fits one curve to all 84 measurements, and the measurements vary around it with a standard deviation of 8.2 μmol/m²/s. Without pooling, each plant has its own asymptote, and the measurements vary around the curves of their own plants with a standard deviation of 1.9 μmol/m²/s. So most of the scatter around the common curve is variation among plants, which complete pooling counts as variation of the measurements. The asymptotes without pooling range from 14.2 μmol/m²/s (plant Mc2) to 45.0 μmol/m²/s (Qn3), and the three chilled plants from Mississippi have the lowest ones. No pooling describes these 12 plants, but it cannot say how much plants of this grass vary in their capacity for uptake, or which curve to expect for a plant that was not measured.

Delivery

The two NLLs are in the chapter ("Three ways to share information") and are run in sections 1 and 2 of the demo; do not show code here. If a student asks: the no-pooling NLL makes `log_asym` a vector with one value per plant and selects the asymptote of each measurement with `a[plant]`.

Slide 12 — Discussion: which pooling answers which question?

This discussion applies the three ways of slide 11 to questions about a survey. The ponds are a hypothetical example for the discussion.

Discussion and expected answers — included in the three minutes

Give students two minutes to match the questions with a neighbour, then take one answer per item.

1. **Which of these 20 ponds has the most newts?** A question about the surveyed ponds, so **no pooling** answers it, and so does **partial pooling**, whose predicted pond values are shrunk towards the typical pond (slide 20) and are less sensitive to chance variation in ponds with few or noisy counts. Complete pooling cannot answer it, because it gives every pond the same density.
2. **How much does the density vary among the ponds of the region? Partial pooling**: the variance among ponds is a parameter of the model. No pooling describes only these 20 ponds and no distribution of ponds; complete pooling assumes that ponds do not differ.
3. **How many newts in a pond that was not surveyed? Partial pooling**: the typical pond and the range of new ponds (slide 24). No pooling has no value for a pond that was not surveyed.

4. **Does the density increase with pond area, and how precisely do we know it?** Area is measured on the ponds, a group-level predictor (slide 17), so **partial pooling with area as a fixed effect**. Complete pooling with area would treat the 100 counts as independent and overstate the precision of the effect (slide 4). No pooling cannot estimate it at all, because the value of each pond absorbs everything that differs among ponds, including their area.

Accept "both no and partial pooling" for the first question if students say why. The fourth answer is the one that students usually miss.

Break (10 minutes).

Slide 13 — The marginal likelihood

In a mixed model the random effects are not observed. The likelihood of the parameters $\boldsymbol{\theta}$, which are the fixed effects and the variances of the model, is therefore the probability of the observations with the random effects integrated out, that is, their marginal distribution (slide 7). For group j, with its observations collected in the vector $\mathbf{y}_j$, this is the integral on the slide. In it, $f\left(\mathbf{y}_j \mid u, \boldsymbol{\theta}\right)$ is the joint density of the observations of group j for a given value u of its random effect (the product of the densities of its observations, which are independent for a given u), and $f\left(u \mid \sigma_{\text{group}}\right)$ is the Normal density of the random effect, with mean zero and standard deviation σ_{group}. The groups are independent, so the likelihood is the product of the L_j over the groups, and the log-likelihood is the sum of their logarithms. This is the **marginal likelihood**, and its maximisation is called **maximum marginal likelihood**. In plain English, the marginal likelihood of a group averages the likelihood of its observations over all the values that its random effect could take, each weighted by its density under the distribution of group effects.

Why not estimate the deviations as parameters? That is no pooling: the u_j would be ordinary parameters, and no distribution would connect them. In the marginal likelihood the deviations are replaced by one parameter, the standard deviation among groups, which describes how groups vary. The number of parameters then stays the same however many groups are observed, which is one of the conditions under which the large-sample properties of maximum likelihood of Chapter 4 hold.

For the CO_2 data, $\boldsymbol{\theta}$ holds a, b, c, σ_{plant} and σ_{obs}, and there are 12 integrals, one for each plant, each over the deviation of that plant.

Slide 14 — The Laplace approximation

The integral of the marginal likelihood has a closed form only in special cases, such as the Beta-Binomial distribution and the Normal linear mixed model, so in general it must be approximated. The **Laplace approximation** uses the shape of the integrand. As a function of u, the integrand is the likelihood of the observations of the group multiplied by the density of u. It has a peak at the value of u that is most probable for these observations, and near the peak its logarithm is close to a parabola, so the integrand itself is close to a Normal-shaped curve. The area under a Normal-shaped curve is known: it depends only on the height of the peak and on the curvature of the logarithm at the peak. So the approximation finds the peak, measures its height and its curvature,

and replaces the integral by the area under the Normal-shaped curve with the same peak and curvature. This is the quadratic approximation of the log-likelihood that gave the standard errors of Chapter 4, applied to the integrand.

The approximation is accurate when the integrand is close to a Normal-shaped curve: when each group has many observations, so that the likelihood of the group dominates the integrand, and when the observations are Normal, or close to Normal, for a given value of the random effect. For Normal linear mixed models it is exact, because there the logarithm of the integrand is a parabola. It is less accurate when the groups have few observations, and when the observations are counts near zero or binary outcomes, for which the integrand can be skewed. The practical fits models of both kinds (one Binomial count per tank of tadpoles, Poisson counts of salamanders with four visits per site).

Figure walkthrough

The figure shows the integrand for plant Mc2, the plant with the lowest uptake, with the parameters at the estimates of the model that slide 15 fits. The horizontal axis is the deviation of Mc2 on the log scale of the asymptote, and both curves are scaled so that their peak is one. The solid line is the integrand; the dashed line is the Normal-shaped curve with the same peak and the same curvature of the logarithm at the peak. The two curves almost coincide; the integrand has a slightly longer tail towards low values of u, which the symmetric approximation cannot follow. As a result the area under the approximation is 0.3% smaller than the area under the integrand, which changes the log-likelihood of this plant by 0.003. The figure is an illustration: the chapter draws it in a hidden chunk, and in practice no one calculates the integral, because RTMB does it (the takeaway).

Optional remark

The chapter gives the formula of the Laplace approximation in a box; mention it only if asked.

Slide 15 — Random effects in RTMB

In RTMB we do not write the marginal likelihood ourselves. We write the **joint NLL** of the observations and the random effects, minus the logarithm of their joint density: minus the log density of the observations for given random effects plus minus the log density of the random effects (the joint density is the conditional density times the marginal one, slide 7). `MakeADFun()` then integrates the random effects out. Three things change compared with the NLLs of Chapters 4 and 6. The random effects are an element of the list of parameters, a vector with one starting value for each group, usually zero. The NLL adds minus the log density of the random effects to minus the log density of the observations, and the line with the density of the random effects comes before the line that uses them; we need this order to simulate from the model (slide 25). And the argument `random` of `MakeADFun()` names the elements of the list of parameters that are random effects. RTMB then integrates them out with the Laplace approximation: the elements `fn` and `gr` of the object are the marginal NLL and its gradient, as functions of the other parameters only, and `par` holds only those other parameters. So the fit with `optim()` works as before.

For the CO_2 data, the NLL is the no-pooling NLL with the vector of log asymptotes replaced by one parameter `log_asym` plus the deviations `u`, and with the density of `u`, whose standard deviation σ_{plant} is optimised as `log_sd_plant`. The deviations start at zero, and σ_{plant} at 0.1 (plants that differ by about 10%). The chunk on the slide writes the standard deviations inline to fit the slide; the chapter names them first. The fit converges, and `par` holds five estimates; the deviations of the plants are not among them.

Delivery

Show the chunk without running it; section 3 of the demo runs it. Point at the first line of the body (the density of `u`), at `a[plant]` and at `random = "u"`. The vector `plant` is an integer index: it gives the group of each observation, so `a[plant]` picks the asymptote of the plant of each measurement (the comment in the chunk says so).

Optional remark

The chapter describes in a box the inner optimisation that RTMB carries out for the random effects at every evaluation of the NLL; mention it only if asked.

Slide 16 — Variance components and predicted effects

The variances σ^2_{group} and σ^2_{obs} are the **variance components** of the model: the first describes the variation among groups, the second the variation of the observations around the curve of their group. Each has the scale and the units of the quantity that it describes, and the NLL takes their square roots, the standard deviations. `sdreport()` works on the marginal NLL as on any other NLL, and two values of the argument `select` of `summary()` give the two kinds of output: `"fixed"` gives the estimates and standard errors of the parameters $\boldsymbol{\theta}$ on the optimised scales, and `"random"` gives the **predicted group effects**, one row for each group, with their standard errors.

The predicted effect of a group is the peak of its integrand (slide 14), the most probable value of its random effect for the observations of the group and the estimates of $\boldsymbol{\theta}$; it is also called the conditional mode. The random effects are not parameters, so the predicted effects are not estimated by maximum likelihood: they are predictions of unobserved quantities. Their standard errors combine the uncertainty about the effect of a group for given parameter values with the uncertainty of the estimates of the parameters.

A standard deviation on the log scale has no units, and for small values it is approximately the coefficient of variation of the quantity on the original scale (its standard deviation divided by its mean). The curve with $u_j = 0$ has the median asymptote of groups in general: u_j is symmetric around zero and the exponential is increasing, so half of the groups lie above it and half below. It is not their mean, which is larger, because the exponential is not a straight line (the box on Jensen's inequality in Chapter 4).

For the CO_2 data, the measurements vary around the curve of their plant with a standard deviation of about 2.0 µmol/m²/s, close to the 1.9 of no pooling and far from the 8.2 of complete pooling: the mixed model, like no pooling, separates the variation among plants from the variation of the measurements. $\hat{\sigma}_{\text{plant}} \approx 0.34$ is not very small, and it means that the asymptotes of plants vary

around their median by roughly a third. The estimate $\hat{a} \approx 30.9$ µmol/m²/s is the median asymptote of plants in general. The Quebec plants have positive predicted effects, and the chilled plants from Mississippi (Mc1, Mc2 and Mc3) the most negative ones, -0.370, -0.755 and -0.422: their asymptotes are about 31%, 53% and 34% below the median ($e^{-0.370} \approx 0.69$, $e^{-0.755} \approx 0.47$, $e^{-0.422} \approx 0.66$). So much of the variation among plants follows their origin and treatment, which the model does not know about yet.

Delivery

Show the chunk without running it; the numbers are in its comments, and section 3 of the demo prints them. The point to stress is the second bullet: a predicted effect is a prediction, not an estimate.

Transition

The next slide gives the model what it does not know yet: the origin and treatment of each plant.

Slide 17 — Group-level predictors

A **group-level predictor** is measured on the group and not on each observation: the origin of a plant, the treatment that it received, the habitat of a site. All the observations of a group share its value. In a mixed model it enters as a fixed effect on the parameter that varies among groups, here on its log scale, where it explains part of the variation among groups. On the log scale the effects add up, so on the original scale they multiply. The random effects then describe the variation that remains among groups with the same values of the predictors, and the variance among groups becomes smaller. That is the takeaway.

When there are two predictors, the effect of one may depend on the level of the other: that is an **interaction**. The chapter writes the effects of origin and chilling as a baseline plus contrasts, with an interaction: one combination of levels is the baseline, and an indicator that is 1 or 0 switches each effect on or off. In the equation of the example, M_j is 1 for a plant from Mississippi and 0 for a plant from Quebec, C_j is 1 for a chilled plant and 0 otherwise, and a is now the typical asymptote (with $u_j = 0$) of the baseline group, the non-chilled plants from Quebec. β_{miss} is the effect of the origin in non-chilled plants, β_{chill} the effect of chilling in plants from Quebec, and β_{int} the difference between the effect of chilling in plants from Mississippi and in plants from Quebec. Chilling multiplies the typical asymptote of a Quebec plant by $e^{\beta_{\text{chill}}}$ and that of a Mississippi plant by $e^{\beta_{\text{chill}}+\beta_{\text{int}}}$. The next slide shows how R builds the indicators. A quantity calculated from the coefficients, such as the logarithm of a chilling factor, is reported by the NLL with `ADREPORT()`; for a positive quantity the Wald interval is calculated on the log scale and its endpoints are transformed back with `exp()`.

With origin and chilling in the model, $\hat{\sigma}_{\text{plant}}$ falls from 0.34 to 0.079, so the variance among plants on the log scale falls by about 94%. Origin and chilling explain most of the variation among plants, and the asymptotes of plants of the same origin and treatment differ by roughly 8%. Chilling multiplies the typical asymptote of plants from Quebec by about 0.90, with a Wald interval from the log scale of 0.79 to 1.03, and that of plants from Mississippi by about 0.59 (0.52 to 0.68). So the plants from the colder origin are hardly affected by a cold night, whereas the plants from the

warmer origin lose about two fifths of their capacity. Slide 23 returns to the uncertainty of these effects.

Delivery

The code is on the chapter page ("Group-level predictors and shrinkage", subsection "Effects of origin and chilling") and runs in section 4 of the demo. Read the equation term by term, and ask which term would be zero if a cold night affected the plants of both origins in the same way (β_{int}).

Transition

We could calculate M_j and C_j ourselves, but R builds them, and the whole sum, from a formula.

Slide 18 — The model matrix

The **model matrix** (also called the design matrix) of a set of predictors has one row for each group, or for each observation when the predictors are measured on the observations, and one column for each coefficient. Row j holds the values by which the coefficients are multiplied for group j: for the equation of slide 17, 1 (for the baseline $\log(a)$), M_j, C_j and M_jC_j. R builds the whole matrix from a **formula** with `model.matrix()`, and from now on the chapter lets it do this bookkeeping in every model with predictors.

`model.matrix()` takes a formula and a data frame. The predictors are properties of the plants, so `plant_info` has one row per plant, with the origin and treatment from its first measurement (the chapter also names its rows after the plants). `Type * Treatment` is short for `Type + Treatment + Type:Treatment`. The result, `X_asym`, has 12 rows and four columns. The first, `(Intercept)`, is 1 for every plant. The name of each other column joins a factor and the level for which the column is 1: `TypeMississippi` is M_j, `Treatmentchilled` is C_j, and `TypeMississippi:Treatmentchilled` is M_jC_j.

The first level of each factor is its **reference level**, the level of the baseline, and it gets no column. `levels()` shows the order of the levels, and `relevel()` makes another level the first one. The model would be the same, but its coefficients would be other differences, so we choose the reference levels against which the question compares the others: here, plants from Quebec, the colder origin, that were not chilled.

In the NLL, `X_asym %*% beta_asym` is a **matrix product**: it multiplies each column by its coefficient and adds up the columns, so row j is the sum of the equation without u_j. For Qn1 the row is $(1, 0, 0, 0)$ and the product is $\log(a)$; for Mc1 the row is $(1, 1, 1, 1)$ and the product is $\log(a) + \beta_{\text{miss}} + \beta_{\text{chill}} + \beta_{\text{int}}$. The starting values, `c(log(30), rep(0, ncol(X_asym) - 1))`, follow the number of columns, so a change of formula needs no change of code.

The functions of R that fit models from a formula, such as `lm()`, `glm()` and `glmmTMB()`, build the same matrix, name their coefficients after its columns and follow the same rules for reference levels. In RTMB we build it ourselves, which lets us put it on the log asymptote of a non-linear curve.

Delivery

Point at the two lines of code: the matrix is built once, outside the NLL, and the NLL holds only the product. If students ask what `%*%` does, write the row of Mc1 times the four coefficients on the board. Mention the warning callout of the chapter in one sentence: check the levels, because `factor()` sorts them alphabetically, and convert an ordered factor with `factor(x, ordered = FALSE)`. Section 4 of the demo prints the levels and four rows of `X_asym`.

Slide 19 — Two ways to encode groups

The effects of groups defined by predictors, such as the four combinations of origin and treatment, can be written in two ways. With one parameter for each group, each group has its own typical value (here, four typical log asymptotes). With a baseline plus effects and interactions, one group is the baseline and the other parameters are differences from it (here, $\log(a)$, β_{miss}, β_{chill} and β_{int}). In `model.matrix()`, the formula `~ 0 + Type:Treatment` gives the first form, one column for each combination and no intercept (the `0`), and `~ Type * Treatment` the second. Both have the same number of parameters and describe the same model: they have the same maximum likelihood and give the same fitted curves, and each set of parameters can be calculated from the other. In the example, both forms reach a minimum NLL of 192.086.

The second form makes the differences between treatments parameters, so their estimates, standard errors and intervals come directly from the fit, and a simpler model is obtained by leaving a term out of the formula (the interaction, for example; slide 28 does that). The first form makes the typical value of each group a parameter. Which one to use depends on the question: for the CO_2 experiment, whose question is how chilling affects the two origins, the differences are what we want to report.

Delivery

The two columns appear one at a time. This is the callout "Two ways to encode groups" of the chapter. The slide writes the formulas with general factors `A` and `B`; the example box gives them for origin and treatment. Both encodings describe the same model, so they reach the same maximum: an NLL of 192.086, minus the log-likelihood of the model with the interaction in the chapter's comparison table (-192.09).

Slide 20 — Shrinkage

The predicted effect of a group is a compromise between the observations of that group and the distribution of groups. Its observations alone would give the no-pooling estimate; the distribution of group effects pulls the prediction towards the typical value of groups with the same predictors. This pull is called **shrinkage**. It is stronger when the group has few observations, when its observations are noisy compared with what they need to determine, and when the variance among groups is small, because the distribution then says that groups are alike. A group with many precise observations stays close to its own estimate. Shrinkage is a form of **regularisation**: the values of many groups are pulled towards a common value, which makes each of them less sensitive to the noise in the observations of its group. In a mixed model the shrinkage follows from the model itself, through the distribution of the group effects; it is not a correction applied after the fit.

Figure walkthrough

For each plant, the open blue circle is its asymptote without pooling and the filled vermillion circle its predicted asymptote under a mixed model; a grey line joins them where they differ visibly. Top panel: the model with a random asymptote only, whose dashed line is the median asymptote of plants in general, 30.9 μmol/m²/s. The two circles nearly coincide for every plant: every plant has seven precise measurements and $\hat{\sigma}_{\text{plant}}$ is large, so the measurements of each plant dominate its prediction. The largest change is that of Mc2, from 14.2 to 14.5 μmol/m²/s. Bottom panel: the model with origin and chilling, whose dashed lines are the typical asymptotes of the four combinations (about 42, 38, 31 and 18.5 μmol/m²/s). Now the distribution of plant effects is narrow, because plants of the same origin and treatment are similar, and the shrinkage is visible: Mc2 moves from 14.2 to 15.6 μmol/m²/s, about a third of the way towards the typical 18.5 of chilled plants from Mississippi. Mc2 moves most because it lies farthest below the typical asymptote of its group, and because its uptake is the lowest of all: the same observation error of about 2 μmol/m²/s is a larger fraction of its uptake, so its measurements determine its asymptote on the log scale less precisely.

Optional remark

The chapter gives the shrinkage weight of a Normal linear model in a box; Practice 2 uses it.

Slide 21 — Practice 2: wing length of chicks in nests

Short activity — allow about two minutes of work and one to discuss

Students use R as a calculator. Expected answers, from the chapter's solution:

1. Nest A is pulled more, because two chicks say less about their nest than eight do.
2. With $\sigma^2_{\text{nest}} = 4$ mm² the weights are $8/12 \approx 0.67$ (A) and $32/36 \approx 0.89$ (B), so the predicted deviations are 2.0 mm and 2.7 mm. In R: `3 * c(2, 8) * 4 / (c(2, 8) * 4 + 4)`.
3. With $\sigma^2_{\text{nest}} = 1$ mm² the weights are $2/6 \approx 0.33$ and $8/12 \approx 0.67$, and the predicted deviations 1.0 mm and 2.0 mm.

When nests vary less, the model expects a deviation of 3 mm to come partly from the chance variation of the few chicks measured, so both nests are pulled more strongly towards the common mean, and the nest with fewer chicks most of all. These are the three conditions of slide 20 in numbers.

Break (10 minutes).

Slide 22 — Live demo: from pooling to partial pooling

The demo runs the steps of slides 11 to 20 for the CO_2 data, from complete pooling to the model with origin and chilling, and ends with the curves of observed and new plants, which slide 24 explains. It has no model comparison: the comparison follows on slides 27 and 28. The script needs the packages RTMB and RTMBdist; the data set `CO2` is built into R. Its code and object names are those of the chapter, so the numbers agree with the chapter and the slides. The whole script runs in a few seconds (1.5 seconds in a check in R 4.6.1), so there is no waiting.

During the break, open R and run section 0, “Packages and data”. It loads RTMB and RTMBdist (with their messages about masked functions), prints the first rows of the data and the names of the 12 plants, builds the list of data `co2_data`, draws the 12 plants in base R and defines `cmb()`. Keep the session and the plot open. Spend the two minutes at the slide reading the five steps, then switch to R and run the sections one at a time.

Walkthrough of the script — 16 minutes

1. **Complete pooling (about 2 minutes).** Start from the plot of section 0: solid lines are non-chilled plants, dashed lines chilled ones, circles Quebec and triangles Mississippi; the dashed lines with triangles lie lowest. Define the NLL with one curve and fit it from the values read from the plot (asymptote 30, zero uptake at 50 µL/L, rate 0.01). Convergence code 0; the standard deviation of the measurements around the common curve is 8.23 µmol/m²/s.
2. **No pooling (about 2 minutes).** The same NLL with `log_asym` as a vector of 12 and `a[plant]`; every plant starts at the same asymptote. Convergence code 0; the standard deviation around the curves of the plants is 1.86. The 12 asymptotes run from 14.2 (Mc2) to 45.0 (Qn3), and the three chilled Mississippi plants have the lowest (21.3, 14.2 and 20.1). Ask the first question below.
3. **Partial pooling (about 4 minutes).** Define the joint NLL and point at the order: the density of `u` first, then `a = exp(log_asym + u)`. Process it with `random = "u"` and print `mixed_obj$par`: five starting values, and no `u`. Fit: convergence code 0; on the original scale a is 30.9, b 0.0096 L/µL, c 48.4 µL/L, `sd_plant` 0.335 (log scale) and `sd_obs` 2.01 µmol/m²/s. `summary(mixed_sd, select = "fixed")` gives the standard errors on the optimised scales (0.21 for `log_sd_plant`). The predicted plant effects: 0.145 to 0.372 for the Quebec plants, close to zero for the non-chilled Mississippi plants (-0.072 to 0.054), and -0.370, -0.755 and -0.422 for Mc1 to Mc3, each with a standard error of about 0.10 to 0.11. Ask the second question below.
4. **Origin and chilling, and shrinkage (about 5 minutes).** Build `plant_info`, one row per plant, and print the levels of the two factors: `Quebec` and `nonchilled` come first, so they are the baseline. Print the rows of `X_asym` for the first plant of each origin and treatment: $(1, 0, 0, 0)$ for Qn1 up to $(1, 1, 1, 1)$ for Mc1, with the column names of slide 18. Build `group_info`, the four combinations, and its model matrix `X_asym_group`, and add both matrices to the list of data. The NLL computes `log_a = X_asym %*% beta_asym + u` and the two log chilling factors from the typical log asymptotes of the four combinations, `X_asym_group %*% beta_asym` (chilled minus non-chilled, for each origin), which it reports with `ADREPORT()`. Point at the starting value of `beta_asym`, one element per column of the model matrix. Convergence code 0. The table of estimates, with the rows of `beta_asym` named after the columns of `X_asym`: `TypeMississippi` -0.31, `Treatmentchilled` -0.10 and `TypeMississippi:Treatmentchilled` -0.42 (standard error 0.10); the rate, the location and `log_sd_obs` hardly change. The standard deviation among plants falls from 0.335 to 0.079. The chilling factors: Quebec 0.90 (0.79 to 1.03), Mississippi 0.59 (0.52 to 0.68). The table of asymptotes shows the typical values of the four groups (42.2, 38.2, 31.1 and 18.5), each plant's row of `X_asym` times the estimates, and Mc2 at 14.2 without pooling and 15.6 predicted. The plot shows the shrinkage in base R: open circles without pooling, filled vermillion circles predicted, dashed lines typical. Ask the third question below.

5. **Observed and new plants (about 3 minutes).** The typical asymptotes of the four combinations come from `X_asym_group` times the estimates; this is what `predict()` with `newdata` does for the models that R fits from a formula. The typical asymptote of chilled Mississippi plants is 18.46 µmol/m²/s, and 95% of new plants of this group have asymptotes between 15.81 and 21.55. The plot draws the measurements of Mc1 to Mc3, the curve of each from its predicted asymptote, the typical curve (thick, vermillion) and the range of new plants (dashed). Point out that the curve of Mc2 lies above its own measurements and just below the range of new plants. End by saying that the curves of the measured plants and the curves of new plants answer different questions (slide 24).

Questions to ask during the demo (about 1 minute, within the sections above)

1. In section 2, "Why is the standard deviation so much smaller without pooling?" Because complete pooling counts the differences among plants as variation of the measurements; most of the 8.2 is variation among plants (slide 11).
2. In section 3, "Why does `par` have five elements and not 17?" The 12 deviations are integrated out of the likelihood; they cost one parameter, σ_{plant} (slides 13 and 27).
3. In section 4, "Why does the standard deviation among plants fall, but not the one of the measurements?" The predictors act on the asymptote, a property of the plants, so they explain variation among plants and nothing within a plant (slide 17).

Slide 23 — Uncertainty of variance components and REML

`sdreport()` and `TMB::tmbprofile()` work on the marginal NLL as on any other NLL; each evaluation of the NLL during a profile contains a Laplace approximation, so a profile takes longer, but it needs no new code. Variance components need care. A variance is bounded below by zero, so we optimise the logarithm of the standard deviation. The observations inform the variance among groups only through the deviations of the groups, and part of the information in these is used up by the fixed effects on the parameter that varies, so a variance among groups is usually estimated much less precisely than the fixed effects. And its log-likelihood is often asymmetric, the more so when the estimate is close to zero. For variance components we therefore use the likelihood profile interval of the log standard deviation, and transform its endpoints back: `exp()` gives the interval of the standard deviation and squaring that gives the interval of the variance, because both functions are increasing.

Maximum likelihood estimates of variances also tend to be too small when the fixed effects are estimated from the same data. Chapter 3 met the simplest case: for i.i.d. Normal observations the maximum likelihood estimate of the variance divides by n and is too small on average, because the deviations are taken from the estimated mean, which lies closer to the observations than the mean of the model; dividing by $n - 1$ removes the bias. In a grouped model the same happens at the level of the groups: the fixed effects of the group-level predictors are fitted to the deviations of the groups, so these look smaller than they are, and the bias grows with the number of fixed effects estimated from the groups and falls with the number of groups. **Restricted maximum likelihood** (REML) estimates the variance components from a likelihood in which the fixed effects have been integrated out as well, as if they were random effects with a flat distribution. For

i.i.d. Normal observations it gives the estimator that divides by $n - 1$, and for Normal linear mixed models it is the standard method. In RTMB, REML is one change to the call of `MakeADFun()`: the fixed effects are added to `random`, and only the variance parameters are left for `optim()`. REML was introduced by Patterson and Thompson (1971), "Recovery of inter-block information when block sizes are unequal", *Biometrika* 58: 545–554.

Two cautions apply. For non-linear curves or observations that are not Normal, REML through the Laplace approximation is itself an approximation, and the reduction of the bias is not guaranteed. And the REML likelihood depends on which fixed effects were integrated out, so models with different fixed effects are compared with their maximum likelihood fits. REML is worth its extra fit when the variance components are the question.

For the CO_2 data, the standard deviation among plants of the same origin and treatment lies between 0.045 and 0.139 (likelihood profile interval), so the asymptotes of such plants differ by between about 5% and 14%. The interval of the variance, from 0.0020 to 0.019, spans almost a factor of ten, although the model was fitted to 84 measurements: the variance among plants is informed by 12 plants. The REML estimate of σ_{plant}, 0.105, is a third larger than the maximum likelihood estimate of 0.079: plants of the same origin and treatment differ by roughly 10% instead of 8%. The estimate of σ_{obs} hardly changes (2.05 against 2.04 µmol/m²/s), because the measurements around the curve of each plant are many compared with the parameters of the curve. The takeaway is the practical consequence: to estimate a variance among groups more precisely, observe more groups.

Delivery

The code is on the chapter page ("Uncertainty in a grouped model"); it is not run live. If a student asks about the interaction: its profile interval is -0.64 to -0.21, so the chilling factor of Mississippi plants is between 0.53 and 0.81 times that of Quebec plants. Exercise 5 of the practical checks the bias of both estimators by simulation.

Slide 24 — Curves for observed and new groups

A fitted grouped model gives three kinds of curve, which answer different questions. The **curve of an observed group** uses the predicted effect of that group, and describes the group that was measured, shrunk towards its typical value (slide 20). The **typical curve** has $u_j = 0$ and describes the median group with given values of the predictors; it is not the mean group (slide 16). The curve of a **new group**, one that was not measured, has an effect that is a new draw from the distribution of group effects, so we can only give the range in which it is expected to lie. With a Normal distribution of deviations, 95% of new groups have deviations within $\pm 1.96\,\sigma_{\text{group}}$, so their asymptotes lie between $a_{\text{typ}} e^{-1.96\,\sigma_{\text{group}}}$ and $a_{\text{typ}} e^{1.96\,\sigma_{\text{group}}}$, where a_{typ} is the typical asymptote for their predictors. The curve increases with u at every value of x, so these two curves are the 2.5% and 97.5% quantiles of the curves of new groups at every x. The range is calculated at the estimates and leaves out the uncertainty of the estimates themselves, which would widen it.

Questions about the measured groups ("how does this plant respond?") use their own curves; questions about groups in general ("which curve should we expect for a plant from Mississippi after a cold night?") use the typical curve and the range of new groups. That is the takeaway.

For the CO_2 data, a new chilled plant from Mississippi is expected to have an asymptote between 15.8 and 21.5 μmol/m²/s around a typical 18.5, and a new non-chilled plant from Quebec one between 36 and 49 around 42. The ranges are narrow because plants of the same origin and treatment are similar. The mean asymptote of plants is $a_{\text{typ}} e^{\sigma^2_{\text{plant}}/2}$ (the mean of a LogNormal distribution), only 0.3% above the median for $\hat{\sigma}_{\text{plant}} = 0.079$; the difference grows quickly with the variance among groups, as Practice 3 shows.

Figure walkthrough

The figure shows the three chilled plants from Mississippi: their measurements as blue points, with a different symbol for each plant, and the curve of each plant from its predicted deviation as a blue line. The thick vermillion line is the typical curve of this origin and treatment, and the light vermillion band the range that contains the curves of 95% of new plants. The curves of Mc1 and Mc3 lie above the typical curve (predicted asymptotes 20.6 and 19.7). The curve of Mc2 lies just below the band and above its own measurements, because its predicted asymptote is shrunk towards the typical one. The band is not a confidence band (Chapter 6): it describes how plants vary, not how precisely the typical curve is estimated.

Slide 25 — Practice 3: sapling growth in forest stands

Short activity — allow about two minutes of work and one to discuss

Students use R as a calculator. Expected answers, from the chapter's solution:

1. $20\, e^{-1.96\times 0.4} \approx 9.1$ to $20\, e^{1.96\times 0.4} \approx 43.8$ cm/year, which is wide because σ_{stand} is large. In R: `20 * exp(qnorm(p = c(0.025, 0.975)) * 0.4)`.
2. $20\, e^{0.5} \approx 33.0$ cm/year. That value describes this stand only. For a stand that was not surveyed we know only the distribution of stand effects, so we expect a typical growth rate of 20 cm/year, anywhere between about 9 and 44 cm/year.
3. No: 20 cm/year is the median. The exponential stretches the stands above the median more than it compresses those below, so the mean is larger. Simulating one million stands with `mean(20 * exp(rnorm(n = 1e6, mean = 0, sd = 0.4)))` gives about 21.6 cm/year (21.65 with seed 702), 8% above the median. The mean of the LogNormal distribution, $20\, e^{0.4^2/2}$, is 21.67 cm/year.

The second answer is the point of slide 24: the curve of an observed group answers a question about that group, and the typical curve with its range a question about groups in general.

Slide 26 — Checking a grouped model

A fitted grouped model is a data generating process with two levels: first the effects of the groups are drawn from their distribution, and then the observations are drawn for these effects. Checking the model means comparing data simulated in this way with the observations, with scaled quantile residuals as in Chapter 6. The observations are marked with `OBS()` in the NLL, and `obj$simulate()` then draws both levels: new effects for all the groups, and new observations for these effects. This is called **unconditional simulation**, because the effects of the groups are not kept at their predicted values. RTMB simulates in the order of the lines of the NLL: at the line with the density of the random effects it draws new values of them, and the lines below use these val-

ues, which is why that line must come first (slide 15). Keeping the effects at their predicted values instead, **conditional simulation**, simulates new observations of the same groups; it checks the curves of the observed groups and not the distribution of group effects.

The residuals of the observations come from `DHARMa::createDHARMa()` as in Chapter 6, with the typical curve of each observation as `fittedPredictedResponse`, because that is its expected value before the effect of its group is known. They share the deviation of their group, so they are not independent values, and a pattern among them can come from a few groups. The residuals of the groups come from `DHARMa::recalculateResiduals()`, whose argument `group` gives the group of each observation: it adds up the observed values of each group, and the simulated values of each group in each hypothetical sample, and calculates one residual per group from these sums.

The chapter's warning is the takeaway. A grouped model can disagree with the data at either level, and the residuals of one level can hide a problem at the other. Residuals of the observations mix the deviation of their group with their own variation, so a curve that misses some groups may look acceptable, and a pattern that looks like a misfit of the curve may come from a few groups. Residuals of the groups summarise each group in one number, so they cannot show a problem with the shape of the curve within groups. Check both. With few groups, a quiet plot of the group residuals says little about whether the distribution of group effects is Normal.

For the CO_2 data, the measurements vary around their typical curves about 80% as much (as a variance) as in the simulated samples: dispersion statistic 0.80. Most of this variation comes from the deviations of 12 plants, which vary a lot from one hypothetical sample to the next, so this is not a clear disagreement. The residual of Mc2 as a group is 0.010: its total uptake is lower than in 99% of the simulated plants of its origin and treatment. Among 12 plants one residual this extreme is not unusual, so this is a pattern to note, not a sign that the model fails. With more plants in this group, several of them in the tails would suggest that the distribution of plant effects should be wider or have a longer tail than the Normal distribution allows.

Figure walkthrough

The figure is DHARMa's own plot of the 84 measurements under the model with origin and chilling, from 1,000 hypothetical samples with new plant deviations (seed 701). On the left, the quantile–quantile plot follows the diagonal. On the right, the residuals against the rank of the typical curve of each measurement, with dashed lines at 0.25, 0.5 and 0.75: the line of the upper quartile lies above 0.75, most of all at low typical values, and DHARMa draws it in red. Many of the measurements with low typical values belong to the chilled plants from Mississippi, and the residuals of Mc1 and Mc3 are mostly high, because both plants lie above the typical curve (slide 24) and their seven measurements share that deviation. So this pattern comes from two plants, not from the shape of the curve. The panels are small at slide size; read them aloud. The printed tests are on the figure because DHARMa draws them; as in Chapter 6, we read the plots and the dispersion statistic as exploratory evidence and do not rest decisions on the p-values.

Slide 27 — AIC for grouped models

Grouped models are compared with AIC calculated from the maximum marginal log-likelihood. K counts the parameters that were estimated, the fixed effects and the variance parameters, but not the random effects, which were integrated out. The vector `par` of a fit holds exactly these parameters, so K is its length.

A model without random effects is the special case $\sigma_{\text{group}} = 0$, which lies at the boundary of the range of the parameter, where the large-sample theory behind AIC does not hold. So a comparison between models with and without random effects is only an approximate guide.

AICc and BIC need the number of independent observations n, and for grouped data it is ambiguous. The observations of a group are not independent, because they share its deviation, but they are not one observation either, because each adds information about the curve of its group. The effective number lies somewhere between the number of groups and the number of observations, and depends on how large the variance among groups is compared with the variance within groups. So we do not use AICc and BIC for grouped models, and compare them with AIC. This is the question that Chapter 6 deferred to this chapter when it chose 16 tanks and not 610 tadpoles as n.

For the CO_2 data, the model with a random asymptote has $K = 5$, not 17: the 12 deviations cost one parameter, σ_{plant}. Ask the question of the example box (84 measurements or 12 plants?) before giving the answer of the last bullet.

The second bullet of the slide is the rule behind the comparison: AIC is meaningful only as a difference between models fitted to the same observations, and it ranks them without showing that the best one fits.

Slide 28 — Comparing the candidates

Only differences between models matter, so the table gives the difference of the AIC of each model from the lowest AIC, which is zero for the preferred model.

With a model matrix, candidates that differ only in their fixed effects share one NLL and differ in their formulas (the first bullet of the slide), and they are fitted by maximum likelihood to the same observations. The three candidates with plant deviations use `treatment_nll` of slide 18: the formula `~ 1` gives a model matrix with the column `(Intercept)` only, which is the random asymptote without predictors, `~ Type + Treatment` gives origin and chilling without their interaction, and `~ Type * Treatment` the model of slide 17. Complete pooling has no plant deviations, so it needs an NLL of its own; the chapter writes it with a model matrix as well, so that with the formula `~ 1` it goes through the same bookkeeping as the other candidates. The chapter builds the list of data and the starting values of a candidate from its formula with two short functions, so the loop over the candidates needs no change of code.

The four candidates answer three questions: do the plants differ at all, do origin and chilling explain part of the differences, and does the effect of chilling depend on the origin? AIC prefers the random asymptote with origin, chilling and their interaction. The model without the interaction

is worse by 8.6 units, so the data support an effect of chilling that differs between the two origins, and the random asymptote without predictors is worse by 25.0 units, so origin and chilling explain a large part of the differences among the plants. Complete pooling comes last, 175 units above the random asymptote without predictors. It lies at the boundary $\sigma_{\text{plant}} = 0$ (slide 27), so this difference is only an approximate guide, but one of this size leaves no doubt that the plants differ. Whether they still differ once their origin and treatment are known is a question that this comparison cannot answer, because every candidate with origin and chilling also has plant deviations. The likelihood profile interval of σ_{plant} on slide 23 answers it: plants of the same origin and treatment differ by between about 5% and 14%, and values near zero lie outside the interval.

No pooling is not in the table. Its AIC, 372.9, is lower than that of every mixed model, but it answers another question. Its likelihood describes new measurements of the same 12 plants, with their 12 asymptotes as parameters, whereas the marginal likelihood of a mixed model describes the measurements of a new plant, whose asymptote is unknown. The two likelihoods are probabilities of different things, so their AICs cannot be compared.

For the ecological question the chapter uses the model with the interaction. Chilling multiplies the typical uptake capacity of plants from Quebec by about 0.90 (0.79 to 1.03) and that of plants from Mississippi by about 0.59 (0.52 to 0.68), and the ratio of the two factors lies between 0.53 and 0.81. So a cold night hardly affects the plants from the colder origin, and lowers the capacity of the plants from the warmer origin by about two fifths. Plants of the same origin and treatment differ by roughly 8% (5% to 14%), or 10% when the variance is estimated by REML. With three plants per combination, these conclusions rest on 12 plants, which is why the intervals are wide.

Delivery

Read the table by rows in pairs: rows 1 and 2 differ only in the plant deviations, rows 2 and 3 in origin and chilling, and rows 3 and 4 only in the interaction. Ask which pair of rows would tell whether plants of the same origin and treatment differ: none, because every model with origin and chilling has plant deviations, so point back to the interval of slide 23. The takeaway is the general point: AIC ranks the candidates, and what the ranking means for the question is a matter of ecological interpretation. Do not present the choice of the model with the interaction as settled by AIC alone.

Slide 29 — A recipe for a grouped response curve

The steps of this chapter extend the recipe of Chapter 6 to observations that come in groups, and the chapter collects them in a note callout before the summary. Read the nine steps with the tool of each: identify the observational unit, the groups and the level at which each predictor is measured, and build the model matrix of the predictors with `model.matrix()`, from a formula and a data frame with one row per group (or per observation), checking the reference level of each factor; choose the curve, the observation model and the parameter that varies among groups, on a scale on which a Normal deviation is plausible (the logarithm of a positive parameter, the logit of a probability); write the joint NLL with the density of the random effects first; process it with `MakeADFun()` and `random`, fit it with `optim()` and check `convergence`; read the variance components and predicted effects from `sdreport()`, with likelihood profile intervals

from `TMB::tmbprofile()` and REML when the variances are the question; draw the curves of observed groups, the typical curve and the range of new groups, each for the question that it answers; mark the response with `OBS()`, simulate with `obj$simulate()` and check the residuals of the observations and of the groups; compare the candidates, fitted by maximum likelihood to the same observations, with AIC, where candidates that differ only in their fixed effects share one NLL and differ in their formulas; and interpret the fixed effects and the variance among groups in units, saying what the number of groups leaves uncertain.

Delivery

About two minutes. Steps 2, 4 and 8 extend steps of Chapter 6; point at the new ones (1, 3, 5 with REML, 6 and 7 with the residuals of the groups). The practical follows the same sequence in its worked exercise.

Slide 30 — The Chapter 7 R practical

The practical fits models with random effects to four data sets in which the observations come in groups; three of them were used in earlier chapters, where the observations were treated as independent. Exercise 1, on the growth of 14 loblolly pines (`Loblolly`, built into R), is worked on the page: the saturating exponential as a growth curve with a random asymptote for each tree, the variance components with a likelihood profile interval, the curves of the trees and of new trees, residuals of measurements and trees, and AIC. Exercise 2 returns to the reed frog tanks of the Chapter 4 practical (`ReedfrogPred`, `emdbook`): a Binomial model with a random effect for each tank on the logit scale, against the Beta-Binomial model of Chapter 6. Exercise 3 uses all the visits to 27 barn owl nests (`Owls`, `glmmTMB`) with a Negative Binomial model and a random effect for each nest, and compares no pooling with partial pooling. Exercise 4 models the salamander counts of Chapter 2 at 23 stream sites (`Salamanders`, `glmmTMB`) with a Poisson model and a random effect for each site. Exercise 5 is extra practice: it compares maximum likelihood and REML estimates of a variance among groups by simulation.

Try each exercise before checking the solution.

Slide 31 — Summary and closing questions

Observations that come in groups share the conditions of their group, so they are not independent. A grouped model describes this with two levels: the effects of the groups follow a distribution, and the observations follow a distribution for given values of these effects, so the observations of a group are independent for a given effect of their group but dependent marginally. Complete pooling ignores the groups, no pooling gives each group parameters of its own, and partial pooling, the mixed model, describes both the measured groups and groups in general. The likelihood is the marginal likelihood, with the random effects integrated out, and RTMB approximates the integrals with the Laplace approximation when we name the random effects in `random`. Group-level predictors explain part of the variation among groups, the variance component describes the variation that remains, and the predicted effects of the groups are shrunk towards the typical value. The predictors enter the NLL through a model matrix, which `model.matrix()` builds from a formula, as the formula interfaces of R do. A variance among groups is informed by the number

of groups, and REML reduces the bias of its estimate. The curve of an observed group, the typical curve and the range of new groups answer different questions. Grouped models are checked at both levels and compared with AIC. On the diagram of slide 3, the lecture added groups to the model and marginalisation to the estimation.

Return to this slide for the two closing questions.

Discussion and expected answers — allow about two minutes

Question 1: the variance among groups is informed by the deviations of the groups, and there is one deviation per group. Measuring each group more often makes each deviation more precise, but the number of deviations stays the same, so the precision of the variance is limited by the number of groups (slide 23). For the CO_2 data, 84 measurements but 12 plants. To estimate it more precisely, observe more groups.

Question 2: for a group that was not measured we do not know its effect, so we use the typical curve for its values of the predictors, with the range of new groups; the curve of a measured group describes only that group (slide 24 and Practice 3). Accept answers that add that the typical curve is the median group, not the mean.

Final transition

In the chapter every fit converged from simple starting values. Chapter 8 turns to the difficulties of finding the maximum of a likelihood: bounds, scaling, starting values, local maxima and problems with the Hessian. Exercise 5 of the practical meets one of them, fits that stop at a variance of practically zero, and Chapter 8 returns to that example. The practical comes first.

Adjustments during delivery

The slides take 116 minutes and leave 14 minutes of slack, which comes last. The demo is kept whole; if it overruns, it takes the slack first. If discussion overruns, use the slack first, then shorten in this order:

- Slide 3 to one sentence: recovers about half a minute.
- Slide 5 to the figure and the example box, with the provenance in one sentence: about one minute.
- Slide 9 to the two definitions: about one minute.
- Slide 19 to the takeaway, with one sentence on the two columns: about one and a half minutes.
- Slide 12 to questions 2 and 4: about one and a half minutes.
- Slide 25 given as homework, with the answers in one sentence: about two and a half minutes.
- Slides 29 and 30 to one sentence each: about two minutes together.

Keep the conditional and marginal distributions (slides 6 and 7), the two-level model with a random effect (slide 8), the three ways to share information (slide 11), the marginal likelihood (slide 13), random effects in RTMB (slide 15), the model matrix (slide 18), which the practical uses in Exercises 2 to 4, shrinkage (slide 20) and the curves for observed and new groups (slide 24).

The breaks are placed between the blocks, after slides 12 and 21 (minutes 35 and 78). If a block runs over, shorten the block rather than the following break.

If time is left: code to run

No code is run live apart from the demo, which stops before the uncertainty of the variance components and the comparison. Students can follow the chunks of the chapter page at home, by section: “Uncertainty in a grouped model” (the likelihood profiles of `log_sd_plant` and of the interaction, the fourth element of `beta_asym`, and the REML fit with the fixed effects in `random`), “Checking a grouped model” (the NLL with `OBS()`, the loop over `treatment_obj$simulate()`, `DHARMa::createDHARMa()` and `DHARMa::recalculateResiduals()`) and “Comparing grouped models” (the NLL of complete pooling with a model matrix, the list `models` of NLLs and formulas, `formula_data()`, `start_values()`, `fit_model()` and the loop that calculates the AIC of each candidate).