

Computing Euler angles from a rotation matrix

Gregory G. Slabaugh

Abstract

This document discusses a simple technique to find all possible Euler angles from a rotation matrix. Determination of Euler angles is sometimes a necessary step in computer graphics, vision, robotics, and kinematics. However, the solution may or may not be obvious.

Rotation matrices

We start off with the standard definition of the rotations about the three principle axes.

A rotation of ψ radians about the x -axis is defined as

$$R_x(\psi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \psi & -\sin \psi \\ 0 & \sin \psi & \cos \psi \end{bmatrix}$$

Similarly, a rotation of θ radians about the y -axis is defined as

$$R_y(\theta) = \begin{bmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{bmatrix}$$

Finally, a rotation of ϕ radians about the z -axis is defined as

$$R_z(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

The angles ψ , θ , and ϕ are the Euler angles.

Generalized rotation matrices

A general rotation matrix can will have the form,

$$R = \begin{bmatrix} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{bmatrix}$$

This matrix can be thought of a sequence of three rotations, one about each principle axis. Since matrix multiplication does not commute, the order of the axes which one rotates about will affect the result. For this analysis, we will rotate first about the x -axis, then the y -axis, and finally the z -axis. Such a sequence of rotations can be represented as the matrix product,

$$\begin{aligned} R &= R_z(\phi)R_y(\theta)R_x(\psi) \\ &= \begin{bmatrix} \cos \theta \cos \phi & \sin \psi \sin \theta \cos \phi - \cos \psi \sin \phi & \cos \psi \sin \theta \cos \phi + \sin \psi \sin \phi \\ \cos \theta \sin \phi & \sin \psi \sin \theta \sin \phi + \cos \psi \cos \phi & \cos \psi \sin \theta \sin \phi - \sin \psi \cos \phi \\ -\sin \theta & \sin \psi \cos \theta & \cos \psi \cos \theta \end{bmatrix} \end{aligned}$$

Given a rotation matrix R , we can compute the Euler angles, ψ , θ , and ϕ by equating each element in R with the corresponding element in the matrix product $R_z(\phi)R_y(\theta)R_x(\psi)$. This results in nine equations that can be used to find the Euler angles.

Finding two possible angles for θ

Starting with R_{31} , we find

$$R_{31} = -\sin \theta.$$

This equation can be inverted to yield

$$\theta = -\sin^{-1}(R_{31}). \quad (1)$$

However, one must be careful in interpreting this equation. Since $\sin(\pi - \theta) = \sin(\theta)$, there are actually *two* distinct values (for $R_{31} \neq \pm 1$) of θ that satisfy Equation 1. Therefore, both the values

$$\begin{aligned} \theta_1 &= -\sin^{-1}(R_{31}) \\ \theta_2 &= \pi - \theta_1 = \pi + \sin^{-1}(R_{31}) \end{aligned}$$

are valid solutions. We will handle the special case of $R_{31} = \pm 1$ later in this report. So using the R_{31} element of the rotation matrix, we are able to determine two possible values for θ .

Finding the corresponding angles of ψ

To find the values for ψ , we observe that

$$\frac{R_{32}}{R_{33}} = \tan(\psi).$$

We use this equation to solve for ψ , as

$$\psi = \text{atan2}(R_{32}, R_{33}), \quad (2)$$

where $\text{atan2}(y, x)$ is arc tangent of the two variables x and y . It is similar to calculating the arc tangent of y/x , except that the signs of both arguments are used to determine the quadrant of the result, which lies in the range $[-\pi, \pi]$. The function atan2 is available in many programming languages.

One must be careful in interpreting Equation 2. If $\cos(\theta) > 0$, then $\psi = \text{atan2}(R_{32}, R_{33})$. However, when $\cos(\theta) < 0$, $\psi = \text{atan2}(-R_{32}, -R_{33})$. A simple way to handle this is to use the equation

$$\psi = \text{atan2}\left(\frac{R_{32}}{\cos \theta}, \frac{R_{33}}{\cos \theta}\right) \quad (3)$$

to compute ψ .

Equation 3 is valid for all cases *except* when $\cos \theta = 0$. We will deal with this special case later in this report. For each value of θ , we compute a corresponding value of ψ using Equation 3, yielding

$$\psi_1 = \text{atan2}\left(\frac{R_{32}}{\cos \theta_1}, \frac{R_{33}}{\cos \theta_1}\right) \quad (4)$$

$$\psi_2 = \text{atan2}\left(\frac{R_{32}}{\cos \theta_2}, \frac{R_{33}}{\cos \theta_2}\right) \quad (5)$$

Finding the corresponding angles of ϕ

A similar analysis holds for finding ϕ . We observe that

$$\frac{R_{21}}{R_{11}} = \tan \phi.$$

We solve for ϕ using the equation

$$\phi = \text{atan2}\left(\frac{R_{21}}{\cos \theta}, \frac{R_{11}}{\cos \theta}\right) \quad (6)$$

Again, this equation is valid for all cases *except* when $\cos \theta = 0$. We will deal with this special case later in this report. For each value of θ , we compute a corresponding value of ϕ using Equation 6,

$$\phi_1 = \text{atan2}\left(\frac{R_{21}}{\cos \theta_1}, \frac{R_{11}}{\cos \theta_1}\right) \quad (7)$$

$$\phi_2 = \text{atan2}\left(\frac{R_{21}}{\cos \theta_2}, \frac{R_{11}}{\cos \theta_2}\right) \quad (8)$$

Two solutions if $\cos \theta \neq 0$

For the case of $\cos \theta \neq 0$, we now have two triplets of Euler angles that reproduce the rotation matrix, namely

$$\begin{aligned} &(\psi_1, \theta_1, \phi_1) \\ &(\psi_2, \theta_2, \phi_2) \end{aligned}$$

Both of these solutions will be valid.

What if $\cos \theta = 0$?

This technique described above does not work if the R_{31} element of the rotation matrix is 1 or -1 , which corresponds to $\theta = -\pi/2$ or $\theta = \pi/2$, respectively, and to $\cos \theta = 0$. When we try to solve for the possible values of ψ and ϕ using the above technique, problems will occur, since the elements R_{11} , R_{21} , R_{32} , and R_{33} will all be zero, and therefore Equations 3 and 6 will become

$$\begin{aligned}\psi &= \text{atan2}\left(\frac{0}{0}, \frac{0}{0}\right) \\ \phi &= \text{atan2}\left(\frac{0}{0}, \frac{0}{0}\right).\end{aligned}$$

In this case R_{11} , R_{21} , R_{32} , and R_{33} do not constrain the values of ψ and ϕ . Therefore, we must use different elements of the rotation matrix to compute the values of ψ and ϕ .

$\theta = \pi/2$ **case:** Consider the case when $\theta = \pi/2$. Then,

$$\begin{aligned}R_{12} &= \sin \psi \cos \phi - \cos \psi \sin \phi = \sin(\psi - \phi) \\ R_{13} &= \cos \psi \cos \phi + \sin \psi \sin \phi = \cos(\psi - \phi) \\ R_{22} &= \sin \psi \sin \phi + \cos \psi \cos \phi = \cos(\psi - \phi) = R_{13} \\ R_{23} &= \cos \psi \sin \phi - \sin \psi \cos \phi = -\sin(\psi - \phi) = -R_{12}\end{aligned}$$

Any ψ and ϕ that satisfy these equations will be a valid solution. Using the equations for R_{12} and R_{13} , we find that

$$\begin{aligned}(\psi - \phi) &= \text{atan2}(R_{12}, R_{13}) \\ \psi &= \phi + \text{atan2}(R_{12}, R_{13})\end{aligned}$$

$\theta = -\pi/2$ **case:** Not surprisingly, a similar result holds for the case when $\theta = -\pi/2$, for which

$$\begin{aligned}R_{12} &= -\sin \psi \cos \phi - \cos \psi \sin \phi = -\sin(\psi + \phi) \\ R_{13} &= -\cos \psi \cos \phi + \sin \psi \sin \phi = -\cos(\psi + \phi) \\ R_{22} &= -\sin \psi \sin \phi + \cos \psi \cos \phi = \cos(\psi + \phi) = -R_{13} \\ R_{23} &= -\cos \psi \sin \phi - \sin \psi \cos \phi = -\sin(\psi + \phi) = R_{12}\end{aligned}$$

Again, using the equations for R_{12} and R_{13} , we find that

$$\begin{aligned}(\psi + \phi) &= \text{atan2}(-R_{12}, -R_{13}) \\ \psi &= -\phi + \text{atan2}(-R_{12}, -R_{13})\end{aligned}$$

```

if ( $R_{31} \neq \pm 1$ )
   $\theta_1 = -\text{asin}(R_{31})$ 
   $\theta_2 = \pi - \theta_1$ 
   $\psi_1 = \text{atan2}\left(\frac{R_{32}}{\cos \theta_1}, \frac{R_{33}}{\cos \theta_1}\right)$ 
   $\psi_2 = \text{atan2}\left(\frac{R_{32}}{\cos \theta_2}, \frac{R_{33}}{\cos \theta_2}\right)$ 
   $\phi_1 = \text{atan2}\left(\frac{R_{21}}{\cos \theta_1}, \frac{R_{11}}{\cos \theta_1}\right)$ 
   $\phi_2 = \text{atan2}\left(\frac{R_{21}}{\cos \theta_2}, \frac{R_{11}}{\cos \theta_2}\right)$ 
else
   $\phi = \text{anything; can set to } 0$ 
  if ( $R_{31} = -1$ )
     $\theta = \pi/2$ 
     $\psi = \phi + \text{atan2}(R_{12}, R_{13})$ 
  else
     $\theta = -\pi/2$ 
     $\psi = -\phi + \text{atan2}(-R_{12}, -R_{13})$ 
  end if
end if

```

Figure 1: Pseudo-code for computing Euler angles from a rotation matrix. See text for details.

Either case: In both the $\theta = \pi/2$ and $\theta = -\pi/2$ cases, we have found that ψ and ϕ are linked. This phenomenon is called *Gimbal lock*. Although in this case, there are an infinite number of solutions to the problem, in practice, one is often interested in finding one solution. For this task, it is convenient to set $\phi = 0$ and compute ψ as described above.

Pseudo-code

We now summarize the method by providing a pseudo-code implementation in Figure 1. The code is very simple.

Example

An example that demonstrates the computation of ψ , θ , and ϕ from a rotation matrix is provided below.

Suppose we are asked to find the Euler angles that produce the matrix

$$R = \begin{bmatrix} .5 & -.1464 & .8536 \\ .5 & .8536 & -.1464 \\ -.7071 & .5 & .5 \end{bmatrix}$$

First, we find the possible values for θ to be

$$\theta_1 = -\sin(-.7071) = \frac{\pi}{4}$$

$$\theta_2 = \pi - \theta_1 = \frac{3\pi}{4}$$

Then, we find the corresponding values of ψ to be

$$\begin{aligned}\psi_1 &= \text{atan2}\left(\frac{.5}{\cos(\pi/4)}, \frac{.5}{\cos(\pi/4)}\right) = \frac{\pi}{4} \\ \psi_2 &= \text{atan2}\left(\frac{.5}{\cos(3\pi/4)}, \frac{.5}{\cos(3\pi/4)}\right) = -\frac{3\pi}{4}\end{aligned}$$

And we find ϕ to be

$$\begin{aligned}\phi_1 &= \text{atan2}\left(\frac{.5}{\cos(\pi/4)}, \frac{.5}{\cos(\pi/4)}\right) = \frac{\pi}{4} \\ \phi_2 &= \text{atan2}\left(\frac{.5}{\cos(3\pi/4)}, \frac{.5}{\cos(3\pi/4)}\right) = -\frac{3\pi}{4}\end{aligned}$$

Therefore, the solutions are

$$\begin{pmatrix} \frac{\pi}{4}, \frac{\pi}{4}, \frac{\pi}{4} \\ -\frac{3\pi}{4}, \frac{3\pi}{4}, -\frac{3\pi}{4} \end{pmatrix}$$

More than one solution?

It is interesting to note that there is *always* more than one sequence of rotations about the three principle axes that results in the same orientation of an object. As we have shown in this report, in the non-degenerate case of $\cos \theta \neq 0$, there are two solutions. For the degenerate case of $\cos \theta = 0$, an infinite number of solutions exist.

As an example, consider a book laying on a table face up in front of you. Define the x -axis as to the right, the y -axis as away from you, and the z -axis up. A rotation of π radians about the y -axis will turn the book so that the back cover is now facing up. Another way to achieve the same orientation would be to rotate the book π radians about the x -axis, and then π radians about the z -axis. Thus, there is more than one way to achieve a desired rotation.

Acknowledgement

I wish to thank Sinisa Segvic, from the University of Zagreb, for providing some insights into how to reduce the number of possible solutions to two for the non-degenerate case. Also, many thanks to Tobias Ziegler from the Fraunhofer Institute for pointing out the need for negative signs on R_{12} and R_{13} in the $\theta = -\pi/2$ case.

References

- [1] Ken Shoemake, Animating Rotation with Quaternion Curves Proc. SIGGRAPH 1985, pp. 245 - 254.