

Title of Report

Libby Brill¹, Sophia Freije¹, Jett Palmer¹, Alea Seifert¹

¹Department of Statistics, Cal Poly - SLO

May 31, 2024

Abstract

Your abstract goes here (max ~250 words).

1 Introduction

According to Tuan et al. (2015), it is said to xxx. Supported by this we found xyz (Wiwanitkit 2010).

2 Data and Methods

2.1 Data Collection

2.2 Statistical Analysis

3 Results




Figure 1: The figure represents the percentage of night-time animal rescues by the London Fire Brigade from 2009 to 2020. Grouped by pets (cat, dog, rabbit) and wild animals (bird, fox, deer), the plot reveals that the percent of pet night rescues is larger than wildlife night rescues across all years.

Table 1: The table depicts the percentage of night-time animal rescues by the London Fire Brigade from 2009 to 2020 of pets and wild animals. The table also includes the total number of calls for each group.

Animal Type	% of Rescues Occuring at Night	Total Number of Animals
Pets	35.0%	4668
Wild Animals	30.0%	1898

4 Discussion

Table 1 Figure 1

5 Conclusion

References

- Tuan, Nguyen Minh, Ho Thi Nhan, Nguyen Van Vinh Chau, Nguyen Thanh Hung, Ha Manh Tuan, Ta Van Tram, Nguyen Le Da Ha, et al. 2015. “Sensitivity and Specificity of a Novel Classifier for the Early Diagnosis of Dengue.” Edited by Scott B Halstead. *PLOS Neglected Tropical Diseases* 9 (4): e0003638. <https://doi.org/10.1371/journal.pntd.0003638>.
- Wiwanitkit, Viroj. 2010. “Dengue Fever: Diagnosis and Treatment.” *Expert Review of Anti-Infective Therapy* 8 (7): 841–45.