

## $\text{Ca}^{2+}$ : a versatile master key for intracellular signaling cascades

Shi-Qiang WANG<sup>1\*</sup>, Michael X ZHU<sup>2\*</sup> & Ernesto CARAFOLI<sup>3\*</sup>

<sup>1</sup>State Key Lab of Biomembrane and Membrane Biotechnology, Peking University College of Life Sciences, Beijing 100871, China;

<sup>2</sup>Department of Integrative Biology and Pharmacology, the University of Texas Health Science Center at Houston, Houston 77030, USA;

<sup>3</sup>Venetian Institute of Molecular Medicine, University of Padova, Padova 35121, Italy

**Citation:** Wang S-Q, Zhu M X, Carafoli E.  $\text{Ca}^{2+}$ : a versatile master key for intracellular signaling cascades. *Sci China Life Sci*, 2011, 54: 683–685, doi: 10.1007/s11427-011-4208-4

$\text{Ca}^{2+}$  is one of the most ancient and versatile intracellular messengers in both animal and plant systems.  $\text{Ca}^{2+}$  interacts with a huge array of signaling proteins, and coordinates the integration of non-signaling proteins into cellular communication systems. In doing so,  $\text{Ca}^{2+}$  plays crucial roles in many biological processes, including gene regulation, fuel generation in the metabolic pathways, substance transport across membranes, hormone and neurotransmitter secretion, cell motility and muscle contraction [1].  $\text{Ca}^{2+}$  also controls the life cycle at various stages, from regulating fertilization and cell growth to modulating programmed cell death (apoptosis).  $\text{Ca}^{2+}$  is unequivocally a master key with the ability to control most cellular processes.

This special issue brings together 12 invited reviews of some of the key functions of  $\text{Ca}^{2+}$  in cells, most of which were presented at the 17th International Symposium on  $\text{Ca}^{2+}$ -binding Proteins and  $\text{Ca}^{2+}$  Function in Health and Disease in Beijing, China, on July 16–20, 2011. As reflected in these reviews, the current era of  $\text{Ca}^{2+}$  research is characterized by intense interest in the structure and functional regulation of  $\text{Ca}^{2+}$  transporters and  $\text{Ca}^{2+}$ -binding proteins, and the roles of  $\text{Ca}^{2+}$  in health and disease processes.

### 1 Mechanisms for maintaining low cytosolic $\text{Ca}^{2+}$

In the early stages of biological evolution, systems were

developed to maintain a low cytosolic  $\text{Ca}^{2+}$  concentration to prevent the precipitation of many phosphates and organic molecules [2]. Because phosphate was evolutionarily established as the essential energy ‘currency’ of cells, the formation of calcium phosphate was a deleterious event that needed minimizing. The intracellular  $\text{Ca}^{2+}$  concentration must be lowered from the environmental level of  $10^{-3}$ – $10^{-2}$  mol L<sup>-1</sup> to around  $10^{-7}$  mol L<sup>-1</sup>. It is believed that  $\text{Ca}^{2+}$ -transporting proteins evolved rapidly to extrude intracellular  $\text{Ca}^{2+}$  from the cell. Indeed, DNA sequencing supports the idea that  $\text{Ca}^{2+}$  pumps arose early in evolution, together with other outwardly directed ion pumps [3].

In eukaryotic cells,  $\text{Ca}^{2+}$  is either pumped out of the cells by plasma membrane  $\text{Ca}^{2+}$ -ATPase (PMCA pump) or sequestered into the endoplasmic reticulum (ER) (or sarcoplasmic reticulum (SR) in muscle cells) by the sarco-/endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase (SERCA pump). More recently, a  $\text{Ca}^{2+}$  pump (the SPCA pump) has also been described in the Golgi membranes. Cell membrane systems, including the plasma, inner mitochondrial and lysosomal membranes, also contain antiporters that transport  $\text{Ca}^{2+}$  at the expense of chemical energy stored in  $\text{Na}^{+}$  or  $\text{H}^{+}$  gradients. The relative contribution of these transport mechanisms varies between cell types. For example, in heart cells, the  $\text{Na}^{+}/\text{Ca}^{2+}$  exchanger plays a more important role than the PMCA pump in extruding  $\text{Ca}^{2+}$  during myocardial relaxation [4].

Under physiological conditions, the export of cytosolic  $\text{Ca}^{2+}$  by these  $\text{Ca}^{2+}$  transporters balances the  $\text{Ca}^{2+}$  entry

\*Corresponding author (email: wsq@pku.edu.cn; michael.x.zhu@uth.tmc.edu; ernesto.carafoli@unipd.it)

(‘leak’) from the extracellular fluid and intracellular organelles. In this special issue, Carafoli [5] and Neyses *et al.* [6] review the roles of different subtypes of the PMCA pumps in maintaining intracellular  $\text{Ca}^{2+}$  homeostasis in the inner ear and the cardiovascular system, respectively. In keeping with the present interest in the “negative” role of  $\text{Ca}^{2+}$ , they also summarize recent findings that dysfunction of the PMCA pumps causes disturbance of the balance between  $\text{Ca}^{2+}$  entry and extrusion, and can lead to deafness and cardiovascular disease.

## 2 Mechanisms for spatiotemporally-specific increases in cytosolic $\text{Ca}^{2+}$

The large  $\text{Ca}^{2+}$  gradient that exists across the plasma membrane and the organellar membranes provides a dynamic force for  $\text{Ca}^{2+}$  to act as an intracellular messenger. As mentioned, the cytosolic  $\text{Ca}^{2+}$  concentration in most eukaryotic cells is four orders of magnitude lower than that in the extracellular fluid or the lumen of the ER. Therefore, even a small amount of  $\text{Ca}^{2+}$  entry into the cytosol causes a dramatic and immediate change against the low- $\text{Ca}^{2+}$  background. While fast, transient  $\text{Ca}^{2+}$  signals are usually generated by  $\text{Ca}^{2+}$ -permeable ion channels residing on the plasma membrane and on the membrane of organelles, slow, tonic  $\text{Ca}^{2+}$  signals tend to arise from reverse-mode operation of  $\text{Ca}^{2+}$  exchangers and from slower operation of  $\text{Ca}^{2+}$  pumps. Different  $\text{Ca}^{2+}$  channels and/or transporters generate  $\text{Ca}^{2+}$  signals with different amplitude, duration, location and kinetics. The variability in the spatiotemporal patterning of different  $\text{Ca}^{2+}$  signals forms the fundamental basis for the unique versatility of  $\text{Ca}^{2+}$  as a signaling messenger [1].

Since the finding by Berridge *et al.* that ER  $\text{Ca}^{2+}$  can be mobilized by inositol 1,4,5-trisphosphate (IP<sub>3</sub>), the mobilization of stored  $\text{Ca}^{2+}$  has become a central concept of  $\text{Ca}^{2+}$  signaling. In the late 1980s, Lee and colleagues (for their review in this issue, see ref. [7]) found that stored  $\text{Ca}^{2+}$  could also be liberated by two other messengers: cyclic ADP-ribose (cADPR) and nicotinic acid adenine dinucleotide phosphate (NAADP). Further studies have shown that cADPR activates the so-called ryanodine receptors (RyRs) in the ER. RyR, the largest known membrane-expressed protein, is another major  $\text{Ca}^{2+}$  release channel of ER/SR, and is particularly important in muscle cells and some neurons. In this special issue, Van Petegem and colleague have built upon their important recent research on the RyR to provide a comprehensive overview of the structural aspects of this giant channel protein [8]. Surprisingly, although cADPR and NAADP are synthesized by the same enzyme, NAADP does not target RyR, instead acting upon two-pore channels (TPCs) in lysosomes. In their review [9], Galione and colleagues discuss how studies of TPCs are enhancing our understanding of NAADP-mediated  $\text{Ca}^{2+}$  signaling. Furthermore, Santella and her colleague also describe a

novel NAADP-gated channel in the plasma membrane of starfish oocytes [10].

The alteration of  $\text{Ca}^{2+}$  signals as a pathological basis of disease is the topic of two papers in this special issue. Bezprozvanny *et al.* [11] and Stutzmann *et al.* [12] discuss presenilin, a putative novel ER  $\text{Ca}^{2+}$  channel. Mutations of presenilin cause mishandling of ER  $\text{Ca}^{2+}$ , and play important role in the pathogenesis of Alzheimer’s disease. The  $\text{Ca}^{2+}$  theory of Alzheimer’s disease is now becoming increasingly popular.

Mitochondria are also important  $\text{Ca}^{2+}$  stores transporting  $\text{Ca}^{2+}$  very efficiently. The discovery of “ $\text{Ca}^{2+}$  marks” has provided direct evidence that  $\text{Ca}^{2+}$  signals can also be generated in mitochondria. The mechanisms that mediate  $\text{Ca}^{2+}$  flux through the mitochondrial inner membrane have long been controversial. In this special issue, Sheu and colleagues [13] discuss the major candidate mechanisms that may be involved in generating mitochondrial  $\text{Ca}^{2+}$  signals.

## 3 Mechanisms that transduce $\text{Ca}^{2+}$ signals to effector cascades

Intracellular  $\text{Ca}^{2+}$  signals originating from  $\text{Ca}^{2+}$  channels act either on local targets within proximal microdomains, or regulate distant cellular events by diffusing through the cytosol. The efficiency and target selectivity of  $\text{Ca}^{2+}$  signaling are thus determined by the affinity and stoichiometry of the interaction between  $\text{Ca}^{2+}$  and  $\text{Ca}^{2+}$ -binding proteins. Therefore, uncovering the mechanisms by which  $\text{Ca}^{2+}$  activates its targets is important in understanding the profound specificity of  $\text{Ca}^{2+}$  signals in activating different signaling cascades. In their review, Maki and colleagues [14] describe how the binding of  $\text{Ca}^{2+}$  induces a conformational change in the penta-EF-hand  $\text{Ca}^{2+}$ -binding protein, ALG-2, and facilitates the binding of ligand.

The recent finding regarding STIM1-Orai1 interaction during store-operated  $\text{Ca}^{2+}$  entry (SOCE) is an elegant example of  $\text{Ca}^{2+}$  signal transduction. The STIM1 protein acts as a  $\text{Ca}^{2+}$  sensor in the lumen of the ER, and can be activated by depletion of ER  $\text{Ca}^{2+}$  during  $\text{Ca}^{2+}$  release. Activated STIM1 in turn stimulates Orai1, a plasma membrane  $\text{Ca}^{2+}$  channel, which delivers  $\text{Ca}^{2+}$  to the cytosol to enhance the refilling of ER. Trebak and his colleague [15] summarize the studies on STIM1-Orai1 signaling in the vascular system, and discuss the prospects for drugs targeting STIM1/Orai1 in the treatment of vascular diseases.

To understand the versatile roles of  $\text{Ca}^{2+}$  signaling, it is important to identify the downstream cascades of  $\text{Ca}^{2+}$ -binding proteins. The combination of bioinformatics with molecular biological experiments has greatly enhanced our ability to find  $\text{Ca}^{2+}$  signaling targets. Here, Naranjo [16] reviews recent advances in the construction of an interactome of DREAM/calsenilin/KChIP3, member(s) of the neuronal  $\text{Ca}^{2+}$  sensor superfamily. Interestingly, DREAM

has an essential role in the nucleus as a  $\text{Ca}^{2+}$ -dependent gene silencer.

In addition to the activation of signaling cascades, another major role of  $\text{Ca}^{2+}$  is to regulate mechanical remodeling of the cytoskeleton, a fundamental mechanism of cell motility. The review by Santella *et al.* shows that  $\text{Ca}^{2+}$  signals, and their interaction with actin, play an important role during egg fertilization [10].

Transporters, channels and signaling targets for  $\text{Ca}^{2+}$  comprise a coordinated system. Given the versatility of  $\text{Ca}^{2+}$  signaling, the exquisite homeostasis of  $\text{Ca}^{2+}$  cycling between the cytosol, organelles and extracellular medium is a prerequisite for the healthy operation of the cell system. As mentioned above, and as shown in a number of reviews in this special issue, disturbance of the homeostasis of  $\text{Ca}^{2+}$  signaling due to genetic or regulatory factors is deleterious to cells. In particular, Neyses [6], Stutzmann [12] and their colleagues have shown that pathogenesis involves multiple errors in  $\text{Ca}^{2+}$  handling. It is thus hoped that better understanding of the cellular and molecular mechanisms underlying  $\text{Ca}^{2+}$  mishandling will yield new targets in the battle against major human diseases.

*This work and the invited reviews in this issue were partially supported by the National High-Tech Research and Development Program of China (Grant No. 2011CB809101), the National Natural Science Foundation of China (Grant Nos. 31110303028 and 30730013) and Science China Press.*

1 Berridge M J, Bootman M D, Roderick H L. Calcium signalling: dy-

- namics, homeostasis and remodelling. *Nat Rev Mol Cell Biol*, 2003, 4: 517–529
- 2 Williams R J P. The evolution of the biochemistry of calcium. In: Krebs J, Michalak M, eds. *Calcium: A Matter of Life or Death*. Amsterdam: Elsevier, 2007
- 3 Ikura M. *Trends Biochem Sci*, 1996, 21: 14–17
- 4 Bers D M. *Excitation-contraction Coupling and Cardiac Contractile force*. 2nd ed. Dordrecht (the Netherlands): Kluwer Academic Publisher, 2001
- 5 Carafoli E. The plasma membrane calcium pump in the hearing process: physiology and pathology. *Sci China Life Sci*, 2011, 54: 686–690
- 6 Cartwright E J, Oceandy D, Austin C, *et al.*  $\text{Ca}^{2+}$  signalling in cardiovascular disease: the role of the plasma membrane calcium pumps. *Sci China Life Sci*, 2011, 54: 691–698
- 7 Lee H C. Cyclic ADP-ribose and NAADP: fraternal twin messengers for calcium signaling. *Sci China Life Sci*, 2011, 54: 699–711
- 8 Kimlicka L, Van Petegem F. The Structural biology of ryanodine receptors. *Sci China Life Sci*, 2011, 54: 712–724
- 9 Galione A, Parrington J, Funnell T. Physiological roles of NAADP-mediated  $\text{Ca}^{2+}$  signaling. *Sci China Life Sci*, 2011, 54: 725–732
- 10 Santella L, Chun J T. Actin, more than just a housekeeping protein at the scene of fertilization. *Sci China Life Sci*, 2011, 54: 733–743
- 11 Supnet C, Bezprozvanny I. Presenilins as endoplasmic reticulum calcium leak channels and Alzheimer's disease pathogenesis. *Sci China Life Sci*, 2011, 54: 744–751
- 12 Chakroborty S, Stutzmann G E. Early calcium dysregulation in Alzheimer's disease: setting the stage for synaptic dysfunction. *Sci China Life Sci*, 2011, 54: 752–762
- 13 Pan S, Ryu S Y, Sheu S S. Distinctive characteristics and functions of multiple mitochondrial  $\text{Ca}^{2+}$  influx mechanisms. *Sci China Life Sci*, 2011, 54: 763–769
- 14 Maki M, Suzuki H, Shibata H. Structure and function of ALG-2, a penta-EF-hand calcium-dependent adaptor protein. *Sci China Life Sci*, 2011, 54: 770–779
- 15 Zhang W, Trebak M. STIM1 and Orai1: novel targets for vascular diseases? *Sci China Life Sci*, 2011, 54: 780–785
- 16 Rivas M, Villar D, González P, *et al.* Building the DREAM interactome. *Sci China Life Sci*, 2011, 54: 786–792

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.