

## Review

# Inter-Organellar $\text{Ca}^{2+}$ Homeostasis in Plant and Animal Systems

Philip Steiner <sup>1,\*</sup>  and Susanna Zierler <sup>1,2,\*</sup> 
<sup>1</sup> Institute of Pharmacology, Faculty of Medicine, Johannes Kepler University Linz, 4020 Linz, Austria

<sup>2</sup> Walther Straub Institute of Pharmacology and Toxicology, Ludwig-Maximilians-Universität München, 80336 Munich, Germany

\* Correspondence: philip.steiner@jku.at (P.S.); susanna.zierler@jku.at (S.Z.)

## Abstract

The regulation of calcium ( $\text{Ca}^{2+}$ ) homeostasis is a critical process in both plant and animal systems, involving complex interplay between various organelles and a diverse network of channels, pumps, and transporters. This review provides a concise overview of inter-organellar  $\text{Ca}^{2+}$  homeostasis, highlighting key regulators and mechanisms in plant and animal cells. We discuss the roles of key  $\text{Ca}^{2+}$  channels and transporters, including  $\text{IP}_3\text{Rs}$ ,  $\text{RyRs}$ ,  $\text{TPCs}$ ,  $\text{MCUs}$ ,  $\text{TRPMLs}$ , and  $\text{P2XR}$ s in animals, as well as their plant counterparts. Here, we explore recent innovations in structural biology and advanced microscopic techniques that have enhanced our understanding of these proteins' structure, functions, and regulations. We examine the importance of membrane contact sites in facilitating  $\text{Ca}^{2+}$  transfer between organelles and the specific expression patterns of  $\text{Ca}^{2+}$  channels and transporters. Furthermore, we address the physiological implications of inter-organellar  $\text{Ca}^{2+}$  homeostasis and its relevance in various pathological conditions. For extended comparability, a brief excursus into bacterial intracellular  $\text{Ca}^{2+}$  homeostasis is also made. This meta-analysis aims to bridge the gap between plant and animal  $\text{Ca}^{2+}$  signaling research, identifying common themes and unique adaptations in these diverse biological systems.

**Keywords:** calcium homeostasis; organelle communication;  $\text{IP}_3$  receptors; ryanodine receptors; two-pore channels; mitochondrial calcium uniporters;  $\text{TRPML}$  channels; membrane contact sites; animal calcium signaling; plant calcium signaling



Academic Editor: Juan A. Rosado

Received: 23 June 2025

Revised: 1 August 2025

Accepted: 4 August 2025

Published: 6 August 2025

**Citation:** Steiner, P.; Zierler, S. Inter-Organellar  $\text{Ca}^{2+}$  Homeostasis in Plant and Animal Systems. *Cells* **2025**, *14*, 1204. <https://doi.org/10.3390/cells14151204>

**Copyright:** © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


## 1. Introduction

Calcium ( $\text{Ca}^{2+}$ ) is a ubiquitous second messenger that plays an important role in numerous cellular processes in both plant and animal systems. What makes  $\text{Ca}^{2+}$  such a powerful second messenger is the enormous chemical gradient between the extracellular free  $\text{Ca}^{2+}$  concentration, ranging between 0.8 and 2 mM, and cytosolic free  $\text{Ca}^{2+}$  concentrations of resting cells, which can be as low as 100 nM, which is ten-thousand times lower [1–5]. The maintenance of  $\text{Ca}^{2+}$  homeostasis is essential for proper cellular function and involves manifold interplay between various organelles [6]. In eukaryotic cells, the endoplasmic reticulum (ER) serves as the primary intracellular  $\text{Ca}^{2+}$  store, while mitochondria, chloroplasts (in plants), and other organelles also contribute to  $\text{Ca}^{2+}$  buffering and signaling. The coordinated efforts of these organelles in regulating  $\text{Ca}^{2+}$  levels are critical for cell survival, growth, and response to environmental stimuli. Calmodulin (CaM) serves as a ubiquitously conserved  $\text{Ca}^{2+}$  sensor in both animal and plant cells, but shows a pronounced functional diversification [7,8]. While animal CaM has a uniform sequence, plants express different CaM isoforms. CaM-like proteins (CMLs) are a plant-specific expansion of  $\text{Ca}^{2+}$  sensors, which are not present in animal cells [9–11].

CMLs comprise unique expression patterns and target specificities, providing plants with specialized  $\text{Ca}^{2+}$  signaling capabilities for environmental adaptation [12,13]. Troponin C (TnC) is predominantly restricted to animal muscle cells, where it regulates contraction through  $\text{Ca}^{2+}$ -dependent conformational changes, with no functional equivalent in plant cells [14].  $\text{Ca}^{2+}$ -dependent protein kinases (CDPKs) are ubiquitously expressed in plant cells and some protozoans but not in animal cells [15,16]. CDPKs combine  $\text{Ca}^{2+}$  sensing and kinase activity within a single polypeptide, enabling direct phosphorylation responses to  $\text{Ca}^{2+}$  signals without requiring separate CaM intermediates [17,18]. Recent advances in imaging techniques, ultra-resolution microscopy and molecular tools have significantly enhanced our understanding of the complex mechanisms underlying interorganellar  $\text{Ca}^{2+}$  homeostasis in both plant and animal systems [19–23]. The diverse network of  $\text{Ca}^{2+}$  signaling pathways involves numerous proteins, including channels, pumps, and exchangers, which work in concert to maintain precise spatiotemporal control over intracellular  $\text{Ca}^{2+}$  concentrations. In animal cells, the sarco/endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase (SERCA) actively pumps  $\text{Ca}^{2+}$  into the ER lumen, while inositol 1,4,5-trisphosphate receptors ( $\text{IP}_3\text{Rs}$ ) and ryanodine receptors (RyRs) mediate  $\text{Ca}^{2+}$  release from the ER [24,25]. Similarly, in plant cells,  $\text{Ca}^{2+}$ -ATPases and intracellular ion channels such as two-pore channels (TPCs) play crucial roles in maintaining  $\text{Ca}^{2+}$  gradients across organelle membranes [26–28]. In this review, we did not focus on voltage-gated  $\text{Ca}^{2+}$  channels as they were extensively reviewed before [29–31]. Mitochondria, with their ability to rapidly uptake and release  $\text{Ca}^{2+}$ , act as important modulators of cytosolic  $\text{Ca}^{2+}$  signals and contribute to the shaping of  $\text{Ca}^{2+}$  microdomains [32,33]. Interorganellar communication, not only between the ER and mitochondria but also between the ER and the endo-lysosomal compartment, has emerged as a critical aspect of  $\text{Ca}^{2+}$  homeostasis. Membrane contact sites (MCS) between these organelles facilitate the efficient transfer of  $\text{Ca}^{2+}$  and other molecules, allowing for fine-tuned regulation of cellular processes [24,34–39]. In recent years, the discovery of proteins such as PDZD8, which were shown to tether ER and mitochondria in mammalian neurons, has shed light on the molecular mechanisms underlying these interactions [40]. Moreover, the store-operated  $\text{Ca}^{2+}$  entry (SOCE) pathway, mediated by STIM and ORAI proteins, plays a vital role in replenishing ER  $\text{Ca}^{2+}$  stores and maintaining long-term  $\text{Ca}^{2+}$  homeostasis and signaling in various cell types, including immune cells [41,42]. In plant systems, the presence of chloroplasts adds another layer of complexity to  $\text{Ca}^{2+}$  signaling networks. These organelles not only participate in photosynthesis but also contribute to  $\text{Ca}^{2+}$  homeostasis and signaling, particularly in response to light and other environmental cues [43]. The evolution of  $\text{Ca}^{2+}$ -based signaling in plants has led to the development of unique mechanisms for integrating information from various cellular compartments and responding to diverse stimuli [44,45]. This review aims to provide a comprehensive overview of the current knowledge regarding the key regulators involved in maintaining  $\text{Ca}^{2+}$  balance across different cellular compartments, highlighting both similarities and differences between plant and animal systems. We will explore the latest findings on interorganellar  $\text{Ca}^{2+}$  homeostasis, discuss the physiological implications of these mechanisms, and identify areas for future research in this rapidly evolving field.

## 2. Key Players in Intracellular $\text{Ca}^{2+}$ Regulation in Animal Cells

The following chapter summarizes the most important intracellular ion channels and transporters regulating  $\text{Ca}^{2+}$  homeostasis and signaling in animal cells. The intracellular localization of the individual channels and transporters is schematically depicted in Figure 1, showing the known predominant position of the channels in the various compartment membranes, including the known protein data bank (PDB) structures.



**Figure 1.** Key regulators for intracellular  $\text{Ca}^{2+}$  homeostasis in animal cells. The illustration indicates key components of intracellular  $\text{Ca}^{2+}$  regulation across various cellular compartments. The endoplasmic reticulum (ER), mitochondria (M), and the endolysosomal (EL) compartment serve as  $\text{Ca}^{2+}$  stores.  $\text{Ca}^{2+}$  release from the ER is mediated by inositol 1,4,5-trisphosphate receptors ( $\text{IP}_3\text{R}$ ) and ryanodine receptors ( $\text{RyR}$ ), while  $\text{Ca}^{2+}$  uptake is facilitated by sarco/endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase (SERCA). The mitochondrial calcium uniporter (MCU) enables rapid  $\text{Ca}^{2+}$  uptake into mitochondria. Two-pore channels (TPC) and transient receptor potential mucolipin (TRPML) channels regulate  $\text{Ca}^{2+}$  flux in EL. P2X receptors (P2XR) act as ligand-gated ion channels in the plasma membrane (PM). PM  $\text{Ca}^{2+}$  ATPase (PMCA),  $\text{Ca}^{2+}$  release-activated  $\text{Ca}^{2+}$  channels (CRACM or Orai), and  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX) maintain cytosolic  $\text{Ca}^{2+}$  levels.  $\text{IP}_3\text{Rs}$  are also present in Golgi bodies (GB), contributing to local  $\text{Ca}^{2+}$  signaling. Created with biorender.com. PDB data were also generated using the PDB tool from biorender.com and show a schematic of the van der Waals structure with color codes corresponding to the different sequences. Localization and orientation correspond to the known nature of the proteins.

### 2.1. Inositol Trisphosphate Receptors

Inositol 1,4,5-trisphosphate receptors ( $\text{IP}_3\text{Rs}$ ) are crucial intracellular  $\text{Ca}^{2+}$  channels that play a vital role in cellular signaling and  $\text{Ca}^{2+}$  homeostasis in animal cells. Recent structural studies have provided new insights into the activation mechanisms of  $\text{IP}_3\text{Rs}$  [46]. These receptors are primarily localized in the ER but can also reside in the biomembrane of Golgi bodies (Figure 1; PDB: 8TL9) and regulate  $\text{Ca}^{2+}$  release in response to  $\text{IP}_3$  and local  $\text{Ca}^{2+}$  increase due to interplay with other ion channels [37,38,46–48].  $\text{IP}_3\text{Rs}$  are involved in various physiological processes, including cell proliferation, smooth muscle contraction, apoptosis, and gene expression [49–51]. Moreover, dysregulation of  $\text{IP}_3\text{Rs}$  has been implicated in several pathological conditions, such as neurodegenerative diseases and cancer [24,51]. The spatial organization of  $\text{IP}_3\text{Rs}$  within cells has been shown to be critical for the propagation of  $\text{Ca}^{2+}$  signals and the selective regulation of cellular responses [47,52].

### 2.2. Ryanodine Receptors

Ryanodine receptors ( $\text{RyRs}$ ) are essential intracellular  $\text{Ca}^{2+}$  release channels located in the sarcoplasmic/endoplasmic reticulum (Figure 1; PDB: 8UQ4) of various animal cells.  $\text{RyRs}$  are regulated by a complex interplay of  $\text{Ca}^{2+}$  concentrations, redox state, small molecules, and protein interactions and primarily conduct  $\text{Ca}^{2+}$  [53–55]. These homotetrameric proteins are important for numerous cellular processes, including excitation–contraction coupling in muscle cells and neurotransmission in neurons [56,57].  $\text{RyR1}$ , for example, directly interacts with dihydropyridine receptors (DHPRs) through physical

coupling mechanisms in skeletal muscle, where tetrads of DHPRs bind to RyR1s in an alternating manner and form two distinct connection sites between the channels without requiring  $\text{Ca}^{2+}$  influx [58]. In contrast, RyR2 is generally activated by  $\text{Ca}^{2+}$  itself through  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (CICR) in cardiac muscle, where  $\text{Ca}^{2+}$  influx via L-type  $\text{Ca}^{2+}$  channels triggers RyR2 opening by binding to  $\text{Ca}^{2+}$ -sensing sites on the receptor [59–61]. Previous investigations have revealed the complex structure–function relationship of RyRs, with super-resolution microscopy techniques identifying distinct subpopulations of RyR clusters in cardiac cells [21,22]. The spatial organization and interactions of these clusters significantly influence  $\text{Ca}^{2+}$  signaling dynamics and cellular function. Furthermore, research has highlighted the involvement of RyRs in pathological conditions, such as their potential role in Alzheimer’s disease [62].

### 2.3. Two-Pore Channels

Two-pore channels (TPCs) are intracellular voltage- and ligand-gated cation channels located in the endo-lysosomal system (Figure 1; PDB: 6NQ0) of animal cells [63]. In most vertebrates, all three TPC variants—TPC1, TPC2, and TPC3—are expressed, except in humans, mice, and rats, where TPC3 is not present [64–67]. These channels play crucial roles in various cellular processes, including  $\text{Ca}^{2+}$  signaling, membrane trafficking, and organelle morphology [39]. Structural and functional studies have revealed that TPCs are regulated by multiple factors, including the second messenger NAADP, indirectly via JPT2 and LSM12, phosphoinositides such as  $\text{PI}(3,5)\text{P}_2$ , and voltage [68–70]. In this context, it was hypothesized that upon activation via NAADP, TPCs conduct primarily  $\text{Ca}^{2+}$  and upon activation via  $\text{PI}(3,5)\text{P}_2$ , they conduct primarily  $\text{Na}^+$  [68–70]. Furthermore, researchers have uncovered novel functions of TPCs in regulating inter-organellar communication, particularly between the ER and other organelles such as the apicoplast in parasites [71] or endo-lysosomes in mice [38]. The interplay between TPC2 and  $\text{IP}_3\text{Rs}$  to coordinate  $\text{Ca}^{2+}$  signals between lysosomes and the ER has also been recently described [38,48] but TPC1 has also been shown to regulate ER–endo-lysosome  $\text{Ca}^{2+}$  homeostasis in certain cell types [38]. TPCs have emerged as important players in several pathophysiological conditions, including Parkinson’s disease, non-alcoholic fatty liver disease, anaphylaxis, Ebola infection, cancer, cardiac dysfunction, and diabetes [38,72]. The complex regulation and diverse functions of TPCs make them promising targets for therapeutic interventions in various diseases.

### 2.4. Stromal Interaction Molecules, Orai Proteins, and Store-Operated Calcium Entry

In animal cells, Stromal Interaction Molecules (STIM) and Orai or CRACM proteins play crucial roles in regulating intracellular  $\text{Ca}^{2+}$  levels and signaling (Figure 1; PDB: 7KR5). STIM proteins (STIM1, STIM2), located in the ER, sense decreases in luminal  $\text{Ca}^{2+}$  and activate Orai channels (Orai1, Orai2, Orai3) in the plasma membrane to facilitate  $\text{Ca}^{2+}$  influx [73–77]. This process, known as store-operated calcium entry (SOCE), is essential for various cellular functions, including cell growth and immune responses [78]. The interaction between STIM and Orai is highly specific, with STIM binding to Orai via conserved domains to activate CRAC channels [79,80]. This interaction is critical for maintaining  $\text{Ca}^{2+}$  homeostasis and modulating downstream signaling pathways [81]. Dysregulation of CRAC channels has been linked to immune deficiencies and other diseases, highlighting their importance in cellular function [73,74,82–84].

### 2.5. Mitochondrial Calcium Uniporters

The mitochondrial calcium uniporter (MCU), located in the inner mitochondrial membrane (Figure 1; PDB: 6XJV), is a highly selective  $\text{Ca}^{2+}$  channel complex that plays a crucial role in regulating intracellular  $\text{Ca}^{2+}$  signaling, bioenergetics and cell death in

animal cells [85]. Previously, it was shown that the MCU complex consists of multiple subunits, including the pore-forming MCU protein, its paralog MCUb, the essential MCU regulator (EMRE), and regulatory MICU proteins in the intermembrane space [86,87]. The complex interplay between these components allows for precise control of mitochondrial  $\text{Ca}^{2+}$  uptake, with MICU1 and MICU2 acting as gatekeepers and regulators by sensing cytosolic  $\text{Ca}^{2+}$  to prevent  $\text{Ca}^{2+}$  overload under resting conditions while facilitating rapid uptake during signaling events [32,88,89]. Interestingly, the activity and expression of MCU varies significantly among different tissues, with cardiac mitochondria exhibiting surprisingly low MCU current density, likely to prevent excessive  $\text{Ca}^{2+}$  uptake in this highly metabolically active tissue [90]. In addition to that, previous studies have highlighted the importance of MCU in various physiological and pathological processes, including T-cell-mediated inflammation, cancer cell migration, and cardiac function, making MCU an essential modulator for future biomedical research [91–93].

## 2.6. Transient Receptor Potential Channels

Transient receptor potential (TRP) channels are a diverse superfamily of cation permeable ion channels, classified into six subfamilies (TRPC, TRPV, TRPM, TRPA, TRPP and TRPML) [94]. Two of these subfamilies, are (also) expressed on organelles and thus described in more detail below: TRPML and TRPM. The transient receptor potential mucolipin (TRPML) subfamily comprises three members (TRPML1, TRPML2, and TRPML3) in mammals, encoded by MCOLN1–3 genes. These non-selective cation channels are primarily localized in the endolysosomal system of animal cells (Figure 1; PDB: 6E7Z), playing pivotal roles in various cellular processes [95]. TRPML channels are regulated by multiple factors, including phosphoinositides, reactive oxygen species (ROS), and pH, with  $\text{PI}(3,5)\text{P}_2$  acting as an endogenous activator and  $\text{PI}(4,5)\text{P}_2$  as an inhibitor [96]. Previous investigations shed light on the complex regulatory mechanisms of TRPMLs, with cryo-electron microscopy providing insights into their molecular architecture [23]. TRPML1, the best-characterized member, functions as an “ROS sensor” and is involved in lysosomal  $\text{Ca}^{2+}$  release, autophagy regulation, and maintenance of cellular redox homeostasis [97]. TRPMLs have been implicated in several pathological conditions, including neurodegenerative disorders such as Alzheimer’s and Parkinson’s diseases, immune diseases as well as various forms of cancer [96,98–104]. Due to their involvement in numerous human diseases, TRPMLs are emerging as potential targets for the development of new therapeutic strategies and treatments [105]. Transient receptor potential melastatin-like (TRPM) channels are classified into TRPM1–7. TRPM2, for example, plays an important role in lysosomal  $\text{Ca}^{2+}$  release in dendritic cells and beta cells [106,107]. Several studies have examined the role of the unique channel-kinase TRPM7 within various vesicular compartments [108–111], highlighting its possible participation in vesicular  $\text{Ca}^{2+}$  signaling. As it also harbors a functional serine/threonine kinase, it may contribute to cellular signaling via modulating other signaling proteins [112–114]. Additional research is required to clarify exact molecular mechanisms.

## 2.7. P2X Receptors

P2X receptors (P2X1–7) are trimeric ATP-gated cation channels that play crucial roles in diverse cellular processes across animal cell types and are mainly located in the plasma membrane (Figure 1; PDB: 8JV8). Recent cryo-electron microscopy studies have depicted detailed structures of P2X receptors, elucidating mechanisms of agonist binding and channel gating [115–117]. P2X receptors are  $\text{Ca}^{2+}$ -permeable and can be modulated by extracellular  $\text{Na}^+$ ,  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ , and  $\text{H}^+$ . P2X receptors, particularly P2X7, are implicated in numerous physiological and pathophysiological conditions, including neurodegenerative diseases, inflammation, and pain [118–121]. The expression and function of P2X receptors vary



across cell types, with P2X7 predominantly expressed in immune and glial cells [122–125]. Recently, the potential of P2X receptors as therapeutic targets was discussed, with the development of subtype-specific modulators showing promise in preclinical models [126,127]. In the study of Poejo et al., it was shown that Preyssler-type polyoxotungstate (P5W30) acts as a novel agonist of purinergic P2 receptors, inducing dose-dependent increases in cytosolic  $\text{Ca}^{2+}$  concentration in mouse hippocampal neuronal cells primarily through metabotropic receptor activation, suggesting its potential as a therapeutic tool for targeting purinergic signaling pathways [128]. In addition to that, the discovery of splice variants, especially for P2X7, has added additional complexity to P2X receptor biology and pharmacology [122,129].

## 2.8. Sarco/Endoplasmic Reticulum $\text{Ca}^{2+}$ -ATPase

The sarco/endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase (SERCA; SERCA1-3 isoforms), located in the ER membrane (Figure 1; PDB: 3B9R), is an important regulator for intracellular  $\text{Ca}^{2+}$  homeostasis in animal cells [130]. Studies have highlighted SERCA's importance in various physiological processes and pathological conditions. In cardiac tissue, SERCA2a gene therapy has shown promise in treating heart failure, improving contractility and survival in animal models [131]. SERCA dysfunction has also been implicated in neurodegenerative diseases [132]. The pump's activity is regulated by phospholamban and sarcolipin in muscle cells [133]. SERCA can be targeted in cancer cells by small molecules and pharmacological compounds such as thapsigargin (and prodrug derivatives) which disrupt  $\text{Ca}^{2+}$  balance and trigger programmed cell death [134–137]. This makes SERCA an important target for cancer research. It was also shown, that polyoxotungstates (POTs), particularly Wells–Dawson POTs and Preyssler-type anions (see above), are potential inhibitors of SERCA [138]. Recent structural studies have provided insights into SERCA's molecular mechanisms, such as investigating the molecular architecture underlying the diverse roles of structurally similar SERCA modulators [139], while its role in cellular processes like autophagy and ER stress response has been largely elucidated [140].

## 2.9. Plasma Membrane Calcium ATPase


The plasma membrane  $\text{Ca}^{2+}$ -ATPase (PMCA; Figure 1; PDB: 6A69) is essential for regulating and maintaining the balance of  $\text{Ca}^{2+}$  ions within animal cells, playing a vital role in intracellular  $\text{Ca}^{2+}$  homeostasis [141]. It was shown that PMCA is involved in various physiological processes, including neuronal signaling, cardiac function, and cancer progression [142]. PMCA isoforms (PMCA1-4) exhibit tissue-specific expression patterns and are regulated by CaM and other signaling molecules [143]. Targeting PMCA function has emerged as a potential therapeutic approach for various diseases such as deafness, neurological disorders, autism, or cardiovascular diseases and cancer [142,144,145].

## 2.10. Sodium–Calcium Exchanger

The sodium–calcium exchanger (NCX; NCX1-3) is important for maintaining intracellular  $\text{Ca}^{2+}$  homeostasis in animal cells. As an antiporter membrane protein, NCX exports one  $\text{Ca}^{2+}$  ion in exchange for three  $\text{Na}^{+}$  ions, utilizing the electrochemical gradient of  $\text{Na}^{+}$  [146]. NCX is essential for various cellular functions, including neurosecretion, cardiac muscle relaxation, and maintaining low intracellular  $\text{Ca}^{2+}$  levels [147,148]. NCXs are primarily found in the plasma membrane [149,150], but isoforms can also be found in the mitochondrial membrane (Figure 1; PDB: 3US9) [151,152]. Also, NCXs' involvement in  $\text{Ca}^{2+}$  signaling pathways and its potential as a therapeutic target in cardiovascular and neurological disorders was highlighted before [148,153].

### 3. Key Players in Intracellular $\text{Ca}^{2+}$ Regulation in Plant Cells

This section provides an overview of the key players involved in intracellular  $\text{Ca}^{2+}$  regulation within plant cells, highlighting the most important  $\text{Ca}^{2+}$  channels and transporters. The intracellular localization of the individual channels and transporters is schematically depicted in Figure 2, showing the known predominant position of the channels in the various compartment membranes, including the known or related protein data bank (PDB) structures.



**Figure 2.** Key regulators for intracellular  $\text{Ca}^{2+}$  homeostasis in plant cells.  $\text{Ca}^{2+}$  homeostasis in plant cells involves the vacuole (V), ER, chloroplast (CHL), and cell wall (CW) as major  $\text{Ca}^{2+}$  stores. SERCA-like pumps (ECA) pump  $\text{Ca}^{2+}$  into the ER. Cyclic nucleotide-gated channels (CNGC) and glutamate receptor-like channels (GLR) facilitate  $\text{Ca}^{2+}$  influx across the plasma (and chloroplast) membrane. Two-pore channels (TPC) mediate  $\text{Ca}^{2+}$  release from the vacuole. Mitochondrial calcium uniporters (MCU) allow  $\text{Ca}^{2+}$  uptake into mitochondria, although their presence is less well-characterized than in animal cells. While not fully characterized in plants, recent studies suggest the potential existence of  $\text{IP}_3$  receptor-like channels in the vacuole ( $\text{vIP}_3\text{R}$ ), contributing to  $\text{Ca}^{2+}$  release. The chloroplast has an important yet special function for intracellular, intraorganellar, and plastidial  $\text{Ca}^{2+}$  homeostasis, with chloroplast MCU (cMCU), GLRs, and  $\text{Ca}^{2+}$ -sensing receptors (CAS) highlighted in this scheme. Created with biorender.com; PDB data were also generated using the PDB tool from biorender.com and show a schematic of the van der Waals structure with color codes corresponding to the different sequences. Localization and orientation correspond to the known nature of the proteins.

#### 3.1. Vacuolar Inositol Trisphosphate Receptor

Vacuolar inositol trisphosphate receptors ( $\text{vIP}_3\text{R}$ ) might play a crucial role in intracellular  $\text{Ca}^{2+}$  homeostasis in plant cells. These receptors are thought to be  $\text{Ca}^{2+}$ -permeable channels located in the vacuolar membrane (tonoplast; Figure 2; PDB: 8TL9, human structure schematically used for visualization due to the lack of plant PDB data), activated by  $\text{IP}_3$  as a second messenger [154–156]. However, their existence and properties in plants remain controversial, with limited homology to animal counterparts [155]. So far, no analogs of mammalian-like  $\text{IP}_3\text{Rs}$  could be characterized in plant cells. Presumably located in the vacuolar membrane,  $\text{vIP}_3\text{Rs}$  are thought to mediate  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (CICR) mechanisms essential for cellular  $\text{Ca}^{2+}$  homeostasis [155,157,158]. They might play a pivotal role in intracellular  $\text{Ca}^{2+}$  regulation by facilitating controlled  $\text{Ca}^{2+}$  efflux from the massive vacuolar  $\text{Ca}^{2+}$  stores—which can contain  $\text{Ca}^{2+}$  concentrations up to

three orders of magnitude higher than cytosolic levels and thereby enabling precise temporal and spatial control of cytosolic  $\text{Ca}^{2+}$  signatures required for diverse physiological processes including stress responses, developmental regulation, and signal transduction pathways [157,159–161]. Despite the potential importance of these receptors, their molecular identity and exact mechanisms in plants are still not fully understood, highlighting the need for further research in this area [155].

### 3.2. Cyclic Nucleotide-Gated Channels

Located in the plasma membrane of plant cells (Figure 2; PDB: 9J34), cyclic nucleotide-gated channels (CNGCs) play an important role in maintaining intracellular  $\text{Ca}^{2+}$  homeostasis [162]. These ion channels, regulated by cyclic nucleotides such as cAMP and cGMP, form a complex network with  $\text{Ca}^{2+}$  signaling pathways [163]. CNGCs are involved in various physiological processes, including defense responses, abiotic stress adaptation, and development [164]. Previous investigations have elucidated that CNGCs contribute to  $\text{Ca}^{2+}$  influx across the plasma membrane, thereby modulating cytosolic  $\text{Ca}^{2+}$  levels [165]. The interaction between CNGCs and  $\text{Ca}^{2+}$  is bidirectional, with  $\text{Ca}^{2+}$ -CaM binding regulating channel activity [166]. Furthermore, the crosstalk between cyclic nucleotides and  $\text{Ca}^{2+}$  signaling has been observed in multiple plant species and cell types, highlighting its importance in plant physiology [163]. Understanding CNGC function and regulation is crucial for deciphering plant responses to environmental stimuli and developmental cues.

### 3.3. Two-Pore Channels

TPCs play a crucial role in intracellular  $\text{Ca}^{2+}$  homeostasis in plant cells. These voltage-dependent ion channels are located in the vacuolar membrane (Figure 2; PDB: 5DQQ) and mediate  $\text{Ca}^{2+}$  release from intracellular stores [27,28]. Compared to its animal counterpart, only TPC1 is relatively well described in plant cells [167,168]. TPCs are involved in various physiological processes, including stress responses and stomatal regulation [169]. Furthermore, TPCs contribute to the generation of specific  $\text{Ca}^{2+}$  signatures that encode information during plant–microbe interactions and abiotic stress responses [170]. Another important innovation is the high-resolution atomic representation of TPC1 in plant cells, which provides insights into the structure but also allows interpretation for potential functionality [171–173]. In Guo et al. [174], it was shown that Arabidopsis TPC (AtTPC1) favors  $\text{Ca}^{2+}$  and is nonselective among monovalent cations, whereas human TPC (HsTPC2) is  $\text{PIP}_2$ -activated and thought to be  $\text{Na}^+$ -selective. There it was shown that replacing three filter residues in AtTPC1 confers  $\text{Na}^+$  selectivity, revealing the structural determinants that drive divergent ion preference in TPCs despite their similar filter sequences. TPCs therefore undoubtedly play an important role in  $\text{Ca}^{2+}$  regulation in plant cells.

### 3.4. Mitochondrial Calcium Uniporters

MCUs are located in the inner mitochondrial membrane (Figure 2; PDB: 6DT0, fungus structure schematically used for visualization due to the lack of plant PDB data) and are critical intracellular  $\text{Ca}^{2+}$  homeostasis regulators in plant cells. Recent studies have demonstrated that plant MCU proteins, similar to their mammalian counterparts, mediate mitochondrial  $\text{Ca}^{2+}$  transport and represent the major route for rapid  $\text{Ca}^{2+}$  uptake [33,175]. The MCU complex in plants consists of pore-forming subunits and regulatory proteins, including MICU, which modulates the channel activity [176]. Research has shown that AtMCU1, an Arabidopsis MCU isoform, forms a  $\text{Ca}^{2+}$ -permeable channel sensitive to known inhibitors and MICU regulation [175]. Impairment of MCU function in plants affects mitochondrial  $\text{Ca}^{2+}$  dynamics, ultrastructure, and root growth under certain conditions [175]. Furthermore, MCU-mediated mitochondrial  $\text{Ca}^{2+}$  transport has been linked to



phytohormone signaling and thigmomorphogenesis, highlighting its importance in plant development and environmental responses [33].

### 3.5. SERCA-like Transporters/ER-Type $\text{Ca}^{2+}$ ATPases

Plant cells contain specialized, SERCA-like  $\text{Ca}^{2+}$  transporters (Figure 2; PDB: 3B9R, rabbit structure schematically used for visualization due to the lack of plant PDB data) known as ER-type  $\text{Ca}^{2+}$  ATPases (ECAs) that play crucial roles in maintaining intracellular  $\text{Ca}^{2+}$  homeostasis [177]. These P-type IIA  $\text{Ca}^{2+}$  ATPases are evolutionarily related to animal SERCA and share approximately 50–53% amino acid sequence identity with their animal counterparts [177,178]. Plant ECAs are primarily localized to the ER membrane and function as ATP-driven  $\text{Ca}^{2+}$  pumps that transport  $\text{Ca}^{2+}$  from the cytosol into the ER lumen, thereby maintaining the low cytosolic  $\text{Ca}^{2+}$  concentrations (0.1–0.2  $\mu\text{M}$ ) essential for cellular function [179]. Unlike animal SERCA pumps, plant ECAs are typically insensitive to  $\text{CaM}$  regulation and show distinct pharmacological properties, being inhibited by cyclopiazonic acid but not by thapsigargin [177,180]. It was shown that plant ECAs, particularly ECA1 in *Arabidopsis*, are essential for stress tolerance, pollen fertility, and proper  $\text{Ca}^{2+}$  signaling responses to environmental stimuli, with their activity being regulated by  $\text{Ca}^{2+}$ -dependent protein kinases (CPKs) that phosphorylate and activate these pumps during osmotic stress [181,182]. However, research on SERCA-like transporters in plant cells has made little progress in recent years compared to their animal counterparts [183].

### 3.6. Glutamate Receptor-like Channels

Plant glutamate receptor-like channels (GLRs) predominantly reside in the plasma membrane but can also be located in the chloroplast membrane (Figure 2; PDB: 6R85) and contribute to the regulation of intracellular  $\text{Ca}^{2+}$  homeostasis and signaling in plant cells [184]. These transmembrane proteins allow the movement of various ions across membranes, particularly  $\text{Ca}^{2+}$ , which act as a key second messenger in plant responses to both endogenous and exogenous stimuli [184]. GLRs are involved in numerous physiological processes, including pollen development, sexual reproduction, root development, stomatal regulation, and pathogen response [184]. Recent studies have demonstrated their importance in long-distance electrical and  $\text{Ca}^{2+}$  signaling [185]. GLRs mediate  $\text{Ca}^{2+}$  influx in response to amino acids and other stimuli, generating specific  $\text{Ca}^{2+}$  signatures that are decoded by  $\text{Ca}^{2+}$ -sensing proteins to initiate downstream signaling cascades [185,186]. Understanding GLR function is crucial for elucidating plant stress responses and developmental adaptations to changing environments [187].

### 3.7. Chloroplast

The chloroplast plays an important and exceptional role as an organelle in intracellular  $\text{Ca}^{2+}$  regulation and signaling in plant cells (Figure 2). It can generate specific stromal  $\text{Ca}^{2+}$  regulation in response to environmental stimuli [43]. Several  $\text{Ca}^{2+}$ -permeable channels and transporters in chloroplast membranes have been identified, including the chloroplast-localized mitochondrial calcium uniporter (cMCU; Figure 2; PDB: 6DT0, fungus structure schematically used for visualization due to the lack of plant PDB data), which mediates  $\text{Ca}^{2+}$  flux across the chloroplast envelope and participates in drought stress response [188]. Other important players include the calcium-sensing receptor (CAS; Figure 2; PDB: 7DD7, animal structure schematically used for visualization due to the lack of plant PDB data) and GLRs (Figure 2; PDB: 6385) [189–191]. Nevertheless, these channels and transporters are mainly involved in forming plastidial  $\text{Ca}^{2+}$  transients and regulating important chloroplast functions such as photosynthesis [43].

#### 4. Similarities of Inter-Organellar $\text{Ca}^{2+}$ Homeostasis in Plants and Animals

Inter-organellar  $\text{Ca}^{2+}$  homeostasis in plant and animal cells exhibits several similarities, despite their distinct evolutionary paths. Both systems rely on a complex network of  $\text{Ca}^{2+}$  channels, pumps, and exchangers to maintain precise spatiotemporal control over intracellular  $\text{Ca}^{2+}$  concentrations. The ER serves as an important intracellular  $\text{Ca}^{2+}$  store in both plant and animal cells [6,192], with  $\text{Ca}^{2+}$ -ATPases actively pumping  $\text{Ca}^{2+}$  into the ER lumen. However, this aspect is not sufficiently understood in plant cells [183]. Mitochondria play a crucial role in modulating cytosolic  $\text{Ca}^{2+}$  signals and shaping  $\text{Ca}^{2+}$  microdomains in both systems [32,33]. Membrane contact sites between organelles, particularly between the ER and mitochondria and between ER and lysosomes, facilitate efficient  $\text{Ca}^{2+}$  transfer and fine-tuned regulation of cellular processes in both plant and animal cells [37,38,40,48,193]. Additionally, both systems utilize voltage-gated  $\text{Ca}^{2+}$  channels for rapid  $\text{Ca}^{2+}$  influx in response to membrane potential changes [27,39,71]. The coordinated efforts of these organelles and their associated proteins are critical for maintaining  $\text{Ca}^{2+}$  homeostasis, which is essential for cell survival, growth, and response to environmental stimuli in both systems [194,195]. As an example, both plant and animal TPCs function as intracellular cation channels that integrate voltage,  $\text{Ca}^{2+}$ , and ligand signals to regulate  $\text{Ca}^{2+}$ -dependent processes, including stress responses and organellar communication, with structural similarities such as dual Shaker-like domains enabling their conserved role in modulating cytosolic  $\text{Ca}^{2+}$  dynamics [71,171,196]. While plant TPCs (e.g., Arabidopsis AtTPC1) primarily mediate vacuolar  $\text{Ca}^{2+}$  waves during abiotic stress and immune signaling, animal TPCs (especially TPC1 and TPC2) similarly govern endolysosomal  $\text{Ca}^{2+}$  release for trafficking and metabolic regulation, highlighting their shared evolutionary origin as  $\text{Ca}^{2+}$ -regulated gatekeepers of intracellular compartments [71,197,198].

#### 5. Differences in Inter-Organellar $\text{Ca}^{2+}$ Homeostasis in Plants and Animals

The regulation of  $\text{Ca}^{2+}$  homeostasis plays a pivotal role in both plant and animal cellular processes, but significant differences exist between these kingdoms. In plants,  $\text{Ca}^{2+}$  signaling has evolved to be more direct and specialized compared to animals [199,200]. While animals possess diverse  $\text{Ca}^{2+}$ -influx mechanisms at the plasma membrane, plants have experienced a loss of diversity in this area [200]. For example, L-type and T-type voltage-gated  $\text{Ca}^{2+}$  channels are crucial for intracellular  $\text{Ca}^{2+}$  regulation in animal cells but are absent in plant cells, which have developed different  $\text{Ca}^{2+}$  signaling mechanisms, utilizing specialized channels like TPC, CNGCs, and GLRs to regulate intracellular  $\text{Ca}^{2+}$  homeostasis [199,200]. Plants have developed unique  $\text{Ca}^{2+}$ -binding proteins, such as CDPKs and CDPK-related kinases (CRKs), which are absent in animals [200,201]. The evolution of the  $\text{Ca}^{2+}$ -storing vacuole in plants provides additional possibilities for regulating cytosolic  $\text{Ca}^{2+}$  influx. Plants lack specialized muscle and nerve tissue-specific  $\text{Ca}^{2+}$  signaling genes found in animals, such as RyRs [200]. However, basic cellular  $\text{Ca}^{2+}$  homeostasis mechanisms such as MCUs and TPCs are conserved between plants and animals [27,32,38,188]. These differences in  $\text{Ca}^{2+}$  homeostasis reflect the distinct evolutionary pressures and environmental adaptations faced by plants and animals, particularly in response to abiotic stresses like temperature fluctuations [186]. As a direct example, animal TPCs are predominantly localized in endolysosomal membranes and are primarily activated by NAADP and  $\text{PI}(3,5)\text{P}_2$  to regulate  $\text{Ca}^{2+}$  signaling for vesicle trafficking and disease pathways, whereas plant TPCs such as AtTPC1 reside in vacuolar membranes and depend on voltage and cytosolic  $\text{Ca}^{2+}$  for stress-induced  $\text{Ca}^{2+}$  waves during immune responses [198,202]. Structural differences include EF-hand domains in plant TPCs enabling

direct Ca<sup>2+</sup> regulation, absent in animal TPCs, which instead exhibit NAADP-mediated activation and amplify signals via Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release from the ER [65,198].

6. Inter-Organellar Ca<sup>2+</sup> Regulation in Animal and Plant Cells

Intracellular Ca<sup>2+</sup> regulation is crucial for cellular signaling in both plants and animals, involving complex inter-organellar relationships. In animals, the ER–mitochondria interface plays a pivotal role in Ca<sup>2+</sup> transfer, with optimal efficiency at a distance of about 20 nm [203]. Similarly, also in plants, Ca<sup>2+</sup> fluxes between organelles like mitochondria, ER, Golgi bodies, and chloroplast are essential for Ca<sup>2+</sup> signaling [204–209]. However, plants lack direct homologues of mammalian RyRs in the ER, suggesting different mechanisms. The plasma membrane–ER interface is critical in both systems for lipid and Ca<sup>2+</sup> homeostasis [210–213]. In plants, the vacuole is a major Ca<sup>2+</sup> store, unlike in animals where the ER is predominant [204]. In particular, TPCs should be highlighted, which are currently discussed as potential secondary regulators of intracellular Ca<sup>2+</sup> via the inter-organellar crosstalk between endo-lysosomes and ER [37,39,48] but also the vacuole and ER in plant cells [214]. In addition, the interaction between mitochondria and (endo)lysosomes plays an important role in intracellular Ca<sup>2+</sup> homeostasis and mitochondrial Ca<sup>2+</sup> dynamics [215,216]. Also, peroxisomes can play an important role in intracellular Ca<sup>2+</sup> regulation by functioning as Ca<sup>2+</sup>-buffering organelles that maintain basal Ca<sup>2+</sup> levels around 600 nM and can increase up to 2.4 μM upon stimulation in both animal and plant cells [217,218]. Only recently it was demonstrated that peroxisomes take up Ca<sup>2+</sup> during cytosolic Ca<sup>2+</sup> increases and exhibit beat-to-beat calcium uptake in cardiomyocytes, positioning them as crucial contributors to cellular Ca<sup>2+</sup> homeostasis and excitation–contraction coupling [217]. The importance of the exact interplay of organelles in a cell was only recently elucidated in the organelle interactome study by Valm et al. [219]. There, a multispectral imaging method to study the interactions among six membrane-bound organelles in live cells was developed, revealing their spatial and temporal relationships and how these change under different conditions. This approach provides a powerful tool and new possibilities for understanding cellular organization and dynamics such as inter-organellar, intracellular Ca<sup>2+</sup> regulation. Overall, while both plant and animal systems utilize Ca<sup>2+</sup> as a secondary messenger, the specific inter-organellar interactions and regulatory mechanisms appear to distinctly differ between plants and animals. Table 1 compares well-characterized inter-organellar Ca<sup>2+</sup> regulation, divided into kingdom, interacting organelles, specific functionality, and representative literature.

Table 1. Inter-organellar Ca<sup>2+</sup> regulation in animal and plant cells.

Kingdom	Organelles	Functionality	References
Animal	ER—mitochondria	Spatial Ca <sup>2+</sup> transfer (IP <sub>3</sub> R)	[203]
Animal	Plasma membrane—ER	Lipid- and Ca <sup>2+</sup> homeostasis (ORAI, IP <sub>3</sub> R)	[210,211]
Animal	Endolysosomes—ER	Spatial Ca <sup>2+</sup> transfer (TPC)	[37,39,48]
Animal	Endolysosomes—mitochondria	Intracellular and mitochondrial Ca <sup>2+</sup> dynamics (TRPML)	[215,216]
Animal	ER—Golgi	Interorganellar Ca <sup>2+</sup> regulation (IP <sub>3</sub> R)	[207–209]
Plant	Chloroplast—mitochondria	Ca <sup>2+</sup> signaling (MCU)	[204,205]
Plant	Plasma membrane—ER	Lipid- and Ca <sup>2+</sup> homeostasis	[212,213]
Plant	Chloroplast—ER	Intracellular Ca <sup>2+</sup> regulation	[206]
Plant	Vacuole—ER	Vacuolar Ca <sup>2+</sup> buffering and spatial Ca <sup>2+</sup> transfer (TPC)	[214,216]

7. Excursus: Intracellular Ca<sup>2+</sup> Regulation in Bacteria

While bacteria lack traditional membrane-bound organelles found in eukaryotes, they possess sophisticated Ca<sup>2+</sup> transport systems that regulate intracellular Ca<sup>2+</sup> homeostasis

between different cellular compartments. Bacterial  $\text{Ca}^{2+}$  regulation involves specialized P-type ATPases such as LMCA1 from *Listeria monocytogenes*, which transports a single  $\text{Ca}^{2+}$  ion across the plasma membrane with unique mechanistic properties distinct from eukaryotic systems [220,221]. Recent studies have identified voltage-gated  $\text{Ca}^{2+}$  channels like CavMr in *Meiothermus ruber* that represent evolutionary links between bacterial and mammalian  $\text{Ca}^{2+}$  channels [222]. Bacteria also utilize  $\text{Ca}^{2+}$ -binding proteins and  $\text{Ca}^{2+}$  leak channels, such as CalC in *Pseudomonas aeruginosa*, that mediate rapid  $\text{Ca}^{2+}$  transients in response to environmental stimuli and regulate gene expression controlling virulence factors and biofilm formation [223,224]. Additionally, some bacteria store  $\text{Ca}^{2+}$  in specialized compartments like acidocalcisomes, which contain  $\text{Ca}^{2+}$ - $\text{PO}_4^{3-}$  complexes that function in  $\text{Ca}^{2+}$  homeostasis and potentially facilitate  $\text{Ca}^{2+}$  deposition for biofilm matrix mineralization [223,225]. These findings demonstrate that bacterial  $\text{Ca}^{2+}$  regulation represents a fundamental signaling mechanism that predates eukaryotic evolution and plays critical roles in bacterial survival, pathogenesis, and environmental adaptation. However, similar to plant cells, the extent of research regarding intracellular  $\text{Ca}^{2+}$  regulation lags behind that of animal cells.

## 8. Discussion and Conclusions

This review highlights the complex mechanisms of inter-organellar  $\text{Ca}^{2+}$  homeostasis in plant and animal systems, emphasizing the critical roles of various channels and transporters. Recent advancements in structural biology, imaging techniques, and advanced microscopic methods have significantly enhanced our understanding of these processes [19,21–23,226,227]. The complex interplay between organelles, particularly the ER, mitochondria, and endo-lysosomes, underscores the importance of membrane contact sites in facilitating efficient  $\text{Ca}^{2+}$  transfer [35–37,39]. The discovery of novel proteins like PDZD8 in mammalian neurons has shed light on the molecular basis of these interactions [40]. Additionally, the store-operated  $\text{Ca}^{2+}$  entry pathway, mediated by STIM and ORAI proteins, has emerged as a crucial mechanism for maintaining long-term  $\text{Ca}^{2+}$  homeostasis and signaling [41,42]. In plant systems, the presence of chloroplasts adds another layer of complexity to  $\text{Ca}^{2+}$  signaling networks, particularly in response to environmental stimuli [43]. Future research should focus on elucidating the precise spatiotemporal dynamics of  $\text{Ca}^{2+}$  signaling and its implications in various physiological and pathological conditions. This knowledge could pave the way for novel therapeutic interventions targeting  $\text{Ca}^{2+}$  homeostasis in both plant and animal systems. Intracellular  $\text{Ca}^{2+}$  regulation in plant cells and bacteria remains significantly more cryptic than in animal cells due to fundamental differences in cellular architecture and signaling complexity. Obviously, bacteria predominantly lack traditional membrane-bound organelles and plants have evolved different  $\text{Ca}^{2+}$  signaling systems compared to animals, with simplified influx mechanisms and yet more complex sensor networks that require two to three times more  $\text{Ca}^{2+}$ -binding protein species to achieve the same signaling specificity [199]. This is based on the predominant sessile nature of plants, which necessitates more sophisticated internal regulatory mechanisms to respond to environmental changes [228]. The current understanding of plant  $\text{Ca}^{2+}$  channels remains remarkably limited, with many key channels still genetically unidentified and their mechanistic basis not fully understood [229]. Unlike animals, where  $\text{Ca}^{2+}$  oscillation systems are well characterized through  $\text{IP}_3\text{Rs}$  and  $\text{RyRs}$ , plants lack these identical, homologous proteins [199], relying instead on less well-studied alternative mechanisms which still leave important scientific questions unanswered, decades after their initial investigation [230,231]. Overall, not only plant and animal cells have developed distinct strategies for intracellular and inter-organellar  $\text{Ca}^{2+}$  regulation, but even different cell types within a single organism have evolved diverse  $\text{Ca}^{2+}$  homeostasis mechanisms.

Nevertheless, the fundamental principles of intracellular  $\text{Ca}^{2+}$  regulation are conserved, revealing surprising similarities from bacterial cells to plants and even human cells.

**Author Contributions:** Conceptualization, P.S. and S.Z.; writing—original draft preparation, P.S. and S.Z.; writing—review and editing, P.S. and S.Z.; visualization, P.S.; supervision, S.Z.; funding acquisition, S.Z. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by Deutsche Forschungsgemeinschaft (DFG), grant number TRR-152 P14. Supported by Johannes Kepler University Open Access Publishing Fund.

**Data Availability Statement:** No new data were created or analyzed in this study.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

- Islam, M.S. Calcium Signaling: From Basic to Bedside. *Adv. Exp. Med. Biol.* **2020**, *1131*, 1–6. [\[CrossRef\]](#) [\[PubMed\]](#)
- Jiang, Y.; Ding, P. Calcium signaling in plant immunity: A spatiotemporally controlled symphony. *Trends Plant Sci.* **2023**, *28*, 74–89. [\[CrossRef\]](#) [\[PubMed\]](#)
- Song, L.; Tang, Y.; Law, B.Y.K. Targeting calcium signaling in Alzheimer’s disease: Challenges and promising therapeutic avenues. *Neural Regen. Res.* **2024**, *19*, 501–502. [\[CrossRef\]](#)
- Berridge, M.J.; Bootman, M.D.; Roderick, H.L. Calcium signalling: Dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol.* **2003**, *4*, 517–529. [\[CrossRef\]](#)
- Clapham, D.E. Calcium signaling. *Cell* **2007**, *131*, 1047–1058. [\[CrossRef\]](#)
- Burgoyne, T.; Patel, S.; Eden, E.R. Calcium signaling at ER membrane contact sites. *Biochim. Biophys. Acta* **2015**, *1853*, 2012–2017. [\[CrossRef\]](#)
- Villalobo, A.; Gonzalez-Munoz, M.; Berchtold, M.W. Proteins with calmodulin-like domains: Structures and functional roles. *Cell. Mol. Life Sci.* **2019**, *76*, 2299–2328. [\[CrossRef\]](#)
- Perochon, A.; Aldon, D.; Galaud, J.P.; Ranty, B. Calmodulin and calmodulin-like proteins in plant calcium signaling. *Biochimie* **2011**, *93*, 2048–2053. [\[CrossRef\]](#) [\[PubMed\]](#)
- Astegno, A.; Bonza, M.C.; Vallone, R.; La Verde, V.; D’Onofrio, M.; Luoni, L.; Molesini, B.; Dominici, P. Arabidopsis calmodulin-like protein CML36 is a calcium ( $\text{Ca}^{2+}$ ) sensor that interacts with the plasma membrane  $\text{Ca}^{2+}$ -ATPase isoform ACA8 and stimulates its activity. *J. Biol. Chem.* **2017**, *292*, 15049–15061. [\[CrossRef\]](#)
- La Verde, V.; Dominici, P.; Astegno, A. Towards Understanding Plant Calcium Signaling through Calmodulin-Like Proteins: A Biochemical and Structural Perspective. *Int. J. Mol. Sci.* **2018**, *19*, 1331. [\[CrossRef\]](#)
- Yadav, M.; Pandey, J.; Chakraborty, A.; Hassan, M.I.; Kundu, J.K.; Roy, A.; Singh, I.K.; Singh, A. A Comprehensive Analysis of Calmodulin-Like Proteins of Glycine max Indicates Their Role in Calcium Signaling and Plant Defense Against Insect Attack. *Front. Plant Sci.* **2022**, *13*, 817950. [\[CrossRef\]](#) [\[PubMed\]](#)
- Heyer, M.; Scholz, S.S.; Reichelt, M.; Kunert, G.; Oelmüller, R.; Mithofer, A. The  $\text{Ca}^{2+}$  sensor proteins CML37 and CML42 antagonistically regulate plant stress responses by altering phytohormone signals. *Plant Mol. Biol.* **2022**, *109*, 611–625. [\[CrossRef\]](#) [\[PubMed\]](#)
- Ogunrinde, A.; Munro, K.; Davidson, A.; Ubaid, M.; Snedden, W.A. Arabidopsis Calmodulin-Like Proteins, CML15 and CML16 Possess Biochemical Properties Distinct from Calmodulin and Show Non-overlapping Tissue Expression Patterns. *Front. Plant Sci.* **2017**, *8*, 2175. [\[CrossRef\]](#) [\[PubMed\]](#)
- Sevrieva, I.R.; Kampourakis, T.; Irving, M. Structural changes in troponin during activation of skeletal and heart muscle determined in situ by polarised fluorescence. *Biophys. Rev.* **2024**, *16*, 753–772. [\[CrossRef\]](#)
- Boudsocq, M.; Droillard, M.J.; Regad, L.; Lauriere, C. Characterization of Arabidopsis calcium-dependent protein kinases: Activated or not by calcium? *Biochem. J.* **2012**, *447*, 291–299. [\[CrossRef\]](#)
- Dekomah, S.D.; Bi, Z.; Dormatey, R.; Wang, Y.; Haider, F.U.; Sun, C.; Yao, P.; Bai, J. The role of CDPKs in plant development, nutrient and stress signaling. *Front. Genet.* **2022**, *13*, 996203. [\[CrossRef\]](#)
- Schulz, P.; Herde, M.; Romeis, T. Calcium-dependent protein kinases: Hubs in plant stress signaling and development. *Plant Physiol.* **2013**, *163*, 523–530. [\[CrossRef\]](#)
- An, L.; Fang, H.; Zhang, X.; Tang, J.; Gong, J.; Yi, Y.; Tang, M. Genome-Wide Identification and Characterization of the CDPK Family of Genes and Their Response to High-Calcium Stress in *Yinshania henryi*. *Genes* **2025**, *16*, 109. [\[CrossRef\]](#)
- Moccia, F.; Fiorio Pla, A.; Lim, D.; Lodola, F.; Gerbino, A. Intracellular  $\text{Ca}^{2+}$  signalling: Unexpected new roles for the usual suspect. *Front. Physiol.* **2023**, *14*, 1210085. [\[CrossRef\]](#)



20. Pirayesh, N.; Giridhar, M.; Ben Khedher, A.; Vothknecht, U.C.; Chigri, F. Organellar calcium signaling in plants: An update. *Biochim. Biophys. Acta* **2021**, *1868*, 118948. [\[CrossRef\]](#)
21. Hou, Y.; Bai, J.; Shen, X.; de Langen, O.; Li, A.; Lal, S.; Dos Remedios, C.G.; Baddeley, D.; Ruygrok, P.N.; Soeller, C.; et al. Nanoscale Organisation of Ryanodine Receptors and Junctophilin-2 in the Failing Human Heart. *Front. Physiol.* **2021**, *12*, 724372. [\[CrossRef\]](#)
22. Maltsev, A.V.; Ventura Subirachs, V.; Monfredi, O.; Juhaszova, M.; Ajay Warriar, P.; Rakshit, S.; Tagirova, S.; Maltsev, A.V.; Stern, M.D.; Lakatta, E.G.; et al. Structure-Function Relationship of the Ryanodine Receptor Cluster Network in Sinoatrial Node Cells. *Cells* **2024**, *13*, 1885. [\[CrossRef\]](#)
23. Schmiede, P.; Fine, M.; Li, X. The regulatory mechanism of mammalian TRPMLs revealed by cryo-EM. *FEBS J.* **2018**, *285*, 2579–2585. [\[CrossRef\]](#)
24. Loncke, J.; Kaasik, A.; Bezprozvanny, I.; Parys, J.B.; Kerkhofs, M.; Bultynck, G. Balancing ER-Mitochondrial  $\text{Ca}^{2+}$  Fluxes in Health and Disease. *Trends Cell Biol.* **2021**, *31*, 598–612. [\[CrossRef\]](#)
25. Kovacs, G.; Reimer, L.; Jensen, P.H. Endoplasmic Reticulum-Based Calcium Dysfunctions in Synucleinopathies. *Front. Neurol.* **2021**, *12*, 742625. [\[CrossRef\]](#)
26. Garcia Bossi, J.; Kumar, K.; Barberini, M.L.; Dominguez, G.D.; Rondon Guerrero, Y.D.C.; Marino-Buslje, C.; Obertello, M.; Muschietti, J.P.; Estevez, J.M. The role of P-type IIA and P-type IIB  $\text{Ca}^{2+}$ -ATPases in plant development and growth. *J. Exp. Bot.* **2020**, *71*, 1239–1248. [\[CrossRef\]](#)
27. Furuichi, T.; Cunningham, K.W.; Muto, S. A putative two pore channel AtTPC1 mediates  $\text{Ca}^{2+}$  flux in Arabidopsis leaf cells. *Plant Cell Physiol.* **2001**, *42*, 900–905. [\[CrossRef\]](#)
28. Hedrich, R.; Muller, T.D.; Marten, I.; Becker, D. TPC1 vacuole SV channel gains further shape—Voltage priming of calcium-dependent gating. *Trends Plant Sci.* **2023**, *28*, 673–684. [\[CrossRef\]](#)
29. Zamponi, G.W.; Striessnig, J.; Koschak, A.; Dolphin, A.C. The Physiology, Pathology, and Pharmacology of Voltage-Gated Calcium Channels and Their Future Therapeutic Potential. *Pharmacol. Rev.* **2015**, *67*, 821–870. [\[CrossRef\]](#)
30. Simms, B.A.; Zamponi, G.W. Neuronal voltage-gated calcium channels: Structure, function, and dysfunction. *Neuron* **2014**, *82*, 24–45. [\[CrossRef\]](#)
31. Dolphin, A.C. Voltage-gated calcium channels and their auxiliary subunits: Physiology and pathophysiology and pharmacology. *J. Physiol.* **2016**, *594*, 5369–5390. [\[CrossRef\]](#)
32. Hasan, P.; Berezhnaya, E.; Rodriguez-Prados, M.; Weaver, D.; Bekeova, C.; Cartes-Saavedra, B.; Birch, E.; Beyer, A.M.; Santos, J.H.; Seifert, E.L.; et al. MICU1 and MICU2 control mitochondrial calcium signaling in the mammalian heart. *Proc. Natl. Acad. Sci. USA* **2024**, *121*, e2402491121. [\[CrossRef\]](#)
33. Ruberti, C.; Feitosa-Araujo, E.; Xu, Z.; Wagner, S.; Grenzi, M.; Darwish, E.; Lichtenauer, S.; Fuchs, P.; Parmagnani, A.S.; Balcerowicz, D.; et al. MCU proteins dominate in vivo mitochondrial  $\text{Ca}^{2+}$  uptake in Arabidopsis roots. *Plant Cell* **2022**, *34*, 4428–4452. [\[CrossRef\]](#)
34. Poggio, E.; Vallese, F.; Hartel, A.J.W.; Morgenstern, T.J.; Kanner, S.A.; Rauh, O.; Giamogante, F.; Barazzuol, L.; Shepard, K.L.; Colecraft, H.M.; et al. Perturbation of the host cell  $\text{Ca}^{2+}$  homeostasis and ER-mitochondria contact sites by the SARS-CoV-2 structural proteins E and M. *Cell Death Dis.* **2023**, *14*, 297. [\[CrossRef\]](#)
35. Sassano, M.L.; Felipe-Abrio, B.; Agostinis, P. ER-mitochondria contact sites; a multifaceted factory for  $\text{Ca}^{2+}$  signaling and lipid transport. *Front. Cell Dev. Biol.* **2022**, *10*, 988014. [\[CrossRef\]](#)
36. Patel, S.; Brailoiu, E. Triggering of  $\text{Ca}^{2+}$  signals by NAADP-gated two-pore channels: A role for membrane contact sites? *Biochem. Soc. Trans.* **2012**, *40*, 153–157. [\[CrossRef\]](#)
37. Kilpatrick, B.S.; Eden, E.R.; Hockey, L.N.; Yates, E.; Futter, C.E.; Patel, S. An endosomal NAADP-sensitive two-pore  $\text{Ca}^{2+}$  channel regulates ER-endosome membrane contact sites to control growth factor signaling. *Cell Rep.* **2017**, *18*, 1636–1645. [\[CrossRef\]](#)
38. Arlt, E.; Fraticelli, M.; Tsvilovskyy, V.; Nadolni, W.; Breit, A.; O'Neill, T.J.; Resenberger, S.; Wennemuth, G.; Wahl-Schott, C.; Biel, M.; et al. TPC1 deficiency or blockade augments systemic anaphylaxis and mast cell activity. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 18068–18078. [\[CrossRef\]](#)
39. Steiner, P.; Arlt, E.; Boekhoff, I.; Gudermann, T.; Zierler, S. Two-pore channels regulate inter-organellar  $\text{Ca}^{2+}$  homeostasis in immune cells. *Cells* **2022**, *11*, 1465. [\[CrossRef\]](#)
40. Hirabayashi, Y.; Kwon, S.K.; Paek, H.; Pernice, W.M.; Paul, M.A.; Lee, J.; Erfani, P.; Raczkowski, A.; Petrey, D.S.; Pon, L.A.; et al. ER-mitochondria tethering by PDZD8 regulates  $\text{Ca}^{2+}$  dynamics in mammalian neurons. *Science* **2017**, *358*, 623–630. [\[CrossRef\]](#)
41. Moccia, F.; Zuccolo, E.; Soda, T.; Tanzi, F.; Guerra, G.; Mapelli, L.; Lodola, F.; D'Angelo, E. Stim and Orai proteins in neuronal  $\text{Ca}^{2+}$  signaling and excitability. *Front. Cell. Neurosci.* **2015**, *9*, 153. [\[CrossRef\]](#)
42. Demaurex, N.; Nunes, P. The role of STIM and ORAI proteins in phagocytic immune cells. *Am. J. Physiol. Cell Physiol.* **2016**, *310*, C496–C508. [\[CrossRef\]](#)
43. Navazio, L.; Formentin, E.; Cendron, L.; Szabo, I. Chloroplast Calcium Signaling in the Spotlight. *Front. Plant Sci.* **2020**, *11*, 186. [\[CrossRef\]](#)

44. Wang, T.; Chen, X.; Ju, C.; Wang, C. Calcium signaling in plant mineral nutrition: From uptake to transport. *Plant Commun.* **2023**, *4*, 100678. [\[CrossRef\]](#)
45. Xu, T.; Niu, J.; Jiang, Z. Sensing Mechanisms: Calcium Signaling Mediated Abiotic Stress in Plants. *Front. Plant Sci.* **2022**, *13*, 925863. [\[CrossRef\]](#)
46. Prole, D.L.; Taylor, C.W. Structure and Function of IP<sub>3</sub> Receptors. *Cold Spring Harb. Perspect. Biol.* **2019**, *11*, a035063. [\[CrossRef\]](#)
47. Thillaiappan, N.B.; Chavda, A.P.; Tovey, S.C.; Prole, D.L.; Taylor, C.W. Ca<sup>2+</sup> signals initiate at immobile IP<sub>3</sub> receptors adjacent to ER-plasma membrane junctions. *Nat. Commun.* **2017**, *8*, 1505. [\[CrossRef\]](#)
48. Yuan, Y.; Arige, V.; Saito, R.; Mu, Q.; Brailoiu, G.C.; Pereira, G.J.S.; Bolsover, S.R.; Keller, M.; Bracher, F.; Grimm, C.; et al. Two-pore channel-2 and inositol trisphosphate receptors coordinate Ca<sup>2+</sup> signals between lysosomes and the endoplasmic reticulum. *Cell Rep.* **2024**, *43*, 113628. [\[CrossRef\]](#)
49. Parys, J.B.; Lemos, F.O. The interplay between associated proteins, redox state and Ca<sup>2+</sup> in the intraluminal ER compartment regulates the IP<sub>3</sub> receptor. *Cell Calcium* **2024**, *117*, 102823. [\[CrossRef\]](#)
50. Narayanan, D.; Adebisi, A.; Jaggar, J.H. Inositol trisphosphate receptors in smooth muscle cells. *Am. J. Physiol. Heart Circ. Physiol.* **2012**, *302*, H2190–H2210. [\[CrossRef\]](#)
51. Rezuchova, I.; Hudecova, S.; Soltysova, A.; Matuskova, M.; Durinikova, E.; Chovancova, B.; Zuzcak, M.; Cihova, M.; Burikova, M.; Penesova, A.; et al. Type 3 inositol 1,4,5-trisphosphate receptor has antiapoptotic and proliferative role in cancer cells. *Cell Death Dis.* **2019**, *10*, 186. [\[CrossRef\]](#)
52. Lock, J.T.; Smith, I.F.; Parker, I. Spatial-temporal patterning of Ca<sup>2+</sup> signals by the subcellular distribution of IP<sub>3</sub> and IP<sub>3</sub> receptors. *Semin. Cell Dev. Biol.* **2019**, *94*, 3–10. [\[CrossRef\]](#)
53. Meissner, G. The structural basis of ryanodine receptor ion channel function. *J. Gen. Physiol.* **2017**, *149*, 1065–1089. [\[CrossRef\]](#)
54. Diercks, B.P.; Werner, R.; Weidemuller, P.; Czarniak, F.; Hernandez, L.; Lehmann, C.; Rosche, A.; Kruger, A.; Kaufmann, U.; Vaeth, M.; et al. ORAI1, STIM1/2, and RYR1 shape subsecond Ca<sup>2+</sup> microdomains upon T cell activation. *Sci. Signal.* **2018**, *11*, eaat0358. [\[CrossRef\]](#)
55. Pessah, I.N. Ryanodine receptor acts as a sensor for redox stress. *Pest. Manag. Sci.* **2001**, *57*, 941–945. [\[CrossRef\]](#)
56. Kobayashi, T.; Kurebayashi, N.; Murayama, T. The Ryanodine Receptor as a Sensor for Intracellular Environments in Muscles. *Int. J. Mol. Sci.* **2021**, *22*, 10795. [\[CrossRef\]](#)
57. Lalo, U.; Pankratov, Y. Astrocyte ryanodine receptors facilitate gliotransmission and astroglial modulation of synaptic plasticity. *Front. Cell. Neurosci.* **2024**, *18*, 1382010. [\[CrossRef\]](#)
58. Xu, J.; Liao, C.; Yin, C.C.; Li, G.; Zhu, Y.; Sun, F. In situ structural insights into the excitation-contraction coupling mechanism of skeletal muscle. *Sci. Adv.* **2024**, *10*, ead11126. [\[CrossRef\]](#)
59. Luo, S.; Benitah, J.P.; Gomez, A.M. Can cardiomyocytes bypass the ‘Ca<sup>2+</sup>-induced Ca<sup>2+</sup>-release’ mechanism? *Cardiovasc. Res.* **2024**, *120*, 6–9. [\[CrossRef\]](#)
60. Porretta, A.P.; Pruvot, E.; Bhuiyan, Z.A. Calcium Release Deficiency Syndrome (CRDS): Rethinking “Atypical” Catecholaminergic Polymorphic Ventricular Tachycardia. *Cardiogenetics* **2024**, *14*, 211–220. [\[CrossRef\]](#)
61. Nikolaienko, R.; Bovo, E.; Zima, A.V. Expression level of cardiac ryanodine receptors dictates properties of Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release. *Biophys. Rep.* **2024**, *4*, 100183. [\[CrossRef\]](#)
62. Chiantia, G.; Hidisoglu, E.; Marcantoni, A. The Role of Ryanodine Receptors in Regulating Neuronal Activity and Its Connection to the Development of Alzheimer’s Disease. *Cells* **2023**, *12*, 1236. [\[CrossRef\]](#) [\[PubMed\]](#)
63. Steiner, P.; Arlt, E.; Boekhoff, I.; Gudermann, T.; Zierler, S. TPC Functions in the Immune System. *Handb. Exp. Pharmacol.* **2023**, *278*, 71–92. [\[CrossRef\]](#) [\[PubMed\]](#)
64. Calcraft, P.J.; Ruas, M.; Pan, Z.; Cheng, X.; Arredouani, A.; Hao, X.; Tang, J.; Rietdorf, K.; Teboul, L.; Chuang, K.T.; et al. NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* **2009**, *459*, 596–600. [\[CrossRef\]](#) [\[PubMed\]](#)
65. Zhu, M.X.; Ma, J.; Parrington, J.; Calcraft, P.J.; Galione, A.; Evans, A.M. Calcium signaling via two-pore channels: Local or global, that is the question. *Am. J. Physiol. Cell Physiol.* **2010**, *298*, C430–C441. [\[CrossRef\]](#)
66. Rahman, T.; Cai, X.; Brailoiu, G.C.; Abood, M.E.; Brailoiu, E.; Patel, S. Two-pore channels provide insight into the evolution of voltage-gated Ca<sup>2+</sup> and Na<sup>+</sup> channels. *Sci. Signal.* **2014**, *7*, ra109. [\[CrossRef\]](#)
67. Cai, X.; Patel, S. Degeneration of an intracellular ion channel in the primate lineage by relaxation of selective constraints. *Mol. Biol. Evol.* **2010**, *27*, 2352–2359. [\[CrossRef\]](#)
68. Roggenkamp, H.G.; Khansahib, I.; Hernandez, C.L.; Zhang, Y.; Lodygin, D.; Kruger, A.; Gu, F.; Mockl, F.; Lohndorf, A.; Wolters, V.; et al. HN1L/JPT2: A signaling protein that connects NAADP generation to Ca<sup>2+</sup> microdomain formation. *Sci. Signal.* **2021**, *14*, eabd5647. [\[CrossRef\]](#)
69. Gunaratne, G.S.; Brailoiu, E.; He, S.; Unterwald, E.M.; Patel, S.; Slama, J.T.; Walseth, T.F.; Marchant, J.S. Essential requirement for JPT2 in NAADP-evoked Ca<sup>2+</sup> signaling. *Sci. Signal.* **2021**, *14*, eabd5605. [\[CrossRef\]](#)
70. Zhang, J.; Guan, X.; Shah, K.; Yan, J. Lsm12 is an NAADP receptor and a two-pore channel regulatory protein required for calcium mobilization from acidic organelles. *Nat. Commun.* **2021**, *12*, 4739. [\[CrossRef\]](#)

71. Li, Z.H.; King, T.P.; Ayong, L.; Asady, B.; Cai, X.; Rahman, T.; Vella, S.A.; Coppens, I.; Patel, S.; Moreno, S.N.J. A plastid two-pore channel essential for inter-organelle communication and growth of *Toxoplasma gondii*. *Nat. Commun.* **2021**, *12*, 5802. [\[CrossRef\]](#) [\[PubMed\]](#)
72. Patel, S.; Kilpatrick, B.S. Two-pore channels and disease. *Biochim. Biophys. Acta* **2018**, *1865*, 1678–1686. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Butorac, C.; Krizova, A.; Derler, I. Review: Structure and Activation Mechanisms of CRAC Channels. *Adv. Exp. Med. Biol.* **2020**, *1131*, 547–604. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Derler, I.; Jardin, I.; Romanin, C. Molecular mechanisms of STIM/Orai communication. *Am. J. Physiol. Cell Physiol.* **2016**, *310*, C643–C662. [\[CrossRef\]](#)
75. Zhang, S.L.; Yu, Y.; Roos, J.; Kozak, J.A.; Deerinck, T.J.; Ellisman, M.H.; Stauderman, K.A.; Cahalan, M.D. STIM1 is a  $\text{Ca}^{2+}$  sensor that activates CRAC channels and migrates from the  $\text{Ca}^{2+}$  store to the plasma membrane. *Nature* **2005**, *437*, 902–905. [\[CrossRef\]](#)
76. Vig, M.; Peinelt, C.; Beck, A.; Koomoa, D.L.; Rabah, D.; Koblan-Huberson, M.; Kraft, S.; Turner, H.; Fleig, A.; Penner, R.; et al. CRACM1 is a plasma membrane protein essential for store-operated  $\text{Ca}^{2+}$  entry. *Science* **2006**, *312*, 1220–1223. [\[CrossRef\]](#)
77. Prakriya, M.; Feske, S.; Gwack, Y.; Srikanth, S.; Rao, A.; Hogan, P.G. Orai1 is an essential pore subunit of the CRAC channel. *Nature* **2006**, *443*, 230–233. [\[CrossRef\]](#)
78. Tedeschi, V.; La Russa, D.; Franco, C.; Vinciguerra, A.; Amantea, D.; Secondo, A. Plasma Membrane and Organellar Targets of STIM1 for Intracellular Calcium Handling in Health and Neurodegenerative Diseases. *Cells* **2021**, *10*, 2518. [\[CrossRef\]](#)
79. Li, Z.; Liu, L.; Deng, Y.; Ji, W.; Du, W.; Xu, P.; Chen, L.; Xu, T. Graded activation of CRAC channel by binding of different numbers of STIM1 to Orai1 subunits. *Cell Res.* **2011**, *21*, 305–315. [\[CrossRef\]](#)
80. Park, C.Y.; Hoover, P.J.; Mullins, F.M.; Bachhawat, P.; Covington, E.D.; Raunser, S.; Walz, T.; Garcia, K.C.; Dolmetsch, R.E.; Lewis, R.S. STIM1 clusters and activates CRAC channels via direct binding of a cytosolic domain to Orai1. *Cell* **2009**, *136*, 876–890. [\[CrossRef\]](#)
81. Berlansky, S.; Humer, C.; Sallinger, M.; Frischauf, I. More Than Just Simple Interaction between STIM and Orai Proteins: CRAC Channel Function Enabled by a Network of Interactions with Regulatory Proteins. *Int. J. Mol. Sci.* **2021**, *22*, 471. [\[CrossRef\]](#)
82. Emrich, S.M.; Yoast, R.E.; Trebak, M. Physiological Functions of CRAC Channels. *Annu. Rev. Physiol.* **2022**, *84*, 355–379. [\[CrossRef\]](#)
83. Vaeth, M.; Kahlfuss, S.; Feske, S. CRAC Channels and Calcium Signaling in T Cell-Mediated Immunity. *Trends Immunol.* **2020**, *41*, 878–901. [\[CrossRef\]](#)
84. Fahrner, M.; Schindl, R.; Romanin, C. Studies of Structure-Function and Subunit Composition of Orai/STIM Channel. In *Calcium Entry Channels in Non-Excitable Cells*; Kozak, J.A., Putney, J.W., Jr., Eds.; CRC Press: Boca Raton, FL, USA, 2018; pp. 25–50.
85. Pathak, T.; Trebak, M. Mitochondrial  $\text{Ca}^{2+}$  signaling. *Pharmacol. Ther.* **2018**, *192*, 112–123. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Balderas, E.; Eberhardt, D.R.; Lee, S.; Pleinis, J.M.; Sommakia, S.; Balynas, A.M.; Yin, X.; Parker, M.C.; Maguire, C.T.; Cho, S.; et al. Mitochondrial calcium uniporter stabilization preserves energetic homeostasis during Complex I impairment. *Nat. Commun.* **2022**, *13*, 2769. [\[CrossRef\]](#) [\[PubMed\]](#)
87. Kamer, K.J.; Mootha, V.K. The molecular era of the mitochondrial calcium uniporter. *Nat. Rev. Mol. Cell Biol.* **2015**, *16*, 545–553. [\[CrossRef\]](#) [\[PubMed\]](#)
88. Fan, M.; Zhang, J.; Tsai, C.W.; Orlando, B.J.; Rodriguez, M.; Xu, Y.; Liao, M.; Tsai, M.F.; Feng, L. Structure and mechanism of the mitochondrial  $\text{Ca}^{2+}$  uniporter holocomplex. *Nature* **2020**, *582*, 129–133. [\[CrossRef\]](#)
89. Gibhardt, C.S.; Riemer, J.; Bogeski, I. Calcium and redox signals at mitochondrial interfaces: A nanoview perspective. *Cell Calcium* **2022**, *103*, 102550. [\[CrossRef\]](#)
90. Fieni, F.; Lee, S.B.; Jan, Y.N.; Kirichok, Y. Activity of the mitochondrial calcium uniporter varies greatly between tissues. *Nat. Commun.* **2012**, *3*, 1317. [\[CrossRef\]](#)
91. Paupe, V.; Prudent, J. New insights into the role of mitochondrial calcium homeostasis in cell migration. *Biochem. Biophys. Res. Commun.* **2018**, *500*, 75–86. [\[CrossRef\]](#)
92. Shumanska, M.; Lodygin, D.; Gibhardt, C.S.; Ickes, C.; Stejerean-Todoran, I.; Krause, L.C.M.; Pahl, K.; Jacobs, L.; Paluschkiwitz, A.; Liu, S.; et al. Mitochondrial calcium uniporter complex controls T-cell-mediated immune responses. *EMBO Rep.* **2024**, *26*, 407–442. [\[CrossRef\]](#)
93. Stejerean-Todoran, I.; Zimmermann, K.; Gibhardt, C.S.; Vultur, A.; Ickes, C.; Shannan, B.; Bonilla Del Rio, Z.; Wolling, A.; Cappello, S.; Sung, H.M.; et al. MCU controls melanoma progression through a redox-controlled phenotype switch. *EMBO Rep.* **2022**, *23*, e54746. [\[CrossRef\]](#)
94. Flockerzi, V.; Nilius, B. TRPs: Truly remarkable proteins. *Handb. Exp. Pharmacol.* **2014**, *222*, 1–12. [\[CrossRef\]](#)
95. Santoni, G.; Maggi, F.; Amantini, C.; Marinelli, O.; Nabissi, M.; Morelli, M.B. Pathophysiological Role of Transient Receptor Potential Mucolipin Channel 1 in Calcium-Mediated Stress-Induced Neurodegenerative Diseases. *Front. Physiol.* **2020**, *11*, 251. [\[CrossRef\]](#) [\[PubMed\]](#)
96. Ouologuem, L.; Bartel, K. Endolysosomal transient receptor potential mucolipins and two-pore channels: Implications for cancer immunity. *Front. Immunol.* **2024**, *15*, 1389194. [\[CrossRef\]](#) [\[PubMed\]](#)

97. Zhang, X.; Cheng, X.; Yu, L.; Yang, J.; Calvo, R.; Patnaik, S.; Hu, X.; Gao, Q.; Yang, M.; Lawas, M.; et al. MCOLN1 is a ROS sensor in lysosomes that regulates autophagy. *Nat. Commun.* **2016**, *7*, 12109. [[CrossRef](#)] [[PubMed](#)]
98. Zhang, L.; Fang, Y.; Zhao, X.; Zheng, Y.; Ma, Y.; Li, S.; Huang, Z.; Li, L. miR-204 silencing reduces mitochondrial autophagy and ROS production in a murine AD model via the TRPML1-activated STAT3 pathway. *Mol. Ther. Nucleic Acids* **2021**, *24*, 822–831. [[CrossRef](#)]
99. Wu, L.-K.; Agarwal, S.; Kuo, C.-H.; Kung, Y.-L.; Day, C.H.; Lin, P.-Y.; Lin, S.-Z.; Hsieh, D.J.; Huang, C.-Y.; Chiang, C.-Y. Artemisia Leaf Extract protects against neuron toxicity by TRPML1 activation and promoting autophagy/mitophagy clearance in both in vitro and in vivo models of MPP+/MPTP-induced Parkinson's disease. *Phytomedicine* **2022**, *104*, 154250. [[CrossRef](#)]
100. Pan, Y.; Zhao, Q.; He, H.; Qi, Y.; Bai, Y.; Zhao, J.; Yang, Y. TRPML1 as a potential therapeutic target for triple-negative breast cancer: A review. *Front. Oncol.* **2023**, *13*, 1326023. [[CrossRef](#)]
101. Krogsaeter, E.; Rosato, A.S.; Grimm, C. TRPMLs and TPCs: Targets for lysosomal storage and neurodegenerative disease therapy? *Cell Calcium* **2022**, *103*, 102553. [[CrossRef](#)]
102. Spix, B.; Castiglioni, A.J.; Remis, N.N.; Flores, E.N.; Wartenberg, P.; Wyatt, A.; Boehm, U.; Gudermann, T.; Biel, M.; Garcia-Anoveros, J.; et al. Whole-body analysis of TRPML3 (MCOLN3) expression using a GFP-reporter mouse model reveals widespread expression in secretory cells and endocrine glands. *PLoS ONE* **2022**, *17*, e0278848. [[CrossRef](#)]
103. Spix, B.; Butz, E.S.; Chen, C.C.; Rosato, A.S.; Tang, R.; Jeridi, A.; Kudrina, V.; Plesch, E.; Wartenberg, P.; Arlt, E.; et al. Lung emphysema and impaired macrophage elastase clearance in mucolipin 3 deficient mice. *Nat. Commun.* **2022**, *13*, 318. [[CrossRef](#)]
104. Spix, B.; Chao, Y.K.; Abrahamian, C.; Chen, C.C.; Grimm, C. TRPML Cation Channels in Inflammation and Immunity. *Front. Immunol.* **2020**, *11*, 225. [[CrossRef](#)] [[PubMed](#)]
105. Rautenberg, S.; Keller, M.; Leser, C.; Chen, C.C.; Bracher, F.; Grimm, C. Expanding the Toolbox: Novel Modulators of Endolysosomal Cation Channels. *Handb. Exp. Pharmacol.* **2023**, *278*, 249–276. [[CrossRef](#)] [[PubMed](#)]
106. Lange, I.; Yamamoto, S.; Partida-Sanchez, S.; Mori, Y.; Fleig, A.; Penner, R. TRPM2 functions as a lysosomal Ca<sup>2+</sup>-release channel in beta cells. *Sci. Signal.* **2009**, *2*, ra23. [[CrossRef](#)] [[PubMed](#)]
107. Sumoza-Toledo, A.; Lange, I.; Cortado, H.; Bhagat, H.; Mori, Y.; Fleig, A.; Penner, R.; Partida-Sanchez, S. Dendritic cell maturation and chemotaxis is regulated by TRPM2-mediated lysosomal Ca<sup>2+</sup> release. *FASEB J.* **2011**, *25*, 3529–3542. [[CrossRef](#)]
108. Abiria, S.A.; Krapivinsky, G.; Sah, R.; Santa-Cruz, A.G.; Chaudhuri, D.; Zhang, J.; Adstamomkonkul, P.; DeCaen, P.G.; Clapham, D.E. TRPM7 senses oxidative stress to release Zn<sup>2+</sup> from unique intracellular vesicles. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E6079–E6088. [[CrossRef](#)]
109. Doyle, C.A.; Busey, G.W.; Lobst, W.H.; Kiessling, V.; Renken, C.; Doppalapudi, H.; Stremska, M.E.; Manjgowda, M.C.; Arish, M.; Wang, W.; et al. Endosomal fusion of pH-dependent enveloped viruses requires ion channel TRPM7. *Nat. Commun.* **2024**, *15*, 8479. [[CrossRef](#)]
110. Krapivinsky, G.; Mochida, S.; Krapivinsky, L.; Cibulsky, S.M.; Clapham, D.E. The TRPM7 ion channel functions in cholinergic synaptic vesicles and affects transmitter release. *Neuron* **2006**, *52*, 485–496. [[CrossRef](#)]
111. Zierler, S.; Sumoza-Toledo, A.; Suzuki, S.; Duill, F.O.; Ryazanova, L.V.; Penner, R.; Ryazanov, A.G.; Fleig, A. TRPM7 kinase activity regulates murine mast cell degranulation. *J. Physiol.* **2016**, *594*, 2957–2970. [[CrossRef](#)]
112. Khajavi, N.; Beck, A.; Ricku, K.; Beyerle, P.; Jacob, K.; Syamsul, S.F.; Belkacemi, A.; Reinach, P.S.; Schreier, P.C.; Salah, H.; et al. TRPM7 kinase is required for insulin production and compensatory islet responses during obesity. *JCI Insight* **2023**, *8*, e163397. [[CrossRef](#)]
113. Nadoln, W.; Immler, R.; Hoelting, K.; Fraticelli, M.; Rippahhn, M.; Rothmiller, S.; Matsushita, M.; Boekhoff, I.; Gudermann, T.; Sperandio, M.; et al. TRPM7 Kinase Is Essential for Neutrophil Recruitment and Function via Regulation of Akt/mTOR Signaling. *Front. Immunol.* **2020**, *11*, 606893. [[CrossRef](#)] [[PubMed](#)]
114. Romagnani, A.; Vettore, V.; Rezzonico-Jost, T.; Hampe, S.; Rottoli, E.; Nadoln, W.; Perotti, M.; Meier, M.A.; Hermanns, C.; Geiger, S.; et al. TRPM7 kinase activity is essential for T cell colonization and alloreactivity in the gut. *Nat. Commun.* **2017**, *8*, 1917. [[CrossRef](#)]
115. Oken, A.C.; Lisi, N.E.; Krishnamurthy, I.; McCarthy, A.E.; Godsey, M.H.; Glasfeld, A.; Mansoor, S.E. High-affinity agonism at the P2X(7) receptor is mediated by three residues outside the orthosteric pocket. *Nat. Commun.* **2024**, *15*, 6662. [[CrossRef](#)] [[PubMed](#)]
116. Sander, S.; Muller, I.; Garcia-Alai, M.M.; Nicke, A.; Tidow, H. New insights into P2X7 receptor regulation: Ca<sup>2+</sup>-calmodulin and GDP bind to the soluble P2X7 ballast domain. *J. Biol. Chem.* **2022**, *298*, 102495. [[CrossRef](#)] [[PubMed](#)]
117. Durner, A.; Durner, E.; Nicke, A. Improved ANAP incorporation and VCF analysis reveal details of P2X7 current facilitation and a limited conformational interplay between ATP binding and the intracellular ballast domain. *eLife* **2023**, *12*, e82479. [[CrossRef](#)]
118. Zheng, H.; Liu, Q.; Zhou, S.; Luo, H.; Zhang, W. Role and therapeutic targets of P2X7 receptors in neurodegenerative diseases. *Front. Immunol.* **2024**, *15*, 1345625. [[CrossRef](#)]
119. Alves, M.; Gil, B.; Villegas-Salmeron, J.; Salari, V.; Martins-Ferreira, R.; Arribas Blazquez, M.; Menendez Mendez, A.; Da Rosa Gerbatin, R.; Smith, J.; de Diego-Garcia, L.; et al. Opposing effects of the purinergic P2X7 receptor on seizures in neurons and microglia in male mice. *Brain Behav. Immun.* **2024**, *120*, 121–140. [[CrossRef](#)]



120. Bianchi, C.; Alvarez-Castelao, B.; Sebastian-Serrano, A.; Di Lauro, C.; Soria-Tobar, L.; Nicke, A.; Engel, T.; Diaz-Hernandez, M. P2X7 receptor inhibition ameliorates ubiquitin-proteasome system dysfunction associated with Alzheimer's disease. *Alzheimers Res. Ther.* **2023**, *15*, 105. [\[CrossRef\]](#)
121. Engel, T.; Nicke, A.; Deussing, J.M.; Sperlagh, B.; Diaz-Hernandez, M. Editorial: P2X7 as Common Therapeutic Target in Brain Diseases. *Front. Mol. Neurosci.* **2021**, *14*, 656011. [\[CrossRef\]](#)
122. Tewari, M.; Michalski, S.; Egan, T.M. Modulation of Microglial Function by ATP-Gated P2X7 Receptors: Studies in Rat, Mice and Human. *Cells* **2024**, *13*, 161. [\[CrossRef\]](#) [\[PubMed\]](#)
123. Jooss, T.; Zhang, J.; Zimmer, B.; Rezzonico-Jost, T.; Rissiek, B.; Felipe Pelczar, P.; Seehusen, F.; Koch-Nolte, F.; Magnus, T.; Zierler, S.; et al. Macrophages and glia are the dominant P2X7-expressing cell types in the gut nervous system—No evidence for the role of neuronal P2X7 receptors in colitis. *Mucosal Immunol.* **2023**, *16*, 180–193. [\[CrossRef\]](#)
124. Sierra-Marquez, J.; Schaller, L.; Sassenbach, L.; Ramirez-Fernandez, A.; Alt, P.; Rissiek, B.; Zimmer, B.; Schredelseker, J.; Hector, J.; Stahler, T.; et al. Different localization of P2X4 and P2X7 receptors in native mouse lung—Lack of evidence for a direct P2X4-P2X7 receptor interaction. *Front. Immunol.* **2024**, *15*, 1425938. [\[CrossRef\]](#)
125. Sluyter, R.; Adriouch, S.; Fuller, S.J.; Nicke, A.; Sophocleous, R.A.; Watson, D. Animal Models for the Investigation of P2X7 Receptors. *Int. J. Mol. Sci.* **2023**, *24*, 8225. [\[CrossRef\]](#)
126. Dunker, C.; Vinnenberg, L.; Isaak, A.; Karabatak, E.; Hundehage, P.; Budde, T.; Murakami, K.; Junker, A. Exploring P2X receptor activity: A journey from cellular impact to electrophysiological profiling. *Biochem. Pharmacol.* **2024**, *229*, 116543. [\[CrossRef\]](#)
127. Cabral-Garcia, G.A.; Cruz-Munoz, J.R.; Valdez-Morales, E.E.; Barajas-Espinosa, A.; Linan-Rico, A.; Guerrero-Alba, R. Pharmacology of P2X Receptors and Their Possible Therapeutic Potential in Obesity and Diabetes. *Pharmaceuticals* **2024**, *17*, 1291. [\[CrossRef\]](#)
128. Poejo, J.; Gumerova, N.I.; Rompel, A.; Mata, A.M.; Aureliano, M.; Gutierrez-Merino, C. Unveiling the agonistic properties of Preyssler-type Polyoxotungstates on purinergic P2 receptors. *J. Inorg. Biochem.* **2024**, *259*, 112640. [\[CrossRef\]](#)
129. Illes, P.; Muller, C.E.; Jacobson, K.A.; Grutter, T.; Nicke, A.; Fountain, S.J.; Kennedy, C.; Schmalzing, G.; Jarvis, M.F.; Stojilkovic, S.S.; et al. Update of P2X receptor properties and their pharmacology: IUPHAR Review 30. *Br. J. Pharmacol.* **2021**, *178*, 489–514. [\[CrossRef\]](#)
130. Primeau, J.O.; Armanious, G.P.; Fisher, M.E.; Young, H.S. The SarcoEndoplasmic Reticulum Calcium ATPase. *Subcell. Biochem.* **2018**, *87*, 229–258. [\[CrossRef\]](#)
131. del Monte, F.; Williams, E.; Lebeche, D.; Schmidt, U.; Rosenzweig, A.; Gwathmey, J.K.; Lewandowski, E.D.; Hajjar, R.J. Improvement in survival and cardiac metabolism after gene transfer of sarcoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase in a rat model of heart failure. *Circulation* **2001**, *104*, 1424–1429. [\[CrossRef\]](#)
132. Britzolaki, A.; Saurine, J.; Klocke, B.; Pitychoutis, P.M. A Role for SERCA Pumps in the Neurobiology of Neuropsychiatric and Neurodegenerative Disorders. *Adv. Exp. Med. Biol.* **2020**, *1131*, 131–161. [\[CrossRef\]](#)
133. Gorski, P.A.; Ceholski, D.K.; Young, H.S. Structure-Function Relationship of the SERCA Pump and Its Regulation by Phospholamban and Sarcolipin. *Adv. Exp. Med. Biol.* **2017**, *981*, 77–119. [\[CrossRef\]](#) [\[PubMed\]](#)
134. Khurram, I.; Khan, M.U.; Ibrahim, S.; Ghani, M.U.; Amin, I.; Falzone, L.; Herrera-Bravo, J.; Setzer, W.N.; Sharifi-Rad, J.; Calina, D. Thapsigargin and its prodrug derivatives: Exploring novel approaches for targeted cancer therapy through calcium signaling disruption. *Med. Oncol.* **2024**, *42*, 7. [\[CrossRef\]](#)
135. Kim, J.; Chang, H.S.; Yun, H.J.; Chang, H.J.; Park, K.C. New Small-Molecule SERCA Inhibitors Enhance Treatment Efficacy in Lenvatinib-Resistant Papillary Thyroid Cancer. *Int. J. Mol. Sci.* **2024**, *25*, 10646. [\[CrossRef\]](#)
136. Kim, S.M.; Park, K.; Yun, H.J.; Kim, J.M.; Choi, K.H.; Park, K.C. Identification of new small molecules for selective inhibition of SERCA 1 in patient-derived metastatic papillary thyroid cancer. *Br. J. Pharmacol.* **2025**, *182*, 2392–2408. [\[CrossRef\]](#)
137. Steiner, P.; Melek, K.; Andosch, A.; Wiesbauer, L.; Madlmayr, A.; Duggan, M.; Kerschbaum, H.H.; Zierler, S. Thapsigargin triggers a non-apoptotic, caspase-independent programmed cell death in basophilic leukaemia cells. *Cell Death Discov.* **2025**, *11*, 313. [\[CrossRef\]](#)
138. Aureliano, M.; Fraqueza, G.; Berrocal, M.; Cordoba-Granados, J.J.; Gumerova, N.I.; Rompel, A.; Gutierrez-Merino, C.; Mata, A.M. Inhibition of SERCA and PMCA  $\text{Ca}^{2+}$ -ATPase activities by polyoxotungstates. *J. Inorg. Biochem.* **2022**, *236*, 111952. [\[CrossRef\]](#)
139. Aguayo-Ortiz, R.; Espinoza-Fonseca, L.M. Linking Biochemical and Structural States of SERCA: Achievements, Challenges, and New Opportunities. *Int. J. Mol. Sci.* **2020**, *21*, 4146. [\[CrossRef\]](#)
140. Rashid, H.O.; Yadav, R.K.; Kim, H.R.; Chae, H.J. ER stress: Autophagy induction, inhibition and selection. *Autophagy* **2015**, *11*, 1956–1977. [\[CrossRef\]](#)
141. Strehler, E.E.; Caride, A.J.; Filoteo, A.G.; Xiong, Y.; Penniston, J.T.; Enyedi, A. Plasma membrane  $\text{Ca}^{2+}$  ATPases as dynamic regulators of cellular calcium handling. *Ann. N. Y. Acad. Sci.* **2007**, *1099*, 226–236. [\[CrossRef\]](#)
142. Stafford, N.; Wilson, C.; Oceandy, D.; Neyses, L.; Cartwright, E.J. The Plasma Membrane Calcium ATPases and Their Role as Major New Players in Human Disease. *Physiol. Rev.* **2017**, *97*, 1089–1125. [\[CrossRef\]](#)



143. Krebs, J. Structure, Function and Regulation of the Plasma Membrane Calcium Pump in Health and Disease. *Int. J. Mol. Sci.* **2022**, *23*, 1027. [[CrossRef](#)] [[PubMed](#)]
144. Themistocleous, S.C.; Yiallouris, A.; Tsioutis, C.; Zaravinos, A.; Johnson, E.O.; Patrikios, I. Clinical significance of P-class pumps in cancer. *Oncol. Lett.* **2021**, *22*, 658. [[CrossRef](#)]
145. Berrocal, M.; Cordoba-Granados, J.J.; Carabineiro, S.A.C.; Gutierrez-Merino, C.; Aureliano, M.; Mata, A.M. Gold Compounds Inhibit the Ca-ATPase Activity of Brain PMCA and Human Neuroblastoma SH-SY5Y Cells and Decrease Cell Viability. *Metals* **2021**, *11*, 1934. [[CrossRef](#)]
146. Yu, S.P.; Choi, D.W. Na<sup>+</sup>-Ca<sup>2+</sup> exchange currents in cortical neurons: Concomitant forward and reverse operation and effect of glutamate. *Eur. J. Neurosci.* **1997**, *9*, 1273–1281. [[CrossRef](#)]
147. Scranton, K.; John, S.; Angelini, M.; Steccanella, F.; Umar, S.; Zhang, R.; Goldhaber, J.I.; Olcese, R.; Ottolia, M. Cardiac function is regulated by the sodium-dependent inhibition of the sodium-calcium exchanger NCX1. *Nat. Commun.* **2024**, *15*, 3831. [[CrossRef](#)]
148. Dong, Y.; Yu, Z.; Li, Y.; Huang, B.; Bai, Q.; Gao, Y.; Chen, Q.; Li, N.; He, L.; Zhao, Y. Structural insight into the allosteric inhibition of human sodium-calcium exchanger NCX1 by XIP and SEA0400. *EMBO J.* **2024**, *43*, 14–31. [[CrossRef](#)]
149. Ren, X.; Philipson, K.D. The topology of the cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger, NCX1. *J. Mol. Cell. Cardiol.* **2013**, *57*, 68–71. [[CrossRef](#)]
150. Liao, J.; Li, H.; Zeng, W.; Sauer, D.B.; Belmares, R.; Jiang, Y. Structural insight into the ion-exchange mechanism of the sodium/calcium exchanger. *Science* **2012**, *335*, 686–690. [[CrossRef](#)]
151. Palty, R.; Silverman, W.F.; Hershfinkel, M.; Caporale, T.; Sensi, S.L.; Parnis, J.; Nolte, C.; Fishman, D.; Shoshan-Barmatz, V.; Herrmann, S.; et al. NCLX is an essential component of mitochondrial Na<sup>+</sup>/Ca<sup>2+</sup> exchange. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 436–441. [[CrossRef](#)]
152. Gobbi, P.; Castaldo, P.; Minelli, A.; Salucci, S.; Magi, S.; Corcione, E.; Amoroso, S. Mitochondrial localization of Na<sup>+</sup>/Ca<sup>2+</sup> exchangers NCX1-3 in neurons and astrocytes of adult rat brain in situ. *Pharmacol. Res.* **2007**, *56*, 556–565. [[CrossRef](#)]
153. Xue, J.; Zeng, W.; Han, Y.; John, S.; Ottolia, M.; Jiang, Y. Structural mechanisms of the human cardiac sodium-calcium exchanger NCX1. *Nat. Commun.* **2023**, *14*, 6181. [[CrossRef](#)]
154. Pottosin, I.I.; Schonknecht, G. Vacuolar calcium channels. *J. Exp. Bot.* **2007**, *58*, 1559–1569. [[CrossRef](#)]
155. Krinke, O.; Novotna, Z.; Valentova, O.; Martinec, J. Inositol trisphosphate receptor in higher plants: Is it real? *J. Exp. Bot.* **2007**, *58*, 361–376. [[CrossRef](#)]
156. Brosnan, J.M.; Sanders, D. Identification and Characterization of High-Affinity Binding Sites for Inositol Trisphosphate in Red Beet. *Plant Cell* **1993**, *5*, 931–940. [[CrossRef](#)]
157. Schonknecht, G. Calcium Signals from the Vacuole. *Plants* **2013**, *2*, 589–614. [[CrossRef](#)]
158. Randall, S.K. Characterization of vacuolar calcium-binding proteins. *Plant Physiol.* **1992**, *100*, 859–867. [[CrossRef](#)]
159. Peiter, E. The plant vacuole: Emitter and receiver of calcium signals. *Cell Calcium* **2011**, *50*, 120–128. [[CrossRef](#)]
160. Ren, H.; Zhao, X.; Li, W.; Hussain, J.; Qi, G.; Liu, S. Calcium Signaling in Plant Programmed Cell Death. *Cells* **2021**, *10*, 1089. [[CrossRef](#)]
161. Costa, A.; Navazio, L.; Szabo, I. The contribution of organelles to plant intracellular Calcium signalling. *J. Exp. Bot.* **2018**, *69*, 4175–4193. [[CrossRef](#)]
162. Jarratt-Barnham, E.; Wang, L.; Ning, Y.; Davies, J.M. The Complex Story of Plant Cyclic Nucleotide-Gated Channels. *Int. J. Mol. Sci.* **2021**, *22*, 874. [[CrossRef](#)]
163. Swiezawska-Boniecka, B.; Duszyn, M.; Kwiatkowski, M.; Szmidt-Jaworska, A.; Jaworski, K. Cross Talk Between Cyclic Nucleotides and Calcium Signaling Pathways in Plants-Achievements and Prospects. *Front. Plant Sci.* **2021**, *12*, 643560. [[CrossRef](#)]
164. Dietrich, P.; Moeder, W.; Yoshioka, K. Plant Cyclic Nucleotide-Gated Channels: New Insights on Their Functions and Regulation. *Plant Physiol.* **2020**, *184*, 27–38. [[CrossRef](#)] [[PubMed](#)]
165. Ma, W.; Berkowitz, G.A. Ca<sup>2+</sup> conduction by plant cyclic nucleotide gated channels and associated signaling components in pathogen defense signal transduction cascades. *New Phytol.* **2011**, *190*, 566–572. [[CrossRef](#)]
166. Pan, Y.; Chai, X.; Gao, Q.; Zhou, L.; Zhang, S.; Li, L.; Luan, S. Dynamic Interactions of Plant CNGC Subunits and Calmodulins Drive Oscillatory Ca<sup>2+</sup> Channel Activities. *Dev. Cell* **2019**, *48*, 710–725.e5. [[CrossRef](#)] [[PubMed](#)]
167. Carpaneto, A.; Gradogna, A. Modulation of calcium and potassium permeation in plant TPC channels. *Biophys. Chem.* **2018**, *236*, 1–7. [[CrossRef](#)] [[PubMed](#)]
168. Hedrich, R.; Marten, I. TPC1-SV channels gain shape. *Mol. Plant* **2011**, *4*, 428–441. [[CrossRef](#)]
169. Tong, T.; Li, Q.; Jiang, W.; Chen, G.; Xue, D.; Deng, F.; Zeng, F.; Chen, Z.H. Molecular Evolution of Calcium Signaling and Transport in Plant Adaptation to Abiotic Stress. *Int. J. Mol. Sci.* **2021**, *22*, 12308. [[CrossRef](#)]
170. Negi, N.P.; Prakash, G.; Narwal, P.; Panwar, R.; Kumar, D.; Chaudhry, B.; Rustagi, A. The calcium connection: Exploring the intricacies of calcium signaling in plant-microbe interactions. *Front. Plant Sci.* **2023**, *14*, 1248648. [[CrossRef](#)]
171. Patel, S.; Penny, C.J.; Rahman, T. Two-pore Channels Enter the Atomic Era: Structure of Plant TPC Revealed. *Trends Biochem. Sci.* **2016**, *41*, 475–477. [[CrossRef](#)]

172. Guo, J.T.; Zeng, W.Z.; Chen, Q.F.; Lee, C.; Chen, L.P.; Yang, Y.; Cang, C.L.; Ren, D.J.; Jiang, Y.X. Structure of the voltage-gated two-pore channel TPC1 from. *Nature* **2016**, *531*, 196–201. [[CrossRef](#)]
173. Kintzer, A.F.; Stroud, R.M. Structure, inhibition and regulation of two-pore channel TPC1 from *Arabidopsis thaliana*. *Nature* **2016**, *531*, 258–262. [[CrossRef](#)]
174. Guo, J.; Zeng, W.; Jiang, Y. Tuning the ion selectivity of two-pore channels. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 1009–1014. [[CrossRef](#)] [[PubMed](#)]
175. Teardo, E.; Carraretto, L.; Wagner, S.; Formentin, E.; Behera, S.; De Bortoli, S.; Larosa, V.; Fuchs, P.; Lo Schiavo, F.; Raffaello, A.; et al. Physiological Characterization of a Plant Mitochondrial Calcium Uniporter in Vitro and in Vivo. *Plant Physiol.* **2017**, *173*, 1355–1370. [[CrossRef](#)] [[PubMed](#)]
176. Tsai, M.F.; Phillips, C.B.; Ranaghan, M.; Tsai, C.W.; Wu, Y.; Williams, C.; Miller, C. Dual functions of a small regulatory subunit in the mitochondrial calcium uniporter complex. *eLife* **2016**, *5*, e15545. [[CrossRef](#)]
177. Liang, F.; Cunningham, K.W.; Harper, J.F.; Sze, H. ECA1 complements yeast mutants defective in  $\text{Ca}^{2+}$  pumps and encodes an endoplasmic reticulum-type  $\text{Ca}^{2+}$ -ATPase in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 8579–8584. [[CrossRef](#)]
178. Wimmers, L.E.; Ewing, N.N.; Bennett, A.B. Higher plant  $\text{Ca}^{2+}$ -ATPase: Primary structure and regulation of mRNA abundance by salt. *Proc. Natl. Acad. Sci. USA* **1992**, *89*, 9205–9209. [[CrossRef](#)]
179. He, J.; Rossner, N.; Hoang, M.T.T.; Alejandro, S.; Peiter, E. Transport, functions, and interaction of calcium and manganese in plant organellar compartments. *Plant Physiol.* **2021**, *187*, 1940–1972. [[CrossRef](#)]
180. Hwang, I.; Ratterman, D.M.; Sze, H. Distinction between Endoplasmic Reticulum-Type and Plasma Membrane-Type  $\text{Ca}^{2+}$  Pumps (Partial Purification of a 120-Kilodalton  $\text{Ca}^{2+}$ -ATPase from Endomembranes). *Plant Physiol.* **1997**, *113*, 535–548. [[CrossRef](#)]
181. Wang, C.; Tang, R.J.; Kou, S.; Xu, X.; Lu, Y.; Rauscher, K.; Voelker, A.; Luan, S. Mechanisms of calcium homeostasis orchestrate plant growth and immunity. *Nature* **2024**, *627*, 382–388. [[CrossRef](#)]
182. Rahmati Ishka, M.; Brown, E.; Rosenberg, A.; Romanowsky, S.; Davis, J.A.; Choi, W.G.; Harper, J.F. *Arabidopsis*  $\text{Ca}^{2+}$ -ATPases 1, 2, and 7 in the endoplasmic reticulum contribute to growth and pollen fitness. *Plant Physiol.* **2021**, *185*, 1966–1985. [[CrossRef](#)]
183. Periasamy, M.; Kalyanasundaram, A. SERCA pump isoforms: Their role in calcium transport and disease. *Muscle Nerve* **2007**, *35*, 430–442. [[CrossRef](#)]
184. Grenzi, M.; Bonza, M.C.; Costa, A. Signaling by plant glutamate receptor-like channels: What else! *Curr. Opin. Plant Biol.* **2022**, *68*, 102253. [[CrossRef](#)]
185. Simon, A.A.; Navarro-Retamal, C.; Feijo, J.A. Merging Signaling with Structure: Functions and Mechanisms of Plant Glutamate Receptor Ion Channels. *Annu. Rev. Plant Biol.* **2023**, *74*, 415–452. [[CrossRef](#)]
186. Kang, X.; Zhao, L.; Liu, X. Calcium Signaling and the Response to Heat Shock in Crop Plants. *Int. J. Mol. Sci.* **2023**, *25*, 324. [[CrossRef](#)] [[PubMed](#)]
187. Yu, B.; Liu, N.; Tang, S.; Qin, T.; Huang, J. Roles of Glutamate Receptor-Like Channels (GLRs) in Plant Growth and Response to Environmental Stimuli. *Plants* **2022**, *11*, 3450. [[CrossRef](#)] [[PubMed](#)]
188. Corti, F.; Festa, M.; Stein, F.; Stevanato, P.; Siroka, J.; Navazio, L.; Vothknecht, U.C.; Alboresi, A.; Novak, O.; Formentin, E.; et al. Comparative analysis of wild-type and chloroplast MCU-deficient plants reveals multiple consequences of chloroplast calcium handling under drought stress. *Front. Plant Sci.* **2023**, *14*, 1228060. [[CrossRef](#)] [[PubMed](#)]
189. Nomura, H.; Komori, T.; Uemura, S.; Kanda, Y.; Shimotani, K.; Nakai, K.; Furuichi, T.; Takebayashi, K.; Sugimoto, T.; Sano, S.; et al. Chloroplast-mediated activation of plant immune signalling in *Arabidopsis*. *Nat. Commun.* **2012**, *3*, 926. [[CrossRef](#)]
190. Teardo, E.; Segalla, A.; Formentin, E.; Zanetti, M.; Marin, O.; Giacometti, G.M.; Lo Schiavo, F.; Zoratti, M.; Szabo, I. Characterization of a plant glutamate receptor activity. *Cell. Physiol. Biochem.* **2010**, *26*, 253–262. [[CrossRef](#)]
191. Frank, J.; Happeck, R.; Meier, B.; Hoang, M.T.T.; Stribny, J.; Hause, G.; Ding, H.; Morsomme, P.; Baginsky, S.; Peiter, E. Chloroplast-localized BICAT proteins shape stromal calcium signals and are required for efficient photosynthesis. *New Phytol.* **2019**, *221*, 866–880. [[CrossRef](#)]
192. Resentini, F.; Grenzi, M.; Ancora, D.; Cademartori, M.; Luoni, L.; Franco, M.; Bassi, A.; Bonza, M.C.; Costa, A. Simultaneous imaging of ER and cytosolic  $\text{Ca}^{2+}$  dynamics reveals long-distance ER  $\text{Ca}^{2+}$  waves in plants. *Plant Physiol.* **2021**, *187*, 603–617. [[CrossRef](#)] [[PubMed](#)]
193. Li, C.; Duckney, P.; Zhang, T.; Fu, Y.; Li, X.; Kroon, J.; De Jaeger, G.; Cheng, Y.; Hussey, P.J.; Wang, P. TraB family proteins are components of ER-mitochondrial contact sites and regulate ER-mitochondrial interactions and mitophagy. *Nat. Commun.* **2022**, *13*, 5658. [[CrossRef](#)] [[PubMed](#)]
194. Arruda, A.P.; Hotamisligil, G.S. Calcium Homeostasis and Organelle Function in the Pathogenesis of Obesity and Diabetes. *Cell Metab.* **2015**, *22*, 381–397. [[CrossRef](#)] [[PubMed](#)]
195. Resentini, F.; Ruberti, C.; Grenzi, M.; Bonza, M.C.; Costa, A. The signatures of organellar calcium. *Plant Physiol.* **2021**, *187*, 1985–2004. [[CrossRef](#)]
196. Deutsch, R.; Kudrina, V.; Freichel, M.; Grimm, C. Two-pore channel regulators—Who is in control? *Front. Physiol.* **2024**, *15*, 1534071. [[CrossRef](#)]

197. Jin, X.; Zhang, Y.; Alharbi, A.; Hanbashi, A.; Alhoshani, A.; Parrington, J. Targeting Two-Pore Channels: Current Progress and Future Challenges. *Trends Pharmacol. Sci.* **2020**, *41*, 582–594. [[CrossRef](#)]
198. Lagostena, L.; Festa, M.; Pusch, M.; Carpaneto, A. The human two-pore channel 1 is modulated by cytosolic and luminal calcium. *Sci. Rep.* **2017**, *7*, 43900. [[CrossRef](#)]
199. Nagata, T.; Iizumi, S.; Satoh, K.; Ooka, H.; Kawai, J.; Carninci, P.; Hayashizaki, Y.; Otomo, Y.; Murakami, K.; Matsubara, K.; et al. Comparative analysis of plant and animal calcium signal transduction element using plant full-length cDNA data. *Mol. Biol. Evol.* **2004**, *21*, 1855–1870. [[CrossRef](#)]
200. Edel, K.H.; Marchadier, E.; Brownlee, C.; Kudla, J.; Hetherington, A.M. The Evolution of Calcium-Based Signalling in Plants. *Curr. Biol.* **2017**, *27*, R667–R679. [[CrossRef](#)]
201. Xiao, X.H.; Yang, M.; Sui, J.L.; Qi, J.Y.; Fang, Y.J.; Hu, S.N.; Tang, C.R. The calcium-dependent protein kinase (CDPK) and CDPK-related kinase gene families in *Hevea brasiliensis*-comparison with five other plant species in structure, evolution, and expression. *FEBS Open Bio* **2017**, *7*, 4–24. [[CrossRef](#)]
202. Patel, S.; Marchant, J.S.; Brailoiu, E. Two-pore channels: Regulation by NAADP and customized roles in triggering calcium signals. *Cell Calcium* **2010**, *47*, 480–490. [[CrossRef](#)]
203. Dematteis, G.; Tapella, L.; Casali, C.; Talmon, M.; Tonelli, E.; Reano, S.; Ariotti, A.; Pessolano, E.; Malecka, J.; Chrostek, G.; et al. ER-mitochondria distance is a critical parameter for efficient mitochondrial  $\text{Ca}^{2+}$  uptake and oxidative metabolism. *Commun. Biol.* **2024**, *7*, 1294. [[CrossRef](#)]
204. Stael, S.; Wurzing, B.; Mair, A.; Mehler, N.; Voithknecht, U.C.; Teige, M. Plant organellar calcium signalling: An emerging field. *J. Exp. Bot.* **2012**, *63*, 1525–1542. [[CrossRef](#)]
205. Carraretto, L.; Checchetto, V.; De Bortoli, S.; Formentin, E.; Costa, A.; Szabo, I.; Teardo, E. Calcium Flux across Plant Mitochondrial Membranes: Possible Molecular Players. *Front. Plant Sci.* **2016**, *7*, 354. [[CrossRef](#)] [[PubMed](#)]
206. Cortese, E.; Moscatiello, R.; Pettiti, F.; Carraretto, L.; Baldan, B.; Frigerio, L.; Voithknecht, U.C.; Szabo, I.; De Stefani, D.; Brini, M.; et al. Monitoring calcium handling by the plant endoplasmic reticulum with a low- $\text{Ca}^{2+}$ -affinity targeted aequorin reporter. *Plant J.* **2022**, *109*, 1014–1027. [[CrossRef](#)] [[PubMed](#)]
207. Ramazanov, B.R.; Parchure, A.; Di Martino, R.; Kumar, A.; Chung, M.; Kim, Y.; Griesbeck, O.; Schwartz, M.A.; Luini, A.; von Blume, J. Calcium flow at ER-TGN contact sites facilitates secretory cargo export. *Mol. Biol. Cell* **2024**, *35*, ar50. [[CrossRef](#)] [[PubMed](#)]
208. Rayl, M.; Truitt, M.; Held, A.; Sargeant, J.; Thorsen, K.; Hay, J.C. Penta-EF-Hand Protein Peflin Is a Negative Regulator of ER-To-Golgi Transport. *PLoS ONE* **2016**, *11*, e0157227. [[CrossRef](#)]
209. Malek, M.; Wawrzyniak, A.M.; Koch, P.; Luchtenborg, C.; Hessenberger, M.; Sachsenheimer, T.; Jang, W.; Brugger, B.; Haucke, V. Inositol triphosphate-triggered calcium release blocks lipid exchange at endoplasmic reticulum-Golgi contact sites. *Nat. Commun.* **2021**, *12*, 2673. [[CrossRef](#)]
210. Yang, Y.; Valencia, L.A.; Lu, C.H.; Nakamoto, M.L.; Tsai, C.T.; Liu, C.; Yang, H.; Zhang, W.; Jahed, Z.; Lee, W.R.; et al. Plasma membrane curvature regulates the formation of contacts with the endoplasmic reticulum. *Nat. Cell Biol.* **2024**, *26*, 1878–1891. [[CrossRef](#)]
211. Gil, D.; Guse, A.H.; Dupont, G. Three-Dimensional Model of Sub-Plasmalemmal  $\text{Ca}^{2+}$  Microdomains Evoked by the Interplay Between ORAI1 and InsP(3) Receptors. *Front. Immunol.* **2021**, *12*, 659790. [[CrossRef](#)]
212. Liu, L.; Li, J. Communications Between the Endoplasmic Reticulum and Other Organelles During Abiotic Stress Response in Plants. *Front. Plant Sci.* **2019**, *10*, 749. [[CrossRef](#)] [[PubMed](#)]
213. Saheki, Y.; De Camilli, P. Endoplasmic Reticulum-Plasma Membrane Contact Sites. *Annu. Rev. Biochem.* **2017**, *86*, 659–684. [[CrossRef](#)]
214. Pottosin, I.; Dobrovinskaya, O. Major vacuolar TPC1 channel in stress signaling: What matters,  $\text{K}^+$ ,  $\text{Ca}^{2+}$  conductance or an ion-flux independent mechanism? *Stress Biol.* **2022**, *2*, 31. [[CrossRef](#)] [[PubMed](#)]
215. Peng, W.; Wong, Y.C.; Krainc, D. Mitochondria-lysosome contacts regulate mitochondrial  $\text{Ca}^{2+}$  dynamics via lysosomal TRPML1. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 19266–19275. [[CrossRef](#)] [[PubMed](#)]
216. Rosencrans, W.M.; Rajendran, M.; Bezrukov, S.M.; Rostovtseva, T.K. VDAC regulation of mitochondrial calcium flux: From channel biophysics to disease. *Cell Calcium* **2021**, *94*, 102356. [[CrossRef](#)]
217. Sargsyan, Y.; Bickmeyer, U.; Gibhardt, C.S.; Streckfuss-Bomeke, K.; Bogeski, I.; Thoms, S. Peroxisomes contribute to intracellular calcium dynamics in cardiomyocytes and non-excitable cells. *Life Sci. Alliance* **2021**, *4*, e202000987. [[CrossRef](#)]
218. Sargsyan, Y.; Kalinowski, J.; Thoms, S. Calcium in peroxisomes: An essential messenger in an essential cell organelle. *Front. Cell Dev. Biol.* **2022**, *10*, 992235. [[CrossRef](#)]
219. Valm, A.M.; Cohen, S.; Legant, W.R.; Melunis, J.; Hershsberg, U.; Wait, E.; Cohen, A.R.; Davidson, M.W.; Betzig, E.; Lippincott-Schwartz, J. Applying systems-level spectral imaging and analysis to reveal the organelle interactome. *Nature* **2017**, *546*, 162–167. [[CrossRef](#)]

220. Faxen, K.; Andersen, J.L.; Gourdon, P.; Fedosova, N.; Morth, J.P.; Nissen, P.; Moller, J.V. Characterization of a *Listeria monocytogenes* Ca<sup>2+</sup> pump: A SERCA-type ATPase with only one Ca<sup>2+</sup>-binding site. *J. Biol. Chem.* **2011**, *286*, 1609–1617. [[CrossRef](#)]
221. Prabudiansyah, I.; Oradd, F.; Magkakis, K.; Pounot, K.; Levantino, M.; Andersson, M. Dephosphorylation and ion binding in prokaryotic calcium transport. *Sci. Adv.* **2024**, *10*, eadp2916. [[CrossRef](#)]
222. Suarez-Delgado, E.; Islas, L.D. A novel origin for calcium selectivity. *eLife* **2020**, *9*, e55216. [[CrossRef](#)]
223. Kolodkin-Gal, I.; Parsek, M.R.; Patrauchan, M.A. The roles of calcium signaling and calcium deposition in microbial multicellularity. *Trends Microbiol.* **2023**, *31*, 1225–1237. [[CrossRef](#)] [[PubMed](#)]
224. King, M.M.; Kayastha, B.B.; Franklin, M.J.; Patrauchan, M.A. Calcium Regulation of Bacterial Virulence. *Adv. Exp. Med. Biol.* **2020**, *1131*, 827–855. [[CrossRef](#)] [[PubMed](#)]
225. Zhou, H.; Hu, Y.-Y.; Tang, Z.-X.; Jiang, Z.-B.; Huang, J.; Zhang, T.; Shen, H.-Y.; Ye, X.-P.; Huang, X.-Y.; Wang, X.; et al. Calcium Transport and Enrichment in Microorganisms: A Review. *Foods* **2024**, *13*, 3612. [[CrossRef](#)]
226. Steiner, P.; Buchner, O.; Andosch, A.; Wanner, G.; Neuner, G.; Lutz-Meindl, U. Fusion of Mitochondria to 3-D Networks, Autophagy and Increased Organelle Contacts are Important Subcellular Hallmarks during Cold Stress in Plants. *Int. J. Mol. Sci.* **2020**, *21*, 8753. [[CrossRef](#)]
227. Steiner, P.; Luckner, M.; Kerschbaum, H.; Wanner, G.; Lutz-Meindl, U. Ionic stress induces fusion of mitochondria to 3-D networks: An electron tomography study. *J. Struct. Biol.* **2018**, *204*, 52–63. [[CrossRef](#)]
228. Edel, K.H.; Kudla, J. Increasing complexity and versatility: How the calcium signaling toolkit was shaped during plant land colonization. *Cell Calcium* **2015**, *57*, 231–246. [[CrossRef](#)]
229. Swarbreck, S.M.; Colaco, R.; Davies, J.M. Plant calcium-permeable channels. *Plant Physiol.* **2013**, *163*, 514–522. [[CrossRef](#)]
230. Kudla, J.; Becker, D.; Grill, E.; Hedrich, R.; Hippler, M.; Kummer, U.; Parniske, M.; Romeis, T.; Schumacher, K. Advances and current challenges in calcium signaling. *New Phytol.* **2018**, *218*, 414–431. [[CrossRef](#)]
231. Plieth, C. Plant calcium signaling and monitoring: Pros and cons and recent experimental approaches. *Protoplasma* **2001**, *218*, 1–23. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.