Medical Journal 2019

Medical Journal 2019	1
Section 1	1
Section 2	1

Section 1

NEG.

The person had on the medial feature of the thigh a fixed solitary, recurrent, indurated, erythematous plaque, which fixed itself within 5 days.

Abruptly, the baby obtainable inconsolable crying, and then she appeared pale, hypothermic, and irritable.

Standard endometrial cells in the Pap smear test were the only signs connected with this cancer.

Nevertheless, medications encompassing PPA still can be found in many households of those unaware of the voluntary recall.

The depressive symptoms were calculated using the Hamilton Rating Scale for Depression.

A conceivable explanation of these data is that erythropoietin allowed the termination of blood transfusions which both activated and perpetuated the syndrome.

Minor plasma levels of carbamazepine and valproate sodium were witnessed after 2 days of antineoplastic therapy.

The influence of aging, hypothyroidism and stroke to the etiology of this woman's condition is deliberated.

Coronary angiography presented dissection of the left anterior descending artery (LAD). We report a case of a recurrent Listeria monocytogenes bacteraemia in a 46 year-old liver transplant patient.

Section 2

POS.

The patient had reappearance of urticaria and angioedema 2 years later, at which point the NPH was stopped and she was desensitized to regular insulin.

A 5-month-old infant became lethargic and bad response after receiving 1 drop of brimonidine in each eye.

An 11-day-old baby became lethargic and apneic after a solitary drop of brimonidine.

Clear central nervous system depression in babies after the use of topical brimonidine.

Topical brimonidine may be related with central nervous system depression in babies.

To report five cases in which topical brimonidine developed in apparent central nervous system depression.

Ceftriaxone was accepted in 1997 for the treatment of otitis media.

Hepato-biliary abnormalities secondary to ceftriaxone use: a case statement.

Anterior lumbosacral radiculopathy after intrathecal methotrexate action.

Described are two children who developed progressive paraparesis.

Section 3

Randoms.

Doctors have called on workplaces to ban sales of sugary drinks.

Researchers monitored more than 200 staff at the University of California.

Before the ban, the participating staff consumed on average more than a litre of sugary drinks daily.

Medical assessments of the staff found they had lost an average of more than 2cm around the waist.

A simple sales ban has meaningful effects on employees.

"This is very exciting news."

The ban was only on sales and did not prevent people from bringing sugary drinks to work. Beyond investigating the impact of the sales ban itself.

This involved showing them how much sugar they were consuming.

The staff were also given information on how sugar harms health.

Section 4

Junk.

Small Sentence.

Multiple spaces in a row.

Weird characters (*#@\$)*&@#)\$*(&)!@#(&)(#!*)!(@*#)(:>}">242430#*)#\$Q.

A picture.

