

Supplement to Strong Tree-Childness Is a Sharp Identifiability Boundary for Level-2 Jukes–Cantor Networks

Alec Kriebel
Independent Researcher

August 2026

1 Purpose and scope

This supplement is a reader-oriented map from the mathematical statements in the article to the exact reproducibility package. It does not broaden the theorem. The positive result concerns binary, LSA-valid, already-simple, reticulation-preserving semi-directed strongly tree-child level-2 networks under the open four-state Jukes–Cantor model

$$0 < x_e < 1, \quad 0 < \lambda_r < 1.$$

Admissible rootings reduce to one fixed mixed graph after the single root suppression in Definition 2.1 of the article; arbitrary hidden rooted refinements are not included.

The relation $N \preceq_{\text{JC}} N'$ means that a source-regular common point has a relatively open neighborhood in the regular source-model locus contained in the target image. The relation $N \bowtie_{\text{JC}} N'$ requires a full-dimensional regular stochastic germ in both directions. Neither relation means equality of complex model closures or equality of the complete open stochastic images.

2 Theorem and proof map

The main theorem states

$$N \preceq_{\text{JC}} N' \iff \mathcal{C}(N, N'),$$

where $\mathcal{C}(N, N')$ means that the labelled reduced bridge trees agree and corresponding nontrivial blobs are labelled-isomorphic or differ by ordinary triangle redirection **T**. The same condition characterizes $N \bowtie_{\text{JC}} N'$, so there is no proper one-sided generic containment. The logical dependencies are:

- P1.** The fixed-mixed-graph convention and no-omnian criterion identify the topology class and imply that a level-2 strong blob has at most one triangle.
- P2.** Exact Fourier flattening minors recover every cut split pointwise, including the crossing case with two active endpoint tensors.
- P3.** Positive rank-one cut factorization recovers the bridge tree’s local tensors only modulo the full incidence-scaling action. It does not recover physical bridge multipliers. The physical fibre

is the ambient incidence orbit intersected with the complete open realization locus, including every local-arm, bridge, and inheritance constraint.

- P4.** Every nontrivial level-2 factor contracts to a cycle or one of four directed theta cores.
- P5.** A decorated graph-to-polynomial atlas classifies every rigid bounded relation in both containment directions. The only nonisomorphic common local germ is ordinary triangle redirection.
- P6.** Marginal submersions and coherent one- and two-port probes restore every arbitrary subdivision word from the bounded support.
- P7.** Projective localization turns global containment into independent local containments; the bridge tree has no scaling holonomy.
- P8.** Conversely, finitely many local triangle germs glue through the common bridge tree, proving the global common germ.

For a fixed source topology, the singular locus, lower-rank images, wrong topology intersections, and certified nonzero witness hypersurfaces form a proper algebraic exceptional subset. Outside it, the exact distribution determines the labelled topology modulo T .

3 Exact convention checks

The independent convention implementation reconstructs rooted bidegrees, acyclicity, the lowest-stable-ancestor condition, one-step root suppression, mixed-graph simplicity, retained arrowheads, admissible rootings, and the tree-child property. Its relevant exact checks are:

Object	exact count or outcome
Primitive support mixed graphs	12
Admissible rootings of those supports	100
Tree-child rootings of the simple double-triangle $K_4 - e$ core	0
Hidden one-step zipper admitted as a rooting of the final tree	no
Nested hidden zipper admitted as a rooting of the final tree	no

The last two regressions prevent the rejected cleanup-fibre convention from re-entering the active theorem. Exhaustive cleanup remains permitted only in explicit restriction or displayed-network operations; it does not enlarge the admissible-rooting set of a fixed mixed graph.

4 Local classification certificates

The finite theorem object is a directed decorated relation, not a target graph identifier. Each normalized record retains both mixed graphs, side colours, independently distinguished incoming boundaries, physical port matching, source-to-target direction, and all vertex, edge, inheritance, label, and Fourier-coordinate transports. The verifier regenerates the chain

graphs $\rightarrow$ switchings $\rightarrow$ descendant masks $\rightarrow$ Fourier tensors $\rightarrow$ pullback $\rightarrow$ sign/rank certificate.

The following counts are checksums of the regenerated universe rather than premises of the proof.

Certificate family	exact count
Raw three-outgoing decorated presentations	10,826
Canonical three-outgoing directed relations	10,466
Strict three-outgoing relations	5,284
Restoration-root raw coverages	5,344
Three-outgoing restoration states	68,584
Three-outgoing isomorphism / T terminals	120 / 24
Direct residual anchors	62
Four-outgoing completion records	6,138
Four-outgoing raw survivors	192
Four-outgoing restoration states	2,106
One-port direct-anchor probes	2,642
Two-port direct-anchor probes	18,224

Every strict relation has a regenerated exact polynomial, flattening minor, or Bernstein-basis sign certificate on the complete open cube. Every equality terminal is independently canonicalized and accepted only as a labelled isomorphism or ordinary T. Mutation tests reject deleted relations, merged source embeddings, altered port matches, source–target reversal, separator reassignment, broken transports, missing hard-cover keys, and incoherent probes.

5 Triangle-free Omega sharpness certificate

The historical four-leaf Omega pair was admitted only after a bounded primary and clean-room audit. Each fixed mixed graph is simple, binary, triangle-free, level two, and has seven admissible rootings, exactly two of which are tree-child. Hence both lie in $W_{TC} \setminus S_{TC}$. The source and target are labelled nonisomorphic: leaf 1 has unordered distances $\{3, 3\}$ to the reticulations in the source and $\{2, 4\}$ in the target.

The exact nine-variable rational correspondence annihilates all 64 zero-sum Fourier differences. At its certified point, all 256 Fourier entries and all 256 inverse-pattern probabilities agree and every parameter is strict. The four exact rank-nine minors are

$$\begin{array}{cc}
\frac{171}{2305843009213693952000000}, & \frac{513}{9223372036854775808000000}, \\
\frac{57}{576460752303423488000000}, & \frac{189}{2305843009213693952000000}.
\end{array}$$

With the four pendant multipliers set to one, the core Jacobian has shape 14×10 and exact rational-function rank six. In the edge order of the paper, the first six nonconstant orbit rows and columns $(a_0, a_1, a_2, a_3, a_4, a_7)$ have determinant

$$-\frac{723}{8589934592} \neq 0$$

at the strict source point. Four pendant torus directions add at most four dimensions, while

$$a_4 \partial_{a_4} c_g + a_7 \partial_{a_7} c_g = \mathbf{1}[g_4 \neq 0] c_g$$

identifies the leaf-4 pendant direction with a core tangent direction. This gives the exact upper bound nine. Thus both complete models and their local intersection have dimension nine. Repeated

identical cherry substitution has a positive analytic inverse, adds two dimensions per leaf, and gives the triangle-free family of dimension $2n + 1$ for every $n \geq 4$.

The clean-room release verifier rejects all twelve mandated mutations, including incomplete rooting enumeration, a changed arrowhead or parent, boundary parameters, a changed regenerated Fourier coordinate, and a replaced rank-nine minor.

6 Triangle-containing Theta sharpness certificate

The independent frozen Theta package verifies a second mechanism. Its two four-leaf theta graphs are nonisomorphic and not T-equivalent, lie in $W_{TC} \setminus S_{TC}$, agree on all 256 Fourier coordinates at a strict quadratic point, and have nonzero rank-eight minors. The common irreducible JC model has dimension eight. The same cherry inverse extends this family to every $n \geq 4$, with common dimension $2n$. Theta is not used as a move in the positive strong-class proof.

7 Deterministic reproduction

From the repository root, in a clean checkout, run

```
bash reproducibility/verify_quick.sh
bash reproducibility/verify_full.sh
bash reproducibility/verify_regenerate_all.sh
```

The quick command checks active scope, hashes, crosswalks, convention regressions, and the independent Omega release record. The full command replays every committed exact component certificate. The regeneration command starts from primitive graph encodings and rebuilds the bounded relation, bridge, restoration, probe, triangle, Omega, and Theta evidence.

The whole-proof checker is a fail-closed consistency gate, not a substitute for the component calculations. The mathematical evidence is supplied by the graph-derived component scripts and their independently implemented replays. Complete commands, environment versions, elapsed times, peak-memory records, output hashes, and clean-status checks are stored in the release transcripts.

8 Theorem-to-artifact crosswalk

Claim	Repository-relative evidence
Convention and no-omnian characterization	<code>s_tc_jc_landmark_closure/reviews/final_standard_convention/</code>
Pointwise cut preservation and bridge fibre	<code>s_tc_jc_landmark_closure/reviews/global_bridge/</code>
Decorated bounded relation atlas	<code>s_tc_jc_landmark_closure/reviews/n3_universe_generator/</code> and <code>s_tc_jc_landmark_closure/reviews/bounded_directed_relation_cleanroom/</code>
Four-outgoing theta gate	<code>s_tc_jc_landmark_closure/reviews/theta2_signature_gate/</code>
Restoration and coherent probes	<code>s_tc_jc_landmark_closure/reviews/final_hard_cover_cleanroom/</code> and <code>s_tc_jc_landmark_closure/reviews/direct_anchor_probe_closure/</code>

Claim	Repository-relative evidence
Ordinary triangle germ	<code>s_tc_jc_landmark_closure/reviews/triangle_redirection_cleanroom/</code>
Triangle-free Omega family	<code>omega_audit/</code>
Triangle-containing Theta family	<code>s_tc_jc_sharp_boundary/</code>
Active release consistency	<code>s_tc_jc_landmark_closure/reproducibility/verify_active_release.py</code>

9 Interpretation limits

The theorem reconstructs topology, not all numerical parameters. In particular, bridge data determine projective local tensor orbits and normalized effective scales, not physical bridge lengths. The ordinary triangle result is a common full-dimensional regular germ; it does not assert equality of complete stochastic images or one orientation compatible with every distribution. No pointwise classification of the exceptional locus, richer substitution model, finite-sample inference result, or complete weak-class classification is claimed.