A Martin kernel with isolated Harnack contact in an annulus

Alec Kriebel
Independent researcher
ORCID: 0009-0001-9320-500X

12 October 2026
Version 1.0.0; unrefereed research note

Abstract

For the annulus $1 < |z| < e$, we exhibit a normalized minimal Martin kernel that is strictly below the envelope of normalized positive harmonic functions at every point of a punctured neighborhood of the base point. Four explicit positive harmonic competitors give a uniform first-order gap in every direction. The result disproves the extension to multiply connected domains of the equality-along-a-Green-line property in Hayman and Lingham's Problem 7.42, attributed to Arsove and Johnson. We give direct constructions of the Green and Martin kernels and a proof of minimality. OpenAI tools were used extensively in developing and independently checking the argument. This note has not undergone human peer review or formal proof-assistant verification.

1 The question and the counterexample

For a plane Green domain D and $a \in D$, define

$$H_a(z) = \sup\{h(z) : h > 0 \text{ harmonic on } D,\ h(a) \le 1\}.$$

A finite Harnack chain shows that $H_a(z) < \infty$ for every $z \in D$; also $H_a(a) = 1$. In the disc centered at a, a normalized boundary Poisson kernel equals H_a on the radius toward its pole. Problem 7.42 of Hayman and Lingham [1] asks whether the corresponding property, equality along a Green line issuing from a for each minimal Martin kernel, persists in multiply connected domains. We use this equality-along-the-line meaning, supplied by the disc example, rather than equality only at a single point. The latter holds automatically at a.

Theorem 1. *Let $D = \{z \in \mathbb{C} : 1 < |z| < e\}$, $a = e^{1/2}$ and $\xi = -1$. The minimal Martin kernel K_ξ at the inner-boundary point ξ, normalized by $K_\xi(a) = 1$, satisfies*

$$K_\xi(z) < H_a(z) \qquad (z \in U \setminus \{a\})$$

for some neighborhood U of a in D.

Thus the kernel cannot equal the envelope along any nontrivial curve issuing from a, and in particular along a Green line. The theorem does not address isolated contact farther from a. A single such kernel in this smooth doubly connected Green domain negates the proposed universal extension.

2 The Green and Martin kernels

Write $z = e^{x+iy}$, where $0 < x < 1$ and y is taken modulo 2π. Harmonic functions on D become periodic harmonic functions on this cylinder. The base point is $A = (1/2, 0)$ and ξ is $(0, \pi)$. For the infinite strip, the Poisson density on the side $x = 0$, relative to its horizontal boundary coordinate, is

$$p(x,t) = \frac{\sin(\pi x)}{2(\cosh(\pi t) - \cos(\pi x))}. \tag{1}$$

Indeed the map $w = y + ix \mapsto e^{\pi w}$ sends the strip to the upper half-plane; transforming its Poisson density yields (1). Periodize it and normalize:

$$P(x,y) = \sum_{k\in\mathbb{Z}} p(x, y - \pi + 2\pi k), \qquad K(x,y) = \frac{P(x,y)}{P(A)}. \tag{2}$$

Every term is positive. On compact subsets of the interior, each fixed finite order of derivatives has an exponentially summable bound in $|k|$. Consequently P is smooth, positive, periodic and harmonic, and termwise differentiation is valid.

We identify K with the required Martin kernel directly. With $w = y + ix$ and $w' = \eta + is$, the strip Green function with logarithmic singularity $-\log|w - w'|$ is

$$\begin{aligned} g_S(w,w') &= \log\left|\frac{e^{\pi w} - e^{\pi\overline{w'}}}{e^{\pi w} - e^{\pi w'}}\right| \\ &= \frac{1}{2}\log\frac{\cosh(\pi(y-\eta)) - \cos(\pi(x+s))}{\cosh(\pi(y-\eta)) - \cos(\pi(x-s))}. \end{aligned} \tag{3}$$

The numerator in the last ratio exceeds its denominator by $2\sin(\pi x)\sin(\pi s)$, so this function is positive in the strip and vanishes on both sides. Its periodization

$$g_C(w,w') = \sum_{k\in\mathbb{Z}} g_S(w, w' + 2\pi k)$$

converges locally with derivatives away from the pole by exponential decay in horizontal separation. It is periodic, has one logarithmic pole on the quotient cylinder and vanishes on its boundary. The maximum principle identifies it with the cylinder Green function.

Differentiating (3) at $s = 0$ gives

$$\partial_s g_S(y + ix, \eta + is)|_{s=0} = 2\pi p(x, y - \eta).$$

The periodized series and its boundary derivatives converge uniformly when the first variable ranges over an interior compact set, η is near π, and $s \geq 0$ is small. For $P_\eta(x,y) = \sum_k p(x, y - \eta + 2\pi k)$, Taylor expansion therefore gives

$$g_C(w, \eta + is) = 2\pi s P_\eta(w) + O(s^2).$$

Since $P_\pi(A) > 0$, division by the same expression at A shows

$$\frac{g_C(w, \eta + is)}{g_C(A, \eta + is)} \longrightarrow K(w) \qquad \text{as } (\eta, s) \to (\pi, 0), \quad s > 0, \tag{4}$$

locally uniformly in the interior. The expansion is uniform in η, so the approach need not be nontangential.

To establish minimality, suppose v is harmonic and $0 \le v \le P$. Away from $(0,\pi)$, P has continuous zero boundary values, and so does v. Near that point put $t = y - \pi$ and $r^2 = x^2 + t^2$. Expansion of (1), with the other summands smooth and vanishing on $x = 0$, gives

$$P(x, \pi + t) = \frac{x}{\pi r^2} + O(x), \tag{5}$$

uniformly as $r \to 0$ in the half-disc. Odd reflection of v across $x = 0$ is harmonic in a punctured disc and has size $O(1/r)$. The single-valued real harmonic Laurent expansion then allows only a logarithmic term, degree-one singular terms and a regular harmonic part: taking circle Fourier coefficients excludes all negative powers of degree at least two. Oddness eliminates the logarithmic and tangential singular terms. Hence

$$v(x, \pi + t) = B\frac{x}{r^2} + v_0(x, t),$$

where v_0 is harmonic across the origin and odd in x. Along $t = 0$, domination and (5) give $0 \le B \le 1/\pi$. The difference $v - \pi BP$ has its singularity canceled and extends continuously with value zero at every point of the compact annular boundary. The maximum principle gives $v = \pi BP$. Thus P is minimal. Together with (4), this proves that K is the normalized minimal Martin kernel K_ξ.

3 An all-direction competitor gap

Set

$$q_k = \operatorname{sech}((2k-1)\pi^2), \qquad S_1 = \sum_k q_k, \qquad S_2 = \sum_k q_k^2.$$

At A, direct differentiation of (1) and pairing k with $1 - k$ give

$$P(A) = \tfrac{1}{2}S_1, \qquad P_x(A) = -\tfrac{\pi}{2}S_2, \qquad P_y(A) = 0.$$

It follows that

$$\nabla K(A) = (-c, 0), \qquad c = \pi S_2/S_1. \tag{6}$$

The ratio S_2/S_1 is a positive weighted mean of numbers at most $\operatorname{sech}(\pi^2)$. Using $3 < \pi < 4$ and $\cosh(\pi^2) > 1 + \pi^4/2 > 83/2$, we obtain the exact bound

$$0 < c \le \pi \operatorname{sech}(\pi^2) < \frac{8}{83} < \frac{1}{10}. \tag{7}$$

No numerical truncation is used here.

Let $b = 1/(2\cosh(1/2))$ and define four competitors

$$h_{x,\pm}(x, y) = 1 \pm (x - 1/2),$$
$$h_{y,\pm}(x, y) = 1 \pm b \sin y \cosh(x - 1/2).$$

They are periodic and harmonic, equal one at A, and are positive throughout the cylinder: the first pair lies between $1/2$ and $3/2$, and the second pair is at least $1/2$. Thus each is admissible for H_a. Their gradients at A are $(\pm 1, 0)$ and $(0, \pm b)$. The series for $\cosh(1/2)$ gives

$$\cosh(1/2) \le \sum_{n \ge 0} (1/4)^n = 4/3 < 2, \qquad b > 1/4.$$

For $d = (u, v) \neq 0$, write $\rho = |d|$ and $M = \max\{|u|, b|v|\}$. Equations (6)–(7) give

$$\max_j (\nabla h_j(A) - \nabla K(A)) \cdot d = M + cu \geq (1 - c)M,$$
$$M \geq \tfrac{1}{2}(|u| + b|v|) \geq \tfrac{b}{2}(|u| + |v|) > \rho/8.$$

Consequently

$$\max_j (\nabla h_j(A) - \nabla K(A)) \cdot d > \frac{75}{664}\rho > \rho/10. \tag{8}$$

This is a uniform bound over all real directions; it is not a sampled inequality.

The finite family $h_j - K$ is smooth near A and vanishes at A. On a sufficiently small closed coordinate disc, Taylor's theorem supplies a common constant $C_0 \geq 1$ with remainders bounded by $C_0\rho^2$. By (8), for $0 < \rho < \min\{r_0, 1/(20C_0)\}$ we have

$$\max_j h_j(A + d) - K(A + d) > \rho/10 - C_0\rho^2 > 0.$$

Since $H_a \geq \max_j h_j$, this proves Theorem 1. No smoothness of the envelope is assumed. A curve carrying equality and having non-base points arbitrarily close to a would contradict the theorem, which yields the stated negative answer for Green lines.

4 Attribution, scope and verification

The question is tied to Problem 7.42 and its update in the inspected 2018 version of [1], printed pages 173–174. That update reports no progress known to its editors at that time. This historical report is not a certification of current worldwide openness. The strip kernels, conformal change of coordinates, harmonic reflection and maximum principle are classical tools. The contribution presented here is their explicit use to negate the fixed-Martin-kernel Green-line property.

Related work on Harnack and other conformally invariant metrics includes Herron [2] and Bear and Smith [3]. The one-sided normalized envelope used here should be distinguished from symmetric Harnack distances. A dated, bounded priority audit accompanies the verification package. The note makes no claim of being the first historical resolution, and absence of an identified earlier counterexample is not proof of priority.

The companion materials retain the original authored proof and its independent AI reviews, identify their exact source commit, and provide reproducible checks with their limits stated explicitly. The theorem is established by the analytic argument above; integrity checks and numerical experiments do not replace that proof. OpenAI tools were used extensively in solving, drafting and adversarially checking the result. The author is responsible for the submission. This is an unrefereed preprint, with no completed human peer review or proof-assistant certification claimed.

References

[1] W. K. Hayman and E. F. Lingham, *Research Problems in Function Theory (New Edition)*, arXiv:1809.07200v2 (2018), Problem 7.42 and Update 7.42, pp. 173–174. arXiv:1809.07200v2.

[2] D. A. Herron, The Harnack and other conformally invariant metrics, *Kodai Math. J.* **10** (1987), 9–19. doi:10.2996/kmj/1138037356.

[3] H. S. Bear and W. Smith, A tale of two conformally invariant metrics, *J. Math. Anal. Appl.* **318** (2006), 498–506. doi:10.1016/j.jmaa.2005.05.065.