

Technical Supplement to Exact Diffusion Design for Maximally Collective Stable Turing Patterns in Binary-Complex Mass-Action Networks

Alec Kriebel

August 2026

Contents

S1	Reaction matrices, flux cone, and semipositive conservation	1
S2	All-spectrum principal-block classification	2
S3	The principal-minor diffusion-ray theorem in full detail	3
S4	Order- $(n - 1)$ omission minors and diffusion design	3
S5	Improved unit-equilibrium critical profile	4
S6	Conservation-compatible stable-mode corrections	6
S7	Equilibrium-scaled stable family	8
S8	Exact coefficient tables	13
S8.1	Coefficientwise-nonnegative modulus certificates	13
S8.2	Signed scalar and rational-function certificates	19
S9	Near-threshold control example	22
S10	Semilinear stability and robustness	23
S11	Numerical protocol and independent verification	24

S1 Reaction matrices, flux cone, and semipositive conservation

Order species as $X_1, \dots, X_m, Z$ and reactions as $R_0, R_2, \dots, R_{m-2}, R_a, R_b, R_+, R_-$. For each indexed reaction, the source and target vectors have total molecularity at most two. The source matrix Y_m and stoichiometric matrix Γ_m are obtained by placing these source vectors and reaction differences in columns.

The balance equations $\Gamma_m v = 0$ read, successively, equality of every flux around the long circuit and equality of the two boundary-pair fluxes. Thus

$$\ker \Gamma_m = \text{span}\{(\mathbf{1}_m, 0, 0), (0_m, 1, 1)\}.$$

The balance equations show that this kernel is exactly two-dimensional. Since Γ_m has $m + 2$ columns, rank-nullity gives $\text{rank } \Gamma_m = m$. As an independent explicit check, retain the rows $X_1, X_3, \dots, X_m, Z$ and the columns $R_2, \dots, R_{m-2}, R_a, R_b, R_+$, with the chain range empty when $m = 3$. The resulting maximal minor has determinant $4(-1)^m$. The vector

$$c = (0, 4, \dots, 4, 2, 1)^T$$

is orthogonal to every reaction vector, so $\text{im } \Gamma_m = c^\perp$. Notice that c is semipositive rather than positive: X_1 is absent.

For flux $(a\mathbf{1}_m, b, b)$ and unit equilibrium, $A_m(a, b) = \Gamma_m \text{diag}(a\mathbf{1}_m, b, b)Y_m^T$ has precisely the entries printed in the main text. At an arbitrary positive equilibrium, right multiplication by $H = \text{diag}(1/x_i^*)$ gives $J = A_m(a, b)H$. Conversely the rate assignment $k_r = v_r/(x^*)^{y_r}$ realizes every $a, b, H > 0$.

S2 All-spectrum principal-block classification

Use the directed convention $X_j \rightarrow X_i$ when the (i, j) Jacobian entry is nonzero. The interior arrows are the one-way chain $X_2 \rightarrow X_3 \rightarrow \cdots \rightarrow X_{m-1}$. The boundary arrows are those incident to X_1, X_m, Z and the returns $X_{m-1} \rightarrow X_1, X_m \rightarrow X_2$. The following case analysis is exhaustive for an induced set with fewer than m vertices.

For $m = 3$, inspect the principal sets of order at most two directly. The sets $\{X_1, X_2\}$ and $\{X_2, X_3\}$ are the two complete long cycles; every other non-singleton strongly connected component is a principal subgraph of $\{X_1, X_3, Z\}$. Hence assume $m \geq 4$ in the chain argument below.

1. If a component contains a chain arrow and returns through $X_{m-1} \rightarrow X_1$, then the unique forward path from X_1 forces every vertex $X_1, \dots, X_{m-1}$. This is the first complete long cycle.
2. If it contains a chain arrow and returns through $X_m \rightarrow X_2$, then the forward path forces every vertex $X_2, \dots, X_m$. This is the second complete long cycle.
3. The alternative chain closure $X_1 \rightarrow X_2 \rightarrow \cdots \rightarrow X_m \rightarrow X_1$ uses m vertices and is excluded. Adding any boundary vertex to either complete long cycle also uses at least m vertices.
4. If neither long cycle is complete, every feedback path not involving a complete chain lies inside $\{X_1, X_m, Z\}$. All other retained chain vertices have no return path and therefore form negative singleton diagonal blocks.

At $b = 2a$, the coefficient A_{m1} vanishes. This deletes the arrow $X_1 \rightarrow X_m$. That arrow lies on neither complete long cycle, so both long-cycle blocks and their characteristic equations are unchanged. Within the boundary graph, edge deletion can split but cannot merge strongly connected components, so the same list remains exhaustive.

For the boundary triad, the Routh–Hurwitz difference expands as

$$\begin{aligned} c_1 c_2 - c_3 = & a(4a^2 h_1^2 h_m + 16a^2 h_1 h_m^2 + 11ab h_1^2 h_m + 4ab h_1^2 h_Z \\ & + 32ab h_1 h_m^2 + 32ab h_1 h_Z h_m + 64ab h_Z h_m^2 + 7b^2 h_1^2 h_m \\ & + 4b^2 h_1^2 h_Z + 7b^2 h_1 h_m^2 + 48b^2 h_1 h_Z h_m + 16b^2 h_1 h_Z^2 \\ & + 16b^2 h_Z h_m^2 + 64b^2 h_Z^2 h_m). \end{aligned}$$

Every monomial is strictly positive. The two long-cycle modulus inequalities are strict because $a + b > a$ and $4a + b > 4a$. Frobenius permutation therefore proves Hurwitz stability of every smaller principal block.

For completeness, the core determinant has a short exact Schur-complement calculation. First factor the column scalings $\prod_{i=1}^m h_i$. For $m \geq 4$, the lower-bidiagonal block on $X_3, \dots, X_{m-1}$ has determinant $(-a)^{m-3}$, and the bottom-left corner of its inverse is $-1/a$. Eliminating that block leaves, in the order (X_1, X_2, X_m) ,

$$B_m^{\text{core}} = \begin{pmatrix} -(a+b) & -a & -b \\ -a & -a & 2a \\ 2a-b & 2a & -(4a+b) \end{pmatrix}, \quad \det B_m^{\text{core}} = 2a^2 b.$$

When $m = 3$ the interior block is empty and B_m^{core} is the original X -species block. Restoring the column factors therefore gives, in every dimension,

$$(-1)^m \det J_{\widehat{Z}, \widehat{Z}} = -2a^{m-1}b \prod_{i=1}^m h_i.$$

55 The complete omission table is derived in Section S4 below.

56 S3 The principal-minor diffusion-ray theorem in full detail

Let $n \geq 2$, $a_I = (-1)^{|I|} \det J_{I,I}$, and $a_\emptyset = 1$. Assume $\det J = 0$, $a_I > 0$ for $|I| \leq n-2$, and $\sum_{|I|=n-1} a_I > 0$. The last condition follows from a simple zero and stable complementary spectrum but is stated directly because it is all the proof uses. Multilinearity in the diagonal entries of sD gives

$$\det(sD - J) = \sum_{I \subseteq [n]} a_I \prod_{j \notin I} s d_j.$$

Regrouping by $k = n - |I|$ yields

$$\det(sD - J) = \sum_{k=1}^n \beta_k(D) s^k, \quad \beta_k(D) = \sum_{|I|=n-k} a_I \prod_{j \notin I} d_j.$$

57 For $k \geq 2$, every contributing set has order at most $n-2$, so the stated coefficient hypothesis gives
58 $a_I > 0$ and hence $\beta_k > 0$.

59 The bracket $q_D(s) = \beta_1 + \beta_2 s + \dots + \beta_n s^{n-1}$ is strictly increasing for $s > 0$. If $\beta_1 < 0$, it has
60 exactly one positive zero and its derivative there is positive. This proves scalar simplicity of the
61 dispersion root. Ordinary algebraic simplicity of zero as an eigenvalue of $J - s_* D$ follows from the
62 characteristic-polynomial derivative below.

For the positive-real-eigenvalue band, differentiate

$$\chi_s(\lambda) = \sum_I a_I \prod_{j \notin I} (\lambda + s d_j).$$

63 Every order-at-most- $(n-2)$ contribution is nonnegative for $\lambda \geq 0$. The order- $(n-1)$ contribution to
64 χ'_s is the constant $\sum_{|I|=n-1} a_I$. This is positive by hypothesis. Under the common spectral corollary
65 it equals the coefficient of s in $\det(sI - J)$, and equal damping sends the simple zero and all stable
66 eigenvalues left. Thus $\chi'_s > 0$ on the nonnegative real axis. The sign of $\chi_s(0)$ then determines
67 whether there is exactly one positive real eigenvalue. At $s = s_*$, $\chi_{s_*}(0) = 0$ and $\chi'_{s_*}(0) > 0$, so zero
68 is an algebraically simple eigenvalue of $J - s_* D$.

69 The theorem says nothing about nonreal unstable eigenvalues when $s > s_*$. The all-mode
70 certificates below are separate, profile-specific results.

71 S4 Order- $(n-1)$ omission minors and diffusion design

72 Right multiplication by H multiplies a principal determinant by the product of its retained h_i , so
73 the combinatorial coefficients may be computed at $H = I$.

Omitting Z leaves the complete X -chain block. The Schur-complement calculation above gives

$$(-1)^m \det J_{\widehat{Z}, \widehat{Z}} = -2a^{m-1}b \prod_{i=1}^m h_i.$$

For $2 \leq j \leq m-1$, omit X_j . A Frobenius permutation orders the right chain fragment $X_{j+1}, \dots, X_{m-1}$, then $\{X_1, X_m, Z\}$, then the left fragment $X_2, \dots, X_{j-1}$; empty fragments are omitted. The resulting matrix is block lower triangular. The $m-3$ chain singletons contribute signed factors ah_i , and the boundary triad contributes

$$(-1)^3 \det J_{\{X_1, X_m, Z\}, \{X_1, X_m, Z\}} = 16a^2b h_1 h_m h_Z.$$

Therefore

$$(-1)^m \det J_{\widehat{X_j}, \widehat{X_j}} = 16a^{m-1}bh_Z \prod_{\substack{1 \leq i \leq m \\ i \neq j}} h_i.$$

74 Omitting X_1 leaves the restricted semipositive conservation vector in the left kernel. Omitting X_m
 75 leaves the restriction of $H^{-1}(2, -2, \dots, -2, 0, 1)^T$ in the right kernel, because the omitted coordinate
 76 is zero. Hence both determinants vanish.

Multiplication by the omitted diffusivities produces

$$\beta_1(D) = 2a^{m-1}b \left(\prod_{i=1}^m h_i \right) \left(8h_Z \sum_{j=2}^{m-1} \frac{d_j}{h_j} - d_Z \right).$$

77 This yields the exact diffusion criterion and explains the coefficient eight as the exact ratio $16/2$.
 Define the stable realization domain

$$\mathfrak{S}_m = \{(a, b, H) : a, b > 0, H \succ 0, A_m(a, b)H|_{c^\perp} \text{ is Hurwitz}\}.$$

The complete region $\mathfrak{S}_m$ is not characterized here. For any $(a, b, H) \in \mathfrak{S}_m$, if $d_{\min} = \min_i d_i$, then

$$d_Z > 8h_Z \sum_{j=2}^{m-1} \frac{d_j}{h_j} \geq d_{\min} T(H),$$

78 so $\chi_D > T(H)$. Setting all X diffusivities to one and $d_Z = T(H) + \varepsilon$ proves the nonattained fixed- H
 79 infimum. Since $h_Z/h_j \geq 1/\chi_H$, one obtains $\chi_D \chi_H > 8(m-2)$. The implication $T(H) > 1$ is
 80 necessary under homogeneous stability; its converse is not proved.

81 S5 Improved unit-equilibrium critical profile

Put $K_i = 91m - 181 - i$. The right vector $r = (r_1, \dots, r_m, r_Z)^T$ and diffusion profile are

$$r_1 = 1, \quad r_i = -\frac{K_i}{63(m-2)}, \quad r_m = -\frac{2}{9}, \quad r_Z = \frac{5}{14},$$

$$d_1 = \frac{23}{63}, \quad d_i = \frac{1}{K_i}, \quad d_m = \frac{1}{7}, \quad d_Z = \frac{16}{45}.$$

Direct substitution gives $(A_m - D_m)r = 0$. With $\ell_Z = 1$, the left vector $\ell = (\ell_1, \dots, \ell_m, \ell_Z)^T$ is

$$\ell_1 = -\frac{266}{815}, \quad \ell_i = \frac{78260(m-2)}{163(91m-180-i)}, \quad \ell_m = \frac{18368}{7335}, \quad \ell_Z = 1.$$

Define

$$\mathfrak{h}_m = \sum_{j=1}^{m-2} \frac{1}{91m-181-j}.$$

Then

$$\begin{aligned} \ell^T r &= -\frac{7043400m - 13600927 - 7043400\mathfrak{h}_m}{924210} < 0, \\ \ell^T D_m r &= -\frac{2(1760850\mathfrak{h}_m + 16559)}{462105} < 0. \end{aligned}$$

⁸² The harmonic bound $\mathfrak{h}_m \leq (m-2)/(90m-179)$ makes the first sign an elementary shifted-polynomial
⁸³ inequality.

⁸⁴ Homogeneous and spatial determinants

The homogeneous determinant has

$$\det(\lambda I - A_m) = \lambda q_m(\lambda), \quad \lambda q_m = (1 + \lambda)^{m-3} P(\lambda) - R(\lambda),$$

with $P = \lambda^4 + 12\lambda^3 + 42\lambda^2 + 47\lambda + 16$ and $R = 5\lambda^2 + 33\lambda + 16$. For $m = 3$,

$$q_3 = (\lambda + 7)(\lambda^2 + 5\lambda + 2)$$

is Hurwitz. For $m \geq 4$, the exact 35-term certificate for $|1 + \lambda|^2 |P|^2 - |R|^2$ is strictly positive away from the origin in the closed right half-plane. Since $|1 + \lambda| \geq 1$ there, a nonzero closed-right-half-plane root of q_m would contradict $(1 + \lambda)^{m-3} P = R$. Finally, $q_m(0) = 16m - 34 > 0$, so the conservation zero is simple and the complementary homogeneous spectrum is stable for every $m \geq 3$. For a mode with damping factor t ,

$$\det(\lambda I - A_m + tD_m) = Q_m F - G,$$

where

$$\begin{aligned} Q_m &= \prod_{i=2}^{m-1} \left(\lambda + 1 + \frac{t}{K_i} \right), \\ g_1 &= \lambda + 2 + \frac{23}{63}t, \quad g_m = \lambda + 5 + \frac{1}{7}t, \quad g_Z = \lambda + 4 + \frac{16}{45}t, \\ F &= g_1 g_m g_Z - 4g_1 - 4g_m + g_Z, \quad G = g_Z(4g_1 + g_m) - 36. \end{aligned}$$

At onset, $\prod_{i=2}^{m-1} (1 + K_i^{-1}) = 91/90$. The exact certificate tables appear in Section S8. The equality point is algebraically simple, not merely isolated from nonzero closed-right-half-plane roots. Indeed, with

$$\Pi_m(\lambda) = \det(\lambda I - A_m + D_m) = Q_m(\lambda, 1)F(\lambda, 1) - G(\lambda, 1),$$

one has the all-dimensional identity

$$\Pi'_m(0) = \frac{7043400m - 13600927 - 7043400\mathfrak{h}_m}{255150} = -\frac{163}{45} \ell^T r > 0.$$

⁸⁵ Thus zero is algebraically simple and the kernel and cokernel of $A_m - D_m$ are one-dimensional.

S6 Conservation-compatible stable-mode corrections

The quadratic tensor is obtained reaction by reaction. Its components are

$$\begin{aligned}(B(u, v))_1 &= -(u_1 v_{m-1} + u_{m-1} v_1) + 2u_Z v_Z - (u_1 v_m + u_m v_1), \\ (B(u, v))_2 &= -(u_1 v_2 + u_2 v_1) + 2u_m v_m, \\ (B(u, v))_i &= u_1 v_{i-1} + u_{i-1} v_1 - u_1 v_i - u_i v_1, \\ (B(u, v))_m &= 2(u_1 v_{m-1} + u_{m-1} v_1) - 4u_m v_m + 2u_Z v_Z - (u_1 v_m + u_m v_1), \\ (B(u, v))_Z &= -4u_Z v_Z + 2(u_1 v_m + u_m v_1).\end{aligned}$$

Here the third line holds for $3 \leq i \leq m-1$. Reaction-wise conservation implies $c^T B(u, v) = 0$.

The zero mode solves

$$A_m w_0 = -\frac{1}{4}B(r, r), \quad c^T w_0 = 0.$$

The exact solution is

$$\begin{aligned}(w_0)_1 &= \frac{182448m - 373417}{31752(8m - 17)}, \\ (w_0)_2 &= \frac{1008m^2 - 20459m + 37138}{31752(m - 2)(8m - 17)}, \\ (w_0)_i &= (w_0)_2 - \frac{i - 2}{126(m - 2)}, \quad (w_0)_m = -\frac{1}{81}, \quad (w_0)_Z = \frac{16861m - 34044}{7938(8m - 17)}.\end{aligned}$$

The second harmonic solves

$$(A_m - 4D_m)w_2 = -\frac{1}{4}B(r, r).$$

With

$$\sigma = \frac{1}{126(m - 2)}, \quad T_i = \frac{K_{i-3}K_{i-2}K_{i-1}K_i}{K_{-1}K_0K_1K_2},$$

the chain recurrence is

$$(w_2)_i = T_i \left((w_2)_2 + \frac{\sigma K_2}{3} \right) - \frac{\sigma K_i}{3}.$$

Its boundary denominator is

$$\mathcal{Q}_m = 589180301m^3 - 3500015940m^2 + 6930529579m - 4574434500.$$

After $m = u + 3$, the coefficients are

$$589180301, \quad 1802606769, \quad 1838302066, \quad 624878904,$$

so it never vanishes for $m \geq 3$. The complete boundary solve is exposed as follows. Write

$$W_1 = (w_2)_1, \quad W_2 = (w_2)_2, \quad W_m = (w_2)_m, \quad W_Z = (w_2)_Z, \quad b^{(2)} = -\frac{1}{4}B(r, r),$$

and set

$$\delta_m = \frac{\sigma}{3} (T_{m-1}K_2 - K_{m-1}).$$

Substitution of the recurrence into the four boundary rows gives the complete system

$$\begin{pmatrix} -218/63 & -T_{m-1} & -1 & 2 \\ -1 & -(1+4/K_2) & 2 & 0 \\ 1 & 2T_{m-1} & -39/7 & 2 \\ 2 & 0 & 2 & -244/45 \end{pmatrix} \begin{pmatrix} W_1 \\ W_2 \\ W_m \\ W_Z \end{pmatrix} = \begin{pmatrix} b_1^{(2)} + \delta_m \\ b_2^{(2)} \\ b_m^{(2)} - 2\delta_m \\ b_Z^{(2)} \end{pmatrix}.$$

Its determinant is exactly

$$\frac{64\mathcal{Q}_m}{6615(91m-183)(91m-181)(91m-180)}.$$

88 All factors in the denominator are positive for $m \geq 3$, while the displayed shifted-positive certificate
 89 proves $\mathcal{Q}_m > 0$. Thus the four boundary values are the unique solution of this displayed system. Their
 90 expanded rational forms are supplied in the machine-readable archive for convenient verification.

The cubic coefficient is

$$c_m = \frac{\ell^T \{B(r, w_0) + \frac{1}{2}B(r, w_2)\}}{\ell^T r}.$$

This is the cubic coefficient of the one-dimensional center-manifold normal form in the adjoint-projection coordinate. Setting the reduced vector field to zero gives the associated stationary Lyapunov–Schmidt equation. The spatial reflection $(\mathcal{R}u)(\xi) = u(\pi - \xi)$ commutes with the Neumann semiflow, preserves the fixed integrated-mass class, and sends $r \cos \xi$ to $-r \cos \xi$. On an equivariant center manifold the adjoint-projection action is $\mathcal{R} : A \mapsto -A$; hence the reduced vector field is odd in A , and reflection exchanges the two nonzero branches. For each fixed m , the branch perturbation tends to zero in $H_N^2 \hookrightarrow C^0$ from the base equilibrium $\mathbf{1}$; hence both reflection-related equilibria remain componentwise strictly positive for sufficiently small $\mu > 0$. For the cubic numerator, introduce

$$\begin{aligned} P_R(m) = & 68605040480814208768m^4 - 550882186169626030957m^3 \\ & + 1658612632937449670852m^2 - 2219226476204103501323m \\ & + 1113379274975809565700, \end{aligned}$$

and

$$\begin{aligned} P_C(m) = & 652054120726848m^4 - 5151971981328467m^3 \\ & + 15265080924982572m^2 - 20102347725659113m \\ & + 9927281930180400. \end{aligned}$$

Then the exact decomposition is

$$N_m = N_m^{\text{ref}} = R_m + C_m \mathfrak{h}_m,$$

where

$$R_m = \frac{P_R(m)}{286118780220(8m-17)\mathcal{Q}_m}, \quad C_m = -\frac{215P_C(m)}{11645046(8m-17)\mathcal{Q}_m}.$$

91 The shifted-positive coefficient certificates in Section S8.2 show that $\mathcal{Q}_m > 0$, $P_R(m) > 0$, and
 92 $P_C(m) > 0$ for $m \geq 3$. Consequently $R_m > 0$ and $C_m < 0$; in the latter conclusion the external
 93 factor -215 reverses the sign of the positive polynomial P_C .

The exact clearing identity is

$$R_m + C_m \frac{m-2}{90m-179} = \frac{L_m}{286118780220(8m-17)(90m-179)\mathcal{Q}_m},$$

where

$$\begin{aligned} L_m = & 2729945147827667886720m^5 - 27755132420474170999952m^4 \\ & + 112813395868533457497683m^3 - 229153280695458887386228m^2 \\ & + 232620996871721820873517m - 94412163900120968220300. \end{aligned} \quad (\text{S6.1})$$

After $m = u + 3$, its coefficients are

$$\begin{aligned} & 2729945147827667886720, 13194044796940847300848, 25446870127333515303059, \\ & 24475321329207325509911, 11736484608366875120374, 2243933770033305838488, \end{aligned}$$

all positive. Since $C_m < 0$ and $\mathfrak{h}_m \leq (m-2)/(90m-179)$, the complete comparison is

$$R_m + C_m \mathfrak{h}_m \geq R_m + C_m \frac{m-2}{90m-179} > 0.$$

94 Thus the cubic numerator is positive and, since $\ell^T r < 0$, $c_m < 0$.

95 S7 Equilibrium-scaled stable family

Let $\nu = m - 2$ and define

$$\kappa_\nu = \begin{cases} 1/\sqrt{3}, & \nu = 1, \\ \sqrt{5}/2, & \nu \geq 2, \end{cases} \quad L_0(\nu) = \frac{\kappa_\nu}{\sqrt{\nu}}, \quad L_1(\nu) = \frac{90\nu}{90\nu+1}.$$

This interval is nonempty. For $\nu = 1$, square both positive sides and use $8281 < 24300$. For $\nu \geq 2$,

$$(90\nu+1)^2 \leq (91\nu)^2 < 6480\nu^3,$$

which is equivalent to $L_0(\nu) < L_1(\nu)$. The lower endpoints are sufficient boundaries of the certificates below, not proved intrinsic dynamical boundaries. For $L \in [L_0, L_1]$, set

$$h_1 = h_m = h_Z = 1, \quad h_i = \frac{K_i}{LK_{i-1}}.$$

Write $\mathbf{H}_m(L) = \text{diag}(h_1, \dots, h_m, h_Z)$. The physical equilibrium is $x^* = \mathbf{H}_m(L)^{-1} \mathbf{1}$ and the physical diffusion matrix is $D^{\text{phys}} = \mathbf{H}_m(L) \Delta$, where, throughout this supplement, $\Delta := \Delta_m$ is the improved unit profile. In normalized variables $\hat{x} = \mathbf{H}_m(L)x$,

$$\partial_t \hat{x} = \mathbf{H}_m(L) \{f_m(\hat{x}) + (1-\mu) \Delta \partial_{\xi\xi} \hat{x}\}.$$

At $\mu = 0$, the stationary bracket $f_m(\hat{x}) + \Delta \partial_{\xi\xi} \hat{x}$ and its critical right vector are unchanged. The dynamic left vector is

$$\tilde{\ell}(L) = \mathbf{H}_m(L)^{-1} \ell,$$

and the physical conservation gauge uses $\mathbf{H}_m(L)^{-1}c$. The crossing numerator is unchanged by the row scaling:

$$\tilde{\ell}(L)^T \mathbf{H}_m(L) \Delta r = \ell^T \Delta r < 0.$$

Together with $\tilde{\ell}(L)^T r < 0$, this gives

$$\eta_m(L) = \frac{\tilde{\ell}(L)^T \mathbf{H}_m(L) \Delta r}{\tilde{\ell}(L)^T r} > 0.$$

Moreover,

$$\ker[\mathbf{H}_m(L)(A_m - \Delta)] = \ker(A_m - \Delta) = \text{span}\{r\},$$

96 because $\mathbf{H}_m(L)$ is invertible and the unit critical zero is simple. If $\mathbf{H}_m(L)(A_m - \Delta)v = r$ for a
 97 generalized eigenvector v , then left multiplication by $\tilde{\ell}(L)^T$ would imply $0 = \tilde{\ell}(L)^T r$, a contradiction.
 98 Hence the scaled critical zero is algebraically simple as well.

For interior chain factors,

$$g_i = \frac{K_{i-1}}{K_i}(1 + L\lambda) + \frac{t-1}{K_i}.$$

When $t \geq 1$ and $\Re\lambda \geq 0$,

$$|Q_m|^2 \geq \left(\frac{91}{90}\right)^2 |1 + L\lambda|^{2\nu} \geq \left(\frac{91}{90}\right)^2 \left(1 + 2\nu L\Re\lambda + \frac{1}{3}(\Im\lambda)^2\right).$$

99 The last inequality holds because $\nu L^2 \geq 1/3$ throughout the certified interval. The 84-term spatial
 100 coefficient table in Section S8 is nonnegative and is strict except at $(\lambda, t) = (0, 1)$.

Homogeneous stability uses a different boundary polynomial. At $t = 0$,

$$\frac{\det(\lambda I - \mathbf{H}_m(L)A_m)}{\det \mathbf{H}_m(L)} = Q(\lambda)F_0(\lambda) - R(\lambda),$$

where

$$Q = \prod_{i=2}^{m-1} \left(1 + L \frac{K_{i-1}}{K_i} \lambda\right), \quad F_0 = \lambda^3 + 11\lambda^2 + 31\lambda + 16, \quad R = 5\lambda^2 + 33\lambda + 16.$$

For $\Re\lambda = x \geq 0$, every factor in Q has modulus at least $|1 + L\lambda|$. If $\nu \geq 2$, then

$$|Q|^2 \geq |1 + L\lambda|^{2\nu} \geq 1 + 2\nu Lx + \nu L^2 (\Im\lambda)^2 \geq 1 + Ax + \frac{5}{4}(\Im\lambda)^2, \quad A = 2\nu L.$$

Here $A \geq \sqrt{5\nu} \geq \sqrt{10} > 1/4$. The exact polynomial

$$\left(1 + Ax + \frac{5}{4}(\Im\lambda)^2\right) |F_0|^2 - |R|^2$$

has 22 nonzero monomials. With $U = A - 1/4$, its coefficient polynomials have nonnegative rational coefficients and the expression is strict away from $\lambda = 0$. Thus there is no nonzero homogeneous eigenvalue in the closed right half-plane. The coefficient $5/4$ is sharp for this coefficientwise certificate. If it is replaced by B , then at $x = 0$ the coefficient of $z = y^2$ in $(1 + Bz)|F_0|^2 - |R|^2$ is $256B - 320$. It

vanishes at $B = 5/4$ and is negative for $B < 5/4$. This does not show that L_0 is an intrinsic dynamical boundary; it identifies only the boundary of the present exact modulus certificate. Moreover

$$(QF_0 - R)'(0) = 16L \sum_{i=2}^{m-1} \frac{K_{i-1}}{K_i} - 2 > 0,$$

where the inequality uses $K_{i-1}/K_i > 1$ and $L \geq \sqrt{5}/(2\sqrt{\nu})$. Thus the conservation zero is simple. If $\nu = 1$, put $\gamma = 91L/90$. Direct division by the conservation factor gives

$$\frac{QF_0 - R}{\lambda} = \gamma\lambda^3 + (1 + 11\gamma)\lambda^2 + (6 + 31\gamma)\lambda + (16\gamma - 2).$$

101 Here $\gamma > 1/8$, and the remaining cubic Routh–Hurwitz difference is $6 + 99\gamma + 325\gamma^2 > 0$. This proves
102 homogeneous relative stability in the exceptional dimension as well.

Let $\rho_m = (2, -2, \dots, -2, 0, 1)^T$. The physical zero-mode gauge is

$$w_0(L) = w_0^{\text{ref}} + \tau_m(L)\rho_m, \quad \tau_m(L) = -\frac{(\mathbf{H}_m(L)^{-1}c)^T w_0^{\text{ref}}}{(\mathbf{H}_m(L)^{-1}c)^T \rho_m}.$$

The second harmonic is unchanged. The cubic numerator satisfies

$$N_m(L) = N_m^{\text{ref}} + \tau_m(L)S_m,$$

where $N_m^{\text{ref}} = R_m + C_m \mathbf{h}_m$. To make its quantitative margin explicit, set

$$\begin{aligned} P_{\text{ref}}(\nu) = & 3790502986637265684840\nu^5 - 974216530468600286489\nu^4 \\ & - 53103567440921218871\nu^3 - 576386186827093561\nu^2 \\ & + 3649732858601219\nu + 55281268032918 \end{aligned}$$

and

$$\begin{aligned} D_{\text{ref}}(\nu) = & 715296950550(8\nu - 1)(90\nu + 1) \\ & \times (589180301\nu^3 + 35065866\nu^2 + 629431\nu + 3306). \end{aligned}$$

Since $C_m < 0$ and $\mathbf{h}_m \leq \nu/(90\nu + 1)$, exact clearing gives

$$N_m^{\text{ref}} - \frac{1}{100} \geq \frac{P_{\text{ref}}(\nu)}{D_{\text{ref}}(\nu)}.$$

The denominator is positive for $\nu \geq 1$. After $\nu = v + 1$, the coefficients of P_{ref} , in descending powers $v^5, \dots, v^0$, are

$$\begin{aligned} & 3790502986637265684840, 17978298402717728137711, 33955060177057334483573, \\ & 31899843595051464379292, 14895188986348368035728, 2762610207555043720056. \end{aligned}$$

103 They are positive, so $P_{\text{ref}}(\nu) > 0$ and $N_m^{\text{ref}} > 1/100$.

Separately,

$$S_m = -\frac{4(1760850\mathbf{h}_m - 10253)}{462105}.$$

The bounds $1/91 \leq \mathfrak{h}_m < 1/90$ give $S_m < 0$ because

$$\frac{1760850}{91} - 10253 > 0,$$

and give $S_m > -1/10$ because

$$\frac{4}{462105} \left(\frac{1760850}{90} - 10253 \right) < \frac{1}{10}.$$

For the remaining gauge estimate, define

$$\begin{aligned} A_\tau &= 1494249120\mathfrak{h}_m L\nu^2 - 69786990\mathfrak{h}_m L\nu + 108738630L\nu^2 \\ &\quad + 1214388L\nu - 8521L - 125249670\nu^2 + 1031940\nu, \\ B_\tau &= 32760\mathfrak{h}_m L\nu + 32760L\nu^2 + 4L - 4095\nu. \end{aligned}$$

Then the closed form is

$$\tau_m(L) = -\frac{A_\tau}{15876(8\nu - 1)B_\tau}.$$

On the broader certified cubic range $L \geq 1/\sqrt{3\nu}$,

$$B_\tau = 4095\nu[8L(\nu + \mathfrak{h}_m) - 1] + 4L > 0,$$

because $L\nu \geq 1/\sqrt{3}$. Exact differentiation then gives

$$\partial_{\mathfrak{h}} \tau_m = -\frac{4225L\nu^2(182448L\nu + 1008L - 7513)}{2B_\tau^2} < 0,$$

$$\partial_L \tau_m = -\frac{65\nu(-61531470\mathfrak{h}_m\nu + 125249670\nu^2 + 1031940\nu - 7513)}{252B_\tau^2} < 0.$$

¹⁰⁴ For the first sign, $L\nu \geq 1/\sqrt{3} > 1/2$ makes the parenthesis positive. For the second, use $\mathfrak{h}_m < 1/90$
¹⁰⁵ and $\nu \geq 1$. Thus the maximum occurs at $\mathfrak{h}_m = 1/91$ and $L = 1/\sqrt{3\nu}$.

Put $y = \sqrt{3\nu}$ and define the unreduced endpoint polynomial

$$\begin{aligned} P_\tau(\nu, y) &= -1040195520\nu^3 + 756272790\nu^2 y - 507201030\nu^2 \\ &\quad - 21412755\nu y - 935658\nu + 58481 \end{aligned}$$

and $D_\tau(\nu, y) = -32760\nu^2 + 4095\nu y - 360\nu - 4$. Direct substitution gives

$$\tau_m(1/91, 1/y) - \frac{1}{20} = -\frac{P_\tau(\nu, y)}{79380(8\nu - 1)D_\tau(\nu, y)}.$$

Since $y < 2\nu$, $D_\tau < 0$. For $\nu = 1$, the coefficient of y in $P_\tau(1, y)$ is positive and $\sqrt{3} < 7/4$, whence

$$P_\tau(1, \sqrt{3}) < P_\tau(1, 7/4) = -\frac{1049074663}{4} < 0.$$

For $\nu \geq 2$, $y \leq 5\nu/4$ and deletion of the negative term $-21412755\nu y$ gives $P_\tau(\nu, y) < P_{\text{up}}(\nu)$, where

$$P_{\text{up}}(\nu) = -\frac{189709065}{2}\nu^3 - 507201030\nu^2 - 935658\nu + 58481.$$

This upper-bound polynomial is human-checkable:

$$P'_{\text{up}}(\nu) = -\frac{569127195}{2}\nu^2 - 1014402060\nu - 935658 < 0 \quad (\nu \geq 0),$$

and

$$P_{\text{up}}(2) = -2789453215 < 0.$$

106 Hence $P_{\text{up}}(\nu) < 0$ for every $\nu \geq 2$, and therefore $P_{\tau}(\nu, y) < 0$ in all dimensions. Because the displayed
107 denominator $79380(8\nu - 1)D_{\tau}$ is negative, the exact endpoint comparison proves $\tau_m(L) < 1/20$.

If $\tau_m(L) \leq 0$, then $\tau_m(L)S_m \geq 0$. If instead $0 < \tau_m(L) < 1/20$, then $\tau_m(L)S_m > -1/200$. Together with $N_m^{\text{ref}} > 1/100$, this yields $N_m(L) > 1/200$. Also

$$\ell^T \mathbf{H}_m(L)^{-1} r = -\frac{485873}{924210} - \frac{11180}{1467} L(m-2) < 0.$$

Thus the scaled cubic coefficient is explicitly

$$c_m(L) = \frac{N_m(L)}{\widetilde{\ell}(L)^T r} = \frac{N_m(L)}{\ell^T \mathbf{H}_m(L)^{-1} r} < 0.$$

108 Therefore the family is uniformly supercritical over the certified interval. Indeed, the cubic comparison
109 was proved on the broader range $L \geq 1/\sqrt{3\nu}$; the present lower endpoint agrees with this value
110 for $\nu = 1$ and is larger for $\nu \geq 2$. The diagonal concentration scaling commutes with the same
111 spatial reflection $\xi \mapsto \pi - \xi$. Hence the scaled center-manifold vector field remains odd in its
112 adjoint-projection amplitude, and reflection exchanges the two nonzero patterned branches. For
113 each fixed m and L , the normalized branch converges to $\mathbf{1}$ in $H_N^2 \hookrightarrow C^0$ and is componentwise
114 strictly positive for sufficiently small $\mu > 0$. Multiplication by the positive diagonal matrix $\mathbf{H}_m(L)^{-1}$
115 preserves componentwise positivity in physical concentrations.

The contrasts are

$$\chi_D(L) = \frac{23}{63}(91\nu L), \quad \chi_H(L) = \frac{91\nu - 1}{91\nu L},$$

where χ_H measures equilibrium concentration-scale separation, not a universal measure of biological expense or implausibility. Their product is fixed:

$$\chi_D(L)\chi_H(L) = \frac{23(91\nu - 1)}{63} = \frac{23(91m - 183)}{63} = \chi_D^{\text{unit}}(m).$$

Thus the family reallocates this fixed contrast product between physical diffusion and equilibrium concentration scales; it does not reduce the product. The first contrast is strictly increasing and the second strictly decreasing in L , while

$$\frac{\chi_D(L_0)}{\chi_H(L_0)} = \begin{cases} 27209/2430, & \nu = 1, \\ \frac{136045\nu}{36(91\nu - 1)}, & \nu \geq 2, \end{cases} > 1.$$

Consequently $\chi_D(L) > \chi_H(L)$ throughout the certified interval, and $\max(\chi_D, \chi_H) = \chi_D(L)$ is uniquely minimized within this family at $L = L_0$. At that endpoint,

$$\chi_D = \frac{2093}{63} \kappa_{\nu} \sqrt{\nu}, \quad \chi_H = \frac{\sqrt{\nu}}{\kappa_{\nu}} \frac{91\nu - 1}{91\nu},$$

116 so both contrasts, and hence their maximum, have square-root growth, compared with linear growth
117 for the unit-equilibrium stable profile.

S8 Exact coefficient tables

S8.1 Coefficientwise-nonnegative modulus certificates

The first four tables below are coefficientwise-nonnegative modulus certificates. Each row records an exponent vector and the exact coefficient of the corresponding polynomial. The later shifted-polynomial tables are signed scalar certificates; for each such table we state the polynomial, any overall sign prefactor, the parameter domain, and the sign conclusion separately. Rational coefficients in the four modulus tables are printed in single-line a/b notation so adjacent exact entries remain visually separated.

By conjugation, take $y \geq 0$ and use the variables

$$\lambda = x + iy = x + i\sqrt{z}, \quad z = y^2, \quad t = 1 + s, \quad x, y, z, s \geq 0.$$

For the unit-profile spatial calculation, define

$$g_1 = \lambda + 2 + \frac{23}{63}t, \quad g_m = \lambda + 5 + \frac{1}{7}t, \quad g_z = \lambda + 4 + \frac{16}{45}t,$$

$$F = g_1 g_m g_z - 4g_1 - 4g_m + g_z, \quad G = g_z(4g_1 + g_m) - 36.$$

The 77-term table is exactly the expansion in (x, z, s) of

$$E_{77}(x, z, s) = \left[\left(\frac{91}{90} \right)^2 + z \right] |F(x + i\sqrt{z}, 1 + s)|^2 - |G(x + i\sqrt{z}, 1 + s)|^2.$$

Every one of its 77 displayed coefficients is strictly positive; there is no constant term. Hence equality occurs only at $x = z = s = 0$, equivalently $(\lambda, t) = (0, 1)$.

For the equilibrium-scaled spatial calculation, put $A = 2\nu L > 0$. The 84-term table is exactly the expansion in (x, z, s) of

$$E_{84}(x, z, s; A) = \left(\frac{91}{90} \right)^2 \left(1 + Ax + \frac{z}{3} \right) |F(x + i\sqrt{z}, 1 + s)|^2 - |G(x + i\sqrt{z}, 1 + s)|^2.$$

Its 84 coefficients are polynomials in A with nonnegative rational coefficients and are positive for $A > 0$. Again there is no constant term, and equality occurs only at $x = z = s = 0$, equivalently $(\lambda, t) = (0, 1)$.

For the equilibrium-scaled homogeneous calculation, define

$$F_0(\lambda) = \lambda^3 + 11\lambda^2 + 31\lambda + 16, \quad R(\lambda) = 5\lambda^2 + 33\lambda + 16, \quad U = A - \frac{1}{4}.$$

The 22-term table is exactly the expansion in (x, z) of

$$E_{22}(x, z; U) = \left[1 + \left(U + \frac{1}{4} \right) x + \frac{5}{4} z \right] |F_0(x + i\sqrt{z})|^2 - |R(x + i\sqrt{z})|^2.$$

For $U \geq 0$, all 22 coefficient polynomials are nonnegative, and the expression vanishes only at $x = z = 0$, equivalently $\lambda = 0$.

Finally, with

$$P(\lambda) = \lambda^4 + 12\lambda^3 + 42\lambda^2 + 47\lambda + 16,$$

the unit-profile homogeneous table is the 35-term expansion in (x, z) of

$$E_{35}(x, z) = |1 + \lambda|^2 |P(\lambda)|^2 - |R(\lambda)|^2, \quad \lambda = x + i\sqrt{z}.$$

133 All 35 coefficients are strictly positive, there is no constant term, and equality occurs only at $x = z = 0$,
 134 equivalently $\lambda = 0$.

135 From the repository root, the command `python independent_verifier/verify_symbolic_certificates.py`
 136 regenerates and checks the tables from exact symbolic expressions.

137 **35-term homogeneous certificate**

$\deg_x$	$\deg_z$	coefficient
10	0	1
9	0	26
8	1	5
8	0	277
7	1	104
7	0	1582
6	2	10
6	1	892
6	0	5356
5	2	156
5	1	4034
5	0	11282
4	3	10
4	2	1014
4	1	10432
4	0	15116
3	3	104
3	2	3322
3	1	15628
3	0	12612
2	4	5
2	3	460
2	2	5804
2	1	13296
2	0	5568
1	4	26
1	3	870
1	2	4858
1	1	5700
1	0	960
0	5	1
0	4	61
0	3	728
0	2	1508
0	1	192

77-term improved-profile spatial certificate

$\deg_x$	$\deg_z$	$\deg_s$	coefficient
6	1	0	1
6	0	0	8281/8100
5	1	1	544/315
5	1	0	7474/315
5	0	1	160888/91125
5	0	0	4420871/182250
4	2	0	3
4	1	2	40058/33075
4	1	1	1147126/33075
4	1	0	9681647/44100
4	0	2	3384901/2733750
4	0	1	96932147/2733750
4	0	0	1073317517/5467500
3	2	1	1088/315
3	2	0	14948/315
3	1	3	2744576/6251175
3	1	2	41276762/2083725
3	1	1	8186375738/31255875
3	1	0	10434543841/10418625
3	0	3	115958336/258339375
3	0	2	3487886389/172226250
3	0	1	39671949511/172226250
3	0	0	315432259813/516678750
2	3	0	3
2	2	2	147968/99225
2	2	1	4065856/99225
2	2	0	112938659/396900
2	1	4	33947257/393824025
2	1	3	2164641406/393824025
2	1	2	1130666508023/9845600625
2	1	1	9041228678666/9845600625
2	1	0	5074377793279/2187911250
2	0	4	5737086433/65101522500
2	0	3	182912198807/32550761250
2	0	2	2142193804481/21700507500
2	0	1	17194597630733/32550761250
2	0	0	34399269232129/65101522500
1	3	1	544/315
1	3	0	7474/315
1	2	3	1817216/6251175
1	2	2	24359162/2083725
1	2	1	5056135954/31255875
1	2	0	47437544293/62511750
1	1	5	3399584/393824025

$\deg_x$	$\deg_z$	$\deg_s$	coefficient
1	1	4	57943048/78764805
1	1	3	1934522477168/88610405625
1	1	2	16308922895827/59073603750
1	1	1	85574440190593/59073603750
1	1	0	62146993882837/25317258750
1	0	5	143632424/16275380625
1	0	4	2448093778/3255076125
1	0	3	98527787884/5425126875
1	0	2	475980699652/3255076125
1	0	1	4488365319374/16275380625
1	0	0	1086412028/10333575
0	4	0	1
0	3	2	27794/99225
0	3	1	624478/99225
0	3	0	1747307/26460
0	2	4	1744613/78764805
0	2	3	437610814/393824025
0	2	2	452220491743/19691201250
0	2	1	4035916860091/19691201250
0	2	0	10861695961357/13127467500
0	1	6	135424/393824025
0	1	5	2964608/78764805
0	1	4	962538972893/637994920500
0	1	3	43240103983367/1594987301250
0	1	2	238300444324673/1063324867500
0	1	1	170939874507539/227855328750
0	1	0	32638690751329/65101522500
0	0	6	5721664/16275380625
0	0	5	125254688/3255076125
0	0	4	6479423564/5425126875
0	0	3	41256502232/3255076125
0	0	2	513831327844/16275380625
0	0	1	148103696/10333575

139 **22-term equilibrium-scaled homogeneous certificate** ($U = A - 1/4$)

$\deg_x$	$\deg_z$	coefficient
7	0	$1/4 + U$
6	1	$5/4$
6	0	$13/2 + 22U$
5	1	$113/4 + 3U$
5	0	$271/4 + 183U$
4	2	$15/4$
4	1	$971/4 + 44U$
4	0	$673/2 + 714U$

$\deg_x$	$\deg_z$	coefficient
3	2	$223/4 + 3U$
3	1	$997 + 242U$
3	0	$2849/4 + 1313U$
2	3	$15/4$
2	2	$311 + 22U$
2	1	$7919/4 + 586U$
2	0	$312 + 992U$
1	3	$111/4 + U$
1	2	$3077/4 + 59U$
1	1	$6593/4 + 609U$
1	0	$256U$
0	4	$5/4$
0	3	$299/4$
0	2	$3181/4$

140 **84-term equilibrium-scaled spatial certificate**

$\deg_x$	$\deg_z$	$\deg_s$	coefficient
7	0	0	$8281/8100A$
6	1	0	$8281/24300$
6	0	1	$160888/91125A$
6	0	0	$8281/8100 + 4420871/182250A$
5	1	1	$160888/273375$
5	1	0	$4420871/546750 + 8281/2700A$
5	0	2	$3384901/2733750A$
5	0	1	$160888/91125 + 96932147/2733750A$
5	0	0	$4420871/182250 + 1210005017/5467500A$
4	2	0	$8281/8100$
4	1	2	$3384901/8201250$
4	1	1	$96932147/8201250 + 321776/91125A$
4	1	0	$315078023/4100625 + 4420871/91125A$
4	0	3	$115958336/258339375A$
4	0	2	$3384901/2733750 + 3487886389/172226250A$
4	0	1	$96932147/2733750 + 45494837011/172226250A$
4	0	0	$1073317517/5467500 + 503404909813/516678750A$
3	2	1	$321776/273375$
3	2	0	$4420871/273375 + 8281/2700A$
3	1	3	$115958336/775018125$
3	1	2	$3487886389/516678750 + 6251648/4100625A$
3	1	1	$47319306931/516678750 + 171782416/4100625A$
3	1	0	$578603925523/1550036250 + 2360113561/8201250A$
3	0	4	$5737086433/65101522500A$
3	0	3	$115958336/258339375 + 182912198807/32550761250A$
3	0	2	$3487886389/172226250 + 2513945531981/21700507500A$
3	0	1	$39671949511/172226250 + 29165273350733/32550761250A$

$\deg_x$	$\deg_z$	$\deg_s$	coefficient
3	0	0	$315432259813/516678750 + 138537167092129/65101522500A$
2	3	0	$8281/8100$
2	2	2	$6251648/12301875$
2	2	1	$171782416/12301875 + 160888/91125A$
2	2	0	$4871148347/49207500 + 4420871/182250A$
2	1	4	$5737086433/195304567500$
2	1	3	$182912198807/97652283750 + 76777376/258339375A$
2	1	2	$2613196695629/65101522500 + 2058349189/172226250A$
2	1	1	$33256099805357/97652283750 + 28172026051/172226250A$
2	1	0	$184975683058783/195304567500 + 388034017813/516678750A$
2	0	5	$143632424/16275380625A$
2	0	4	$5737086433/65101522500 + 2448093778/3255076125A$
2	0	3	$182912198807/32550761250 + 119438423884/5425126875A$
2	0	2	$2142193804481/21700507500 + 878963880652/3255076125A$
2	0	1	$17194597630733/32550761250 + 21944279796374/16275380625A$
2	0	0	$34399269232129/65101522500 + 21842305628/10333575A$
1	3	1	$160888/273375$
1	3	0	$4420871/546750 + 8281/8100A$
1	2	3	$76777376/775018125$
1	2	2	$2058349189/516678750 + 2348593/8201250A$
1	2	1	$29084261011/516678750 + 52768391/8201250A$
1	2	0	$212816762834/775018125 + 1090212071/16402500A$
1	1	5	$143632424/48826141875$
1	1	4	$2448093778/9765228375 + 294839597/13020304500A$
1	1	3	$124275398572/16275380625 + 36978113783/32550761250A$
1	1	2	$9956722796683/97652283750 + 503148512477/21700507500A$
1	1	1	$11312104230233/19530456750 + 6606581721077/32550761250A$
1	1	0	$241757054977/221433750 + 52308962411329/65101522500A$
1	0	6	$5721664/16275380625A$
1	0	5	$143632424/16275380625 + 125254688/3255076125A$
1	0	4	$2448093778/3255076125 + 8242156364/5425126875A$
1	0	3	$98527787884/5425126875 + 86436565592/3255076125A$
1	0	2	$475980699652/3255076125 + 3438519744244/16275380625A$
1	0	1	$4488365319374/16275380625 + 7094103536/10333575A$
1	0	0	$1086412028/10333575 + 4999696/6561A$
0	4	0	$8281/24300$
0	3	2	$2348593/24603750$
0	3	1	$52768391/24603750$
0	3	0	$570259573/24603750$
0	2	4	$294839597/39060913500$
0	2	3	$36978113783/97652283750$
0	2	2	$521791643711/65101522500$
0	2	1	$3617447476357/48826141875$
0	2	0	$30203751676613/97652283750$
0	1	6	$5721664/48826141875$
0	1	5	$125254688/9765228375$
0	1	4	$11480941147/21700507500$

$\deg_x$	$\deg_z$	$\deg_s$	coefficient
0	1	3	975299997269/97652283750
0	1	2	17163191001769/195304567500
0	1	1	10010796373877/32550761250
0	1	0	650029271329/65101522500
0	0	6	5721664/16275380625
0	0	5	125254688/3255076125
0	0	4	6479423564/5425126875
0	0	3	41256502232/3255076125
0	0	2	513831327844/16275380625
0	0	1	148103696/10333575

141 S8.2 Signed scalar and rational-function certificates

142 Each generated row below states the exact scalar polynomial or rational expression, the descending
143 coefficient order after its indicated shift, the sign of the polynomial on its parameter domain, and
144 whether an external negative prefactor reverses that sign. The shifts use $u = m - 3$ and $v = \nu - 1$,
145 leaving $z = y^2$ reserved for the modulus tables. In particular, the P_C row certifies $C_m < 0$ through
146 its external factor -215 ; the sign of S_m is certified separately.

Unit-profile boundary denominator. The polynomial being certified is

$$Q_m = 589180301m^3 - 3500015940m^2 + 6930529579m - 4574434500.$$

After $m = u + 3$, the following coefficients are listed in descending powers of u :

$$589180301, \quad 1802606769, \quad 1838302066, \\ 624878904$$

147 Every coefficient is positive, so $Q_m > 0$ for $m \geq 3$. There is no external sign prefactor.

Unit-profile critical-denominator bound. The auxiliary polynomial is

$$633906000m^2 - 2491895430m + 2448652733.$$

After $m = u + 3$, the following coefficients are listed in descending powers of u :

$$633906000, \quad 1311540570, \quad 678120443$$

148 Every coefficient is positive, so the polynomial is positive for $m \geq 3$. There is no external sign
149 prefactor.

Unit-profile cubic numerator lower bound. The polynomial being certified is

$$L_m = 2729945147827667886720m^5 - 27755132420474170999952m^4 \\ + 112813395868533457497683m^3 - 229153280695458887386228m^2 \\ + 232620996871721820873517m - 94412163900120968220300.$$

This is the exact clearing polynomial defined in (S6.1). Its rational identity is

$$R_m + C_m \frac{m-2}{90m-179} = \frac{L_m}{286118780220(8m-17)(90m-179)\mathcal{Q}_m}.$$

After $m = u + 3$, the following coefficients are listed in descending powers of u :

$$\begin{aligned} &2729945147827667886720, \quad 13194044796940847300848, \quad 25446870127333515303059, \\ &24475321329207325509911, \quad 11736484608366875120374, \quad 2243933770033305838488 \end{aligned}$$

150 Every coefficient is positive, so $L_m > 0$ for $m \geq 3$. It enters the clearing identity with a positive
151 denominator and no external negative prefactor.

Reference coefficient R_m . Define

$$\begin{aligned} P_R(m) = &68605040480814208768m^4 - 550882186169626030957m^3 \\ &+ 1658612632937449670852m^2 - 2219226476204103501323m \\ &+ 1113379274975809565700, \end{aligned} \quad R_m = \frac{P_R(m)}{286118780220(8m-17)\mathcal{Q}_m}.$$

After $m = u + 3$, the following coefficients of P_R are listed in descending powers of u :

$$\begin{aligned} &68605040480814208768, \quad 272378299600144474259, \quad 405345143374782665711, \\ &267974666768626234894, \quad 66402795166594173768 \end{aligned}$$

152 Every coefficient is positive. The denominator is positive for $m \geq 3$, and there is no external negative
153 prefactor; hence $R_m > 0$.

Unit-profile coefficient C_m . Define

$$\begin{aligned} P_C(m) = &652054120726848m^4 - 5151971981328467m^3 \\ &+ 15265080924982572m^2 - 20102347725659113m + 9927281930180400, \\ C_m = &-\frac{215P_C(m)}{11645046(8m-17)\mathcal{Q}_m}. \end{aligned}$$

After $m = u + 3$, the following coefficients of P_C are listed in descending powers of u :

$$\begin{aligned} &652054120726848, \quad 2672677467393709, \quad 4108255612276161, \\ &2806739366867294, \quad 719107361052288 \end{aligned}$$

154 Every coefficient is positive. Since the denominator is positive and the external prefactor is -215 ,
155 this proves $C_m < 0$.

Reference cubic margin. Let $\nu = m - 2$ and define

$$\begin{aligned} P_{\text{ref}}(\nu) = &3790502986637265684840\nu^5 - 974216530468600286489\nu^4 \\ &- 53103567440921218871\nu^3 - 576386186827093561\nu^2 \\ &+ 3649732858601219\nu + 55281268032918, \\ D_{\text{ref}}(\nu) = &715296950550(8\nu-1)(90\nu+1)(589180301\nu^3 + 35065866\nu^2 + 629431\nu + 3306). \end{aligned}$$

The exact comparison is

$$N_m^{\text{ref}} - \frac{1}{100} \geq \frac{P_{\text{ref}}(\nu)}{D_{\text{ref}}(\nu)}.$$

After $\nu = v + 1$, the following coefficients of P_{ref} are listed in descending powers of v :

$$\begin{aligned} &3790502986637265684840, \quad 17978298402717728137711, \quad 33955060177057334483573, \\ &31899843595051464379292, \quad 14895188986348368035728, \quad 2762610207555043720056 \end{aligned}$$

156 Every coefficient is positive, and $D_{\text{ref}}(\nu) > 0$ for $\nu \geq 1$. There is no external sign prefactor. Hence
157 $N_m^{\text{ref}} > 1/100$.

Gauge upper-bound tail. Let $\nu = m - 2$. The exact rational expression being bounded is

$$\tau_m(L) = -\frac{A_\tau}{15876(8\nu - 1)B_\tau},$$

where

$$\begin{aligned} A_\tau &= 1494249120\mathfrak{h}_m L\nu^2 - 69786990\mathfrak{h}_m L\nu + 108738630L\nu^2 \\ &\quad + 1214388L\nu - 8521L - 125249670\nu^2 + 1031940\nu, \\ B_\tau &= 32760\mathfrak{h}_m L\nu + 32760L\nu^2 + 4L - 4095\nu. \end{aligned}$$

The signed upper-bound polynomial is

$$P_{\text{up}}(\nu) = -\frac{189709065}{2}\nu^3 - 507201030\nu^2 - 935658\nu + 58481.$$

The following coefficients are listed in descending powers of ν :

$$\begin{aligned} &-189709065/2, \quad -507201030, \quad -935658, \\ &58481 \end{aligned}$$

This is a signed coefficient list, not a coefficientwise positivity certificate. Direct differentiation gives

$$P'_{\text{up}}(\nu) = -\frac{569127195}{2}\nu^2 - 1014402060\nu - 935658 < 0 \quad (\nu \geq 0),$$

158 and $P_{\text{up}}(2) = -2789453215 < 0$. Thus $P_{\text{up}}(\nu) < 0$ for $\nu \geq 2$. There is no external negative prefactor
159 reversing this conclusion.

The signed scalar S_m . This sign is certified separately from P_C . With

$$S_m = -\frac{4(1760850\mathfrak{h}_m - 10253)}{462105}, \quad \frac{1}{91} \leq \mathfrak{h}_m < \frac{1}{90},$$

the exact comparisons

$$\frac{1760850}{91} - 10253 > 0, \quad \frac{4}{462105} \left(\frac{1760850}{90} - 10253 \right) < \frac{1}{10},$$

160 give $-1/10 < S_m < 0$. The displayed expression itself carries the external negative prefactor -4 ,
161 which reverses the positive bracket and proves the upper sign. Combining $N_m^{\text{ref}} > 1/100$, $S_m > -1/10$,
162 and $\tau_m(L) < 1/20$ gives $N_m(L) > 1/200$.

S9 Near-threshold control example

Return here to $a = b = 1$ and $H = I$. For scalars p, u, v, q , with u the Latin letter, define the affine-chain critical vector $r^{\text{aff}} = (r_1, \dots, r_m, r_Z)^T$ by

$$r_1 = 1, \quad r_i = -\left(u + v \frac{m-1-i}{m-2}\right) \quad (2 \leq i \leq m-1), \quad r_m = -p, \quad r_Z = q.$$

The diagonal matrix $D = \text{diag}(d_1, \dots, d_m, d_Z)$ determined by $(A_m - D)r^{\text{aff}} = 0$ has entries

$$\begin{aligned} d_1 &= -2 + u + p + 2q, \\ d_2 &= \frac{1 + 2p - u - v(m-3)/(m-2)}{u + v(m-3)/(m-2)}, \quad d_i = \frac{v}{(m-2)u + v(m-1-i)} \quad (3 \leq i \leq m-1), \\ d_m &= \frac{2u - 5p - 2q - 1}{p}, \quad d_Z = \frac{2 - 2p - 4q}{q}. \end{aligned}$$

For design parameters ω, θ, M , the near-threshold affine path is

$$\begin{aligned} p &= \varepsilon, \quad u = 1 + (2 - \omega)\varepsilon + \theta\varepsilon^2, \quad v = \omega\varepsilon - \theta\varepsilon^2, \\ q &= \frac{1}{2} - \left(\frac{1}{2} + \omega\right)\varepsilon + \left(\theta - \frac{M}{2}\right)\varepsilon^2. \end{aligned}$$

Then the critical vector tends to one half of the homogeneous kernel vector. The diffusion excess over the exact linear threshold is

$$d_Z - 8 \sum_{j=2}^{m-1} d_j = 4 \left(M + 6\omega + 3\omega^2 - \frac{\omega^2}{m-2} \right) \varepsilon^2 + O(\varepsilon^3).$$

For $(\omega, \theta, M) = (2/9, 1/2, 1)$ and $m = 3$, the selected path gives

$$d_1 = \frac{\varepsilon(3\varepsilon + 8)}{6}, \quad d_2 = \frac{\varepsilon(4 - 9\varepsilon)}{9\varepsilon^2 + 32\varepsilon + 18}, \quad d_3 = \varepsilon, \quad d_Z = \frac{16\varepsilon}{9 - 13\varepsilon}.$$

All four entries are positive for $0 < \varepsilon \leq 10^{-3}$, and

$$d_Z - 8d_2 = \frac{8\varepsilon^2(197 - 99\varepsilon)}{(9 - 13\varepsilon)(9\varepsilon^2 + 32\varepsilon + 18)} > 0.$$

Let $\ell_\varepsilon^T(A_3 - D_\varepsilon) = 0$, and obtain w_0, w_2 from the reaction-derived Hessian tensor B_3 through the equations of Section S6. Substitution in

$$c_3(\varepsilon) = \frac{\ell_\varepsilon^T \{B_3(r, w_0) + \frac{1}{2}B_3(r, w_2)\}}{\ell_\varepsilon^T r}$$

gives the exact rational expression reconstructed by the public verifier and

$$c_3(\varepsilon) = \frac{6}{1379} + \frac{421985}{11409846}\varepsilon + O(\varepsilon^2),$$

with $c_3(\varepsilon) > 0$ throughout $0 < \varepsilon \leq 10^{-3}$ by an exact coefficient certificate. For the complete onset check, write

$$\det(\lambda I - A_3 + tD_\varepsilon) = \lambda^4 + a_1\lambda^3 + a_2\lambda^2 + a_3\lambda + a_4, \quad H_2 = a_1a_2 - a_3, \quad H_3 = a_3H_2 - a_1^2a_4.$$

After cancelling common factors and substituting $t = 1 + s$ and $\varepsilon = [1000(1 + w)]^{-1}$, the exact numerator and denominator polynomials for $a_1, a_2, a_3, a_4/(t - 1), H_2, H_3$ have strictly positive coefficients for $s, w \geq 0$. At $t = 1$, $a_4 = 0$ while $a_1, a_2, a_3, H_2 > 0$; hence zero is simple and the cubic complement is Hurwitz. Moreover $\partial_t a_4|_{t=1} > 0$, so $\partial_t \lambda|_{t=1} = -(\partial_t a_4)/a_3 < 0$, equivalently the $\mu = 1 - t$ crossing coefficient is positive. For every $t > 1$, the strict quartic Routh–Hurwitz inequalities prove stability, including every higher Neumann mode. Finally, the fixed-mass homogeneous factor is $q_3(\lambda) = (\lambda + 7)(\lambda^2 + 5\lambda + 2)$. Thus, for each fixed ε in the stated interval, this is a primary transverse subcritical control example, not a universal nonlinear lower bound; no dimension- or ε -uniform bifurcation neighborhood is claimed.

S10 Semilinear stability and robustness

At onset for the unit-equilibrium family, write

$$\mathcal{L}_0 = A_m + D_m \partial_{\xi\xi} : X_c \longrightarrow Y_c, \quad X_c = \left\{ u \in H_N^2 : \int_0^\pi c^T u = 0 \right\}, \quad Y_c = \left\{ g \in L^2 : \int_0^\pi c^T g = 0 \right\}.$$

The identity $c^T A_m = 0$ and the Neumann boundary conditions make these fixed-mass subspaces invariant under the linearized evolution. The Neumann cosine transform sends this operator to $A_m - k^2 D_m$. On the homogeneous coefficient the mass restriction gives the invertible map $A_m : c^\perp \rightarrow c^\perp$. The $k = 1$ block has kernel $\text{span}\{r\}$ and range $\ell^\perp$, while every $k \geq 2$ block is invertible by the all-mode certificate. For $k^2 \geq 2\|D_m^{-1} A_m\|$, the factorization

$$A_m - k^2 D_m = -k^2 D_m (I - k^{-2} D_m^{-1} A_m)$$

and the Neumann series give

$$\|(A_m - k^2 D_m)^{-1}\| \leq 2\|D_m^{-1}\|k^{-2}.$$

Thus compatible L^2 Fourier data have an H_N^2 preimage, and

$$\ker \mathcal{L}_0 = \text{span}\{r \cos \xi\}, \quad \text{ran } \mathcal{L}_0 = \left\{ g \in Y_c : \int_0^\pi \ell^T g \cos \xi \, d\xi = 0 \right\}.$$

This proves directly that $\mathcal{L}_0$ is Fredholm of index zero and that its cokernel is $\text{span}\{\ell \cos \xi\}$. The same cosine-block argument applies to the equilibrium-scaled family after replacing the mode matrices by $H_m(L)(A_m - k^2 \Delta)$ and the left vector by $\tilde{\ell}(L)$.

On the real fixed-integrated-mass L^2 space, the positive diagonal Neumann diffusion operator has domain H_N^2 , is sectorial, and generates an analytic semigroup with compact resolvent. Its half-order fractional domain is H^1 . The reaction Jacobian is a bounded multiplication perturbation. Moreover,

$$\alpha(A_m - k^2 D_m) \leq \left\| \frac{A_m + A_m^T}{2} \right\|_2 - k^2 d_{\min} \rightarrow -\infty,$$

so the onset complementary spectral gap is determined by finitely many low modes. Positive diffusion lower bounds and bounded reaction Jacobians give the corresponding uniform high-frequency bound along a sufficiently small patterned branch; analytic perturbation preserves the finitely many low-mode gaps. In one space dimension, H^1 is a Banach algebra and embeds into C^0 , so the quadratic mass-action Nemytskii map is smooth from H^1 to L^2 . The patterned branch lies in H_N^2 . The fixed-mass restriction removes the homogeneous conservation zero, and Neumann boundary conditions on

an interval leave no continuous translation-neutral mode. Strict spectral negativity therefore implies local exponential asymptotic stability in H^1 .

For robustness, perturb positive steady fluxes and equilibrium coordinates within the positive-equilibrium realization manifold, reconstructing the rate constants accordingly; also allow small perturbations of diffusion ratios and interval length. Use one scalar diffusion multiplier as the implicit variable. The critical eigenvalue derivative is nonzero, so the implicit function theorem continues the crossing. Low-mode gaps persist by analytic perturbation. A positive diffusion lower bound and a reaction-Jacobian upper bound give one neighborhood controlling all high modes. Smooth dependence of the reduced normal-form coefficients preserves the negative cubic sign and the branch spectral gap. This is a retuned codimension-one statement, not a claim that arbitrary nearby parameter points are already critical, and no dimension-uniform radius is claimed.

S11 Numerical protocol and independent verification

Cosine–Galerkin simulations use exact modal convolution for the quadratic reaction terms and an adaptive stiff integrator. Initial data are small perturbations of **1** in the critical direction. The first-mode amplitude is measured by the adjoint projection. Spatial and temporal refinement tables, open-format profile data, exact parameters, and all solver tolerances are included in the data archive. No numerical output is used in a proof.

The release contains independent programs for the reaction family, flux cone, all-spectrum block theorem, omission minors, principal-minor diffusion-ray theorem, diffusion criterion, contrast bounds, improved profile, Pareto family, mode certificates, harmonic corrections, cubic sign, and branch stability. Mutation tests change signs, endpoints, the factor eight, Fourier factors, and certificate coefficients and must be rejected.