

# Quiver Varieties and Symmetric Pairs

Yiqiang Li

University at Buffalo, The State University of New York

REPRESENTATION THEORY  
COMBINATORICS AND GEOMETRY  
October 2018



- Motivations: Schur dualities  
—based on joint works with H. Bao, J. Kujawa and W. Wang.

# Overview:

- Motivations: Schur dualities  
—based on joint works with H. Bao, J. Kujawa and W. Wang.
- Results on nilpotent Slodowy slices

# Overview:

- Motivations: Schur dualities  
—based on joint works with H. Bao, J. Kujawa and W. Wang.
- Results on nilpotent Slodowy slices
- The construction of iQuiver Varieties (iQV)

# Overview:

- Motivations: Schur dualities  
—based on joint works with H. Bao, J. Kujawa and W. Wang.
- Results on nilpotent Slodowy slices
- The construction of iQuiver Varieties (iQV)
- Connection with real classical groups

Schur duality and its generalizations.

- Schur duality:  $U(\mathfrak{gl}_n) \curvearrowright (\mathbb{C}^n)^{\otimes d} \curvearrowright \mathbb{C}[S_d]$ . (Schur  $\sim 1901$ )

# Background: Schur duality and beyond

Schur duality and its generalizations.

- Schur duality:  $U(\mathfrak{gl}_n) \curvearrowright (\mathbb{C}^n)^{\otimes d} \curvearrowright \mathbb{C}[S_d]$ . (Schur  $\sim$  1901)
- qSchur duality:  $U_q(\mathfrak{gl}_n) \curvearrowright T_{n,d} \curvearrowright \mathbf{H}_d$ , with  $T_{n,d} = (\mathbb{C}(q)^n)^{\otimes d}$ . (Jimbo, 1986)


# Background: Schur duality and beyond

Schur duality and its generalizations.


- Schur duality:  $U(\mathfrak{gl}_n) \curvearrowright (\mathbb{C}^n)^{\otimes d} \curvearrowleft \mathbb{C}[S_d]$ . (Schur  $\sim$  1901)
- qSchur duality:  $U_q(\mathfrak{gl}_n) \curvearrowright T_{n,d} \curvearrowleft \mathbf{H}_d$ , with  $T_{n,d} = (\mathbb{C}(q)^n)^{\otimes d}$ . (Jimbo, 1986)
- iSchur duality:  $U_q^\sigma(\mathfrak{gl}_n) \curvearrowright T_{n,d} \curvearrowleft \mathbf{H}_d^{B/C/D}$ . (Green 1997, Bao-Wang 2013)



# Compatibility and quantum symmetric pairs




## Compatibility and quantum symmetric pairs


$$U_q^\sigma(\mathfrak{gl}_n) \text{ is a coideal: } \Delta(U_q^\sigma(\mathfrak{gl}_n)) \subseteq U_q^\sigma(\mathfrak{gl}_n) \otimes U_q(\mathfrak{gl}_n).$$



## Compatibility and quantum symmetric pairs


$$U_q^\sigma(\mathfrak{gl}_n) \text{ is a coideal: } \Delta(U_q^\sigma(\mathfrak{gl}_n)) \subseteq U_q^\sigma(\mathfrak{gl}_n) \otimes U_q(\mathfrak{gl}_n).$$

The pair  $(U_q(\mathfrak{gl}_n), U_q^\sigma(\mathfrak{gl}_n))$  is a quantum symmetric pair.

These algebras were studied previously by Noumi, Letzter, Kolb, etc.


- A symmetric pair  $(\mathfrak{g}, \mathfrak{g}^\theta)$  is a complex semisimple Lie algebra  $\mathfrak{g}$  together with its fixed-point subalgebra  $\mathfrak{g}^\theta$  under an involution  $\theta$ .

# Symmetric pairs

- A symmetric pair  $(\mathfrak{g}, \mathfrak{g}^\theta)$  is a complex semisimple Lie algebra  $\mathfrak{g}$  together with its fixed-point subalgebra  $\mathfrak{g}^\theta$  under an involution  $\theta$ .
- Classification of symmetric pairs goes back to É Cartan, and is equivalent to the classification of *real simple* Lie algebras, via Satake diagrams= bicolor Dynkin diagrams with diagram involution.


# Symmetric pairs

- A symmetric pair  $(\mathfrak{g}, \mathfrak{g}^\theta)$  is a complex semisimple Lie algebra  $\mathfrak{g}$  together with its fixed-point subalgebra  $\mathfrak{g}^\theta$  under an involution  $\theta$ .
- Classification of symmetric pairs goes back to É Cartan, and is equivalent to the classification of *real simple* Lie algebras, via Satake diagrams = bicolor Dynkin diagrams with diagram involution.
- When specialized to  $q = 1$ , the pair  $(U_q(\mathfrak{gl}_n), U_q^\sigma(\mathfrak{gl}_n))$  is of type  $A_{iii}/A_{iv}$ , with Satake diagram (all vertices are white):



# Symmetric pairs

- A symmetric pair  $(\mathfrak{g}, \mathfrak{g}^\theta)$  is a complex semisimple Lie algebra  $\mathfrak{g}$  together with its fixed-point subalgebra  $\mathfrak{g}^\theta$  under an involution  $\theta$ .
- Classification of symmetric pairs goes back to É Cartan, and is equivalent to the classification of *real simple* Lie algebras, via Satake diagrams = bicolor Dynkin diagrams with diagram involution.
- When specialized to  $q = 1$ , the pair  $(U_q(\mathfrak{gl}_n), U_q^\sigma(\mathfrak{gl}_n))$  is of type  $A_{iii}/A_{iv}$ , with Satake diagram (all vertices are white):



- Caution:  $U_q^\sigma(\mathfrak{gl}_n)$  NOT a fixed-point subalgebra of  $U_q(\mathfrak{gl}_n)$ .



# Applications/Why do we care?

- iSchur duality is used by Bao-Wang to solve Kazhdan-Lusztig problem for  $\mathfrak{osp}_{m|n}$ , following Brundan's approach to  $\mathfrak{gl}_{m|n}$ .

# Applications/Why do we care?

- iSchur duality is used by Bao-Wang to solve Kazhdan-Lusztig problem for  $\mathfrak{osp}_{m|n}$ , following Brundan's approach to  $\mathfrak{gl}_{m|n}$ .
- iHowe duality is used by Ehrig-Stroppel to solve problems for  $\mathfrak{so}_{2m}$ .

# Applications/Why do we care?

- iSchur duality is used by Bao-Wang to solve Kazhdan-Lusztig problem for  $\mathfrak{osp}_{m|n}$ , following Brundan's approach to  $\mathfrak{gl}_{m|n}$ .
- iHowe duality is used by Ehrig-Stroppel to solve problems for  $\mathfrak{so}_{2m}$ .
- Bao-Wang's work contains an icanonical basis (iCB for short) for representations of coideal subalgebras of quantum  $\mathfrak{sl}_n$ .

# Applications/Why do we care?

- iSchur duality is used by Bao-Wang to solve Kazhdan-Lusztig problem for  $\mathfrak{osp}_{m|n}$ , following Brundan's approach to  $\mathfrak{gl}_{m|n}$ .
- iHowe duality is used by Ehrig-Stroppel to solve problems for  $\mathfrak{so}_{2m}$ .
- Bao-Wang's work contains an icanonical basis (iCB for short) for representations of coideal subalgebras of quantum  $\mathfrak{sl}_n$ .
- Bao-Wang's work on iCB has been extended by themselves to any coideal subalgebras of quantum groups of finite type.

- qSchur duality was realized geometrically by Beilinson, Lusztig and MacPherson (BLM) and Grojnowski and Lusztig ~1990.

# Geometric Schur duality

- qSchur duality was realized geometrically by Beilinson, Lusztig and MacPherson (BLM) and Grojnowski and Lusztig  $\sim 1990$ .
- Geometric iSchur duality by Bao, Kujawa, L. and Wang (BKLW) in 2014.

# Geometric Schur duality

- qSchur duality was realized geometrically by Beilinson, Lusztig and MacPherson (BLM) and Grojnowski and Lusztig  $\sim 1990$ .
- Geometric iSchur duality by Bao, Kujawa, L. and Wang (BKLW) in 2014.
- BKLW obtained an iCB for quantum symmetric pairs of type Aiii/iv. A slight modification leads to remarkable positivity (L.-Wang, Fan-L.). (weak categorification)

# Geometric Schur duality

- qSchur duality was realized geometrically by Beilinson, Lusztig and MacPherson (BLM) and Grojnowski and Lusztig  $\sim 1990$ .
- Geometric iSchur duality by Bao, Kujawa, L. and Wang (BKLW) in 2014.
- BKLW obtained an iCB for quantum symmetric pairs of type Aiii/iv. A slight modification leads to remarkable positivity (L.-Wang, Fan-L.). (weak categorification)
- Genuine categorification of (part of ) BKLW's work has been done by H. Bao, P. Shan, W. Wang and B. Webster.



# *i*Quiver varieties?

There is a ‘type  $A$ ’ line of research:

$$\begin{array}{ccccc} \mathcal{F}_{n,d} & \rightsquigarrow & T^*\mathcal{F}_{n,d} & \rightsquigarrow & \text{Nakajima varieties} \\ \mathfrak{sl}_n & & \text{dual} & & \mathfrak{g}_{ADE} \end{array}$$

# *i*Quiver varieties?

There is a ‘type A’ line of research:

$$\begin{array}{ccccc} \mathcal{F}_{n,d} & \rightsquigarrow & T^* \mathcal{F}_{n,d} & \rightsquigarrow & \text{Nakajima varieties} \\ \mathfrak{sl}_n & & \text{dual} & & \mathfrak{g}_{ADE} \end{array}$$

In light of the previous work, We should have a line of research, based on “classical type” geometry:

$$\begin{array}{ccccc} \mathcal{F}_{n,d}^\sigma & \rightsquigarrow & T^* \mathcal{F}_{n,d}^\sigma & \rightsquigarrow & \text{iQV????} \\ \mathfrak{sl}_n^\sigma & & \text{dual} & & \mathfrak{g}_{ADE}^\theta \end{array}$$

The existence of iQV is also conjectured by Weiqiang Wang.

# A simple answer

The simple answer to construct iQV:

## Analogue

$$\mathfrak{M}_\zeta(\mathbf{w}) \rightsquigarrow \mathfrak{g}$$

# A simple answer

The simple answer to construct iQV:

## Analogue

$$\begin{array}{lll} \mathfrak{M}_\zeta(\mathbf{w}) & \rightsquigarrow & \mathfrak{g} \\ iQV?? & \rightsquigarrow & \mathfrak{g}^\theta, \text{ for } \theta \in \text{Aut}(\mathfrak{g}) \end{array}$$

# A simple answer

The simple answer to construct iQV:

## Analogue

$$\mathfrak{M}_\zeta(\mathbf{w}) \rightsquigarrow \mathfrak{g}$$

$$iQV?? \rightsquigarrow \mathfrak{g}^\theta, \text{ for } \theta \in \text{Aut}(\mathfrak{g})$$

Answer:  $iQV = \mathfrak{M}_\zeta(\mathbf{w})^\sigma$ : the fixed point locus of  $\mathfrak{M}_\zeta(\mathbf{w})$  under an  $\sigma$ .

# Comparison, I

We have the following comparison of results in QV and iQV.

Nakajima varieties=QV	iQV
symplectic resolution	symplectic <b>partial</b> resolution

# Comparison, I

We have the following comparison of results in QV and iQV.

Nakajima varieties=QV	iQV
symplectic resolution	symplectic <b>partial</b> resolution
$\pi$ semismall	$\pi^\sigma$ semismall

# Comparison, I

We have the following comparison of results in QV and iQV.

Nakajima varieties=QV	iQV
symplectic resolution	symplectic <b>partial</b> resolution
$\pi$ semismall	$\pi^\sigma$ semismall
nilpotent Slodowy slices of $GL_n$	$GL_n, Sp_{2n}, O_n, \dots$



# Comparison, I

We have the following comparison of results in QV and iQV.

Nakajima varieties=QV	iQV
symplectic resolution	symplectic <b>partial</b> resolution
$\pi$ semismall	$\pi^\sigma$ semismall
nilpotent Slodowy slices of $GL_n$	$GL_n, Sp_{2n}, O_n, \dots$
Weyl groups action of type $A_\ell/D_\ell/E_{6,7,8}$	$ADE, B_\ell/C_\ell/F_4/G_2$

# Comparison, I

We have the following comparison of results in QV and iQV.

Nakajima varieties=QV	iQV
symplectic resolution	symplectic <b>partial</b> resolution
$\pi$ semismall	$\pi^\sigma$ semismall
nilpotent Slodowy slices of $GL_n$	$GL_n, Sp_{2n}, O_n, \dots$
Weyl groups action of type $A_\ell/D_\ell/E_{6,7,8}$	$ADE, B_\ell/C_\ell/F_4/G_2$
$\mathfrak{M}_\zeta$ connected, $\zeta$ generic	<b>not connected in general</b>

# Comparison, I

We have the following comparison of results in QV and iQV.

Nakajima varieties=QV	iQV
symplectic resolution	symplectic <b>partial</b> resolution
$\pi$ semismall	$\pi^\sigma$ semismall
nilpotent Slodowy slices of $GL_n$	$GL_n, Sp_{2n}, O_n, \dots$
Weyl groups action of type $A_\ell/D_\ell/E_{6,7,8}$	$ADE, B_\ell/C_\ell/F_4/G_2$
$\mathfrak{M}_\zeta$ connected, $\zeta$ generic	<b>not connected in general</b>
$\mathfrak{M}_0$ normal	<b>not normal in general</b>

# Comparison, I

We have the following comparison of results in QV and iQV.

Nakajima varieties=QV	iQV
symplectic resolution	symplectic <b>partial</b> resolution
$\pi$ semismall	$\pi^\sigma$ semismall
nilpotent Slodowy slices of $GL_n$	$GL_n, Sp_{2n}, O_n, \dots$
Weyl groups action of type $A_\ell/D_\ell/E_{6,7,8}$	$ADE, B_\ell/C_\ell/F_4/G_2$
$\mathfrak{M}_\zeta$ connected, $\zeta$ generic	<b>not connected in general</b>
$\mathfrak{M}_0$ normal	<b>not normal in general</b>
Unitary instantons	$Sp/SO$ instantons (Nakajima)



**Caution:** the automorphism  $\sigma$  is not always an involution. For the Weyl group action of type  $G_2$ ,  $\sigma$  is of order 6.

# Application I: Rectangular symmetry

One of the rectangular symmetries reads as follows

$$\begin{array}{ccc}
 \tilde{S}_{\mu', \lambda}^{\text{sp}} & \xrightarrow{\cong} & \tilde{S}_{\nu', \hat{\lambda}}^{\text{o}} \\
 \pi^{\sigma} \downarrow & & \downarrow \pi^{\tilde{\sigma}} \\
 S_{\mu', \lambda}^{\text{sp}} & \xrightarrow{\cong} & S_{\nu', \hat{\lambda}}^{\text{o}}
 \end{array}$$

where each pair  $(\mu', \tilde{\mu}')$  and  $(\lambda, \tilde{\lambda})$  can be fit into a rectangle:



# Special case: two-row Slodowy slices

As a special case of the rectangular symmetry, we recover

Henderson-Licata, 2013

$$S_{n^1, k^1 (n-k)^1}^{\mathfrak{sp}_n} \cong S_{1^1 (n+1)^1, (k+1)^1 (n+1-k)^1}^{\mathfrak{o}_{n+2}}.$$

# Special case: two-row Slodowy slices

As a special case of the rectangular symmetry, we recover

Henderson-Licata, 2013

$$S_{n^1, k^1(n-k)^1}^{\mathfrak{sp}_n} \cong S_{1^1(n+1)^1, (k+1)^1(n+1-k)^1}^{\mathfrak{o}_{n+2}}$$

Simultaneously, we also deduce

Wilbert, 2015; Ehrig-Stroppel, 2013

$$\mathcal{B}_{e_0}^{\mathfrak{sp}_n} \cong \mathcal{B}_{\hat{e}_o}^{\mathfrak{o}_{n+2}} \text{ — Springer fibers of the associated Slodowy slices.}$$

# Special case: two-row Slodowy slices

As a special case of the rectangular symmetry, we recover

Henderson-Licata, 2013

$$\mathcal{S}_{n^1, k^1(n-k)^1}^{\text{sp}_n} \cong \mathcal{S}_{1^1(n+1)^1, (k+1)^1(n+1-k)^1}^{o_{n+2}}.$$

Simultaneously, we also deduce

Wilbert, 2015; Ehrig-Stroppel, 2013

$$\mathcal{B}_{e_0}^{\text{sp}_n} \cong \mathcal{B}_{\hat{e}_0}^{so_{n+2}} \text{ — Springer fibers of the associated Slodowy slices.}$$

And we solve a conjecture of Henderson-Licata for free:



$$\tilde{\mathcal{S}}_{n^1, k^1(n-k)^1}^{\text{sp}_n} \cong \tilde{\mathcal{S}}_{1^1(n+1)^1, (k+1)^1(n+1-k)^1}^{o_{n+2}}.$$



# Application II: Column removal reduction


## Enhancement of Kraft-Procesi's column removal reduction

We have  $\mathcal{S}_{\check{\mu}', \check{\lambda}}^{\text{sp}|\check{\lambda}|} \cong \mathcal{S}_{\mu', \lambda}^{\text{o}|\lambda|}$ , if the partitions are related as follows.




# Application II: Row removal reduction

We have  $\mathcal{S}_{\ddot{\mu}', \ddot{\lambda}}^{\text{sp}|\ddot{\lambda}|} \cong \mathcal{S}_{\mu', \lambda}^{\text{sp}|\lambda|}$ , if




# An introduction to Nakajima varieties


One starts with a quiver  $Q$  with underlying graph  $\Gamma$ , and its framed version  $Q^f$  by adding an extra copy of vertex set and arrow connecting to the original vertices. For example



Consider the geometries:



Consider the geometries:



Each step yields rich geometries and contains much representation theoretical information for the Lie algebra  $\mathfrak{g}_\Gamma$  associated to  $\Gamma$ .

# Hamiltonian reduction

Specifically, consider the cotangent space  $T^*\text{rep}Q_{\mathbf{v},\mathbf{w}}^f$  of representations of  $Q_{\mathbf{v},\mathbf{w}}^f$  of fixed dimension vectors  $\mathbf{v}, \mathbf{w}$ . There is a (reductive/gauge) group  $G_{\mathbf{v}}$  acts nicely on  $T^*\text{rep}Q_{\mathbf{v},\mathbf{w}}^f$ . General machinery in symplectic geometry says that there is a moment map

$$\mu : T^*\text{rep}Q_{\mathbf{v},\mathbf{w}}^f \rightarrow (\text{Lie } G_{\mathbf{v}})^*.$$

# Hamiltonian reduction

Specifically, consider the cotangent space  $T^*\text{rep}Q_{\mathbf{v},\mathbf{w}}^f$  of representations of  $Q_{\mathbf{v},\mathbf{w}}^f$  of fixed dimension vectors  $\mathbf{v}, \mathbf{w}$ . There is a (reductive/gauge) group  $G_{\mathbf{v}}$  acts nicely on  $T^*\text{rep}Q_{\mathbf{v},\mathbf{w}}^f$ . General machinery in symplectic geometry says that there is a moment map

$$\mu : T^*\text{rep}Q_{\mathbf{v},\mathbf{w}}^f \rightarrow (\text{Lie}G_{\mathbf{v}})^*.$$

Nakajima (quiver) variety is defined to be the Hamiltonian reduction

$$\mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) = \mu^{-1}(\zeta_{\mathbb{C}}) //_{\zeta_{\mathbb{R}}} G_{\mathbf{v}}, \quad \zeta = (\zeta_{\mathbb{C}}, \zeta_{\mathbb{R}}).$$

## Rank one: $\zeta = (0, 1)$ or $(0, 0)$

In rank one case,

$$T^*_{\text{rep}} Q^f_{\mathbf{v}, \mathbf{w}} = \text{Hom}(\mathbb{C}^{\mathbf{w}}, \mathbb{C}^{\mathbf{v}}) \oplus \text{Hom}(\mathbb{C}^{\mathbf{v}}, \mathbb{C}^{\mathbf{w}})$$

and the fiber at 0 of the moment map is given by

$$\mu^{-1}(0) = \{(p, q) | pq = 0\}.$$



## Rank one: $\zeta = (0, 1)$ or $(0, 0)$

In rank one case,

$$T^*_{\text{rep}} Q^f_{\mathbf{v}, \mathbf{w}} = \text{Hom}(\mathbb{C}^{\mathbf{w}}, \mathbb{C}^{\mathbf{v}}) \oplus \text{Hom}(\mathbb{C}^{\mathbf{v}}, \mathbb{C}^{\mathbf{w}})$$

and the fiber at 0 of the moment map is given by

$$\mu^{-1}(0) = \{(p, q) | pq = 0\}.$$

### Rank one Nakajima variety

Nakajima varieties are

$$\mathfrak{M}_{(0,1)}(\mathbf{v}, \mathbf{w}) = \{(p, q) \in \mu^{-1}(0) | q \text{ injective}\} / GL(\mathbb{C}^{\mathbf{v}}), \text{ (GIT quotient)}$$

$$\mathfrak{M}_{(0,0)}(\mathbf{v}, \mathbf{w}) = \mu^{-1}(0) // GL(\mathbb{C}^{\mathbf{v}}) \quad (\text{categorical quotient})$$

# Rank one: Cotangent bundle of Grassmannian (Nakajima)

## Nakajima varieties and cotangent bundle of Grassmannian

The assignment  $(p, q) \mapsto (qp, \text{im}(q))$  identifies Nakajima varieties with the cotangent bundle of Grassmannian and its affinization.

$$\begin{array}{ccc} \mathfrak{M}_{(0,1)}(\mathbf{v}, \mathbf{w}) & \xrightarrow{\cong} & T^*\text{Gr}(\mathbf{v}, \mathbf{w}) \\ \downarrow & & \downarrow \text{Springer resolution} \\ \mathfrak{M}_{(0,0)}(\mathbf{v}, \mathbf{w}) & \longrightarrow & \{x \in \text{End}(\mathbb{C}^{\mathbf{w}}) \mid x^2 = 0, \dots\} \end{array}$$

## Ginzburg's setting

In general, the cotangent bundle  $T^*\mathcal{F}_{n,d}$  used in Ginzburg's construction is a very special case of Nakajima varieties of type A.

# Isomorphisms on quiver varieties

Let  $a : \Gamma \rightarrow \Gamma$  be a diagram automorphism.

Naive diagram isomorphism

$$a : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{a(\zeta)}(a(\mathbf{v}), a(\mathbf{w})).$$

# Isomorphisms on quiver varieties

Let  $a : \Gamma \rightarrow \Gamma$  be a diagram automorphism.

## Naive diagram isomorphism

$$a : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{a(\zeta)}(a(\mathbf{v}), a(\mathbf{w})).$$

## Rank one

$$a = 1.$$

# Isomorphisms on quiver varieties

Let  $a : \Gamma \rightarrow \Gamma$  be a diagram automorphism.

## Naive diagram isomorphism

$$a : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{a(\zeta)}(a(\mathbf{v}), a(\mathbf{w})).$$

## Rank one

$$a = 1.$$

Associate to each  $\mathbb{C}^{\mathbf{v}_i}$  and  $\mathbb{C}^{\mathbf{w}_i}$  a non-degenerate bilinear form.

# Isomorphisms on quiver varieties

Let  $a : \Gamma \rightarrow \Gamma$  be a diagram automorphism.

## Naive diagram isomorphism

$$a : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{a(\zeta)}(a(\mathbf{v}), a(\mathbf{w})).$$

## Rank one

$$a = 1.$$

Associate to each  $\mathbb{C}^{\mathbf{v}_i}$  and  $\mathbb{C}^{\mathbf{w}_i}$  a non-degenerate bilinear form. One can define automorphisms, via taking adjoints, on Nakajima's varieties:

## Isomorphism $\tau_{\zeta}$

$$\tau_{\zeta} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{-\zeta}(\mathbf{v}, \mathbf{w}).$$

# Isomorphisms on quiver varieties

Let  $a : \Gamma \rightarrow \Gamma$  be a diagram automorphism.

## Naive diagram isomorphism

$$a : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{a(\zeta)}(a(\mathbf{v}), a(\mathbf{w})).$$

## Rank one

$$a = 1.$$

Associate to each  $\mathbb{C}^{\mathbf{v}_i}$  and  $\mathbb{C}^{\mathbf{w}_i}$  a non-degenerate bilinear form. One can define automorphisms, via taking adjoints, on Nakajima's varieties:

## Isomorphism $\tau_{\zeta}$

$$\tau_{\zeta} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{-\zeta}(\mathbf{v}, \mathbf{w}).$$

## Rank one

$$\tau_{\zeta} : (p, q) \mapsto (-q^*, p^*) \text{ (modulo } G_{\mathbf{v}}).$$

Recall  $\mathcal{W}_\Gamma$  be the Weyl group of  $\Gamma$ .

Reflection functors  $S_\omega$  of Nakajima, Lusztig and Maffei

$$S_\omega : \mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{\omega(\zeta)}(\omega *_{\mathbf{w}} \mathbf{v}, \mathbf{w}), \forall \omega \in \mathcal{W}_\Gamma.$$



# Isomorphisms cont'd

Recall  $\mathcal{W}_\Gamma$  be the Weyl group of  $\Gamma$ .

Reflection functors  $S_\omega$  of Nakajima, Lusztig and Maffei

$$S_\omega : \mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{\omega(\zeta)}(\omega *_{\mathbf{w}} \mathbf{v}, \mathbf{w}), \forall \omega \in \mathcal{W}_\Gamma.$$

Rank one

$$S_i : (p, q) \mapsto (p', q'), \mathbb{C}^{\mathbf{v}} \xrightarrow{q} \mathbb{C}^{\mathbf{w}} \xrightarrow{p'} \mathbb{C}^{\mathbf{w}-\mathbf{v}} \text{ is exact and } qp = q'p'.$$

Taking the composition of the above three isomorphisms yields:

### Isomorphism $\sigma$

$$\sigma \equiv \sigma_{a,\zeta,\omega} := aS_{\omega}\tau_{\zeta} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{-a\omega(\zeta)}(a(\omega *_{\mathbf{w}} \mathbf{v}), a\mathbf{w}).$$

Taking the composition of the above three isomorphisms yields:

### Isomorphism $\sigma$

$$\sigma \equiv \sigma_{a,\zeta,\omega} := aS_{\omega}\tau_{\zeta} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{-a\omega(\zeta)}(a(\omega *_{\mathbf{w}} \mathbf{v}), a\mathbf{w}).$$

### Definition of iQV

$$\mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w})^{\sigma}, \quad \text{if } \zeta = -a\omega(\zeta), \mathbf{v} = a(\omega *_{\mathbf{w}} \mathbf{v}) \text{ and } \mathbf{w} = a\mathbf{w}.$$

Taking the composition of the above three isomorphisms yields:

### Isomorphism $\sigma$

$$\sigma \equiv \sigma_{a,\zeta,\omega} := aS_{\omega}\tau_{\zeta} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{-a\omega(\zeta)}(a(\omega *_{\mathbf{w}} \mathbf{v}), a\mathbf{w}).$$

### Definition of iQV

$$\mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w})^{\sigma}, \quad \text{if } \zeta = -a\omega(\zeta), \mathbf{v} = a(\omega *_{\mathbf{w}} \mathbf{v}) \text{ and } \mathbf{w} = a\mathbf{w}.$$

### The projective morphism $\pi^{\sigma}$

It comes equipped with a projective morphism:

$$\pi^{\sigma} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w})^{\sigma} \rightarrow \mathfrak{M}_0(\mathbf{v}, \mathbf{w})^{\sigma}.$$

Taking the composition of the above three isomorphisms yields:

### Isomorphism $\sigma$

$$\sigma \equiv \sigma_{a,\zeta,\omega} := aS_{\omega}\tau_{\zeta} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w}) \rightarrow \mathfrak{M}_{-a\omega(\zeta)}(a(\omega *_{\mathbf{w}} \mathbf{v}), a\mathbf{w}).$$

### Definition of iQV

$$\mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w})^{\sigma}, \quad \text{if } \zeta = -a\omega(\zeta), \mathbf{v} = a(\omega *_{\mathbf{w}} \mathbf{v}) \text{ and } \mathbf{w} = a\mathbf{w}.$$

### The projective morphism $\pi^{\sigma}$

It comes equipped with a projective morphism:

$$\pi^{\sigma} : \mathfrak{M}_{\zeta}(\mathbf{v}, \mathbf{w})^{\sigma} \rightarrow \mathfrak{M}_0(\mathbf{v}, \mathbf{w})^{\sigma}.$$

### Quiver variety is an iQV

$$\mathfrak{M}_{\zeta}^{\Gamma \times \Gamma}(\mathbf{w})^{\sigma} \cong \mathfrak{M}_{\zeta}(\mathbf{w}) \text{ for } \sigma = a\tau_{\zeta}.$$

## Rank one iQV

The assignment  $(p, q) \mapsto (qp, \text{im}(q))$  identifies iQV with the cotangent bundles of maximal isotropic Grassmannians.

$$\begin{array}{ccc} \mathfrak{M}_{(0,1)}(\mathbf{v}, \mathbf{w})^\sigma & \xrightarrow{\cong} & T^*\text{Gr}(\mathbf{v}, \mathbf{w})^{\sigma'} \\ \downarrow & & \downarrow \\ \mathfrak{M}_{(0,0)}(\mathbf{v}, \mathbf{w})^\sigma & \longrightarrow & \{x \in \text{End}(\mathbb{C}^{\mathbf{w}}) \mid x^2 = 0, \dots\}^{\sigma'} \end{array}$$

here  $\sigma'$  depends on the form on  $\mathbb{C}^{\mathbf{w}}$ .

The action of  $\sigma$  on  $(p, q)$  is  $(p, q) \xrightarrow{\tau_\zeta} (-q^*, p^*) \xrightarrow{S_i} (-(q^*)', (p^*)')$ .

The action of  $\sigma$  on  $(p, q)$  is  $(p, q) \xrightarrow{\tau_\zeta} (-q^*, p^*) \xrightarrow{S_i} (-(q^*)', (p^*)')$ .  
So it sends

$$qp \mapsto -(p^*)'(q^*)' = -p^*q^* = -(qp)^* \longleftrightarrow x \mapsto -x^*$$



The action of  $\sigma$  on  $(p, q)$  is  $(p, q) \xrightarrow{\tau_\zeta} (-q^*, p^*) \xrightarrow{S_i} (-(q^*)', (p^*)')$ .  
So it sends


$$qp \mapsto -(p^*)'(q^*)' = -p^*q^* = -(qp)^* \longleftrightarrow x \mapsto -x^*$$

$$\operatorname{im} q \mapsto \operatorname{im}(p^*)' = \ker q^* \longleftrightarrow F \mapsto F^\perp.$$

$\perp$  is taken with respect to the form on  $\mathbb{C}^{\mathbf{w}}$ .


# Fixed points vs Kraft-Procesi

Kraft-Procesi considered an  $A_n$  quiver with alternating forms, say:



# Fixed points vs Kraft-Procesi


Kraft-Procesi considered an  $A_n$  quiver with alternating forms, say:



Consider the fixed-points  $\mu^{-1}(0)^\sigma$ ,  $G_v^\sigma = O_{v_1} \times Sp_{v_2} \times O_{v_3} \times \dots$ .

# Fixed points vs Kraft-Procesi

Kraft-Procesi considered an  $A_n$  quiver with alternating forms, say:




Consider the fixed-points  $\mu^{-1}(0)^\sigma$ ,  $G_{\mathbf{v}}^\sigma = O_{\mathbf{v}_1} \times Sp_{\mathbf{v}_2} \times O_{\mathbf{v}_3} \times \dots$ .

**Theorem (Kraft-Procesi, 1982): classical nilpotent orbits**

$\mu^{-1}(0)^\sigma // G_{\mathbf{v}}^\sigma \cong \overline{\mathcal{O}_{\mu'}^{sp_{w_1}}}$ , for certain  $\mathbf{v}, \mathbf{w}$  and the associated  $\mu'$ .

# Fixed points vs Kraft-Procesi

Kraft-Procesi considered an  $A_n$  quiver with alternating forms, say:



Consider the fixed-points  $\mu^{-1}(0)^\sigma$ ,  $G_{\mathbf{v}}^\sigma = O_{\mathbf{v}_1} \times Sp_{\mathbf{v}_2} \times O_{\mathbf{v}_3} \times \cdots$ .

**Theorem (Kraft-Procesi, 1982): classical nilpotent orbits**

$\mu^{-1}(0)^\sigma // G_{\mathbf{v}}^\sigma \cong \overline{\mathcal{O}_{\mu'}^{sp_{w_1}}}$ , for certain  $\mathbf{v}, \mathbf{w}$  and the associated  $\mu'$ .

**Theorem: iQV and Kraft-Procesi**

In Kraft-Procesi's setting,  $\mathfrak{M}_0(\mathbf{v}, \mathbf{w})^\sigma \cong \mu^{-1}(0)^\sigma // G_{\mathbf{v}}^\sigma$ , for  $a = 1$

# Cotangent bundles of flag varieties of classical groups

GIT in Kraft-Procesi's approach (one of my mental blocks)

Geometric invariant theory does not seem to apply to Kraft-Procesi's approach: essentially no non-trivial character of  $G_V^\sigma$ .

# Cotangent bundles of flag varieties of classical groups

## GIT in Kraft-Procesi's approach (one of my mental blocks)

Geometric invariant theory does not seem to apply to Kraft-Procesi's approach: essentially no non-trivial character of  $G_V^\sigma$ .

## Theorem: iAnalogue of Ginzburg: Cotangent bundles of isotropic flag varieties


In Kraft-Procesi's setting, we get

$$\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma \cong T^* \mathcal{F}_{\mathbf{v}, \mathbf{w}}^{\text{sp}_{\mathbf{w}_1}}, \quad \text{if } \zeta = (1, 0), \omega = \omega_0.$$

where  $\omega_0$  is the longest element in  $\mathcal{W}_\Gamma$  and  $a = 1$ .

# Nakajima's generalization: nilpotent Slodowy slices


In more general type-A setting of Kraft-Procesi, say:





# Nakajima's generalization: nilpotent Slodowy slices

In more general type-A setting of Kraft-Procesi, say:




Nakajima asserted at several places that

$$\mu^{-1}(0)^\sigma // G_{\mathbf{v}}^\sigma \rightsquigarrow S_{\mu', \lambda}^{\text{sp}} \text{ or } S_{\mu', \lambda}^{\text{o}},$$

where  $S_{\mu', \lambda}^{\text{sp}} = \overline{\mathcal{O}_{\mu'}} \cap S_\lambda \cap \text{sp}_{\tilde{\mathbf{w}}_1}$  is a nilpotent Slodowy slice in  $\text{sp}_{\tilde{\mathbf{w}}_1}$ .

# Nakajima's generalization: nilpotent Slodowy slices

In more general type-*A* setting of Kraft-Procesi, say:



Nakajima asserted at several places that

$$\mu^{-1}(0)^\sigma // G_{\mathbf{v}}^\sigma \rightsquigarrow S_{\mu', \lambda}^{\text{sp}} \text{ or } S_{\mu', \lambda}^{\text{o}},$$

where  $S_{\mu', \lambda}^{\text{sp}} = \overline{\mathcal{O}_{\mu'}} \cap S_\lambda \cap \mathfrak{sp}_{\tilde{\mathbf{w}}_1}$  is a nilpotent Slodowy slice in  $\mathfrak{sp}_{\tilde{\mathbf{w}}_1}$ .

## Proposition

In the above setting, there is a closed immersion (isomorphism expected):

$$\mu^{-1}(0)^\sigma // G_{\mathbf{v}}^\sigma \hookrightarrow S_{\mu', \lambda}^{\text{sp}},$$

which relies on a result of quiver-analogue of classical invariants.

# Partial Resolutions of nilpotent Slodowy slices

## Theorem: Partial Resolutions of nilpotent Slodowy slices

Moreover, in the above type  $A$  Dynkin diagram setting, we have

$$\begin{array}{ccccc}
 \mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma & \xrightarrow{\cong} & \tilde{\mathcal{S}}_{\mu', \lambda}^{\text{sp}} & \hookrightarrow & T^* \mathcal{F}_{\tilde{\mathbf{v}}, \tilde{\mathbf{w}}}^{\text{sp}} \\
 \downarrow \pi^\sigma & & \downarrow \Pi & & \downarrow \Pi \\
 \text{im}(\pi^\sigma) & \xrightarrow{\cong} & \mathcal{S}_{\mu', \lambda}^{\text{sp}} & \hookrightarrow & \mathfrak{sp}_{\tilde{\mathbf{w}}}
 \end{array}$$

(where  $\tilde{\mathbf{w}} = \sum_{i \in I} i \mathbf{w}_i$ ,  $\tilde{\mathbf{v}} = \mathbf{v}_i + \sum_{j \geq i} (j - i) \mathbf{w}_j$ .)

This result leads to previous applications in classical geometries.

# Instantons on ALE space and Nakajima varieties of type A, due to Nakajima

## Unitary

Regular part of Nakajima varieties = unitary instantons on ALE spaces.

It is known to Nakajima that

## Classical type

Regular part of iQV (of some  $\sigma$ ) =  $SP/SO$  instantons on ALE spaces.  
(arxiv: 1801.06286.)

# General properties of iQV

Now we return to the general setting:

**Proposition: Independence of forms on  $V$**

The  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma$  is independent of choices of forms on  $V$ .

# General properties of iQV

Now we return to the general setting:

**Proposition: Independence of forms on  $V$**

The  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma$  is independent of choices of forms on  $V$ .

From now on, forms on  $W$  are either *orthogonal* or *symplectic*.

**Propositioin: Symplectic structure, semismallness**

(1) The  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma$  is a symplectic submanifold of  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})$ .

# General properties of iQV

Now we return to the general setting:

**Proposition: Independence of forms on  $V$**

The  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma$  is independent of choices of forms on  $V$ .

From now on, forms on  $W$  are either *orthogonal* or *symplectic*.

**Propositioin: Symplectic structure, semismallness**

- (1) The  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma$  is a symplectic submanifold of  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})$ .
- (2) The map  $\pi^\sigma : \mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma \rightarrow \text{im}(\pi^\sigma)$  is semismall.

# General properties of iQV

Now we return to the general setting:

## Proposition: Independence of forms on $V$

The  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma$  is independent of choices of forms on  $V$ .

From now on, forms on  $W$  are either *orthogonal* or *symplectic*.

## Proposition: Symplectic structure, semismallness

- (1) The  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma$  is a symplectic submanifold of  $\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})$ .
- (2) The map  $\pi^\sigma : \mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma \rightarrow \text{im}(\pi^\sigma)$  is semismall.

## Proposition: Weyl group action

Let  $\mathcal{W}_\Gamma^{a\omega} = \{x \in \mathcal{W}_\Gamma \mid x\omega = \omega x, a(x) = x\}$ . There exists a  $\mathcal{W}_\Gamma^{a\omega}$ -action:

$$\mathcal{W}_\Gamma^{a\omega} \curvearrowright H^*(\mathfrak{M}_\zeta(\mathbf{v}, \mathbf{w})^\sigma), \quad \mathbf{w} - \mathbf{C}_\Gamma \mathbf{v} = 0.$$

$\mathcal{W}_\Gamma^{a\omega}$  includes Weyl groups of  $B_\ell/C_\ell/F_4/G_2$  types.



## Conjecture

There is an action

$$\mathfrak{g}^\theta \curvearrowright H^*(\mathfrak{M}_\zeta(\mathbf{w})^\sigma), \quad (\zeta \text{ generic})$$

where  $(\mathfrak{g}, \mathfrak{g}^\theta)$  for a symmetric pair of type  $Ai, Aiii, Di, Ei, Eii, Ev, Eviii$ , whose Satake diagram has no black dots. Note  $\mathfrak{g}^\theta$  of type  $Ai$  is  $\mathfrak{so}_n$ .

It holds for  $Aiii/Aiv$ . There are several supporting evidence.

# Connection to real simple groups

Symmetric pairs have been pervasive in the study of representations of real simple/reductive groups.

## QV and real simple groups

Does QV/iQV have more direct connections with real classical groups?

# Cartan decomposition for quiver varieties

To any symmetric pair  $(\mathfrak{g}, \mathfrak{g}^\theta)$ , it yields a complex Cartan decomposition

$$\mathfrak{g} = \mathfrak{g}^\theta \oplus \mathfrak{p},$$

$\mathfrak{p}$  the eigenspace of eigenvalue  $-1$ .

# Cartan decomposition for quiver varieties

To any symmetric pair  $(\mathfrak{g}, \mathfrak{g}^\theta)$ , it yields a complex Cartan decomposition

$$\mathfrak{g} = \mathfrak{g}^\theta \oplus \mathfrak{p},$$

$\mathfrak{p}$  the eigenspace of eigenvalue  $-1$ . The geometry surrounding the nilcone  $\mathcal{N}(\mathfrak{p})$  also has significant representation theoretic information, especially in the orbit method of real reductive groups (Kirillov, Kostant, Vogan, etc.).

# Cartan decomposition for quiver varieties

To any symmetric pair  $(\mathfrak{g}, \mathfrak{g}^\theta)$ , it yields a complex Cartan decomposition

$$\mathfrak{g} = \mathfrak{g}^\theta \oplus \mathfrak{p},$$

$\mathfrak{p}$  the eigenspace of eigenvalue  $-1$ . The geometry surrounding the nilcone  $\mathcal{N}(\mathfrak{p})$  also has significant representation theoretic information, especially in the orbit method of real reductive groups (Kirillov, Kostant, Vogan, etc.).

## Quiver model of $\mathcal{N}(\mathfrak{p})$ : Lagrangian version of iQV

The anti-symplectic version, say  $\mathfrak{M}_\zeta(\mathbf{w})^{\hat{\sigma}}$ , of iQV yields a quiver/linear model of  $\mathcal{N}(\mathfrak{p})$  and associated Slodowy slices. Almost all results from symplectic version have an anti-symplectic counterpart, such as rectangular symmetry etc. (But not the semismallness of projection from  $\pi$ .)

# Cartan decomposition for quiver varieties

## A correspondence

$\mathfrak{g}$	$\mathfrak{g}^\theta$	$\mathfrak{p}$
$\updownarrow$	$\updownarrow$	$\updownarrow$
Nakajima varieties	Symplectic subvarieties	Lagrangian subvarieties

This correspondence works for all symmetric pairs  $(\mathfrak{g}, \mathfrak{g}^\theta)$  of classical type.

# Cartan decomposition for quiver varieties

## A correspondence

$\mathfrak{g}$	$\mathfrak{g}^\theta$	$\mathfrak{p}$
$\updownarrow$	$\updownarrow$	$\updownarrow$
Nakajima varieties	Symplectic subvarieties	Lagrangian subvarieties

This correspondence works for all symmetric pairs  $(\mathfrak{g}, \mathfrak{g}^\theta)$  of classical type. Note that  $\mathcal{N}(\mathfrak{p}) \cong \mathcal{N}(G_{\mathbb{R}})$ , the Kostant-Sekiguchi homeomorphism of Chen-Nadler. It is reasonable to expect the same holds in quiver setting.

## Kostant-Sekiguchi correspondence for quiver varieties

There should be a homeomorphism  $\mathfrak{M}_0(\mathbf{w})_{\mathbb{R}} \cong \mathfrak{M}_0(\mathbf{w})^{\hat{\sigma}}$ .

# Representations of symmetric pairs

Through the work of Harish-Chandra, the following two classes of representations are the same (very roughly speaking).

- Unitary representations of  $G_{\mathbb{R}}$ , a real simple group.



# Representations of symmetric pairs

Through the work of Harish-Chandra, the following two classes of representations are the same (very roughly speaking).

- Unitary representations of  $G_{\mathbb{R}}$ , a real simple group.
- Harish-Chandra  $(\mathfrak{g}, K)$ -modules where  $\mathfrak{g} = \mathbb{C} \otimes \text{Lie} G_{\mathbb{R}}$  and  $(G_{\mathbb{C}}, K)$  is a symmetric pair on group level.

# Representations of symmetric pairs

Through the work of Harish-Chandra, the following two classes of representations are the same (very roughly speaking).

- Unitary representations of  $G_{\mathbb{R}}$ , a real simple group.
- Harish-Chandra  $(\mathfrak{g}, K)$ -modules where  $\mathfrak{g} = \mathbb{C} \otimes \text{Lie} G_{\mathbb{R}}$  and  $(G_{\mathbb{C}}, K)$  is a symmetric pair on group level.

# Representations of symmetric pairs

Through the work of Harish-Chandra, the following two classes of representations are the same (very roughly speaking).

- Unitary representations of  $G_{\mathbb{R}}$ , a real simple group.
- Harish-Chandra  $(\mathfrak{g}, K)$ -modules where  $\mathfrak{g} = \mathbb{C} \otimes \text{Lie} G_{\mathbb{R}}$  and  $(G_{\mathbb{C}}, K)$  is a symmetric pair on group level.

A shadow of the  $(\mathfrak{g}, K)$ -module: Harish-Chandra  $(\mathfrak{g}, \text{Lie} K)$ -modules.  $\text{Lie} K = \mathfrak{g}^{\theta}$  for some  $\theta$ , studied by Dixmier, Lepowsky, Zuckerman, etc. On the other hand, there is a Yangian  $\mathcal{Y} \equiv \mathcal{Y}(\mathfrak{g}_{\Gamma})$  action on the torus equivariant cohomology  $H_{\mathbb{T}}^*(\mathfrak{M}_{\zeta}(\mathbf{w}))$  of Nakajima variety, due to Varagnolo (via correspondence), Maulik-Okounkov (via R-matrix).

# Representations of symmetric pairs

Through the work of Harish-Chandra, the following two classes of representations are the same (very roughly speaking).

- Unitary representations of  $G_{\mathbb{R}}$ , a real simple group.
- Harish-Chandra  $(\mathfrak{g}, K)$ -modules where  $\mathfrak{g} = \mathbb{C} \otimes \text{Lie} G_{\mathbb{R}}$  and  $(G_{\mathbb{C}}, K)$  is a symmetric pair on group level.

A shadow of the  $(\mathfrak{g}, K)$ -module: Harish-Chandra  $(\mathfrak{g}, \text{Lie} K)$ -modules.  $\text{Lie} K = \mathfrak{g}^{\theta}$  for some  $\theta$ , studied by Dixmier, Lepowsky, Zuckerman, etc. On the other hand, there is a Yangian  $\mathcal{Y} \equiv \mathcal{Y}(\mathfrak{g}_{\Gamma})$  action on the torus equivariant cohomology  $H_{\mathbb{T}}^*(\mathfrak{M}_{\zeta}(\mathbf{w}))$  of Nakajima variety, due to Varagnolo (via correspondence), Maulik-Okounkov (via R-matrix).

## $(\mathfrak{g}, \text{Lie} K)$ -structure in Nakajima varieties

There is an action of an (affine) symmetric pair  $(\mathcal{Y}, \mathcal{Y}_{\sigma}) \curvearrowright H_{\mathbb{T}}^*(\mathfrak{M}_{\zeta}(\mathbf{w}))$ , where  $\mathcal{Y}_{\sigma}$  is a twisted Yangian, i.e., an affinization of  $U(\mathfrak{g}_{\Gamma}^{\theta})$ .

# Representations of symmetric pairs

Through the work of Harish-Chandra, the following two classes of representations are the same (very roughly speaking).

- Unitary representations of  $G_{\mathbb{R}}$ , a real simple group.
- Harish-Chandra  $(\mathfrak{g}, K)$ -modules where  $\mathfrak{g} = \mathbb{C} \otimes \text{Lie} G_{\mathbb{R}}$  and  $(G_{\mathbb{C}}, K)$  is a symmetric pair on group level.

A shadow of the  $(\mathfrak{g}, K)$ -module: Harish-Chandra  $(\mathfrak{g}, \text{Lie} K)$ -modules.  $\text{Lie} K = \mathfrak{g}^{\theta}$  for some  $\theta$ , studied by Dixmier, Lepowsky, Zuckerman, etc. On the other hand, there is a Yangian  $\mathcal{Y} \equiv \mathcal{Y}(\mathfrak{g}_{\Gamma})$  action on the torus equivariant cohomology  $H_{\mathbb{T}}^*(\mathfrak{M}_{\zeta}(\mathbf{w}))$  of Nakajima variety, due to Varagnolo (via correspondence), Maulik-Okounkov (via R-matrix).

## $(\mathfrak{g}, \text{Lie} K)$ -structure in Nakajima varieties

There is an action of an (affine) symmetric pair  $(\mathcal{Y}, \mathcal{Y}_{\sigma}) \curvearrowright H_{\mathbb{T}}^*(\mathfrak{M}_{\zeta}(\mathbf{w}))$ , where  $\mathcal{Y}_{\sigma}$  is a twisted Yangian, i.e., an affinization of  $U(\mathfrak{g}_{\Gamma}^{\theta})$ .

How to lift  $(\mathfrak{g}, \text{Lie} K)$ -structure to a  $(\mathfrak{g}, K)$ -structure remains to be done.

- The above result is obtained via Maulik-Okounkov's R-matrix approach.

- The above result is obtained via Maulik-Okounkov's R-matrix approach.
- We show that certain fixed-point subvarieties of  $iQV$  is a product of  $QV$  of smaller ranks.

- The above result is obtained via Maulik-Okounkov's R-matrix approach.
- We show that certain fixed-point subvarieties of  $iQV$  is a product of  $QV$  of smaller ranks.
- **Stable envelopes exist.**



- The above result is obtained via Maulik-Okounkov's R-matrix approach.
- We show that certain fixed-point subvarieties of iQV is a product of QV of smaller ranks.
- Stable envelopes exist.
- The  $R$ -matrix (or rather  $K$ -matrix) satisfies the reflection equation  $RKRK = KRKR$ .

- The above result is obtained via Maulik-Okounkov's  $R$ -matrix approach.
- We show that certain fixed-point subvarieties of  $iQV$  is a product of  $QV$  of smaller ranks.
- Stable envelopes exist.
- The  $R$ -matrix (or rather  $K$ -matrix) satisfies the reflection equation  $RKRK = KRKR$ .
- The twisted Yangian  $\mathcal{Y}_\sigma$  is then constructed using  $K$ -matrix via Faddeev, Reshetikhin, and Takhtajan's construction.

# Comparison, II

Nakajima varieties	iQV / lgrngn version
symp. resolution	symp. partial resolution/ lgrngn
$\pi$ semismall	$\pi^\sigma$ semismall / proper
Slodowy slices of type $GL_n$	symmetric pairs of classical type
Weyl groups action of type $ADE$	$ADE, B_\ell/C_\ell/F_4/G_2$
Rectangular symmetry of $GL_n$	symmetric pairs of classical type
Column/row removal reduct. of $GL_n$	symmetric pairs of classical type

# Comparison, II

Nakajima varieties	iQV / <b>lgrngn version</b>
symp. resolution	symp. partial resolution/ <b>lgrngn</b>
$\pi$ semismall	$\pi^\sigma$ semismall / proper
Slodowy slices of type $GL_n$	<b>symmetric pairs of classical type</b>
Weyl groups action of type $ADE$	$ADE, B_\ell/C_\ell/F_4/G_2$
Rectangular symmetry of $GL_n$	<b>symmetric pairs of classical type</b>
Column/row removal reduct. of $GL_n$	<b>symmetric pairs of classical type</b>
$\mathfrak{g} \curvearrowright H_{top}(\mathfrak{M}_\zeta(\mathbf{w}))$	$\mathfrak{g}^\theta \curvearrowright H^*(\mathfrak{M}_\zeta(\mathbf{w})^\sigma)$ (conj.)

# Comparison, II

Nakajima varieties	iQV / <b>lgrngn version</b>
symp. resolution	symp. partial resolution/ <b>lgrngn</b>
$\pi$ semismall	$\pi^\sigma$ semismall / proper
Slodowy slices of type $GL_n$	<b>symmetric pairs of classical type</b>
Weyl groups action of type $ADE$	$ADE, B_\ell/C_\ell/F_4/G_2$
Rectangular symmetry of $GL_n$	<b>symmetric pairs of classical type</b>
Column/row removal reduct. of $GL_n$	<b>symmetric pairs of classical type</b>
$\mathfrak{g} \curvearrowright H_{top}(\mathfrak{M}_\zeta(\mathbf{w}))$	$\mathfrak{g}^\theta \curvearrowright H^*(\mathfrak{M}_\zeta(\mathbf{w})^\sigma)$ (conj.)
Geometric Rep( $\mathfrak{g}$ )	Geometric Rep( $\mathfrak{g}^\theta$ ) (conj.)

# Comparison, II

Nakajima varieties	iQV / <b>lgrngn version</b>
symp. resolution	symp. partial resolution/ <b>lgrngn</b>
$\pi$ semismall	$\pi^\sigma$ semismall / proper
Slodowy slices of type $GL_n$	<b>symmetric pairs of classical type</b>
Weyl groups action of type $ADE$	$ADE, B_\ell/C_\ell/F_4/G_2$
Rectangular symmetry of $GL_n$	<b>symmetric pairs of classical type</b>
Column/row removal reduct. of $GL_n$	<b>symmetric pairs of classical type</b>
$\mathfrak{g} \curvearrowright H_{top}(\mathfrak{M}_\zeta(\mathbf{w}))$	$\mathfrak{g}^\theta \curvearrowright H^*(\mathfrak{M}_\zeta(\mathbf{w})^\sigma)$ (conj.)
Geometric Rep( $\mathfrak{g}$ )	Geometric Rep( $\mathfrak{g}^\theta$ ) (conj.)
Maulik-Okounkov $\mathcal{R}$ -matrix	<b><math>\mathcal{K}</math>-matrix</b>

# Comparison, II

Nakajima varieties	iQV / lgrngn version
symp. resolution	symp. partial resolution/ lgrngn
$\pi$ semismall	$\pi^\sigma$ semismall / proper
Slodowy slices of type $GL_n$	symmetric pairs of classical type
Weyl groups action of type $ADE$	$ADE, B_\ell/C_\ell/F_4/G_2$
Rectangular symmetry of $GL_n$	symmetric pairs of classical type
Column/row removal reduct. of $GL_n$	symmetric pairs of classical type
$\mathfrak{g} \curvearrowright H_{top}(\mathfrak{M}_\zeta(\mathbf{w}))$	$\mathfrak{g}^\theta \curvearrowright H^*(\mathfrak{M}_\zeta(\mathbf{w})^\sigma)$ (conj.)
Geometric Rep( $\mathfrak{g}$ )	Geometric Rep( $\mathfrak{g}^\theta$ ) (conj.)
Maulik-Okounkov $\mathcal{R}$ -matrix	$\mathcal{K}$ -matrix
Yang-Baxter equation	Reflection equation

# Comparison, II

Nakajima varieties	iQV / <b>lgrngn version</b>
symp. resolution	symp. partial resolution/ <b>lgrngn</b>
$\pi$ semismall	$\pi^\sigma$ semismall / proper
Slodowy slices of type $GL_n$	<b>symmetric pairs of classical type</b>
Weyl groups action of type $ADE$	$ADE, B_\ell/C_\ell/F_4/G_2$
Rectangular symmetry of $GL_n$	<b>symmetric pairs of classical type</b>
Column/row removal reduct. of $GL_n$	<b>symmetric pairs of classical type</b>
$\mathfrak{g} \curvearrowright H_{top}(\mathfrak{M}_\zeta(\mathbf{w}))$	$\mathfrak{g}^\theta \curvearrowright H^*(\mathfrak{M}_\zeta(\mathbf{w})^\sigma) \text{ (conj.)}$
Geometric Rep( $\mathfrak{g}$ )	Geometric Rep( $\mathfrak{g}^\theta$ ) (conj.)
Maulik-Okounkov $\mathcal{R}$ -matrix	<b><math>\mathcal{K}</math>-matrix</b>
Yang-Baxter equation	<b>Reflection equation</b>
RTT formalism of Yangian	<b>Twisted Yangian</b>
$(\mathcal{Y}, \mathcal{Y}_\sigma) \curvearrowright H_{\mathbb{T}}^*(\mathfrak{M}_\zeta(\mathbf{w}))$	
<b>Kostant-Sekiguchi homeomorphism in quiver varieties (conj.)</b>	



Thank you very much!