

Does Frequent Breathing Help When Exercising?

Ali Fathi

Abstract

In this paper, we model the diffusion rate of Oxygen within lungs and compare the absorption capacity in two states, resting and exercising.

1 Model of Lung

Here, we model the lung as a simple membrane permeable to Oxygen:




Figure 1: Diffusion

The oxygen is diffused from the high-concentrated lung sac into low-concentrated blood capillaries.

On the other hand, we model the breathing behavior with a two steps action; *Holding* after a fast inhaling, and *tranquillity* after a fast exhaling (Fig 2).

We ignore those fast rising and falling transitions in our simplified modeling. Consider α to be the percentage of time for holding or inhaling (*i.e.*, $\frac{\alpha}{1-\alpha}$ as the inhale to exhale ratio), N_L as the number of Oxygen molecules inside a full lung, and N_B as the whole number of Oxygen molecules inside the capillaries in direct contact with the sacs, before absorbing new molecules. We define $N = N_L + N_B$. The diffusion rate from one side with n molecules to the another, is (linearly)




Figure 2: Breathing steps

related to the number of molecules in this side:

$$-\beta n$$

Besides, we assume that diffusion merely happens during the holding period, when the lungs have no interaction with outside. If $n(t)$ is the number of Oxygen molecules inside lungs, $N - n(t)$ will be the number of Oxygen molecules inside capillaries, and we will have the following equation:

$$\frac{dn}{dt} = -\beta n + \beta(N - n) = -2\beta\left(n - \frac{N}{2}\right) \quad (1)$$

and the answer is as follows ($n(0) = N_L$):

$$n(t) = \frac{N}{2} + \left(N_L - \frac{N}{2}\right)e^{-2\beta t}. \quad (2)$$

After Δt seconds holding the breath, the total amount of Oxygen absorbed by the capillaries is as follows:

$$\Delta n = N_L - n(\Delta t) = \left(N_L - \frac{N}{2}\right)(1 - e^{-2\beta\Delta t}). \quad (3)$$

2 Comparison Between Normal and Frequent Breathing

We compare a normal breathing with the period of T , and a fast breathing with the period of $\frac{T}{k}$ (breathing increased with a rate of k). In the first case, we absorb the following amount of Oxygen during T seconds ($\Delta t = \alpha T$):

$$\Delta n_1 = \left(N_L - \frac{N}{2}\right)(1 - e^{-2\alpha\beta T}) \quad (4)$$

and for the second case, we absorb the following amount of Oxygen during T seconds:

$$\Delta n_2 = k(N_L - \frac{N}{2})(1 - e^{-2\alpha\beta\frac{T}{k}}). \quad (5)$$

To have a better comparison, we approximate the exponential term with $e^x \approx 1 + x + \frac{x^2}{2}$. So we get:

$$\Delta n_1 = (N_L - \frac{N}{2})(1 - e^{-2\alpha\beta T}) \approx (N_L - \frac{N}{2})(2\alpha\beta T - 2\alpha^2\beta^2 T^2) \quad (6)$$

$$\Delta n_2 = k(N_L - \frac{N}{2})(1 - e^{-2\alpha\beta\frac{T}{k}}) \approx (N_L - \frac{N}{2})(2\alpha\beta T - 2\alpha^2\beta^2\frac{T^2}{k}) \quad (7)$$

that shows Δn_2 is slightly greater than Δn_1 , supporting the advantage for breathing faster during intense exercising!