


## Fusion: MCF - Homework #6

- 1 For improved performance, tokamak plasmas are almost always operated in x-point configurations with elongated plasmas. In this problem we look at the stability of a simplified version of this situation, shown in the figure below.



Here we approximate the plasma in a large aspect ratio tokamak ( $R \gg a$ ) as a toroidal wire at  $R = R_0$ ,  $z = 0$ , carrying a current  $I_p$ . This produces a poloidal field  $B_\theta = \mu_0 I_p / (2\pi r)$ , shown as solid circles in the figure. Coils 1 and 2 are placed at  $R = R_0$ ,  $z = \pm d$ , each carrying a current  $I_C$  in the same direction as  $I_p$ . The forces on the plasma due to these coils balance so it is in vertical equilibrium, and the net magnetic field has an x-point configuration (dashed lines).

- (a) By considering a motion of the plasma in the  $z$  direction by a small amount  $\delta z$ , show that the total vertical force on the plasma is approximately given by

$$\delta F \simeq 2\mu_0 I_p I_C \frac{R_0}{d^2} \delta z$$

[5]

- (b) This force amplifies the original perturbation, and so the configuration is unstable. Show that the growth rate of the instability is

$$\gamma_0 = \left( 2\mu_0 I_p I_C \frac{R_0}{d^2 M} \right)^{1/2}$$

continued

where  $M$  is the mass of the plasma. For a typical tokamak  $\gamma_0 \sim 10^6 \text{s}^{-1}$ , so x-point plasma should only exist for a few  $\mu\text{s}$  – which is clearly not the case in real experiments. [5]

- (c) So let's add two new coils at the same locations as coils 1 and 2, but let these new coils not be attached to a power supply. Moving the plasma at a velocity  $v = \partial_t \delta z$  will induce a current in these coils. Use Faraday's and Ohm's laws to show that this current is

$$I_i = -v \frac{\mu_0 I_p A}{2\pi d \eta}$$

where the total resistance of a coil of cross sectional area  $A$  and resistivity  $\eta$  is  $2\pi R_0 \eta / A$ . [Hint: consider the flux cut by the coils. No non-trivial integration should be needed.] [5]

- (d) By considering the force exerted on the plasma by these currents, show that the growth rate of the vertical instability is now modified to

$$\gamma_{\text{mod}} = \frac{\gamma_0^2}{2\gamma_i} \left[ \left( 1 + \frac{4\gamma_i^2}{\gamma_0^2} \right)^{1/2} - 1 \right]; \quad \gamma_i = \frac{2\pi\eta}{\mu_0 A} \frac{I_C}{I_p}$$

This  $\gamma_i$  is a characteristic rate of resistive changes in the passive coils. Typical values (e.g. copper  $\eta \sim 10^{-8} \Omega\text{m}$ , steel  $\eta \sim 10^{-6} \Omega\text{m}$ ) give  $\gamma_i \sim 1 - 10^2 \text{s}^{-1}$ .

- (i) Show that for small  $\gamma_i/\gamma_0$ , the growth rate is  $\gamma \simeq \gamma_i$ .

- (ii) Comment on the vertical stability of tokamak x-point configurations. [5]

**End of paper**