

Direct proof of a theorem about compact funcoids

OPG-59900	L1 - Tractable	Partiellement résolu	Topology
-----------	----------------	----------------------	----------

Énoncé original (anglais, reproduction mot pour mot)

Conjecture Let f is a T_1 -separable (the same as T_2 for symmetric transitive) compact funcoid and g is a uniform space (reflexive, symmetric, and transitive endoreloid) such that $(\mathsf{tmap}\{FCD\}) g = f$. Then $g = \langle f \times f \rangle^{\ast} \Delta$.

The main purpose here is to find a direct proof of this conjecture. It seems that this conjecture can be derived from the well known theorem about existence of exactly one uniformity on a compact set. But that would be what I call an indirect proof, we need a direct proof instead.

The direct proof may be constructed by correcting all errors and omissions in this draft article.

Direct proof could be better because with it we would get a little more general statement like this:

Conjecture Let f be a T_1 -separable compact reflexive symmetric funcoid and g be a reloid such that

- $(\mathsf{tmap}\{FCD\}) g = f$;
- $g \circ g^{-1} \sqsubseteq g$.

Then $g = \langle f \times f \rangle^{\ast} \Delta$.

Source et attribution. Énoncé reproduit depuis UnsolvedMath, jeu de données officiel sous licence CC BY 4.0.

Archive: <https://www.unsolvedmath.com/problems?difficulty=1>

Jeu de données: <https://huggingface.co/datasets/ulamai/UnsolvedMath>