

What are hyperfunctors isomorphic to?

OPG-59973	L1 - Tractable	Partiellement résolu	Topology
-----------	----------------	----------------------	----------

Énoncé original (anglais, reproduction mot pour mot)

Let $\mathfrak{A}$ be an indexed family of sets.

Products are $\prod A$ for $A \in \prod \mathfrak{A}$.

Hyperfunctors are filters $\mathfrak{F} \in \Gamma$ on the lattice Γ of all finite unions of products.

Problem Is $\bigcap \{\text{tmap}\{FCD\}\}$ a bijection from hyperfunctors $\mathfrak{F} \in \Gamma$ to:

- prestaroids on $\mathfrak{A}$;
- staroids on $\mathfrak{A}$;
- completary staroids on $\mathfrak{A}$?

If yes, is up^{Γ} defining the inverse bijection? If not, characterize the image of the function $\bigcap \{\text{tmap}\{FCD\}\}$ defined on $\mathfrak{F} \in \Gamma$.

Consider also the variant of this problem with the set Γ replaced with the set Γ^{last} of complements of elements of the set Γ .

Source et attribution. Énoncé reproduit depuis UnsolvedMath, jeu de données officiel sous licence CC BY 4.0.

Archive: <https://www.unsolvedmath.com/problems?difficulty=1>

Jeu de données: <https://huggingface.co/datasets/ulamai/UnsolvedMath>