

What is the smallest number of disjoint spanning trees made a graph Hamiltonian

OPG-567	L1 - Tractable	Partiellement résolu	Graph Theory
---------	----------------	----------------------	--------------

Énoncé original (anglais, reproduction mot pour mot)

We are given a complete simple undirected weighted graph $G_1=(V,E)$ and its first arbitrary shortest spanning tree $T_1=(V,E_1)$. We define the next graph $G_2=(V,E\setminus E_1)$ and find on G_2 the second arbitrary shortest spanning tree $T_2=(V,E_2)$. We continue similarly by finding $T_3=(V,E_3)$ on $G_3=(V,E\setminus \cup_{i=1}^2 E_i)$, etc. Let k be the smallest number of disjoint shortest spanning trees as defined above and let $T^k=(V,\cup_{i=1}^k E_i)$ be the graph obtained as union of all k disjoint trees.

Question 1. What is the smallest number of disjoint spanning trees creates a graph T^k containing a Hamiltonian path.

Question 2. What is the smallest number of disjoint spanning trees creates a graph T^k containing a shortest Hamiltonian path?

Questions 3 and 4. Replace in questions 1 and 2 a shortest spanning tree by a 1-tree. What is the smallest number of disjoint 1-trees creates a Hamiltonian graph? What is the smallest number of disjoint 1-trees creates a graph containing a shortest Hamiltonian cycle?

Source et attribution. Énoncé reproduit depuis UnsolvedMath, jeu de données officiel sous licence CC BY 4.0.

Archive: <https://www.unsolvedmath.com/problems?difficulty=1>

Jeu de données: <https://huggingface.co/datasets/ulamai/UnsolvedMath>