

# factorMerger: A Set of Tools to Support Results From Post Hoc Testing

by Agnieszka Sitko and Przemysław Biecek

**Abstract** ANOVA-like statistical tests for differences among groups are available for almost a hundred years. However, for large number of groups the results from commonly used *post-hoc* tests are often hard to interpret. To deal with this problem, the **factorMerger** package constructs and plots the hierarchical relation for the considered factor. Such a hierarchical structure is derived based on the *Likelihood Ratio Test* and is presented with the *Factor Merger Tree* created with the **ggplot2** package (Wickham, 2009). The current implementation handles one-dimensional and multi-dimensional Gaussian models as well as binomial and survival models. In this article the methodological background is outlined and the main functionalities of the package are illustrated using real-data examples.

## Introduction

In this article we present the **factorMerger** package that enriches results from ANOVA tests. The ANOVA method verifies the null hypothesis that a variable of interest  $y$  has the same distribution in all groups that are being compared. If this null hypothesis is rejected a more detailed analysis of differences among categorical variable levels might be needed. The traditional approach is to perform *pairwise post hoc tests* in order to verify which groups differ significantly.

One may find implementations of the traditional *post hoc tests* in many R packages. Package **agricolae** (de Mendiburu, 2016) offers a wide range of them. It gives one of the most popular *post hoc test*, Tukey HSD test (`HSD.test`), its less conservative version — Student-Newman-Keuls test (`SNK.test`) or Scheffe test (`scheffe.test`) which is robust to factor imbalance. These parametric tests are based on Student's t-distribution, thus, are reduced to Gaussian models only. In contrasts, **multcomp** package (Hothorn et al., 2008) can be used with generalized linear models (function `glht`) as it uses general linear hypothesis. Similarly to the **multcomp**, some implementations that accept `glm` objects are also given in **car** (LinearHypothesis, Fox and Weisberg, 2011) and **lsmeans** (Lenth, 2016).

However, an undeniable disadvantage of single-step *post hoc tests* is the inconsistency of their results. For a fixed significance level, it is possible that mean in group A does not differ significantly from the one in group B, similarly with groups B and C. At the same time the difference between group A and C is detected. Then data partition is unequivocal and, as a consequence, impossible to put through.

The problem of clustering categorical variable into non-overlapping groups has already been present in the literature. First, J. Tukey proposed an iterative procedure of merging factor levels based on the studentized range distribution (Tukey, 1949). However, statistical test used in this approach made it limited to Gaussian models. *Collapse And Shrinkage in ANOVA* (CAS-ANOVA, Bondell and Reich, 2008) is an algorithm that extends categorical variable partitioning for generalized linear models in testing. It is based on the Tibshirani's *Fused LASSO* (Tibshirani et al., 2005) with the constraint taken on the pairwise differences within a factor, which yields to their smoothing.

*Delete or Merge Regressors* algorithm (Prochenka, 2016, p. 37) is also adjusted to generalized linear models. It directly uses the hierarchical clustering to gain hierarchical structure of a factor. At the beginning, *DMR4glm* calculates the likelihood ratio test statistics for models arising from pairwise merging of factor levels or deleting factor levels against the full model (the one with all groups included). Then it performs agglomerative clustering taking LRT statistic as a distance — each step of clustering is associated with a model with different factor structure. Experimental studies (Prochenka, 2016, p. 44–91) showed that the *Delete or Merge Regressors*'s performance is better than CAS-ANOVA's when it comes to the model accuracy. *Delete or Merge Regressors*'s implementation may be found in the **DMR** package (Maj et al., 2013).

In this article we present a more direct approach to the problem of merging groups that are being compared. The **factorMerger** package offers an algorithm of hierarchical clustering of factors based on a backward iterative procedure. In each step it chooses a model with the highest *Likelihood Ratio Test* test p-value or, in other words, the highest likelihood. While this algorithm is more complex than *DMR4glm*, it maximizes the likelihood on the merging path<sup>1</sup>. What is more, it is easily expandable for non-parametric models (using permutation tests instead of LRTs).

In addition to the comprehensive algorithm which tries to merge all possible pairs of levels in

<sup>1</sup> Although it may be shown that the *DMR* algorithm is a consistent model selection method, its performance on smaller datasets is undefined. TODO....

a step, also a *successive version* is provided. In the *successive version* only levels which are relatively close can be merged (levels distance is dependent on the model chosen). While the basic approach (all vs. all comparisons) may result in a slightly better partition from the statistical point of view, proposed extension (all vs. subsequent comparisons) seems to be more graceful when it comes to the interpretation. Moreover, the former algorithm is more computationally expensive.

Furthermore, the `factorMerger` package gives an approximate implementation of *DMR4glm* (skipping the deleting procedure) with its *successive version*.

More detailed description of all algorithms implemented in `factorMerger` is given in the section [Methodology](#).

## Methodology

Merging procedures implemented in the `factorMerger` package begin with the full model — with all levels of a given factor included — and iteratively merge one pair of levels until the factor is constant. Uniting two groups reduces by one the number of subsets, so, as initially we have finite number of levels, the procedure will eventually obtain one-level-factor and terminate. In a single iteration *all possible* pairs are considered and the one which optimizes some objective function is joined. Objective functions use likelihood-based statistics we will describe later on.

The `factorMerger` package gives the ability to perform analysis for the wide family of models and choose from the broad spectrum of merging approaches. Depending on the problem statement, some parts of the merging procedure may differ. The general sketch of the algorithm is described below.

---

### Algorithm 1 The outline of the merging procedure

---

```

function MERGEFACTORS(response, factor, family, successive, method)
2:   pairsSet := generatePairs(response, factor, successive)
   M := createModel(response, factor, family)
4:   while |levels(factor)| > 1 do
       toBeMerged := argmaxpair ∈ pairsSet objectiveFunction(pair)
6:   M := updateModel(M0, toBeMerged)
       factor := mergeLevels(factor, pair)
8:   pairsSet := removePair(pairsSet, pair)
   end while
10: end function

```

---

## Model family

In the current version the package supports parametric models:

- single dimensional Gaussian (with the argument family = "gaussian"),
- multi dimensional Gaussian — Gaussian model with multiple  $y$  outputs (with the argument family = "gaussian")<sup>2</sup>,
- binomial (with the argument family = "binomial"),
- survival (with the argument family = "survival").

Each case has its own method of estimating model parameters and a specific likelihood formula. Here we will introduce notation used in the article. We will denote:

- the sample size as  $n$ ,
- the initial number of factor levels as  $k$ ,
- the binary matrix representing group membership as  $X = \{x_{ij}\}_{i,j=1}^{n,k}$ .

$$x_{ij} = \begin{cases} 1 & \text{if } i\text{-th observation belongs to group } j \\ 0 & \text{otherwise.} \end{cases}$$

We assume that groups do not overlap.

- the effects vector of a length  $k$  as  $\beta$ .
- the response vector as  $y = (y_1, \dots, y_n)$  or the response matrix as  $Y = \{y_{ij}\}_{i,j=1}^{n,m}$ .

---

<sup>2</sup>Both single dimensional and multi dimensional Gaussian models use family = "gaussian". However, multi dimensional model uses different functions for likelihood estimation and may require additional preprocessing, thus, it is considered as a separate category.

**Single dimensional Gaussian model** A convenient and commonly made assumption in the analysis of variance is the normality of the errors. Here we will consider a linear model in which beta coefficients represent group means. The model may be written in vector form as

$$y = X\beta + \epsilon, \quad \epsilon \sim \mathcal{N}(0, \sigma^2).$$

Under the above assumptions we may formulate the likelihood of the Gaussian linear model (Friedman et al., 2001, p.31)

$$L(\beta, \sigma | y) = (2\pi\sigma^2)^{-\frac{n}{2}} \exp\left(-\frac{1}{2}(y - X\beta)^T (y - X\beta) / \sigma^2\right)$$

and the corresponding logarithm of the likelihood

$$l(\beta, \sigma | y) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(\sigma^2) - \frac{1}{2}(y - X\beta)^T (y - X\beta) / \sigma^2.$$

To calculate the loglikelihood in the package we use `logLik.lm{stats}`.

**Multi dimensional Gaussian model** In the multi dimensional generalization of the Gaussian model we will observe multiple outputs  $Y$ . It takes the following form

$$Y = X\beta + E, \quad E \sim \mathcal{N}(0, \Sigma),$$

where  $E = (\epsilon_1, \epsilon_2, \dots, \epsilon_m)$  is an  $m$ -dimensional error.

Now, we may write the likelihood

$$L(\beta, \Sigma | Y) = (|2\pi\Sigma|)^{-\frac{1}{2}} \exp\left(-\frac{1}{2}(Y - X\beta)^T \Sigma^{-1} (Y - X\beta)\right)$$

and its logarithm

$$l(\beta, \Sigma | Y) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \log(|\Sigma|) - \frac{1}{2}(Y - X\beta)^T \Sigma^{-1} (Y - X\beta).$$

Unfortunately, `stats` or any commonly used R package do not support multiple responses in the loglikelihood calculation for linear Gaussian models. In the package we use `logLik.mlm` implementation introduced in the **Atools** package (B<b6>ck, 2014) and the `dmvnorm{mvtnorm}` (Genz and Bretz, 2009) implementation for multivariate normal density estimation.

**Binomial model** In the binomial case the observed response  $y$  will take one of two possible disjoint outcomes (success or failure). We will be interested whether probabilities of success in given subpopulations are statistically the same. Let us introduce the assumption

$$y \sim \mathcal{B}(p, n)$$

where  $\mathcal{B}(p, n)$  is the binomial distribution with the probability of success  $p$  and the number of trials  $n$ . We consider the logit model

$$\ln\left(\frac{p}{1-p}\right) = X\beta.$$

Let  $z = \sum_{i=1}^n y_i$ . Now the likelihood may be written as follows (Czepiel, 2002)

$$L(\beta | y) = \frac{n!}{z!(n-z)!} p^z (1-p)^{n-z}.$$

Thus, the logarithm of the likelihood is expressed as

$$l(\beta | y) = zX\beta - n \log(1 + \exp^{X\beta}).$$

TODO: Policzyć loglik, czy na pewno dobrze.

To calculate the loglikelihood for the binomial model we use `logLik.glm{stats}`.

**Survival model** Survival analysis is a branch of statistics for analyzing the time until a specified event takes to happen (such as a patient death or a machine failure). Usually we assume that survival time of an individual depends on two factors: the underlying baseline hazard function and a set of explanatory variables. In the package for survival analysis applications the *Cox proportional hazard regression* introduced by D. R. Cox (Cox, 1992) is used. In this model we assume that the baseline hazard function is the same for the whole population and the effects of predictors are multiplicatively related to the individual hazard.

Let  $\lambda(t)$  be the hazard function and let  $\lambda_0(t)$  be the baseline hazard function, where  $t$  denotes time. Then the Cox model has the following form

$$\lambda(t) = \lambda_0(t) \cdot \exp(X\beta).$$

Let  $y = (y_1, \dots, y_n)$  be a vector of observed times in the sample. For  $i$ -th observation  $y_i$  can be either event time or censoring time. We say that an observation is censored if we do not have complete information about its status in the context of our interest (for example, the patient died of causes other than the disease of our consideration or was lost to follow-up). Let  $C = (C_1, \dots, C_n)$  be the vector of the indicators that the time corresponds to an event. To be more precise,

$$C_i = \begin{cases} 0 & \text{if } i\text{-th observation was censored,} \\ 1 & \text{if for } i\text{-th observation the event occurred.} \end{cases}$$

Then we may construct the partial likelihood (as a function of  $\beta$  only) in the following way

$$L(\beta|y) = \prod_{i:C_i=1} \left[ \frac{\exp(X_i\beta)}{\sum_{j:y_j \geq y_i} \exp(X_j\beta)} \right]$$

and the corresponding loglikelihood

$$l(\beta|y) = \sum_{i:C_i=1} \left( X_i\beta - \log \left( \sum_{j:y_j \geq y_i} \exp(X_j\beta) \right) \right).$$

TODO: Dlaczego możemy korzystać z częściowej wiarygodności?

The Cox regression is implemented in the **survival** package (`coxph{survival}`, Terry M. Therneau and Patricia M. Grambsch, 2000) and the loglikelihood of the model one may find by accessing the field `loglik` of a `coxph` object.

## Pairs considered

Set of hypotheses that are tested during merging may be either comprehensive or limited. This gives two possibilities:

- *all-to-all* (with the argument `successive = FALSE`),
- *successive* (with the argument `successive = TRUE`).

The version *all-to-all* considers all possible pairs of factor levels. In the *successive* approach factor levels are preliminarily sorted and then only consecutive groups are tested for means equality.

## Defining levels similarity

The **factorMerger** package for each model family and merging strategy implements two types of a single iteration of the algorithm. They use one of the following:

- *Likelihood Ratio Test* (with the argument `method = "LRT"`),
- *agglomerative clustering with constant distance matrix* (based on the *DMR4glm* algorithm, with the argument `method = "hclust"`).

## The successive merging

In the *successive* version of the algorithm levels of a categorical variable are sorted. The order depends on the model chosen family chosen.

model	metric
one-dimensional Gaussian	average
multi-dimensional Gaussian	average of the isoMDS transformation
binomial	proportion of successes
survival	relative survival rate

**Table 1:** Factor ordering by model family

For one-dimensional Gaussian and binomial models groups are sorted by means and proportions of success, respectively. In the survival case we estimate survival model, which takes all factor levels separately. Then beta coefficient approximations specify levels order (base level gets coefficient equal to zero).

Multi dimensional Gaussian model needs additional preprocessing. First, group means are computed. Then they are projected into one dimension with the use of the Kruskal's non-metric multidimensional scaling. The `factorMerger` uses isoMDS implementation from the package **MASS** (Venables and Ripley, 2002).

Having set the factor order, we may limit number of comparisons in each step.

### The *all-to-all* merging

Short description...

### The Likelihood Ratio Test statistics

The substantial part of `factorMerger`'s algorithms is calculating the *Likelihood Ratio Test* statistics. In this section we define *LRT* statistic used in merging.

Let us assume  $y$  is a response variable and  $C$  is a factor with  $k$  levels ( $C \in \{1, 2, \dots, k\}$ ). We denote as  $h$  some linear hypothesis on the levels of  $C$ ,  $M_0$  the initial model (taking all factor levels independently) and  $M_h$  — the model under  $h$ . Then,  $LRT(M_h|M_0)$  statistic based on the *Likelihood Ratio Test* is defined as below.

$$LRT(M_h|M_0) = 2 \cdot l(M_0) - 2 \cdot l(M_h),$$

where  $l(\cdot)$  is log-likelihood function.

As  $M_h$  is nested in  $M_0$ , the likelihood of  $M_h$  is not greater than the  $M_0$ 's likelihood. Therefore, if  $\mathcal{H}$  is a set of considered linear hypothesis, hypothesis

$$\operatorname{argmin}_{h \in \mathcal{H}} LRT(M_h|M_0) = \operatorname{argmax}_{h \in \mathcal{H}} l(M_h)$$

will reduce likelihood the least.

**Asymptotic behaviour of the *LRT* statistic** A convenient result by Samuel S. Wilks (Wilks, 1938) shows that  $LRT(M_h|M_0)$  tends asymptotically to chi-squared distribution with degrees of freedom equal to the difference in degrees of freedom between  $M_0$  and  $M_h$  as number of observations approaches infinity. This convergence will be used to evaluate model's 'statistical correctness'.

### The Likelihood Ratio Test-based merging

The *Likelihood Ratio Test*-based approach minimizes likelihood reduction in the merging path. It may be summarized as follow.

TODO: Rozwinać... (Analogia do LRT testów, ale można uprościć do samego loglik)

**Algorithm 2** Merging with the LRT

---

```

function MERGEFACTORS(response, factor, successive)
2:   pairsSet := generatePairs(response, factor, successive)
      M0 := full model
4:   while levels(factor) > 1 do
      toBeMerged := argmaxpair ∈ pairsSet l(updateModel(M0, pair))
6:   M0 := updateModel(M0, toBeMerged)
      factor := mergeLevels(factor, pair)
8:   pairsSet := pairsSet \ pair
      end while
10: end function

```

---

**The DMR4glm-based merging**

TODO: Wstępny opis

**Algorithm 3** Merging with agglomerative clustering

---

```

function MERGEFACTORS(response, factor, successive)
2:   pairsSet := generatePairs(response, factor, successive)
      dist := set of distances
4:   for all pair ∈ pairsSet do
      h := {μpair1 = μpair2}           ▷ hypothesis under which pair is merged
6:   dist[pair] = LRT(Mh|M0)
      end for
8:   if successive then
      hClust(dist, method = "single")
10:  else
      hClust(dist, method = "complete")
12:  end if
end function

```

---

**An R package [factorMerger](#)**

The [factorMerger](#) package provides easy-to-use functions for factor merging and visualizing obtained results.

TODO: Tutaj da się opis głównych funkcji - co robią i tabelki z opisem parametrów

**Merging and getting results****Visualizations**

TODO: Tutaj dojdzie jeszcze opis każdego z paneli osobno

**CASE STUDY: PISA2012****Summary**

Some summary.

**Acknowledgements**

The authors thank to ... .

Work on this package was financially supported by the NCN *Opus grant* 2016/21/B/ST6/02176.

## Bibliography

- A. B<b6>ck. *Atools: Atools*, 2014. URL <https://R-Forge.R-project.org/projects/biostat/>. R package version 0.2/r191. [p3]
- H. D. Bondell and B. J. Reich. Simultaneous factor selection and collapsing levels in ANOVA. *Department of Statistics, North Carolina State University*, 2008. [p1]
- D. R. Cox. Regression models and life-tables. In *Breakthroughs in statistics*, pages 527–541. Springer, 1992. [p4]
- S. A. Czepiel. Maximum likelihood estimation of logistic regression models: theory and implementation. *Available at czep. net/stat/mlelr. pdf*, 2002. [p3]
- F. de Mendiburu. *agricolae: Statistical Procedures for Agricultural Research*, 2016. URL <https://CRAN.R-project.org/package=agricolae>. R package version 1.2-4. [p1]
- J. Fox and S. Weisberg. *An R Companion to Applied Regression*. Sage, Thousand Oaks CA, second edition, 2011. URL <http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>. [p1]
- J. Friedman, T. Hastie, and R. Tibshirani. *The elements of statistical learning*, volume 1. Springer series in statistics Springer, Berlin, 2001. [p3]
- A. Genz and F. Bretz. *Computation of Multivariate Normal and t Probabilities*. Lecture Notes in Statistics. Springer-Verlag, Heidelberg, 2009. ISBN 978-3-642-01688-2. [p3]
- T. Hothorn, F. Bretz, and P. Westfall. Simultaneous inference in general parametric models. *Biometrical Journal*, 50(3):346–363, 2008. [p1]
- R. V. Lenth. Least-squares means: The R package lsmeans. *Journal of Statistical Software*, 69(1):1–33, 2016. doi: 10.18637/jss.v069.i01. [p1]
- A. Maj, A. Prochenka, and P. Pokarowski. *DMR: Delete or Merge Regressors for linear model selection.*, 2013. URL <https://CRAN.R-project.org/package=DMR>. R package version 2.0. [p1]
- A. Prochenka. *Delete or Merge Regressors algorithm*, chapter 4.3, 5–6, pages 37, 44–91. 2016. [p1]
- Terry M. Therneau and Patricia M. Grambsch. *Modeling Survival Data: Extending the Cox Model*. Springer, New York, 2000. ISBN 0-387-98784-3. [p4]
- R. Tibshirani, M. Saunders, S. Rosset, J. Zhu, and K. Knight. Sparsity and smoothness via the fused lasso. *Journal of the Royal Statistical Society*, pages 01–108, 2005. [p1]
- J. Tukey. Comparing Individual Means in the Analysis of Variance. *BIOMETRICS*, pages 99–114, 1949. [p1]
- W. N. Venables and B. D. Ripley. *Modern Applied Statistics with S*. Springer, New York, fourth edition, 2002. URL <http://www.stats.ox.ac.uk/pub/MASS4>. ISBN 0-387-95457-0. [p5]
- H. Wickham. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, 2009. ISBN 978-0-387-98140-6. URL <http://ggplot2.org>. [p1]
- S. S. Wilks. The large-sample distribution of the likelihood ratio for testing composite hypotheses. *The Annals of Mathematical Statistics*, 9(1):60–62, 1938. [p5]

Agnieszka Sitko  
University of Warsaw  
Faculty of Mathematics, Informatics and Mechanics  
Poland  
[ag.agnieszka.sitko@gmail.com](mailto:ag.agnieszka.sitko@gmail.com)

Przemysław Biecek  
University of Warsaw  
Institute of Applied Mathematics and Mechanics  
Poland  
[P.biecek@mimuw.edu.pl](mailto:P.biecek@mimuw.edu.pl)