

4.4 Ordering Relations

Alejandro Ruiz

April 2021

Exercise 5

Proof. To prove that $R \cap (B \times B)$ is a partial order on B we need to prove that $R \cap (B \times B)$ is reflexive, transitive, and antisymmetric. To prove reflexivity, let $x \in B$. Since $B \subseteq A$, Then $x \in A$. Since R is a partial order on A , then R is reflexive on A . Thus, xRx . Since $x \in B$, then $(x, x) \in (B \times B)$. Since x was an arbitrary element of B , then $R \cap (B \times B)$ is reflexive on B .

To prove transitivity, let $x, y, z \in B$, and suppose that $(x, y) \in R \cap (B \times B)$ and $(y, z) \in R \cap (B \times B)$. Then, $(x, y) \in R$ and $(y, z) \in R$. Since R is a partial order on A , then R is transitive on A . Thus, since xRy and yRz , then xRz . Since $x, z \in B$, then $(x, z) \in B \times B$. Hence, $(x, z) \in R \cap (B \times B)$ and so $R \cap (B \times B)$ is transitive on B .

To prove antisymmetry, let $x, y \in B$, and suppose that $(x, y) \in R \cap (B \times B)$ and $(y, x) \in R \cap (B \times B)$. Then, $(x, y) \in R$ and $(y, x) \in R$. Since R is a partial order on A , then R is antisymmetric on A , and given that $(x, y) \in R$ and $(y, x) \in R$, we conclude that $y = x$. Thus, $R \cap (B \times B)$ is antisymmetric on B . \square

Exercise 8

Proof. To prove that T is a partial order on $A \times B$ we need to prove that T is reflexive, transitive, and antisymmetric. To prove reflexivity, let $(x, y) \in A \times B$. Thus, $x \in A$ and $y \in B$. Since R is a partial order on A , then R is reflexive on A , and since $x \in A$, then xRx . Similarly, since S is a partial order on B , then S is reflexive on B , and since $y \in B$, then ySy . Since xRx and ySy , then $(x, y)T(x, y)$. Since (x, y) was an arbitrary element of $A \times B$, then T is reflexive on $A \times B$.

To prove that T is transitive, let $(x_1, y_1), (x_2, y_2), (x_3, y_3) \in A \times B$, and suppose that $(x_1, y_1)T(x_2, y_2)$ and $(x_2, y_2)T(x_3, y_3)$. Since $(x_1, y_1)T(x_2, y_2)$, then x_1Rx_2 and y_1Sy_2 . Similarly, since $(x_2, y_2)T(x_3, y_3)$, then x_2Rx_3 and y_2Sy_3 . Given that R is a partial order on A , then R is transitive on A , so x_1Rx_3 . Similarly, since S is a partial order on B , then S is transitive on B , so y_1Sy_3 . Since x_1Rx_3 and y_1Sy_3 , then $(x_1, y_1)T(x_3, y_3)$, since $(x_1, y_1), (x_2, y_2), (x_3, y_3)$ were arbitrary elements of $A \times B$, then T is transitive on $A \times B$.

To prove that T is antisymmetric, let $(x_1, y_1), (x_2, y_2) \in A \times B$, and suppose that $(x_1, y_1)T(x_2, y_2)$ and $(x_2, y_2)T(x_1, y_1)$. Thus, x_1Rx_2 and y_1Sy_2 , and x_2Rx_1 and y_2Sy_1 , respectively. Since R is a partial order on A , then R is antisymmetric so $x_1 = x_2$. Similarly, since S is a partial order on B , then S is antisymmetric so $y_1 = y_2$. Thus, $(x_1, y_1) = (x_2, y_2)$. Given that $(x_1, y_1), (x_2, y_2)$ were arbitrary elements of $A \times B$, then T is antisymmetric on $A \times B$. \square

To answer the second question, try proving it. Doing the cases that will appear, there will be a contradiction. Thus, T does not have to be a total order even if both R and S are.

Exercise 12

Proof. Suppose B has a minimal element B_1 . By definition, then $B_1 \in B$, so that $B_1 \neq \emptyset$ and $\forall x \in \mathbb{R} \forall y \in \mathbb{R} [(x \in B_1 \wedge x < y) \rightarrow y \in B_1]$. Since $B_1 \neq \emptyset$, then there is some element $b \in B_1$. Since $\forall x \in \mathbb{R} \forall y \in \mathbb{R} [(x \in B_1 \wedge x < y) \rightarrow y \in B_1]$ and $b \in B_1$, then $\forall y \in \mathbb{R} [(b \in B_1 \wedge b < y) \rightarrow y \in B_1]$. Let

$Y = \{y \in \mathbb{R} | y > b\}$, so $Y \subseteq B_1$. Note that since $b \in B_1$ and $b \notin Y$, then $B_1 \neq Y$. Given that $Y \subset B_1$ and $B_1 \subseteq B$, then $Y \subseteq B$. Thus, there is some set $Y \in B$ so that $Y \subseteq B_1$, but $Y \neq B_1$, which is a contradiction of our definition of the S-minimal element of B . Thus, B has no minimal element. \square

Exercise 14

Proof. (\rightarrow). Suppose that b is the R -largest element of B and let $x \in B$ be an arbitrary element. It follows that xRb , so it is true that $(b, x) \in R^{-1}$.

(\leftarrow). Suppose that b is the R^{-1} -smallest element of B and let $x \in B$ be an arbitrary element. It follows that $bR^{-1}x$, so it is true that $(x, b) \in R$. \square

Proof. (\rightarrow). Suppose b is the R -maximal element of B . Also, let $x \in B$ and suppose that $xR^{-1}b$, so it follows that bRx . Since $x \in B$ and bRx , and given that b is the R -maximal element of B , it follows that $x = b$.

(\leftarrow). Suppose that b is an R^{-1} -minimal element of B . Furthermore, let $x \in B$ and suppose that bRx , so it is true that $xR^{-1}b$. Since $x \in B$ and $xR^{-1}b$, and given that b is an R^{-1} -minimal element of B , it follows that $x = b$. \square

Exercise 16

Proof. Suppose that b is the R -largest element of B . First, we prove that b is also the R -maximal element of B . Let $x \in B$ and suppose that bRx . Thus, since b is the R -largest element of B , it follows that xRb . Since $b, x \in B$ and $B \subseteq A$, then $b, x \in A$. Since R is a partial order, then it is antisymmetric, so we conclude that $x = b$. Thus, b is a largest element of B . To prove that b is the only R -maximal element of B , suppose c is an R -maximal element of B , so $c \in B$. Thus, since $c \in B$ and b is an R -maximal element of B , it follows that cRb . Since $b \in B$ and cRb , and given that c is R -maximal, it follows that $c = b$. Thus, b is the only maximal element of B . \square

Exercise 17

If I have any mistakes, please let me know. I am solving it in the way I understand the question that is being posed. We can try proving this statement, so let's you see how our scratch work would look like. The goal would be that c is R -smallest element of C , which we know that it is represented as $\forall x \in C cRx$. Our list of givens would be the following:

- R is a partial order on A
- $B \subseteq A$
- $C \subseteq R : \forall c_1 \in C \forall c_2 \in C [(c_1, c_2) \in C \rightarrow (c_1, c_2) \in R]$
- $C \subseteq A$
- c is R -minimal element of $C : \forall x \in C (xRc \rightarrow x = c)$ and $c \in C$
- $x \in C$, where x is arbitrary (this comes from the logical form of the goal we are trying to prove)

Based on this list, we can get to the following:

1. $x, c \in C$. Since $C \subseteq A$, then $x, c \in A$
2. Since $C \subseteq R$, then xRc or cRx
 - It is true that xRc . However, since c is R -minimal, then $x = c$. Thus, $(x, x) = (c, c) = (x, c) \in R$ and we would be stuck

Thus, our reasoning tells us that the statement is not necessarily true

Exercise 18

Proof. (\rightarrow). Let $x \in A$, and suppose that x is an upper bound of B_1 . Also, suppose that $z \in B_2$. Since $\forall x \in B_2 \exists y \in B_1 (xRy)$ and $z \in B_2$, then there is some element $y_1 \in B_1$ so that zRy_1 . Since $B_2 \subseteq A$ and $z \in B_2$, then $z \in A$. Similarly, since $B_1 \subseteq A$ and $y_1 \in B_1$, then $y_1 \in A$. Given that R is a partial order, R is transitive. Hence, since zRy_1 and y_1Rx , it follows that zRx . Since z was an arbitrary element of B_2 , if x is an upper bound of B_1 , then x is an upper bound of B_2 .

(\leftarrow). Let $x \in A$, and suppose that x is an upper bound of B_2 . Also, suppose that $x \in A$. Since $\forall x \in B_1 \exists y \in B_2 (xRy)$ and $z \in B_1$, then there is some element $y_2 \in B_2$ so that zRy_2 . Given that x is an upper bound of B_2 and $y_2 \in B_2$, then y_2Rx . Note that $B_1 \subseteq A$ and $B_2 \subseteq A$, so $z, y_2 \in A$. Since R is a partial order, it is transitive, and given that zRy_2 and y_2Rx , it follows that zRx . Given that z was an arbitrary element of B_1 , if x is an upper bound of B_2 , then x is an upper bound of B_1 . □

To prove (b), we can actually use the contrapositive or contradiction strategies, since they share something in common

Proof. Suppose that $B_1 \cap B_2 = \emptyset$. We will prove by contradiction, so suppose that either B_1 or B_2 has a maximal element.

Case 1: B_1 has a maximal element. Let $b_1 \in B_1$ be the maximal element of B_1 . Since $\forall x \in B_1 \exists m \in B_2 (xRm)$ and $b_1 \in B_1$, then there is some element $m_1 \in B_2$ and b_1Rm_1 . Since $\forall x \in B_2 \exists n \in B_1 (xRn)$ and $m_1 \in B_2$, there is some $n_1 \in B_1$ and m_1Rn_1 . Since $B_1 \subseteq A$ and $B_2 \subseteq A$, and $b_1, n_1 \in B_1$ and $m_1 \in B_2$, then $b_1, m_1, n_1 \in A$. Since R is a partial order, then it is transitive and since b_1Rm_1 and m_1Rn_1 , then b_1Rn_1 . Since b_1 is a maximal element and $n_1 \in B_1$, then $n_1 = b_1$, so $m_1Rn_1 = m_1Rb_1$. Given that m_1Rb_1 and b_1Rm_1 , and since R is antisymmetric on A , then $m_1 = b_1$. Since $B_1 \cap B_2 = \emptyset$ and $b_1 \in B_1$, then $b_1 \notin B_2$. But $m_1 = b_1$ and $m_1 \in B_2$, so this is a contradiction. Thus, neither B_1 nor B_2 has a maximal element.

Case 2: The logic is similar to Case 1. □

Exercise 20

Proof. Suppose b is the smallest element of B . This implies that since $B \subseteq A$ and $b \in B$, then $b \in A$, and so b is a lower bound of B . Let L be the set of all lower bounds of B , so we can conclude that $L \neq \emptyset$. Now, let $l \in L$ be an arbitrary element. In particular, we know that $L \subseteq A$. We can think of cases:

Case 1: $L = \{b\}$. Since $b \in B$, and $b \in A$, and since R is reflexive on A , then bRb and b is the g.l.b.

Case 2: L has more than one element. All other elements of L must only be in A and be smaller than b , otherwise there would be no smallest element in B . Thus, b is the g.l.b. □

The proof for part (b) follows a similar reasoning.

Exercise 21

Proof. Suppose $x \in U$ and xRy . We need to prove that $y \in U$, so let $z \in B$ be an arbitrary element. Since $x \in U$, then $\forall m \in B mRx$ and $x \in A$. Using the latter and since $z \in B$, then zRx . Since $B \subseteq A$ and $z \in B$, then $z \in A$. Given that R is a partial order, then it is transitive, so since zRx and xRy , it follows that zRy . Since z was an arbitrary element of B , we conclude that $y \in U$. □

Proof. Let $x \in B$ an arbitrary element. We now have to prove that x is a lower bound for U . Then, suppose that $u \in U$ so $\forall m \in B mRu$. Since $x \in B$, then xRu . Given that u was an arbitrary element of U , then x is a lower bound for U . Since x was an arbitrary element of B , then $\forall x \in B (x \text{ is a lower bound for } U)$. □

Part (c) is kind of tricky. We have to be able to distinguish that if x is a l.u.b. of B , we are saying that x is the smallest element of U , not B !

Proof. Suppose that x is g.l.b. of U . Suppose $u_2 \in U$, and so we have to prove that xR_2 and $x \in U$. Suppose $b \in B$, so our new goals are xRu_2 , bRx , and $x \in A$. Let L_U be the set of l.b. of U . Since x is g.l.b. of U , then it is the largest element of L_U . Thus, $\forall m_1 \in U(xRm_1)$ and $x \in A$. Given that $\forall m_1 \in U(xRm_1)$ and $u_2 \in U$, then xRu_2 . Also, since $u_2 \in U$, it follows that $\forall b \in B(bRu_2)$ and $u_2 \in A$. Since $b \in B$ and $\forall b \in B(bRu_2)$, then bRu_2 . Given that $x, u_2 \in A$, then u_2Rx . Since R is a partial order, then it is transitive on A . Given that bRu_2 and u_2Rx , then bRx . Since u_2 was an arbitrary element of U , then $x \in U$. Since u_2 was an arbitrary element of U , then x is a l.u.b. of B . □

Exercise 23

This proof is quite tricky. We really need to know our definitions. I will first prove the l.u.b. and then the g.u.b.

Proof. Suppose $\mathcal{F} \subseteq \mathcal{P}(A)$ and $\mathcal{F} \neq \emptyset$. Let $M \in Up$, where Up is the set of all upper bounds of \mathcal{F} . Suppose that $x \in \cup \mathcal{F}$. Thus, there is some set $N \in \mathcal{F}$ so that $x \in N$. Since $M \in Up$, this means that M is an upper bound for \mathcal{F} , and given that $N \in \mathcal{F}$, it follows that $N \subseteq M$. We know that $x \in N$, so $x \in M$. Since x was an arbitrary element of $\cup \mathcal{F}$, then $\cup \mathcal{F} \subseteq M$. We would like to prove that $\cup \mathcal{F} \in Up$. Suppose that $T \in \mathcal{F}$ and let $y \in T$. Also, suppose that $z \in \cup \mathcal{F}$. Thus, there is some set $N \in \mathcal{F}$ so that $z \in N$. Since $\mathcal{F} \subseteq \mathcal{P}(A)$, and $N \in \mathcal{F}$, it follows that $N \subseteq A$. Since $z \in N$, then $z \in A$. Also, since $T \in \mathcal{F}$ and $y \in T$, then $y \in \cup \mathcal{F}$. Given that z was an arbitrary element of $\cup \mathcal{F}$, then $\cup \mathcal{F} \subseteq A$. Also, given that y was an arbitrary element of T , then $T \subseteq \cup \mathcal{F}$. Furthermore, since T was an arbitrary element of \mathcal{F} , then $\forall T \in \mathcal{F}(T \subseteq \cup \mathcal{F})$. And given that $\cup \mathcal{F} \subseteq A$, which means that $\cup \mathcal{F} \in \mathcal{P}(A)$, we conclude that $\cup \mathcal{F}$ is an upper bound. Hence, $\cup \mathcal{F}$ is the l.u.b. for \mathcal{F} . □

Proof. Suppose $\mathcal{F} \subseteq \mathcal{P}(A)$ and $\mathcal{F} \neq \emptyset$. We want to prove that $\cap \mathcal{F}$ is a the g.l.b. of \mathcal{F} . Thus, two conditions must be satisfied: 1) $\forall M \in L(M \subseteq \cap \mathcal{F} \text{ and } \cap \mathcal{F} \in L$, where L is the set of lower bounds for \mathcal{F} . To prove the second condition, $F \in \mathcal{F}$ and suppose that $x \in \cap \mathcal{F}$. Thus, since $F \in \mathcal{F}$ and $x \in \cap \mathcal{F}$, then $x \in F$. Since $F \subseteq \mathcal{P}(A)$ and $x \in F$, then $x \in A$. This also shows that $\cap \mathcal{F} \subseteq A$. To prove the second condition, suppose $M \in L$ and let $x \in M$. Also, suppose that $F \in \mathcal{F}$. Thus, since $M \in L$ means that M is a lower bound for \mathcal{F} , then $M \subseteq F$. Since $x \in M$, then $x \in F$. Given that x was an arbitrary element of M , then $M \subseteq \cap \mathcal{F}$. Since F was an arbitrary element of \mathcal{F} , then $\forall M \in L(M \subseteq \cap \mathcal{F})$. Thus, we conclude that $\cap \mathcal{F}$ is the g.l.b. of \mathcal{F} . □