

Assignment 3: Mutual Activation

Two proteins A and B mutually activate their own expression. Their dynamics is described by the following coupled differential equations:


$$\frac{dA}{dt} = \frac{\beta B^n}{k_b^n + B^n} - \delta A, \quad \frac{dB}{dt} = \frac{\beta A^n}{k_a^n + A^n} - \delta B$$

where β is the maximum expression and δ is the decay rate (assumed to be the same for the two proteins), and k_a and k_b are the activation thresholds.

- Assuming that $k_a = k_b = k$ and $n \gg 1$, represent graphically the nullclines of the system in the plane (A, B), for $\beta / \delta > k$. Identify the equilibrium points.
- Determine graphically the stability of all the equilibria for the case of section (a), drawing the slope field of the system in the different regions in which the nullclines divide the phase space. What is the biological meaning of this behaviour?
- Assume now that k_a and k_b are different. Fixing k_a , show (also graphically) that if k_b increases a bifurcation occurs. Draw the bifurcation diagram approximately. What is the type of this bifurcation?


My solution:

- The nullclines for $\frac{dA}{dt}$ and $\frac{dB}{dt}$ for the given parameter values are two identical *activation curves*, symmetric about the line $y = x$, representing the points in which the two coupled differential equations are each equal to zero. The equilibrium points are indicated by their three points of intersection: (0, 0), (k, k), (β/δ , β/δ).




- b) In this system there are *two stable equilibria* and *one unstable equilibrium*, displayed by the two full dots and the empty dot respectively. This is clearly shown by the slope field in the diagram: the arrows cross the dA/dt curve vertically, while pointing towards the dB/dt curve, and vice versa. So, the overall slope field points towards the two stable equilibria of the system, while the unstable equilibrium is at the centre of a *separatrix* which divides the phase plane in two basins of attraction.

This means that when the concentration of A and/or B is not sufficiently high (below the separatrix), the degradation term will prevail over the activation loop, the concentrations of our two proteins will both go to zero and the two genes will be in their **OFF state**; on the other hand, when A and/or B are high enough, on the other side of the separatrix in the plane, the activation loop will show its effect, bringing the concentrations of the two proteins to a high steady-state value: the genes are now in their **ON state**.



- c) When k_a is fixed and k_b is increasing, for the dB/dt nullcline we have that *the step region of the curve is shifting to the right*. The concentration of A at which the unstable equilibrium exists increases and increases with the increase in k_a , until it makes contact with the higher fix point; any further increase in k_a will result in the disappearance of both the unstable equilibrium and the higher stable equilibrium: a **saddle node bifurcation** occurs and only the stable equilibrium at zero is remaining, so the genes will be in their OFF states for such high values of k_a .



Here's my initial draft for this assignment.

