

# Probability & Statistics: Assignment 7

Due on 20 December, 2018 at 8:30am

*Prof. D. Eynard*

**A. Romanelli / A. Vicini**

## Problem 1

1. We know from the nature of politics that the distribution that we are dealing with will be a binomial one. As seen in class we can get an estimation of  $\hat{p}$  by computing:

$$\hat{p} = \left( \sum_{i=1}^n x_i \right) \cdot \frac{1}{n}$$

Which yields the following proportion for candidate A:

$$\frac{185}{351} \approx 0.527 = 52.7\% \text{ of the voters}$$

Against candidate B:

$$\frac{351 - 185}{351} \approx 0.473 = 47.3\% \text{ of the voters}$$

We can see that candidate A will be slightly favoured with a 2.7% margin of vantage.

2. We want to find a 95% confidence interval for the  $\hat{p}$  that we just computed:

$$se = \sqrt{\hat{p}(1 - \hat{p})/n} = 0.027$$

We can thus attempt to find an interval so that our confidence results:

$$P(\hat{p} - \delta < \hat{P} < \hat{p} + \delta) = 0.95$$

The estimation of the proportion can be treated as a random variable binomially distributed, since each sample would either satisfy our confidence interval or not. We can then treat this distribution as a normal one according to the **Central Limit Theorem**:

$$\begin{aligned} P(\hat{p} - \delta < \hat{P} < \hat{p} + \delta) &= P\left(-\frac{\delta}{se} < \frac{\hat{P} - \hat{p}}{se} < \frac{\delta}{se}\right) \\ &= 2 \cdot \Phi\left(\frac{\delta}{se}\right) - 1 \end{aligned}$$

We now need to find  $2\Phi\left(\frac{\delta}{se}\right) - 1 = 0.95$ , thus:

$$\begin{aligned} \Phi\left(\frac{\delta}{se}\right) &= \frac{1 + 0.95}{2} \\ \frac{\delta}{se} &= \Phi^{-1}\left(\frac{1 + 0.95}{2}\right) \\ \delta &= 1.96 \cdot se = 0.053 \end{aligned}$$

We have now found the bounds of our interval to be:  $[\hat{p} - \delta, \hat{p} + \delta] = [0.474, 0.58]$

Since our  $\hat{p}$  lies within this interval we can be pretty confident that the estimated proportion is favourable to A.

3. We want to have a margin of error  $\delta = 0.02$ , thus we'd need to increase our sample population  $n$  such that:

$$\begin{aligned} \delta &= 1.96 \cdot se = 0.02 \\ se &= \frac{0.02}{1.96} \\ \sqrt{\hat{p}(1 - \hat{p})/n} &= 0.0102 \\ \frac{0.249271}{0.00010404} &= n \\ n &= 2396 \end{aligned}$$

## Problem 2

1. Given the following data: (13.0, 18.5, 16.4, 14.8, 20.8, 19.3, 18.8, 23.1, 25.1, 16.8, 20.4, 17.4, 16.0, 21.7, 15.2, 21.3, 19.4, 17.3, 23.2, 24.9, 15.2, 19.9, 19.1, 18.1, 25.2, 23.1, 15.3, 19.4, 21.5, 16.8, 15.6, 17.6) we computed the sample mean  $\bar{x}$  and the standard deviation  $\sigma$  with the following code:

```
samples <- c(13.0, 18.5, ..., 15.6, 17.6)
mu <- mean(samples)
print(mu)
sigma <- sqrt(var(samples))
print(sigma)
```

Which gives us  $\bar{x} = 19.068, \sigma = 3.255$

2. Under the assumption of an underlying normal distribution and variance  $\sigma^2 = 3.12$ , we can attempt to find a 95% confidence interval in the following way:

$$[\bar{x} - \delta, \bar{x} + \delta]$$

$$\delta = z^* \frac{\sigma}{\sqrt{n}}$$

Where the z value for our 95% confidence interval will be  $z^* = 1.96$ :

$$\delta_{0.95} = 1.96 \cdot \frac{\sqrt{3.12}}{\sqrt{32}} = 1.96 \cdot \frac{1.766}{5.657} = 0.612$$

Thus we get the resulting :

$$CI_{0.95} = [\bar{x} - \delta_{0.95}, \bar{x} + \delta_{0.95}] = [19.068 - 0.612, 19.068 + 0.612] = [18.46, 19.68]$$

3. In case of a 99% confidence interval our  $z^*$  would be 2.58 instead, which would result in the following margin of error:

$$\delta_{0.99} = 2.58 \cdot \frac{\sqrt{3.12}}{\sqrt{32}} = 0.806$$

And it's clearly visible that  $\delta_{0.95} < \delta_{0.99}$  and thus we'd have a larger confidence interval than before. This comes from the fact that a wider interval grants us more confidence that our sample lies within the interval than not.

Increasing the sample size  $n$  will reduce the margin of error and also tighten the interval.

### Problem 3

- a) Under the assumption of a biased coin with  $\hat{p} = 0.6$ , we can compute the probability as the one of a binomial distribution:

$$P(S_n = n \cdot \hat{p}) = \binom{n}{n \cdot \hat{p}} \hat{p}^{n\hat{p}} (1 - \hat{p})^{n(1-\hat{p})}$$

$$P\left(S_n = \frac{3}{5}n\right) = \binom{n}{\frac{3}{5}n} \left(\frac{3}{5}\right)^{\frac{3}{5}n} \left(\frac{2}{5}\right)^{\frac{2}{5}n}$$

For  $n = 10$ :

$$P(S_{10} = 6) = \binom{10}{6} \left(\frac{3}{5}\right)^6 \left(\frac{2}{5}\right)^4 \approx 210 \cdot 0.0466 \cdot 0.0256 = 0.25$$

For  $n = 100$ :

$$P(S_{100} = 60) = \binom{100}{60} \left(\frac{3}{5}\right)^{60} \left(\frac{2}{5}\right)^{40} \approx 0.081$$

For  $n = 1000$ :

$$P(S_{1000} = 600) = \binom{1000}{600} \left(\frac{3}{5}\right)^{600} \left(\frac{2}{5}\right)^{400} \approx 0.026$$

## Problem 4

## Problem 5

To accept or reject  $H_0$  we need to compute the z-value:

$$z = \frac{x - \mu}{\sigma/\sqrt{n}} \Rightarrow z = \frac{113.5 - 110}{10/4} = 1.4$$

If the resulting  $z$  is greater than what we just computed, then we'll accept  $H_0$ , else we reject it.

- a) A 5% significance level corresponds to  $\alpha = 0.05$ :  $z_{5\%} = \Phi^{-1}(0.95) \stackrel{?}{>} 1.4$

$$z_{5\%} = 1.65 > 1.4, \text{ thus we accept } H_0$$

- b) Given  $\alpha = 0.10$ :  $z_{10\%} = \Phi^{-1}(0.90) \stackrel{?}{>} 1.4$

$$z_{10\%} = 1.28 < 1.4, \text{ thus we reject } H_0 \text{ and accept } H_1 \text{ instead}$$

- c) The p-value corresponds to the right-tail-end probability of the null hypothesis  $H_0$ :

$$p = 1 - \Phi(z) = 1 - \Phi(1.4) \approx 1 - 0.919 = 0.081$$

## Problem 6