



POLITECNICO
MILANO 1863

SCUOLA DI INGEGNERIA INDUSTRIALE
E DELL'INFORMAZIONE

Title

TESI DI LAUREA MAGISTRALE IN
XXXXXXX ENGINEERING - INGEGNERIA XXXXXXX

Author: **Name Surname**

Student ID: 000000

Advisor: Prof. Name Surname

Co-advisors: Name Surname, Name Surname

Academic Year: 20XX-XX

Abstract

Here goes the Abstract in English of your thesis followed by a list of keywords. The Abstract is a concise summary of the content of the thesis (single page of text) and a guide to the most important contributions included in your thesis. The Abstract is the very last thing you write. It should be a self-contained text and should be clear to someone who hasn't (yet) read the whole manuscript. The Abstract should contain the answers to the main scientific questions that have been addressed in your thesis. It needs to summarize the adopted motivations and the adopted methodological approach as well as the findings of your work and their relevance and impact. The Abstract is the part appearing in the record of your thesis inside POLITesi, the Digital Archive of PhD and Master Theses (Laurea Magistrale) of Politecnico di Milano. The Abstract will be followed by a list of four to six keywords. Keywords are a tool to help indexers and search engines to find relevant documents. To be relevant and effective, keywords must be chosen carefully. They should represent the content of your work and be specific to your field or sub-field. Keywords may be a single word or two to four words.

Keywords: here, the keywords, of your thesis

Abstract in lingua italiana

Qui va l'Abstract in lingua italiana della tesi seguito dalla lista di parole chiave.

Parole chiave: qui, vanno, le parole chiave, della tesi

Contents

Abstract	i
Abstract in lingua italiana	iii
Contents	v
Introduction	1
1 Nuclear structure and deformations	3
1.1 Nuclear structure models	3
1.1.1 Phenomenology of the NN interaction	3
1.1.2 Nuclear models	5
1.1.3 Liquid drop model	5
1.1.4 Shell corrections	6
1.2 Nuclear pairing	9
1.3 Nuclear deformations	9
1.3.1 Quadrupole deformation	9
1.3.2 Nilsson model	11
1.3.3 Octupole deformations and parity breaking	13
1.4 Nuclear fission	13
1.4.1 Octupole deformations	15
2 State of the art, objective and motivation	17
2.1 State of the art and motivation	17
2.2 Objectives	18
2.3 Methods	18
3 Energy functional	21
3.1 Hartree-Fock theory	21
3.1.1 Variational principle	21

3.1.2	Hartree-Fock equations	22
3.1.3	Symmetries in Hartree-Fock	24
3.1.4	Density Functional Theory	26
3.2	Pairing in Hartree-Fock theory	27
3.2.1	BCS theory	28
3.2.2	Hartree-Fock-Bogoliubov theory	30
3.3	Skyrme	34
3.3.1	Skyrme force	34
3.3.2	Energy density functional	36
3.3.3	Functionals	40
3.4	Coulomb interaction	40
3.5	Energy calculation	41
4	Numerical methods	45
4.1	Finite differences	45
4.1.1	3D mesh	45
4.1.2	Schrödinger equation	46
4.1.3	Poisson equation	48
4.2	Eigenvalue problem	49
4.2.1	Conjugate Gradient and numerical techniques	50
4.2.2	Iterative eigensolvers	54
4.2.3	General Conjugate Gradient	57
4.3	Code implementation details	59
4.3.1	Constraints	59
4.3.2	Details on the implementation of the code	61
4.3.3	Optimal parameters choice	61
4.3.4	Numerical stability	63
5	Results for spherical nuclei	67
5.1	Physical quantities	67
5.1.1	Mean square radii	67
5.1.2	Deformation parameters	68
5.2	Parameters and mesh choice	69
5.3	Results for ^{16}O	69
5.3.1	Results neglecting Coulomb interaction	69
5.4	Results for heavier nuclei	70
5.4.1	Comparison with experimental binding energies	70

6	Results for deformed nuclei	77
6.1	^{24}Mg	77
6.1.1	HFBTH0 code and calculation details	77
	Bibliography	81
A	Appendix	89
A.1	Spherical harmonics	89
A.1.1	Algorithm	90
A.2	5-point derivatives	90
A.3	Functional derivatives	90
A.4	\bar{U} , \bar{V} matrix structure	92
	List of Figures	93
	List of Tables	95
	List of Symbols	97
	Acknowledgements	99

Introduction

The study of atomic nuclei forms the bridge between fundamental areas of theoretical and computational physics and nuclear engineering. While experimental data has provided over the years invaluable information about nuclear structure, reactions and other important nuclear processes, the predictions provided by a theoretical understanding although still unsatisfactory due to the extremely vast landscape of nuclear physics, are essential for nuclear engineering. In particular, fission reactions, which are of the utmost importance in nuclear engineering, are still poorly understood, models relying on mean-fields and empirical descriptions of the nucleus are too coarse to help us understand the exact physical process, let alone make numerically accurate predictions, which is a major challenge for the simulation of new generation nuclear reactors, which use neutron rich nuclei and fuel materials much less understood than the traditional thermal reactors. In this regard, the most successful approach to the microscopic description of nuclei, is certainly the exact many-body system, which starting from the interactions among nucleons, aims at building a complete description of the nucleus. At the moment, there are two competing frameworks that try to tackle the problem,

- i the *ab-initio* approach, where the interaction is in principle exact, derived from controlled approximations of quantum chromodynamics; and
- ii the use of effective interactions and nuclear Density Functional Theory.

Ab-initio methods, while technically speaking more rigorous, are still limited as of now, since they can only account for light, spherical nuclei. Energy density functionals and effective interactions, such as the Skyrme force, on the other hand are more flexible and less computationally expensive enabling a much wider representation of nuclei across the whole chart, including heavy nuclei, which are of crucial importance in nuclear engineering. D Vautherin and D M Brink laid the foundations of the nuclear Hartree-Fock theory using the Skyrme interaction in 1972, through spherically symmetric calculations, which are unable to account for nuclear deformations, essential for nuclei far from magic numbers and in the heavy region. Over the years, thanks to the increase in computational performance of modern hardware, codes that are able to represent more coordinates have

been written, mainly using basis expansions of the harmonic oscillator, which presents many downsides, such as the inability to represent superdeformed states or nuclei near drip lines. In the past twenty years, the use of meshes to better account for such extremal cases has been introduced, still assuming certain simplifications, such as plane reflections, cylindrical symmetries and so on. The use of fully unconstrained Hartree-Fock methods, of critical importance for exotic deformations, is still a novel endeavour that only a handful of implementations have tackled, due to the high computational cost.

The aim of this work is to explore a new computational approach, the General Conjugate Gradient method, to efficiently solve spatially unconstrained Skyrme functionals. This thesis is organised as follows:

- In chapter 1, a short, comprehensive introduction to nuclear physics is given, as to prime the reader on the essential physical properties of atomic nuclei, starting from phenomenological facts and empirical models. A formal description of nuclear deformations and fission is also given, to highlight the importance of symmetry breaking.
- In chapter 2, a short summary of the methods used in this thesis is given, both theoretical and numerical, as well as a state-of-the-art comparison with other codes.
- In chapter 3, the theoretical framework used in the present work is reviewed, by introducing aspects of Hartree-Fock theory, Density Functional Theory and the effective interaction used in this work.
- In chapter 4, the numerical methods used in this work are presented, along with actual implementations of them in writing the code.
- In chapter 5, results for the spherically symmetric case are presented as a way of benchmarking the new implementation of this thesis, along with a description of the main physical quantities we compare.
- In chapter 6, benchmarks for the deformed nucleus ^{24}Mg are shown, after which novel results regarding... are presented.

1 | Nuclear structure and deformations

In this chapter, a concise introduction to nuclear physics is provided, as a way to understand the essential physical properties of the system under study. First, in section 1.1, we will review the main empirical facts about nuclides, such as particle density distribution and binding energies and the simple phenomenological models historically employed to describe them. Moving on to more advanced topics, that are able to complete the general description of nuclear structure, which are nuclear pairing in section 1.2 and nuclear deformations in section 1.3.

Lastly, in section 1.4, we will overview the nuclear fission process, by deriving a simple model to describe it and discussing the importance of exotic deformations to accurately describe it.

1.1. Nuclear structure models

The study of low energy hadron physics, has always been a challenging task. This is due to the known fact that the strong force, which is responsible for the attraction between the nucleons in a nuclei, is not perturbative at low energies, as opposed to the atomic case for the Coulomb interaction.

1.1.1. Phenomenology of the NN interaction

It is possible to obtain a good insight on the nuclear structure, by using empirical data obtained experimentally on the more macroscopic properties of nuclei, such as the binding energy and the particle density.

Binding energies

Let us start by the omnipresent physical quantity that is the binding energy of a nucleus. We can define it as the mass defect of the nucleus with respect to the constituents –

protons and neutrons – isolated from each other. If Z is the number of protons, N the number of neutrons, and $A = N + Z$ the nuclear mass, then the binding energy E_B is given by

$$E_B = (Zm_p + Nm_n - M)c^2 \quad (1.1)$$

where m_p is the proton mass, m_n the neutron mass, and M the nucleus mass.

In figure 1.1, the binding energy per nucleon E_B/A of nuclei as a function of A is plotted. As shown in the figure, the binding energy per nucleon rapidly saturates and stalls around 7 MeV just after $A = 4$, this striking behaviour is due to nucleons interacting only with near neighbours, since the strong force is a short-range interaction, otherwise, the trend would follow a behaviour of $A(A - 1)$ as in the Coulomb case.




Figure 1.1: Binding energy per nucleon as a function of A . Due to the short range of the strong force, this value saturates around 7 MeV, with a steady, dim decrease after ^{56}Fe .

Nuclear density

An important aspect of nuclear phenomenology which can be easily obtained through electron scattering experiments [27] is the nuclear density. It can be very well represented by a Fermi-like distribution, which reads

$$\rho(r) = \frac{\rho_0}{1 + e^{\frac{r-R_0}{a}}}, \quad (1.2)$$

where R_0 is the nuclear radius, which can be parametrized as $R_0 \approx 1.2A^{1/3}$, and a is the diffusivity, whose value determines how sharp the density drops from its saturation value $\approx \rho_0$ to ≈ 0 . The saturation density ρ_0 is generally universal for all nuclei, amounting to $\approx 0.16 \text{ fm}^{-3}$.

1.1.2. Nuclear models

The formal description of nuclear structure has been proven to be a difficult task over the years. Due to the extremely rich phenomenology of nuclei and the challenges brought by the strong force, as we shall see, many models and further approximations to give a satisfactory description of all nuclides have been proposed.

1.1.3. Liquid drop model

One, if not the first successful model, is the liquid drop model. It is based on the assumption that the nucleus behaves as a liquid droplet, where forces among constituents tend to saturate. This hypothesis, formulated by G. Gamow, culminated in the formalization of the semi-empirical mass formula (SEMF) by N. Bohr and C. F. von Weizsäcker in 1935 [62], which reads

$$E_B = a_V A - a_S A^{2/3} - a_C \frac{Z(Z-1)}{A^{1/3}} - a_A \frac{(N-Z)^2}{A} + \delta_P \quad (1.3)$$

where E_B is the binding energy of the nucleus. Each term has a different physical meaning:

- $a_V A$ is the volume energy of the nucleus, given by the approximately constant binding energy per nucleon, which makes the total energy roughly proportional to A ;
- $a_S A^{2/3}$ is the surface energy, a correction to the volume energy due to outer nucleons – on the surface – interacting with fewer nucleons than those in the inner bulk;
- $a_C Z(Z-1)/A^{1/3}$ is the approximation to the Coulomb energy repulsion of the nucleus, assuming the protons are uniformly distributed;
- $a_A (N-Z)^2/A$ is the asymmetry energy, which is due to the Pauli exclusion principle, since protons and neutrons occupy their respective states, a high imbalance of one species or the other implies loosely bound nucleons, thus a higher energy contribution of those states; and
- δ_P refers to the pairing energy of the nucleus, whose parametrization and physical significance will be later discussed in section 1.2.

The SEMF can be fitted on current data to get a good estimate of binding energies [4], but it still lacks the ability of describing many aspects of nuclear structure, mainly, the nuclear shell structure, which can account for magic numbers and nuclear deformations.




Figure 1.2: Visual representation of the liquid drop model from [23]

1.1.4. Shell corrections

Describing the nucleus through a shell model would account for the quantum mechanical nature of the system, unfortunately, unlike the ‘atomic’ case, we don’t have a source of the field to which nucleons are subjected to, since it’s generated by the nucleons themselves; nonetheless, the formulation of an empirical potential which reproduces experimental data has been proven to be successful in providing useful corrections to the liquid drop model. The so called Woods-Saxon potential is an empirical field used for modelling the average field to which an independent nucleon would feel in a nucleus. It can take different parametrizations depending on the data that one wants to reproduce. It is formulated as to follow the shape of the nuclear density (1.2), and it reads

$$U(\mathbf{r}) = -\frac{U_0(A, N)}{1 + e^{\frac{r-R}{a}}} \quad (1.4)$$

where U_0 is the potential depth

$$U_0(A, N) = U_0 \left(1 \pm \kappa \frac{2N - A}{A} \right), \quad (1.5)$$

the $+$ and $-$ signs refer to protons and neutrons respectively. R refers to the radius of the nuclear surface, generally parametrized as

$$R = r_0 A^{1/3} \quad (1.6)$$

and a is the surface diffuseness, as in the density expression (1.2).

Spin-orbit coupling The success of the shell model is mainly due to the possibility of accounting for spin-orbit coupling, which is included through a term that reads

$$U_{\text{LS}}(\mathbf{r}) = U_0^{\text{LS}} \left(\frac{r_0}{\hbar} \right)^2 \frac{1}{r} \frac{d}{dr} \left(\frac{1}{1 + e^{\frac{r-R}{a}}} \right). \quad (1.7)$$

Coulomb interaction In the spherical case, the coulomb interaction can be taken as the energy potential produced by a sphere of charge Z and radius R , which reads

$$U_C(r) = Ze^2 \begin{cases} \frac{3-(r/R)^2}{2R} & r \leq R, \\ \frac{1}{r} & r > R. \end{cases} \quad (1.8)$$

The complete Hamiltonian then reads

$$\hat{H} = \hat{T} + U + U_{\text{LS}} + U_C, \quad (1.9)$$

where U_C is present only when solving for the proton shells. The solution to the eigenvalue problem $\hat{H}\psi = E\psi$ is of the form

$$\psi_{nljm_j} = \frac{u_{nl}(r)}{r} [Y_l(\hat{\mathbf{r}}) \otimes \chi_{1/2}]_{jm_j} \quad (1.10)$$

where $Y_{nl}(\hat{\mathbf{r}})$ is the spherical harmonic function of degree l and order m , the $\hat{\mathbf{r}}$ is used to denote dependence on the azimuthal and polar angles of \mathbf{r} and \otimes takes the meaning of the angular momentum coupling and $u_{nl}(r)$ satisfies the reduced Schrödinger equation

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dr^2} + \frac{\hbar l(l+1)}{2mr^2} + U(r) \right) \psi_{nl} = E \psi_{nl}. \quad (1.11)$$

The effect of the spin-orbit coupling U_{LS} and the Coulomb repulsion U_C can be accounted for by using first order perturbation theory.

Harmonic oscillator

A small digression on the harmonic oscillator is in order. The solution of the spherical potential

$$U_{\text{HO}}(\mathbf{r}) = \frac{1}{2} m \omega^2 r^2, \quad (1.12)$$

produces the spherical harmonic oscillator basis, which is very similar to the basis one would get solving for the Woods-Saxon potential, provided that ω is taken as $41/A^{1/3}$ MeV. As a matter of fact, the harmonic oscillator basis is often used to perform calculations

in nuclear physics. We will see in section 4.3 that a harmonic oscillator basis is used as starting guess for the numerical solution of a Woods-Saxon potential.

Shell structure

A graphical representation of the shells for a harmonic oscillator is shown in figure 1.3, where the contribution of the spin-orbit coupling is also accounted for; unlike the atomic case, shells whose total angular momentum is higher are lowered in energy, viceversa for lower total angular momentum.




Figure 1.3: Graphical representation of a harmonic oscillator shells, together with the spin-orbit coupling. Shells whose total angular momentum is higher are lowered in energy, viceversa for lower total angular momentum.

1.2. Nuclear pairing

In the semi-empirical mass formula (1.3), the δ_p term can be parametrised as

$$\delta_p = \begin{cases} +\delta_0 & \text{if N and Z are even,} \\ 0 & \text{if A is odd,} \\ -\delta_0 & \text{if N and Z are odd,} \end{cases} \quad (1.13)$$

hence having an even number of neutrons and/or protons increases the binding energy of the nucleus. A common choice for δ_0 is

$$\delta_0 = 12A^{1/2} \text{ MeV.}$$

This is a phenomena closely related to superconductivity, as nucleons of the same type form pairs that lie in higher energy states. An experimental evidence of this fact is known as odd-even staggering, where the separation energy

$$S_n = E_B(A, Z) - E_B(A - 1, Z), \quad (1.14)$$

is higher for even A , an increase that corresponds to the energy necessary to break a pair. We will see in section 3.2 the two main methods to account for pairing at a microscopic level.

1.3. Nuclear deformations

We shall now give a description of the nuclear shape in a formal framework. We will start by expanding the nuclear radius in terms of spherical harmonics, so that we can truncate and omit terms to describe certain configurations of the nucleus, so that we are able to illustrate the simple case of an axial quadrupole deformation. After that, we will briefly discuss the more general case of triaxial, octupole, and parity breaking configurations.

1.3.1. Quadrupole deformation

Assuming the nuclear volume to be evaluated as

$$V(A) = \frac{4}{3}\pi R^3 \quad (1.15)$$

Let us suppose to consider variations of the nuclear radius R in terms of spherical harmonics

$$R(\theta, \phi) = R_0 \left[1 + \sum_{\lambda\mu} \alpha_{\lambda\mu} Y_{\lambda\mu}(\theta, \phi) \right] \quad (1.16)$$

where the moments $\alpha_{\lambda\mu}$ defined as

$$\alpha_{\lambda\mu} = \int Y_{\lambda\mu}^*(\theta, \phi) R(\theta, \phi) d\Omega, \quad (1.17)$$

are considered small, in the sense that $|\alpha_{\lambda\mu}|^2 \ll |\alpha_{\lambda\mu}|$ as to conserve the volume in equation (1.15). We have that Y_{00} is constant, so its moment does not provide additional information to the constant radius. We can set $\alpha_{00} = 0$. Since Y_{10} , Y_{11} and Y_{1-1} are odd for $\theta + \pi$ and $\phi + \pi$, we have that $\alpha_{1\mu}$ vanishes in a reference frame in which the centre of mass is at the origin.

Now, let us consider only $\alpha_{2\mu}$ coefficients and neglect higher degree terms, so that the deformation is purely quadrupolar, then the radius reads

$$R(\theta, \phi) = R_0 \left[1 + \sum_{\mu=-2}^2 \alpha_{2\mu} Y_{2\mu}(\theta, \phi) \right]. \quad (1.18)$$

If we assume to be in the reference frame in which the inertia tensor, proportional to the coefficients $\alpha_{2\mu}$, is diagonal, which is known as intrinsic frame, then the sum

$$\alpha_{21} Y_{21}^* + \alpha_{2-1} Y_{2-1}^*$$

vanishes. Since R is a real valued function, we have the relation

$$\alpha_{\lambda\mu} Y_{\lambda\mu} + \alpha_{\lambda-\mu} Y_{\lambda-\mu} = 2 \operatorname{Re}\{\alpha_{\lambda\mu} Y_{\lambda\mu}\}, \quad (1.19)$$

as a consequence, the resulting expansion reads

$$\begin{aligned} R(\theta, \phi) &= R_0 \left[1 + a_{20} Y_{20} + 2 \operatorname{Re}\{a_{22} Y_{22}\} \right] \\ &= R_0 \left[1 + \sqrt{\frac{5}{16\pi}} \left(a_{20} (3 \cos^2 \theta - 1) + 2a_{22} \sqrt{3} \sin^2 \theta (\cos^2 \phi - \sin^2 \phi) \right) \right]. \end{aligned} \quad (1.20)$$

If we perform the substitution

$$a_{20} = \beta \cos(\gamma) \quad (1.21)$$

$$a_{22} = \beta \sin(\gamma) \quad (1.22)$$

and express the variation of R along the cartesian axes, we get

$$R_x - R_0 = \delta R_x = \sqrt{\frac{5}{4\pi}} \beta R_0 \cos \left(\gamma - \frac{2\pi}{3} \right), \quad (1.23)$$

$$R_y - R_0 = \delta R_y = \sqrt{\frac{5}{4\pi}} \beta R_0 \cos \left(\gamma + \frac{2\pi}{3} \right), \quad (1.24)$$

$$R_z - R_0 = \delta R_z = \sqrt{\frac{5}{4\pi}} \beta R_0 \cos \gamma. \quad (1.25)$$

Assuming the value of β to always be positive, in the case $\gamma = 0$, $\delta R_x = \delta R_y < \delta R_z$, meaning the nucleus is in a *prolate* configuration; while in the case of $\gamma = \pi/3$, $\delta R_x = \delta R_y > \delta R_z$, meaning the nucleus has an *oblate* shape. A general convention is to write β with a negative sign in the oblate case, and a positive sign in the prolate case.

By using trigonometric identities, it is trivial to show that unique shapes are found only for $\gamma \in [0; \pi/3]$, if γ takes a value different from 0 or $\pi/3$, the shape is said to be triaxial, meaning $\delta R_z \neq \delta R_x \neq \delta R_y$, the nucleus has no more rotational symmetries and is only symmetric for reflections along the (x, y) , (x, z) and (y, z) planes, which also induces parity symmetry.

1.3.2. Nilsson model

To understand the effect on single-particle motion of a deformed potential, we can consider the case of an axially deformed harmonic oscillator potential, for which $\omega_z \neq \omega_x = \omega_y = \omega_\perp$, meaning the oscillator frequency takes on a different value on the z axis than in the x and y axes.

To treat the deformation perturbatively, we can assume that the various frequencies deviate from the unperturbed $\omega_0 = 41/A^{1/3}$ MeV, in which case they may read

$$\omega_z = \omega_0 - \frac{2}{3}\varepsilon, \quad (1.26)$$

$$\omega_\perp = \omega_0 + \frac{1}{3}\varepsilon, \quad (1.27)$$

this definition of the frequencies satisfies the conservation of volume, at lowest order in ε , assumed to hold for

$$\omega_0^3 = \omega_z \omega_\perp^2. \quad (1.28)$$

We can thus write the single-particle Hamiltonian in the deformed potential as

$$H = H_0 + \varepsilon H_1, \quad (1.29)$$

$$H_0 = -\frac{\hbar^2}{2m} \nabla^2 + \frac{1}{2} m \omega_0^2 r^2, \quad (1.30)$$

$$\varepsilon H_1 = \frac{1}{3} \omega_0^2 \varepsilon (x^2 + y^2 - 2z^2) = -\frac{1}{3} \sqrt{\frac{16\pi}{5}} m \omega_0^2 \varepsilon r^2 Y_{20}. \quad (1.31)$$

H_0 is the usual spherical harmonic potential, for which the eigenfunctions, expressed through the usual quantum numbers $|nljm_j\rangle$ are known. Assuming ε to be small, we can evaluate the first order correction of H_1 to the system, which reads

$$\Delta E = \langle nljm_j | \varepsilon H_1 | nljm_j \rangle, \quad (1.32)$$

$$= -\frac{1}{3} \sqrt{\frac{16\pi}{5}} \varepsilon m \omega_0^2 \int r^2 u_{nl}(r) \langle jm_j | Y_{20} | jm_j \rangle dr, \quad (1.33)$$

$$= \frac{\varepsilon}{6} m \omega_0^2 \int r^2 u_{nl}(r) \frac{3m_j^2 - j(j+1)}{j(j+1)} dr, \quad (1.34)$$

thus in the limit of large j , states with the maximum total angular momentum projection m_j are shifted upwards, while states with the minimum m_j are shifted downwards; moreover, eigenstates with $\pm m_j$ are degenerate, as expected by the reflection symmetry of the Hamiltonian if the z axis is inverted. Adding further empirical terms to reproduce experimental data, and the spin-orbit coupling, results in the formulation of the Nilsson model [43]. In figure 1.4, a graphical representation of the energy levels in the Nilsson model is shown [63].

Deformed Woods-Saxon

Recent studies of deformed nuclei have been carried out using empirical potentials such as deformed Woods-Saxon potentials [20, 29]. In these models, the nuclear shape is expanded as

$$R(\theta) = R_0 \left[1 + \sum_{\lambda}^L \beta_{\lambda} Y_{\lambda 0} \right], \quad (1.35)$$

so that the solution is axially symmetric and the problem is reduced to just the (r, θ) coordinates, in which we can write the potential as

$$U_{\text{WS}}(r, \theta) = -\frac{U_0(A, N)}{1 + e^{\frac{r-R(\theta)}{a}}}. \quad (1.36)$$




Figure 1.4: Nilsson model energy levels trends, as a function of ε .

1.3.3. Octupole deformations and parity breaking

While quadrupole deformations concern nuclei across the whole chart, octupole deformations are much less common, being found only in heavier nuclei. Under the parity operation $\mathcal{P} : \mathbf{r} \mapsto -\mathbf{r}$, coefficients of the spherical harmonics transform as

$$\mathcal{P}\alpha_{\lambda\mu} = (-1)^\lambda \alpha_{\lambda\mu}, \quad (1.37)$$

hence a nuclear octupole deformation, whose degree $\lambda = 3$, would break parity symmetry. In figure 1.5 a graphical representation of the spherical harmonics for $\lambda = 3$ and $\mu = 0, 2$ is shown.

1.4. Nuclear fission

Nuclear fission is the process by which a nucleus splits into two – sometimes three – nuclei, whether spontaneously, or induced by a reaction. The physics that governs nuclear fission is the one of many-body, large amplitude collective modes, which elongate the the nuclear shape, until the so called *fission barrier* is surmounted and the path to the energy



Figure 1.5: Graphical representation of possible octupole deformations. On the left, the axially symmetric Y_{30} deformation, on the right, the non-axial octupole deformation Y_{32} .

minimum is one where the nuclei fragments itself.

Spontaneous fission model

It should be obvious that a formal treatment of deformations and collective modes is necessary to give a theoretical description of fission reactions. We can derive a simple spontaneous fission model by studying the effect of a simple axial quadrupole deformation on the semiempirical mass formula 1.3.

Let us assume that the nuclear radius may be expanded, as previously done in section 1.3, as

$$R = R_0[1 + \alpha_{20}Y_{20}]. \quad (1.38)$$

Assuming the nuclear volume is conserved across the fission path, the volume energy will not change. As for the surface energy, its variation can be expressed at the lowest order in α_{20} as

$$\Delta E_{\text{surf}} = E_{\text{surf}} - E_{0,\text{surf}} = E_{0,\text{surf}} \frac{2}{5} \alpha_{20}^2. \quad (1.39)$$

Regarding the Coulomb energy, the variation is given by

$$\Delta E_{\text{coul}} = E_{\text{coul}} - E_{0,\text{coul}} = -E_{0,\text{coul}} \frac{1}{5} \alpha_{20}. \quad (1.40)$$

Since the neutron and proton count does not change, the surface and Coulomb energies are the only contributions to the total energy difference. We can write

$$\Delta E = \frac{2}{5}\alpha_{20}^2 a_s A^{2/3} - \frac{1}{5}\alpha_{20}^2 a_c Z^2 A^{-1/3}, \quad (1.41)$$

if we set equation (1.41) to zero, we get, other than the undeformed solution for $\alpha_{20} = 0$,

$$\frac{Z^2}{A} = \frac{2a_s}{a_c}, \quad (1.42)$$

where the ratio $2a_s/a_c$ amounts to ≈ 50 in typical parametrizations of the SEMF. Equation (1.41), shows that for values of the so called *fissility parameter* Z^2/A larger than 50, the energy change becomes negative, favouring a configuration in which the nucleus fragments due to the spontaneous fission.

1.4.1. Octupole deformations

2 | State of the art, objective and motivation

2.1. State of the art and motivation

The need to account for nuclear deformations has been highlighted in chapter 1, particularly in regard to heavy nuclei and the fission process in section 1.4. The solution of the many-body problem has always been a computational challenge, mainly mitigated by using expansions on bases or assuming certain symmetries to reduce the dimensionality of the problem.

The problems that arise when using either of the aforementioned approaches call for the implementation of unconstrained codes.

Basis expansions An efficient method to solve the many-body problem is to use basis expansions of the harmonic oscillator, similar to the nuclear system as mentioned in section 1.1. Several implement this kind of procedure, one of which is used in the present work to benchmark our own implementation in section 6.1.1. The main limitation of such approach is that weakly bound states and large deformations are not well represented, the former due fundamentally different asymptotic behaviours for $r \rightarrow \infty$, between the HO $\sim e^{-r^2}$ – and loosely bound or quasi-resonant states $\sim e^{-r}$ [19, 59], while the latter needs a large number of HO shells to converge, leading to an exponential increase of the computational cost [40].

Symmetry assumptions Another approach to reduce the complexity of the procedure is to assume certain symmetries, as to reduce the dimensionality of the problem to save computational time. The main example of such codes is the spherical solution [15, 61] of the Hartree-Fock equations, but axially symmetric ones exist as well [46]. The limitations of such codes are obvious, as they systematically prevent the representation of certain broken symmetries.

Unconstrained codes In recent years, codes that solve the Hartree-Fock or Hartree-Fock-Bogoliubov problem on an unconstrained 3D mesh have been developed [14, 42, 51, 52]. These codes are able to represent broken symmetries, but they require a huge amount of computational power to be run at an acceptable accuracy, unless certain assumptions are made, such as plane reflection [51, 52]. Hence, the need to explore new, more computationally efficient methods to solve the many-body problem, as done in this thesis through the use of the General Conjugate Gradient method, detailed in section 4.2.3.

2.2. Objectives

The aim of this work is to develop a new implementation of the Hartree-Fock method on an unconstrained 3D mesh, by the use of the General Conjugate Gradient method. The goals addressed by this work are the following:

- assess the feasibility of the General Conjugate Gradient for the solution of large-scale eigenvalue problems;
- solve the self-consistent Hartree-Fock equations on an unconstrained 3D mesh;
- verify the numerical accuracy of the new implementation against existing spherical codes;
- gauge the numerical accuracy of deformations, comparing results with well established deformed codes; and
- attempt to produce novel results that specifically require an unconstrained implementation of this kind, and establish the advance brought to the field by this work.

2.3. Methods

The methods used in this work are the following:

Skyrme energy functional The many-body problem is treated within the well established Hartree-Fock framework, detailed in section 3.1; as mentioned in the introduction in chapter 1, it is not sufficient, as a more general energy density functional approach has to be taken, which is developed in section 3.3 for the Skyrme functional used in this work.

Finite differences and General Conjugate Gradient After the main equations to be solved have been derived, the numerical methods used to solve them are detailed in chapter 4, starting with the numerical discretization of the equations in section 4.1 and

then solving the eigenvalue problem using an implementation of the General Conjugate Gradient method in section 4.2.3. Finally closing with some remarks about specific details about the code and the algorithm parameters in section 4.3.

3 | Energy functional

3.1. Hartree-Fock theory

While a phenomenological description of bulk nuclear structure properties can be carried out using the liquid drop model or empirical mean-field potentials like Woods-Saxon or Nilsson, as we have seen in section 1.1, this is not sufficient to accurately reproduce all experimental ground state observables, systematically throughout the chart of nuclei, in an accurate manner.

A more rigorous approach needs to take into account the fact that the mean field which the nucleons interact with, is generated by the nucleons themselves. Starting from the many-body hamiltonian of the system, we will be able to extract a single particle Hamiltonian, including an effective mean field potential generated by the effective microscopic force, through the use of the Hartree-Fock method.

We start by writing the many-body hamiltonian of the system, which is a collection of A interacting fermions, given by

$$\hat{H} = \hat{T} + \hat{V} = \sum_i -\frac{\hbar^2}{2m} \nabla_i^2 + \sum_{i<j} v_{ij}^{(2)} + \sum_{i<j<k} v_{ijk}^{(3)} + \dots \quad (3.1)$$

The corresponding Schrödinger equation reads

$$\hat{H}\Psi = E\Psi. \quad (3.2)$$

3.1.1. Variational principle

Since \hat{H} is a many body operator, finding its eigenstates would be a rather challenging task. To our aid, comes the variational principle, from which we can show that equation (3.2) is equivalent to

$$\delta E[\Psi] = \delta \frac{\langle \Psi | \hat{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = 0. \quad (3.3)$$

The variation (3.3) can be obtained from an arbitrary variation of Ψ , which can be done independently on $\langle\Psi|$ and $|\Psi\rangle$, since Ψ is complex, yielding

$$\langle\delta\Psi|\hat{H}-E|\Psi\rangle+\langle\Psi|\hat{H}-E|\delta\Psi\rangle=0 \quad (3.4)$$

since the variation is arbitrary, we can multiply by a phase factor $|\delta\Psi\rangle\mapsto i|\delta\Psi\rangle$ and get

$$-i\langle\delta\Psi|\hat{H}-E|\Psi\rangle+i\langle\Psi|\hat{H}-E|\delta\Psi\rangle=0. \quad (3.5)$$

Combining equations (3.4) and (3.5), we get

$$\langle\delta\Psi|\hat{H}-E|\Psi\rangle=0. \quad (3.6)$$

Again, since the variation is arbitrary, equation (3.6) satisfies equation (3.2).

Ground state

Since we always restrict ourselves to a certain subspace of the full Hilbert space, we can only find an approximate solution to the eigenvalue problem. Expanding this solution on the complete set of exact eigenstates of \hat{H} , we have

$$|\Psi\rangle=\sum_n a_n |\Psi_n\rangle \quad (3.7)$$

the total energy amounts to

$$E[\Psi]=\frac{\sum_{nn'}\langle a_{n'}\Psi_{n'}|\hat{H}|a_n\Psi_n\rangle}{\sum_{nn'}\langle a_{n'}\Psi_{n'}|a_n\Psi_n\rangle}=\frac{\sum_n E_n|a_n|^2}{\sum_n |a_n|^2}\geq\frac{\sum_n E_0|a_n|^2}{\sum_n |a_n|^2}\geq E_0, \quad (3.8)$$

where E_0 is the ground state energy of the system. The orthonormality of the Hamiltonian eigenfunctions $\langle\Psi_{n'}|\Psi_n\rangle=\delta_{nn'}$ has been used. Equation (3.8) tells us that the minimum of the functional $E[\Psi]$ in any variationl subspace we are considering is bound from below by the true ground state energy.

3.1.2. Hartree-Fock equations

The Hartree-Fock method is the application of the variational principle (3.6) to a system of independent particles, whose wavefunction takes the form of a Slater determinant,

which reads

$$\Psi = \frac{1}{\sqrt{A!}} \sum_{\{p\}} (-1)^p \varphi_{p(1)}(\mathbf{r}_1) \dots \varphi_{p(A)}(\mathbf{r}_A) \quad (3.9)$$

where φ_i are the single-particle orthonormal states, which serve the role of variational parameters in Hartree-Fock. The Slater determinant sums over all possible permutations of the A fermions on the single particle states, with a $-$ sign according to the parity of the permutation. The Slater determinant satisfies the permutation symmetry of fermions, so that the Pauli exclusion principle is not violated.

To ensure the orthonormality of the single-particle states, we need to add a Lagrange multiplier to the variation (3.6) of E , which ends up reading

$$\delta \left(E[\Psi] - \sum_i \lambda_i \int \varphi_i^* \varphi_i d\mathbf{r} \right) = 0. \quad (3.10)$$

The total energy of the system is given by

$$E[\Psi] = \langle \Psi | \hat{T} + \hat{V} | \Psi \rangle = \langle \Psi | \hat{T} | \Psi \rangle + \langle \Psi | \hat{V} | \Psi \rangle, \quad (3.11)$$

which can be expressed through the single-particle states $\{\varphi_i\}$, yielding

$$\langle \Psi | \hat{T} | \Psi \rangle = \sum_i^A -\frac{\hbar^2}{2m} \int \varphi_i^*(\mathbf{r}) \nabla^2 \varphi_i(\mathbf{r}) d\mathbf{r} = \sum_i \langle i | t | i \rangle \quad (3.12)$$

$$\langle \Psi | \hat{V} | \Psi \rangle = \frac{1}{2} \sum_{ij} \int \varphi_i^*(\mathbf{r}) \varphi_j^*(\mathbf{r}') v_{ij}(\mathbf{r}, \mathbf{r}') \varphi_i(\mathbf{r}) \varphi_j(\mathbf{r}') d\mathbf{r} d\mathbf{r}' \quad (3.13)$$

$$- \frac{1}{2} \sum_{ij} \int \varphi_i^*(\mathbf{r}) \varphi_j^*(\mathbf{r}') v_{ij}(\mathbf{r}, \mathbf{r}') \varphi_i(\mathbf{r}') \varphi_j(\mathbf{r}) d\mathbf{r} d\mathbf{r}' \quad (3.14)$$

$$= \frac{1}{2} \langle ij | \bar{v} | ij \rangle. \quad (3.15)$$

If we use $\varphi_i^*(\mathbf{r})$ as the quantity to be varied in equation (3.10), we get the *Hartree-Fock equations*

$$- \frac{\hbar^2}{2m} \nabla^2 \varphi_i \quad (3.16)$$

$$+ \frac{1}{2} \sum_j^A \int \varphi_j^*(\mathbf{r}') v_{ij}(\mathbf{r}, \mathbf{r}') \varphi_j(\mathbf{r}') \varphi_i(\mathbf{r}) d\mathbf{r}' \quad (3.17)$$

$$- \frac{1}{2} \sum_j^A \int \varphi_j^*(\mathbf{r}') v_{ij}(\mathbf{r}, \mathbf{r}') \varphi_j(\mathbf{r}) \varphi_i(\mathbf{r}') d\mathbf{r}' = h\varphi_i = \varepsilon_i \varphi_i \quad (3.18)$$

here the Lagrange multipliers λ_i have been replaced by ε_i , since they can be interpreted as the energy of the single-particle states.

Now, a couple of remarks are in order.

Exchange interaction The first interaction term (3.17), called Hartree term, arises from considering independent particles, and is also routinely found in classical physics. The second one in (3.18), called Fock term, or exchange term, takes the form of an integral operator and is present when considering quantum mechanical indistinguishable particles. For what concerns the solution of equation (3.16), the Fock term is problematic, and can be avoided using zero-range forces like the Skyrme one [56], which is used in the present work, since it renders the exchange term as a local one.

Self-consistent solution Even if the interaction terms are local, the equation is still highly non-linear, since the mean field potential will be a function of the eigenfunctions themselves. The consequence is that the solution needs to be found *self-consistently*, that is, by solving for the set of eigenfunctions $\{\varphi_i\}$, using them to build the new mean field, and solving again, repeating the process until convergence.

3.1.3. Symmetries in Hartree-Fock

Since the objective of this work is to solve the Hartree-Fock equations without spatial symmetry assumptions, it is useful to first understand how symmetries propagate along the self-consistent calculation.

We start by defining the creation and annihilation operators of the single particle hamiltonian eigenstates, a_i^\dagger, a_i , which abide the usual anticommutation relations of fermions

$$\{a_i, a_j^\dagger\} = \delta_{ij} \quad (3.19)$$

If we expand on a different, orthonormal complete basis $\{\chi_l\}$, we can write the corresponding creation and annihilation operators c_l^\dagger, c_l as

$$\varphi_k = \sum_l D_{lk} \chi_l \quad (3.20)$$

$$a_k^\dagger = \sum_l D_{lk} c_l^\dagger \quad (3.21)$$

$$a_k = \sum_l D_{lk}^\dagger c_l \quad (3.22)$$

Since orthonormality is guaranteed for both sets, taking $\langle \varphi_j | \varphi_k \rangle$ yields

$$\delta_{jk} = \langle \varphi_j | \varphi_k \rangle = \sum_{l'} D_{l'j}^\dagger D_{lk} \langle \chi_l | \chi_l' \rangle \implies DD^\dagger = 1, \quad (3.23)$$

meaning that D is a unitary transformation. We can define the density matrix as

$$\rho_{ll'} = \langle \Psi | c_{l'}^\dagger c_l | \Psi \rangle, \quad (3.24)$$

whose trace is equal to the particle number A , as per equation (3.25)

$$\text{Tr } \rho = \sum_l \langle \Psi | c_l^\dagger c_l | \Psi \rangle = \sum_l \langle \Psi | a_l^\dagger a_l | \Psi \rangle = \sum_l^A \langle \Psi | \Psi \rangle = A. \quad (3.25)$$

Writing the many body hamiltonian (3.2) in the arbitrary basis of second quantization operators c_l^\dagger, c_l , we get

$$\hat{H} = \sum_{l_1 l_2} t_{l_1 l_2} c_{l_1}^\dagger c_{l_2} + \frac{1}{4} \sum_{l_1 l_2 l_3 l_4} \bar{v}_{l_1 l_2 l_3 l_4} c_{l_1}^\dagger c_{l_2}^\dagger c_{l_3} c_{l_4} \quad (3.26)$$

where $t_{l_1 l_2}$ and $\bar{v}_{l_1 l_2 l_3 l_4}$ are defined as

$$t_{l_1 l_2} = \langle - | c_{l_1} c_{l_2} t c_{l_1}^\dagger c_{l_2}^\dagger | - \rangle = \langle l_1 l_2 | t | l_1 l_2 \rangle \quad (3.27)$$

$$\bar{v}_{l_1 l_2 l_3 l_4} = \langle l_1 l_2 l_3 l_4 | v | l_1 l_2 l_3 l_4 \rangle - \langle l_1 l_2 l_4 l_3 | v | l_1 l_2 l_4 l_3 \rangle. \quad (3.28)$$

The minimization (3.10) can be restated as the variation of $\langle \Psi | \hat{H} | \Psi \rangle$, with respect to the density matrix $\rho_{ll'}$, which yields

$$h_{ll'} = \frac{\partial E[\rho]}{\partial \rho_{ll'}} = t + \sum_{kk'} \bar{v}_{lk' l' k} \rho_{kk'} = t + \Gamma_{ll'} \quad (3.29)$$

where $\Gamma_{ll'}$ is the mean field potential in the arbitrary basis. Being h diagonal in the Hartree-Fock basis, the self-consistent solution is the one for which

$$[h, \rho] = 0 \quad (3.30)$$

holds.

Symmetry propagation

Let us suppose to start a Hartree-Fock calculation with an initial guess for which the corresponding density matrix $\rho^{(0)}$ is symmetric under the action of a many-body symmetry operator S which commutes with the Hamiltonian

$$[S, \hat{H}] = 0. \quad (3.31)$$

It can be shown [49] that

$$S\Gamma[\rho]S^\dagger = \Gamma[S\rho S^\dagger]. \quad (3.32)$$

The single particle Hamiltonian h will then display the same property

$$Sh[\rho]S^\dagger = h[S\rho S^\dagger] = h[\rho] \quad (3.33)$$

meaning that h will be symmetric under the action of S , as well as the next iteration's density matrix $\rho^{(1)}$. The symmetry S gets propagated self-consistently until the minimum is found.

This has profound numerical implications, since the minimum energy configuration of a deformed nuclei can be reached only by starting guesses with the same broken symmetries. It can be the case that numerical noise allows to explore the full energy surface, but if one has to take into consideration the numerical cost of a bad guess, then it's still advantageous to start from a correct one in terms of symmetries.

3.1.4. Density Functional Theory

It shall be evident shortly, in section 3.3, that a more general approach to microscopic models has to be taken in order to give a satisfactory description of nuclei and nuclear matter. The method that we will briefly outline here is called Density Functional Theory (DFT).

DFT was introduced by P. Hohenberg and W. Kohn in 1964 [28], by proving two theorems. The **first** Hohenberg Kohn (HK) theorem states that the energy of a fermion system, subject to an external potential V_{ext} can be expressed solely as a functional of the particle density ρ of the system

$$E[\rho] = F[\rho] + \int V_{\text{ext}}\rho(\mathbf{r})d\mathbf{r}, \quad (3.34)$$

where $F[\rho]$ is a universal functional given by the type of fermions considered, while V_{ext} term is the external potential to which the system is subject to; when treating atomic nuclei, the potential is generated by the nucleons themselves, so this term will be omitted

in the following. The **second** HK theorem states that the ground state of the system is found by minimizing the functional (3.34) with respect to ρ .

HK theorems are fundamental but not constructive [16], since they do not provide a form for the functional F , which is intrinsic to the physics of the fermions at hand. A pragmatic approach to using DFT was outlined by Kohn and Sham in 1965 [32]. They proposed expressing the system as a set of non-interacting particles occupying auxiliary orbitals φ_i , which yield the particle density

$$\rho(\mathbf{r}) = \sum_i |\varphi_i(\mathbf{r})|^2 \quad (3.35)$$

and an energy functional of the form

$$E[\rho] = T[\rho] + E_H[\rho] + E_{xc}[\rho]. \quad (3.36)$$

where T is the kinetic energy, which reads

$$T[\rho] = -\frac{\hbar^2}{2m} \sum_i \varphi_i^*(\mathbf{r}) \nabla^2 \varphi_i(\mathbf{r}) \quad (3.37)$$

and E_H is the classical Hartree term, which in an electronic system may read

$$E_H[\rho] = \iint \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r} d\mathbf{r}' \quad (3.38)$$

while E_{xc} is an unknown exchange term. In electronic systems, the Hartree term is known (3.17) and the exchange term can be approximated thanks to the compensation of its error with the one of particles correlation neglect [41]. In nuclear physics, things are more complicated, since both terms are unknown; historically, effective interactions in HF have been used to extract an effective Hamiltonian density from which an energy density functional (EDF) can be formulated, whenever a pure interaction is not sufficient to describe nuclear systems, as we shall see in section 3.3.

3.2. Pairing in Hartree-Fock theory

In this section, we will discuss the two common approaches to include nuclear pairing in the HF theory. The aim of these few pages is to provide a brief overview of how the BCS equations are derived and understand the basics of the more general Hartree-Fock-Bogoliubov theory. The former method is the most widely implemented thanks to its low complexity [9, 15, 33], while the latter, more sophisticated and advanced, is the standard

in modern codes [13, 39, 54]. We will touch on it so that the reader may appreciate in the numerical chapter the natural extension of this work to the more general Bogoliubov ansatz.

3.2.1. BCS theory

The BCS approximation, from Bardeen-Cooper-Schrieffer, is the same theory used to describe Cooper pairs in superconductivity, applied to the nuclear case. The ansatz of BCS is that nucleons are paired in states whose total angular momentum is zero, such a wavefunction can be expressed as $|JM\rangle = |00\rangle$ and reads

$$|00\rangle = \sum_{m_j} \langle jm_j j - m_j | 00 \rangle |jm_j\rangle |j - m_j\rangle \quad (3.39)$$

Introducing the time-reversal operator $\hat{\mathcal{T}} : t \mapsto -t$, it acts on $|00\rangle$ as

$$\hat{\mathcal{T}} |jm_j\rangle = \widetilde{|jm_j\rangle} = (-1)^{j+m_j} |j - m_j\rangle, \quad (3.40)$$

using this relation, equation (3.39) becomes

$$|00\rangle = -\frac{1}{\sqrt{2j+1}} \sum_{m_j} |jm_j\rangle \widetilde{|jm_j\rangle}. \quad (3.41)$$

Hence BCS amounts to replacing the Slater determinant with a more general wavefunction to describe the ground state, which reads

$$|\text{BCS}\rangle = \prod_{k>0} (u_k + v_k a_k^\dagger a_{\tilde{k}}^\dagger) |-\rangle \quad (3.42)$$

where u_k and v_k are real parameters whose meaning will shortly be clear, k is short-hand for $|jm\rangle$, and $\tilde{k} = -k$ denotes the time-reversal state of k ; the product runs over positive k only. The BCS wavefunction is the creation in the vacuum of quasi-particles made of time-reversal paired particles, instead of individual ones. The normalization condition on the BCS wavefunction reads

$$1 = \langle \text{BCS} | \text{BCS} \rangle = \prod_{k>0} \langle - | (u_k + v_k a_k a_{\tilde{k}}) (u_k + v_k a_{\tilde{k}}^\dagger a_k^\dagger) | - \rangle = \prod_{k>0} (u_k^2 + v_k^2) = 1 \quad (3.43)$$

which implies, for every pair k , the condition

$$u_k^2 + v_k^2 = 1. \quad (3.44)$$

Taking the expectation value of the particle number operator $\hat{N} = \sum_k a_k^\dagger a_k$ yields [6]

$$\langle \text{BCS} | \hat{N} | \text{BCS} \rangle = 2 \sum_{k>0} v_k^2, \quad (3.45)$$

while the expectation value of the particle number dispersion reads

$$\langle \Delta \hat{N}^2 \rangle = \langle \hat{N}^2 \rangle - \langle \hat{N} \rangle^2 = 4 \sum_{k>0} v_k^2 u_k^2. \quad (3.46)$$

The consequence of this result is profound. The BCS ansatz does not assume a fixed number of particles, rather it becomes an observable of the system, with an expectation value that depends on how the parameters v_k^2 are set, which represent the probability of finding a particle in the k -th state. We can now write the many body Hamiltonian of the system as in equation (3.26)

$$\hat{H} = \sum_{k_1 k_2} t_{k_1 k_2} a_{k_1}^\dagger a_{k_2} + \frac{1}{4} \sum_{k_1 k_2 k_3 k_4} \bar{v}_{k_1 k_2 k_3 k_4} a_{k_1}^\dagger a_{k_2}^\dagger a_{k_3} a_{k_4} \quad (3.47)$$

and replace it with the Routhian

$$\langle \text{BCS} | \hat{H} - \lambda \hat{N} | \text{BCS} \rangle \quad (3.48)$$

so that the expected number of particles may be fixed, under the appropriate choice of λ , by the relation

$$\frac{\partial}{\partial N} \langle \text{BCS} | \hat{H} | \text{BCS} \rangle = \lambda. \quad (3.49)$$

the Lagrange multiplier λ takes on the meaning of the Fermi energy. We can now apply the variational principle (3.6) to (3.48) using the v_k as variational quantities, which yields

$$4\tilde{\varepsilon}_k^2 u_k^2 v_k^2 = \Delta_k^2 - 4\Delta_k^2 u_k^2 v_k^2, \quad (3.50)$$

where the pairing gap Δ_k is defined as

$$\Delta_k = - \sum_{k'} \bar{v}_{k\bar{k}k'\bar{k}'} v_{k'} u_{k'} \quad (3.51)$$

and the quantity $\tilde{\varepsilon}_k$ is defined as

$$\tilde{\varepsilon}_k = \frac{1}{2} \left[t_{kk} + t_{\bar{k}\bar{k}} - 2\lambda + \sum_{k'} (\bar{v}_{k\bar{k}k'\bar{k}'} v_{k'} u_{k'} + \bar{v}_{\bar{k}k'\bar{k}k'} v_{k'}^2) \right] \quad (3.52)$$

$$= \frac{1}{2} [h_{kk} + h_{\bar{k}\bar{k}}] - \lambda. \quad (3.53)$$

Introducing the quasi-particle energy

$$E_k = \sqrt{\tilde{\varepsilon}_k^2 + \Delta_k^2} \quad (3.54)$$

we can combine definitions (3.51) and (3.54) with equation (3.52), under the normalization condition (3.44), to get an equation for v_k^2

$$v_k^2 = \frac{1}{2} \pm \frac{|\tilde{\varepsilon}_k|}{2E_k}. \quad (3.55)$$

Since in the Hartree-Fock limit, where the occupations v_k^2 are equal to one below the fermi energy and zero above, and the gaps Δ_k vanish, rendering $E_k = \tilde{\varepsilon}_k$, we only select the solution

$$v_k^2 = \frac{1}{2} - \frac{\tilde{\varepsilon}_k}{2E_k}. \quad (3.56)$$

Using the normalization condition to write $u_k^2 = 1 - v_k^2$, and plugging it into the gaps definition (3.51), we arrive to the gap equation

$$\Delta_k = - \sum_{k'} \frac{\Delta_{k'} \bar{v}_{k\bar{k}k'\bar{k}'}}{2E_{k'}} \quad (3.57)$$

The system of equations (3.57, 3.44, 3.55, 3.54, 3.52), together with the condition on \hat{N} – ie $\langle \hat{N} \rangle = N$ – is closed and can be solved numerically, usually through an effective pairing interaction.

3.2.2. Hartree-Fock-Bogoliubov theory

The most general ansatz to account for pairing interactions in Hartree-Fock theory is the Hartree-Fock-Bogoliubov (HFB) theory, it allows a treatment of the mean-field and pairing interactions in a unified way, the quasi-particles created on the vacuum are the most general ones, instead of being time-reversal paired particles. Let us start by writing

a Bogoliubov transformation from the particle basis c_i to a quasi-particle one

$$\beta_k^\dagger = \sum_l U_{lk} c_l^\dagger + V_{lk} c_l. \quad (3.58)$$

If we take the Hermitian conjugate of the relation (3.58), we get the transformation for β_k , we are then able to write in matrix form

$$\begin{pmatrix} \beta \\ \beta^\dagger \end{pmatrix} = \begin{pmatrix} U^\dagger & V^\dagger \\ V^T & U^T \end{pmatrix} \begin{pmatrix} c \\ c^\dagger \end{pmatrix} = \mathcal{W}^\dagger \begin{pmatrix} c \\ c^\dagger \end{pmatrix}, \quad (3.59)$$

where the matrix of matrices \mathcal{W} reads

$$\mathcal{W} = \begin{pmatrix} U & V^* \\ V & U^* \end{pmatrix}. \quad (3.60)$$

Taking the product $\mathcal{W}^\dagger \mathcal{W}$ and imposing separate fermionic commutation relations of the operators $\beta, \beta^\dagger, c, c^\dagger$, we get that \mathcal{W} is unitary, hence

$$\mathcal{W}^\dagger \mathcal{W} = \mathcal{W} \mathcal{W}^\dagger = I. \quad (3.61)$$

We can now invert equation (3.59) by multiplying both sides on the left by \mathcal{W} , which yields

$$\mathcal{W} \begin{pmatrix} \beta \\ \beta^\dagger \end{pmatrix} = \begin{pmatrix} c \\ c^\dagger \end{pmatrix}.$$

Using the Messiah-Bloch decomposition [18], we can write the unitary matrix \mathcal{W} as

$$\mathcal{W} = \begin{pmatrix} D & 0 \\ 0 & D^* \end{pmatrix} \begin{pmatrix} \bar{U} & \bar{V} \\ \bar{V} & \bar{U} \end{pmatrix} \begin{pmatrix} C & 0 \\ 0 & C^* \end{pmatrix} \quad (3.62)$$

where D and C are unitary matrices and \bar{U} and \bar{V} are real matrices, which have a particular blocked form, expressed through the coefficients u_k, v_k ; the reader may refer to appendix A.4 for the explicit representation. We can also define the matrices U, V as

$$U = D \bar{U} C, \quad V = D^* \bar{V} C. \quad (3.63)$$

Using the decomposition (3.62) we can define the *canonical basis* as

$$a_k^\dagger = \sum_l D_{lk}^\dagger c_l^\dagger, \quad (3.64)$$

a *special Bogoliubov transformation* between *paired* levels as

$$\alpha_k^\dagger = u_k a_k^\dagger - v_k a_{\bar{k}}, \quad (3.65)$$

$$\alpha_{\bar{k}}^\dagger = u_k a_{\bar{k}}^\dagger + v_k a_k, \quad (3.66)$$

and *blocked* levels

$$\alpha_i = a_i, \quad \alpha_n^\dagger = a_n^\dagger \quad (3.67)$$

$$\alpha_i = a_i^\dagger, \quad \alpha_n = a_n, \quad (3.68)$$

where $u_k = u_{\bar{k}}$, $v_k = -v_{\bar{k}}$, and a unitary transformation of the quasi-particle operators α_k^\dagger among themselves

$$\beta_k^\dagger = \sum_{k'} C_{k'k} a_{k'}^\dagger. \quad (3.69)$$

We are now able to define the Bogoliubov ground state $|\text{HFB}\rangle$, as the one for which

$$\beta_k |\text{HFB}\rangle = 0 \quad \forall k = 1, \dots, M \quad (3.70)$$

where M is determined by the physical situation [49]. The wavefunction that satisfies this condition reads

$$|\text{HFB}\rangle = \prod_k^M \beta_k |-\rangle. \quad (3.71)$$

We can define the pairing tensor as

$$\kappa_{ll'} = \langle \text{HFB} | c_{l'} c_l | \text{HFB} \rangle, \quad (3.72)$$

which in matrix form reads, alongside the density matrix

$$\kappa = UV^\dagger, \quad \rho = V^* V^T. \quad (3.73)$$

We can now apply the variational principle (3.6)

$$\delta \frac{\langle \text{HFB} | \hat{H} - \lambda \hat{N} | \text{HFB} \rangle}{\langle \text{HFB} | \text{HFB} \rangle} = 0. \quad (3.74)$$

which yields the eigenvalue problem

$$\begin{pmatrix} h - \lambda & \Delta \\ -\Delta^* & -(h - \lambda)^* \end{pmatrix} \begin{pmatrix} U_k \\ V_k \end{pmatrix} = \mathcal{H}_{\text{HFB}} \begin{pmatrix} U_k \\ V_k \end{pmatrix} = E_k \begin{pmatrix} U_k \\ V_k \end{pmatrix}, \quad (3.75)$$

Here, h is the single-particle Hamiltonian, which reads

$$h_{kk'} = t_{kk'} + \Gamma_{kk'}, \quad (3.76)$$

where $\Gamma_{kk'}$ is the mean field potential, given by

$$\Gamma_{kk'} = \sum_{ll'} \bar{v}_{kl'k'l} \rho_{ll'} \quad (3.77)$$

and the pairing field Δ reads

$$\Delta_{kk'} = \sum_{ll'} \bar{v}_{kk'll'} \kappa_{ll'}. \quad (3.78)$$

In the canonical basis, we are able to solve for the occupation numbers

$$u_k^2 = \frac{1}{2} \left(1 + \frac{h_{kk} + h_{\bar{k}\bar{k}}}{\sqrt{(h_{kk} + h_{\bar{k}\bar{k}})^2 + 4\Delta_{k\bar{k}}^2}} \right) \quad (3.79)$$

where $v_k^2 = 1 - u_k^2$ is guaranteed by the unitarity of the matrices. Starting from an initial guess, we solve the eigenvalue problem (3.75), we extract the occupation numbers (3.79), use them to build the new mean field (3.77) and pairing field (3.78), and repeat the process until convergence.

HFB quasi-particle spectrum Let us assume that $\Psi = (U, V)^T$ is a solution of equation (3.75) with eigenvalue E

$$\mathcal{H}_{\text{HFB}} \Psi = E \Psi. \quad (3.80)$$

Let the particle-hole matrix \mathcal{C} be defined as

$$\mathcal{C} = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad (3.81)$$

it's trivial to show that

$$\mathcal{C} \mathcal{H}_{\text{HFB}} \mathcal{C} = -\mathcal{H}_{\text{HFB}}^*, \quad (3.82)$$

and

$$\mathcal{C} = \mathcal{C}^{-1} \implies \mathcal{C} \mathcal{H}_{\text{HFB}} = -\mathcal{H}_{\text{HFB}}^* \mathcal{C}. \quad (3.83)$$

If we take the complex conjugate of equation (3.80), we get

$$\mathcal{H}_{\text{HFB}}^* \Psi^* = E \Psi^*, \quad (3.84)$$

if we multiply both sides on the left by \mathcal{C} and use (3.83), we get

$$-\mathcal{H}_{\text{HFB}} \mathcal{C} \Psi^* = E \mathcal{C} \Psi^*, \quad (3.85)$$

$$\mathcal{H}_{\text{HFB}} \mathcal{C} \Psi^* = -E \mathcal{C} \Psi^*, \quad (3.86)$$

meaning that $\mathcal{C} \Psi^*$ is a solution of the eigenvalue problem (3.75) as well, with eigenvalue $-E$, hence for every quasi-particle energy we have a corresponding opposite-sign one; moreover, it can be proven that the HFB hamiltonian is unbounded, both from below and above [47]. This feature poses a challenge for numerical solutions of the HFB problem, as we shall see in chapter 4.

3.3. Skyrme

Now that the theoretical framework is clear, we can use the Skyrme microscopic effective interaction to do nuclear structure calculations.

3.3.1. Skyrme force

It was first proposed by Tony Skyrme in 1958 [56] as a zero range force between nucleons, comprising a two body attractive term that reads

$$\begin{aligned} v^{(2)}(\mathbf{r}_1, \mathbf{r}_2) = & t_0 (1 + x_0 P_\sigma) \delta(\mathbf{r}) \\ & + \frac{1}{2} t_1 (1 + x_1 P_\sigma) [\mathbf{P}'^2 \delta(\mathbf{r}) + \delta(\mathbf{r}) \mathbf{P}^2] \\ & + t_2 P_\sigma \mathbf{P}' \cdot \delta(\mathbf{r}) \mathbf{P} \\ & + i W_0 \boldsymbol{\sigma} \cdot [\mathbf{P}' \times \delta(\mathbf{r}) \mathbf{P}] \end{aligned}$$

and a three body term, that is

$$v^{(3)}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = t_3 \delta(\mathbf{r}_1 - \mathbf{r}_2) \delta(\mathbf{r}_2 - \mathbf{r}_3)$$

which mimics the repulsive three-body force; without it, a collapse of the nuclear density would occur. The different operators are defined as

$$\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2 \quad \mathbf{R} = \frac{\mathbf{r}_1 + \mathbf{r}_2}{2} \quad (3.87)$$

which are respectively the relative position of two particles and their center of mass coordinate, assuming equal masses.

$$\mathbf{P} = \frac{-i(\nabla_1 - \nabla_2)}{2} \quad (3.88)$$

which is the so called relative wave-number operator, corresponding to the relative momentum of the two interacting particles.

$$\boldsymbol{\sigma} = \boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2 \quad (3.89)$$

being the total spin of the two interacting particles, and lastly

$$\mathbf{P}_\sigma = \frac{(1 + \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2)}{2} \quad (3.90)$$

which represents the spin-exchange operator. Primed operators refer to the adjoint acting on the left.

It's trivial to show that the three-body term, in the case of even-even nuclei, is equivalent to a two-body, density-dependent interaction [61]

$$v^{(3)}(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{6} t_3 (1 + P_\sigma) \delta(\mathbf{r}) \rho(\mathbf{R}) \quad (3.91)$$

The zero-range feature is apparent in the choice of the Dirac delta $\delta(\mathbf{r})$, which allows the writing of the Fock term detailed in (3.18) as a purely local one.

Modern parametrization

The Skyrme force has evolved from the original one to accomodate new nuclei and properties, done through the addition of a few parameters, yielding the following form of the

interaction [10]

$$\begin{aligned}
v^{(2)}(\mathbf{r}_1, \mathbf{r}_2) = & t_0 (1 + x_0 P_\sigma) \delta(\mathbf{r}) \\
& + \frac{1}{2} t_1 (1 + x_1 P_\sigma) [\mathbf{P}'^2 \delta(\mathbf{r}) + \delta(\mathbf{r}) \mathbf{P}^2] \\
& + t_2 (1 + x_2 P_\sigma) \mathbf{P}' \cdot \delta(\mathbf{r}) \mathbf{P} \\
& + \frac{1}{6} t_3 (1 + x_3 P_\sigma) [\rho(\mathbf{R})]^\sigma \delta(\mathbf{r}) \\
& + i W_0 \boldsymbol{\sigma} \cdot [\mathbf{P}' \times \delta(\mathbf{r}) \mathbf{P}] \\
& + \frac{1}{6} t_3 (1 + x_3 P_\sigma) [\rho(\mathbf{R})]^\sigma \delta(\mathbf{r})
\end{aligned}$$

Here, the boundary between Hartree-Fock and DFT starts to thin out, as the exponent σ of the density makes that piece of the force a true three-body interaction only for the value $\sigma = 1$ [21]. On top of that, additional, empirical tuning of the resulting energy density needed to reach satisfactory physical accuracy, such as the case for the spin-orbit couplings [48], prompts for the following, well established proceeding: use the Skyrme interaction to obtain the Hamiltonian density

$$\langle H \rangle = \langle \Psi | H | \Psi \rangle = \int \mathcal{H}(\mathbf{r}) d\mathbf{r} \quad (3.92)$$

and use it as a starting point to build an energy density functional and employ DFT.

3.3.2. Energy density functional

The energy functional to be minimized is of the form [3]

$$E_{\text{HF}} = E_{\text{Kin}} + E_{\text{Skyrme}} + E_{\text{Coul}} = \int (\mathcal{E}_{\text{Kin}} + \mathcal{E}_{\text{Skyrme}} + \mathcal{E}_{\text{Coul}}) d\mathbf{r}. \quad (3.93)$$

Densities

Functional (3.93) can be expressed through a series of particle densities. Let us define them and express them on the spin coordinates up (\uparrow) and down (\downarrow) for the convenience in a mesh representation.

The starting point is the density matrix, defined as

$$\rho_q(\mathbf{r}\sigma, \mathbf{r}\sigma') = \sum_{\alpha} \phi_{\alpha,\sigma}(\mathbf{r}) \phi_{\alpha,\sigma'}^*(\mathbf{r}') \quad (3.94)$$

where the index α goes through all single particle states of the particles of type q (protons, neutrons) and the index σ refers to the spin coordinate. The particle density is defined

as

$$\begin{aligned}\rho_q(\mathbf{r}) &:= \rho_q(\mathbf{r}, \mathbf{r}') \Big|_{\mathbf{r}=\mathbf{r}'} := \sum_{\sigma} \rho(\mathbf{r}\sigma, \mathbf{r}'\sigma) \Big|_{\mathbf{r}=\mathbf{r}'} = \sum_{\alpha} \phi_{\uparrow}(\mathbf{r})\phi_{\uparrow}^*(\mathbf{r}') + \phi_{\downarrow}(\mathbf{r})\phi_{\downarrow}^*(\mathbf{r}') \Big|_{\mathbf{r}=\mathbf{r}'} \\ &= \sum_{\alpha} |\phi_{\uparrow}(\mathbf{r})|^2 + |\phi_{\downarrow}(\mathbf{r})|^2.\end{aligned}\quad (3.95)$$

The kinetic density reads

$$\begin{aligned}\tau_q(\mathbf{r}) &:= \sum_{\alpha} \nabla' \cdot \nabla \rho_q(\mathbf{r}, \mathbf{r}') \Big|_{\mathbf{r}'=\mathbf{r}} \\ &= \sum_{\sigma, \alpha} \nabla \phi_{\sigma}(\mathbf{r}) \cdot \nabla \phi_{\sigma}^*(\mathbf{r}') \Big|_{\mathbf{r}=\mathbf{r}'} = \sum_{\sigma, \alpha} |\nabla \phi_{\sigma}(\mathbf{r})|^2 \\ &= \sum_{\alpha} |\nabla \phi_{\uparrow}(\mathbf{r})|^2 + |\nabla \phi_{\downarrow}(\mathbf{r})|^2.\end{aligned}\quad (3.96)$$

The spin density reads

$$s_q(\mathbf{r}, \mathbf{r}') := \sum_{\sigma\sigma', i} \rho_q(\mathbf{r}\sigma, \mathbf{r}'\sigma') \langle \sigma' | \hat{\sigma} | \sigma \rangle = \sum_{\alpha} \begin{bmatrix} \phi_{\uparrow}^*(\mathbf{r}') & \phi_{\downarrow}^*(\mathbf{r}') \end{bmatrix} \hat{\sigma} \begin{bmatrix} \phi_{\uparrow}(\mathbf{r}) \\ \phi_{\downarrow}(\mathbf{r}) \end{bmatrix} \quad (3.97)$$

and lastly, the spin-orbit density tensor reads

$$\begin{aligned}J_{q, \mu\nu} &:= \frac{1}{2i} (\partial_{\mu} - \partial'_{\mu}) s_{q, \nu}(\mathbf{r}, \mathbf{r}') \Big|_{\mathbf{r}'=\mathbf{r}} \\ &= \frac{1}{2i} \left(\begin{bmatrix} \phi_{\uparrow}^*(\mathbf{r}') & \phi_{\downarrow}^*(\mathbf{r}') \end{bmatrix} \partial_{\mu} \hat{\sigma}_{\nu} \begin{bmatrix} \phi_{\uparrow}(\mathbf{r}) \\ \phi_{\downarrow}(\mathbf{r}) \end{bmatrix} - \begin{bmatrix} \phi_{\uparrow}(\mathbf{r}) & \phi_{\downarrow}(\mathbf{r}) \end{bmatrix} \partial'_{\mu} \hat{\sigma}_{\nu} \begin{bmatrix} \phi_{\uparrow}^*(\mathbf{r}') \\ \phi_{\downarrow}^*(\mathbf{r}') \end{bmatrix} \right) \Big|_{\mathbf{r}'=\mathbf{r}} \\ &= \sum_{\alpha} \text{Im} \left\{ \begin{bmatrix} \phi_{\uparrow}^*(\mathbf{r}) & \phi_{\downarrow}^*(\mathbf{r}) \end{bmatrix} \partial_{\mu} \hat{\sigma}_{\nu} \begin{bmatrix} \phi_{\uparrow}(\mathbf{r}) \\ \phi_{\downarrow}(\mathbf{r}) \end{bmatrix} \right\}\end{aligned}\quad (3.98)$$

which also defines the spin-orbit current vector \mathbf{J} that reads

$$J_{q, \kappa}(\mathbf{r}) = \sum_{\mu\nu} \epsilon_{\kappa\mu\nu} J_{q, \mu\nu}(\mathbf{r}). \quad (3.99)$$

Kinetic functional

The kinetic term can be expressed as

$$\mathcal{E}_{\text{Kin}} = \frac{\hbar^2}{2m} \tau \quad (3.100)$$

which is found integrating by parts (3.37).

Skyrme functional

Since this work only deals with even-even nuclei, only time-even densities, which are the ones previously defined, are non-vanishing, due to the ground state being time-reversal invariant [3]. This reduces the Skyrme functional to the following form [58]

$$\mathcal{E}_{\text{Skyrme}} = \sum_{t=0,1} \left\{ C_t^\rho [\rho_0] \rho_t^2 + C_t^{\Delta\rho} \rho_t \nabla^2 \rho_t + C_t^{\nabla \cdot J} \rho_t \nabla \cdot \mathbf{J}_t + C_t^\tau \rho_t \tau_t \right\} \quad (3.101)$$

where

$$C_0^\rho = +\frac{3}{8}t_0 + \frac{3}{48}t_3\rho_0^\sigma \quad (3.102)$$

$$C_1^\rho = -\frac{1}{8}t_0(1+2x_0) - \frac{1}{48}t_3(1+x_3)\rho_0^\sigma \quad (3.103)$$

$$C_0^\tau = +\frac{3}{16}t_1 + \frac{1}{16}t_2(5+4x_2) \quad (3.104)$$

$$C_1^\tau = -\frac{1}{16}t_1(1+2x_1) + \frac{1}{16}t_2(1+2x_2) \quad (3.105)$$

$$C_0^{\Delta\rho} = -\frac{9}{64}t_1 + \frac{1}{64}t_2(5+4x_2) \quad (3.106)$$

$$C_1^{\Delta\rho} = +\frac{3}{64}t_1(1+2x_1) + \frac{1}{64}t_2(1+2x_2) \quad (3.107)$$

$$C_0^{\nabla \cdot J} = -\frac{3}{4}W_0 \quad (3.108)$$

$$C_1^{\nabla \cdot J} = -\frac{1}{4}W_0. \quad (3.109)$$

Here, $t = 0, 1$ refers to the isoscalar and isovector components of the densities, eg

$$\rho_0 = \rho_p + \rho_n$$

$$\rho_1 = \rho_p - \rho_n.$$

We can now derive the Kohn-Sham equations, by minimizing the functional under the constraint

$$\langle \varphi_i | \varphi_j \rangle = \delta_{ij}. \quad (3.110)$$

The resulting Kohn-Sham equations are of the form

$$\left[-\nabla \left(\frac{\hbar^2}{2m_q^*(\mathbf{r})} \nabla \right) + U_q(\mathbf{r}) + \delta_{q,\text{proton}} U_C(\mathbf{r}) - i\mathbf{B}_q(\mathbf{r}) \cdot (\nabla \times \boldsymbol{\sigma}) \right] \varphi_\alpha = \varepsilon_\alpha \varphi_\alpha \quad (3.111)$$

where an effective mass field arises, which is defined as

$$\frac{\hbar^2}{2m_q^*(\mathbf{r})} = \frac{\delta \mathcal{E}}{\delta \tau_q} \quad (3.112)$$

a mean field potential, which reads

$$U_q(\mathbf{r}) = \frac{\delta \mathcal{E}}{\delta \rho_q} \quad (3.113)$$

and a spin-orbit field, given by

$$\mathbf{B}_q(\mathbf{r}) = \frac{\delta \mathcal{E}}{\delta \mathbf{J}_q}. \quad (3.114)$$

The coulomb field U_C , which is present only in the single particle equation for protons, doesn't come from the Skyrme interaction, rather from the Coulomb part of the whole functional. It will be properly derived in section 3.4.

Following the rules for functional derivatives, outlined in the appendix A.3 we get

$$\begin{aligned} \frac{\hbar^2}{2m_q^*(\mathbf{r})} = & + \frac{\hbar^2}{2m} \\ & + \frac{1}{8}[t_1(2 + x_1) + t_2(2 + x_2)]\rho(\mathbf{r}) \\ & - \frac{1}{8}[t_1(1 + 2x_1) + t_2(1 + 2x_2)]\rho_q(\mathbf{r}) \end{aligned} \quad (3.115)$$

$$\begin{aligned} U_q(\mathbf{r}) = & + \frac{1}{8}[t_1(2 + x_1) + t_2(2 + x_2)]\rho \\ & + \frac{1}{8}[t_2(1 + 2x_2) - t_1(1 + 2x_1)]\rho_q \\ & + \frac{1}{8}[t_1(2 + x_1) + t_2(2 + x_2)]\tau \\ & + \frac{1}{8}[t_2(1 + 2x_2) - t_1(1 + 2x_1)]\tau_q \\ & + \frac{1}{16}[t_2(2 + x_2) - 3t_1(2 + x_1)]\nabla^2\rho \\ & + \frac{1}{16}[3t_1(2x_1 + 1) + t_2(2x_2 + 1)]\nabla^2\rho_q \end{aligned} \quad (3.116)$$

$$\begin{aligned} \mathbf{B}_q(\mathbf{r}) = & + \frac{1}{2}W_0[\nabla\rho + \nabla\rho_q] \\ & - \frac{1}{8}(t_1x_1 + t_2x_2)\mathbf{J} + \frac{1}{8}(t_1 - t_2)\mathbf{J}_q. \end{aligned} \quad (3.117)$$

Unless otherwise specified, unlabelled densities denote isoscalar quantities (sum of neutron and proton).

3.3.3. Functionals

The set of parameters $(t_0, t_1, t_2, t_3, x_0, x_1, x_2, \sigma, W_0)$ in the Skyrme functional (3.101) is not fixed and may vary for different parametrizations. These parameters are fitted on experimental data so that the EDF may produce accurate results. The first parametrization was given by D Vautherin and D M Brink [61], known as S I, was fitted on double magic nuclei and nuclear matter data. This was sufficient for their calculations but not enough to describe other nuclei. Fitting these parameters on further data over the years led to the introduction of other parameter sets such as S I, S II, S III [10]. These were good functionals to reproduce ground states and certain aspects of nuclear structure, but they lacked the ability to account with good accuracy for deformation properties and fission barriers. The first modern functional to address this issue has been the SkM* [2]. Nowadays functionals such as the SLy family [11], try to be as broad as possible, taking into account results from ab-initio theories, exotic nuclei, drip lines, nuclei far from stability and so on.

3.4. Coulomb interaction

Unlike the Skyrme interaction, the Coulomb force is finite-range, giving rise to an unwanted integral operator in the single-particle Hamiltonian. A well known and widely used device is the Slater approximation [25], which gives a local exchange interaction. In this approximation, the Coulomb energy reads

$$E_{\text{Coul}} = \int \mathcal{E}_{\text{Coul}}(\mathbf{r}) d\mathbf{r}$$

where the energy density is given by

$$\mathcal{E}_{\text{Coul}}(\mathbf{r}) = \frac{e^2}{2} \left[\int \frac{\rho_p(\mathbf{r})\rho_p(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}' - \frac{3}{2} \left(\frac{3}{\pi} \right)^{\frac{1}{3}} \rho_p^{4/3}(\mathbf{r}) \right]. \quad (3.118)$$

which results in the Coulomb potential field

$$U_C(\mathbf{r}) = \frac{\delta \mathcal{E}_{\text{Coul}}}{\delta \rho_p} = \frac{e^2}{2} \left[\int \frac{\rho_p(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d^3\mathbf{r}' - 2 \left(\frac{3}{\pi} \right)^{\frac{1}{3}} \rho_p^{1/3}(\mathbf{r}) \right] \quad (3.119)$$

where the first term is the direct Coulomb interaction, which simply is the Coulomb energy generated by the proton density, while the second term is the exchange Coulomb interaction, which is local and depends on the proton density through a power factor of $1/3$. From a computational standpoint, the exchange part is trivial, while the direct one is more involved. One could compute the integral, but the complexity on a 3D mesh grows as $\mathcal{O}(N^6)$, where N is the total number of points on the mesh, rendering it unfeasible for fine calculations.

An alternative approach is to solve the Poisson equation (from now on, V_c refers to the direct part only)

$$\nabla^2 V_c = 4\pi e^2 \rho_p. \quad (3.120)$$

Given the proton density, we can impose Dirichlet boundary conditions, which can be extracted from a quadrupole expansion of the charge density [30]

$$V_c(\mathbf{r}) = 4\pi e^2 \sum_{\lambda=0}^2 \sum_{\mu=-\lambda}^{\lambda} \frac{\langle Q_{\lambda\mu} \rangle Y_{\lambda\mu}}{r^{1+\lambda}} \text{ on } \partial\Omega \quad (3.121)$$

where $\langle Q_{\lambda\mu} \rangle$ is defined as

$$\langle Q_{\lambda\mu} \rangle = \int r^\lambda Y_{\lambda\mu}^*(\mathbf{r}) \rho_p(\mathbf{r}) d^3\mathbf{r} \quad (3.122)$$

Since we expect a charge density confined to the nuclear shape, higher order terms in the expansion can be neglected, provided that the box is sufficiently large.

In a reference frame where the nucleus center of mass is at the origin, the expansion reduces to

$$V_c(\mathbf{r}) = \frac{Ze^2}{r} + e^2 \sum_{\mu=-2}^2 \frac{\langle Q_{2\mu} \rangle Y_{2\mu}}{r^3} \text{ on } \partial\Omega. \quad (3.123)$$

The reader can refer to appendix A.1 for the definition and numerical evaluation of the spherical harmonics $Y_{\lambda\mu}$.

3.5. Energy calculation

One, if not the most important physical quantity we want to compute is the total energy of the system.

Integrated energy

The obvious way would be to evaluate the functional for a given density. We will call this *integrated energy*.

$$E_{\text{int}} = E[\rho, \tau, J_{\mu\nu}] = \int \mathcal{E} d\mathbf{r}$$

Hartree-Fock energy

An alternative approach can be used, as in a stationary point $\delta E = 0$, the single particle eigenvalue equation (3.111) stands true, summarized as

$$(t + U)\varphi_k = \varepsilon_k \varphi_k \quad (3.124)$$

We can multiply (3.124) on the left by φ_k^* and integrate to get

$$\int -\varphi_k^* \frac{\hbar^2}{2m} \nabla^2 \varphi_k d\mathbf{r} + \int \varphi_k^* U \varphi_k d\mathbf{r} = \int \varphi_k^* \varepsilon_k \varphi_k d\mathbf{r} \quad (3.125)$$

The integral on the right hand side of (3.125) evaluates to ε_k due to the orthonormality constraint. If we sum over all states k we get

$$\sum_k \left\{ \int -\varphi_k^* \frac{\hbar^2}{2m} \nabla^2 \varphi_k d\mathbf{r} + \int \varphi_k^* U \varphi_k d\mathbf{r} \right\} = \sum_k \varepsilon_k \quad (3.126)$$

$$\sum_k t_k + \int \rho U = \sum_k \varepsilon_k \quad (3.127)$$

Since U is calculated as the functional derivative with respect to the density ρ (3.113), assuming that the functional has a power dependence from ρ of the form $\mathcal{E}_{\text{Skyrme}} = A\rho^{\sigma+1}$ as in our case, we get the *rearrangement energy*

$$\rho U = \rho \frac{\delta \mathcal{E}_{\text{Skyrme}}}{\delta \rho} = \rho(1 + \sigma)A\rho^\sigma = (1 + \sigma)A\rho^{\sigma+1} = \mathcal{E}_{\text{Skyrme}} + \sigma \mathcal{E}_{\text{Skyrme}} = \mathcal{E}_{\text{Skyrme}} - \mathcal{E}_{\text{rea}} \quad (3.128)$$

If we explicit ρU in equation (3.127) using (3.128), we get to

$$\sum_k t_k + \int (\mathcal{E}_{\text{Skyrme}} - \mathcal{E}_{\text{rea}}) d\mathbf{r} = \sum_k \varepsilon_k$$

Isolating the Skyrme energy density

$$\int \mathcal{E}_{\text{Skyrme}} d\mathbf{r} = \sum_k (\varepsilon_k - t_k) + \int \mathcal{E}_{\text{rea}} d\mathbf{r} \quad (3.129)$$

and given the total energy of the system from (3.11)

$$E = \sum_k t_k + \frac{1}{2} \int \mathcal{E}_{\text{Skyrme}} d\mathbf{r} \quad (3.130)$$

substituting (3.129) in (3.130) yields

$$E_{\text{HF}} = \frac{1}{2} \sum_k (\varepsilon_k + t_k) + \int \mathcal{E}_{\text{rea}} d\mathbf{r} = \frac{1}{2} \left(T + \sum_k \varepsilon_k \right) + E_{\text{rea}} \quad (3.131)$$

which is called *Hartree-Fock energy*.

Sidenote The actual functional, including the Coulomb exchange term, has different ρ terms, which can be summarized as

$$\mathcal{E}_{\text{Skyrme}} = \sum_j A_j \rho^{\sigma_j+1} \implies E_{\text{rea}} = - \sum_j \sigma_j A_j \rho^{\sigma_j+1}$$

This means that only terms with a $\sigma_j \neq 0, -1$ actually contribute to the rearrangement energy.

Since equation (3.131) is valid only for $\delta E = 0$, it's useful to check its equivalence with the integrated energy at convergence, so one can be sure to actually be in a stationary point.

4 | Numerical methods

This chapter is devoted to describing the practical implementation of the nuclear Hartree-Fock method of this work. In section 4.1, the two partial differential equations (PDEs) of interest, the Kohn-Sham equation and the Poisson equation, are numerically approximated through finite differences. In section 4.2.2, a breakdown of numerical solvers for the large-scale eigenvalue problem posed by the KS equation is presented, to pedagogically illustrate the rationale and implementation of the GCG algorithm. Finally, in section 4.3, the self-consistent calculation is presented, along with the implementation of spatial constraints and the optimization of the numerical parameters of GCG.

4.1. Finite differences

The framework used to numerically solve the relevant PDEs of the problem, is the one of finite differences. The core idea is to discretize the domain on a 3D mesh, use Taylor expansions to approximate differential operators and then solve the resulting system of linear equations.

4.1.1. 3D mesh

The first step of the process is representing the different fields in a numerical, discretized fashion. Generally speaking, we deal at most with 2-rank tensors, which depend on three space coordinates and one spin coordinate.

Discretizing the 3D cartesian space with a 3-index mesh, choosing a box whose size along x , y , z is respectively $[-a_x, a_x]$, $[-a_y, a_y]$, $[-a_z, a_z]$, and a number of points n_x, n_y, n_z , the resulting lattice will be given by

$$V = \{(-a_x + ih_x, -a_y + ih_y, -a_z + ih_z)\} = \{(x_i, y_j, z_k)\}$$

Where the indices and step sizes are

$$\begin{aligned} i = 0, \dots, n_x - 1 \quad h_x &= \frac{2a_x}{n_x - 1} \\ j = 0, \dots, n_y - 1 \quad h_y &= \frac{2a_y}{n_y - 1} \\ k = 0, \dots, n_z - 1 \quad h_z &= \frac{2a_z}{n_z - 1} \end{aligned}$$

The following implementation assumes $a = a_x = a_y = a_z = a$ and $n = n_x = n_y = n_z = n$, without losing generality.

Including the spin degree of freedom, we can finally represent the fields in a numerical way through

$$\varphi(\mathbf{r}, \sigma) \mapsto \varphi(x_i, y_j, z_k, s) = \varphi_{ijks} \quad (4.1)$$

Differential operators discretization

By using Taylor series, it's possible to write approximations to derivatives [65], in any point of the lattice, of any (reasonable) order of accuracy, involving only near neighbouring points. In the present work, 5-points derivatives are used, meaning Taylor expansions are written for $\varphi(x \pm h)$ and $\varphi(x \pm 2h)$ to compute the differential operators. Formulae for first and second derivatives are given in appendix A.2.

From the theory background of chapter 3, we discern two main kinds of PDEs: the Schrödinger-like KS equation, and the Poisson equation.

4.1.2. Schrödinger equation

Starting from the Schrödinger equation (3.111), reported here for clarity

$$\left[-\nabla \left(\frac{\hbar^2}{2m_q^*(\mathbf{r})} \nabla \right) + U_q(\mathbf{r}) + \delta_{q,\text{proton}} U_C(\mathbf{r}) - i\mathbf{B}_q(\mathbf{r}) \cdot (\nabla \times \boldsymbol{\sigma}) \right] \varphi = \varepsilon \varphi$$

it can be compactly written as

$$f(\nabla^2 \varphi, \nabla \varphi, \varphi, \mathbf{r}, s) = \varepsilon \varphi. \quad (4.2)$$

If f is linear in φ , it is possible to rewrite it as a linear combination of the values of φ on the mesh, after which we can use linear algebra methods to solve the problem.

Linearity Breaking down each part of the equation, the kinetic term

$$\nabla \left(\frac{\hbar^2}{2m_q^*(\mathbf{r})} \nabla \right) \varphi = \frac{\hbar^2}{2m_q^*(\mathbf{r})} \nabla^2 \varphi + \nabla \left(\frac{\hbar^2}{2m_q^*(\mathbf{r})} \right) \cdot \nabla \varphi \quad (4.3)$$

is evidently linear in φ .

The spin-orbit term of (3.111), which we write as

$$\begin{aligned} \hat{h}_{\text{SO}} &= -i \mathbf{B}_q(\mathbf{r}) \cdot (\nabla \times \boldsymbol{\sigma}) \\ &= -i [\mathbf{B}_{q,x}(\mathbf{r})(\sigma_z \partial_y - \sigma_y \partial_z) + \mathbf{B}_{q,y}(\mathbf{r})(\sigma_x \partial_z - \sigma_z \partial_x) + \mathbf{B}_{q,z}(\mathbf{r})(\sigma_y \partial_x - \sigma_x \partial_y)] \end{aligned}$$

is also linear in φ .

Finally, the mean field terms U_q, U_c

$$(U_q + \delta_{q,\text{proton}} U_c) \varphi$$

are just multiplicative, hence linear.

Given that the whole equation is linear in φ , we can evaluate it on the chosen mesh, using finite differences to approximate the differential operators, yielding a linear eigenvalue problem of the form

$$\sum_{\alpha=1}^N A_{\alpha\beta} \varphi_\beta = E \varphi_\alpha \quad (4.4)$$

where the shorthand notation $N = 2 \cdot N_x \cdot N_y \cdot N_z$ is used to denote the size of the matrix A , which is $N \times N$.

Boundary conditions

We expect the nucleus to be a localized object, leading to vanishing Dirichlet boundary conditions for the Schrödinger equation. Near the boundaries, the derivatives will involve points outside the box and setting these points to zero, is equivalent to solving

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & A & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \varphi_{-2} \\ \varphi_{-1} \\ \varphi \\ \varphi_N \\ \varphi_{N+1} \end{bmatrix} = E \begin{bmatrix} \varphi_{-2} \\ \varphi_{-1} \\ \varphi \\ \varphi_N \\ \varphi_{N+1} \end{bmatrix} \quad (4.5)$$

From this system of equations, we get for points outside the boundary:

$$\begin{cases} \varphi_{-2} = 0 \\ \varphi_{-1} = 0 \\ \dots \end{cases} \quad (4.6)$$

Meaning that φ outside the box will automatically be set to zero if the A matrix is built assuming those points to be zero when computing its coefficients.

4.1.3. Poisson equation

The other fundamental PDE we need to solve is the Poisson equation encountered in section 3.4. Dropping the c and p subscripts, it reads

$$\nabla^2 V = 4\pi e^2 \rho$$

It's simpler than the Schrödinger equation, as it only involves a Laplacian and it's not an eigenvalue problem. The right side is given, and the solution is found by inverting the coefficients' matrix.

Boundary conditions

Unlike the Schrödinger equation, we do not expect the solution to rapidly decay near the boundaries; as reported in section 3.4, we have fixed, non-vanishing boundary conditions, which we have to properly impose on the system.

We can choose a direction, say x , and look at the discretized equation at the boundaries $x = \pm a$. Since the indices j, k won't vary, we can omit them, and ignore the other derivatives in the following equations.

$$\begin{aligned} \nabla^2 V &= \partial_{xx} V + \partial_{yy} V + \partial_{zz} V \\ &= \frac{-V_{i-2} + 16V_{i-1} - 30V_i + 16V_{i+1} - V_{i+2}}{12h^2} + \dots = 4\pi e^2 \rho_i \end{aligned} \quad (4.7)$$

Near a boundary, say $i = 0$, the formula calls for points outside the box, known as *ghost points*. Since they are not part of the linear system, but they are known, we can bring them on the right side of equation (4.7).

$$\frac{-30V_0 + 16V_1 - V_2}{12h^2} = 4\pi e^2 \rho_0 + \frac{V_{-2} - 16V_{-1}}{12h^2} = \tilde{\rho}_0 \quad (4.8)$$

The same procedure must be applied to all equations involving ghost points, e.g. for $i = 1$

$$\frac{+16V_0 - 30V_1 + 16V_2 - V_3}{12h^2} = 4\pi e^2 \rho_1 + \frac{V_{-1}}{12h^2} = \tilde{\rho}_1. \quad (4.9)$$

The proper system to solve will then be

$$AV = \tilde{\rho} \quad (4.10)$$

Where A is constructed as previously specified. Solving with $\tilde{\rho}$ on the right hand side will force the solution to obey boundary conditions.

On higher order approximations and performance

Higher and higher order approximations for derivatives involve more points that are further away. This increases accuracy by reducing the finite differences error, but it also decreases the matrix sparseness.

4.2. Eigenvalue problem

This section is devoted to the approximate solution of the eigenvalue problem, needed for the Schrödinger equation (3.111).

Eigenvalue problems are ubiquitous in physics and engineering, and while solving one for a small matrix is trivial, it still requires roughly $O(n^3)$ operations [24] to do so. More often than not, real computational applications result in large-scale matrices, which are completely out of question for exact eigenvalues calculations, thus requiring the use of approximate algorithms.

We will begin by describing common building blocks of iterative eigensolvers in section 4.2.1, namely:

- the approximate solutions of linear systems by the use of the Conjugate Gradient method;
- matrix preconditioning to speed up convergence;
- the Rayleigh-Ritz procedure to find good approximations to the eigenpairs in a certain subspace; and
- the shift-and-invert method, to select the desired portion of the eigenvalue spectrum.

After describing these building blocks, some of the most commonly used eigensolvers are described in section 4.2.2, focusing on the core ideas and stating their limitations, to

finally address the General Conjugate Gradient method, whose implementation in the present work is detailed in section 4.2.3.

4.2.1. Conjugate Gradient and numerical techniques

Conjugate Gradient method

Solving linear systems of the form

$$Ax = b \quad (4.11)$$

is crucial in many eigensolvers. The Conjugate Gradient (CG) is perhaps the most famous iterative solver in this sense, especially in connection with sparse matrices, as we will see in a moment. CG applies to cases where A is a real, $n \times n$, positive-definite, symmetric matrix, and x and b are n -dimensional vectors.

Many generalizations to this method exist, which relax the requirements on the matrix, like BiCGSTAB, CGRES and so on [53]. We will describe the working principle of CG, but the same applies to all the others, with slight variations.

Steepest descent method The quadratic form $f(x)$ derived from the system (4.11) is

$$f(x) = \frac{1}{2}x^T Ax - b^T x \quad (4.12)$$

If A is symmetric, positive-definite, the shape of $f(x)$ is convex and has a global minimum for

$$\nabla_x f(x) = Ax_m - b = 0 \implies Ax_m = b, \quad (4.13)$$

hence the extremum of the quadratic form is the also the solution of the linear system (4.11).

We can employ the well-known gradient descent technique [55] to find such point: starting from a guess x_0 , we compute the direction d_i where f decreases the most (the residual r_i), compute the step size that gives the largest decrease, and update x_i at each iteration accordingly, repeating until convergence.

$$d_i = r_i = b - Ax_i \quad (4.14)$$

$$x_{i+1} = x_i + \alpha_i r_i \quad (4.15)$$

$$\text{with } \alpha_i \text{ such that } \frac{df}{d\alpha_i} = 0 \implies \alpha_i = \frac{r_i^T r_i}{r_i^T A r_i} \quad (4.16)$$

This is a powerful but highly inefficient procedure. We are not ensuring that the search direction doesn't end up with components in subspaces that were explored already.

It can be proven [55] that the norm of the error $e_i = x_i - x_m$ is minimal at each iteration if the search directions d_i are chosen to be A -orthogonal to the next error, i.e. $d_i^T A e_{i+1} = 0$. This makes the algorithm converge at the exact solution in n steps, but most importantly it allows to truncate the iterations without a large error on the approximation x_i .

In this case, the algorithm is called Conjugate Gradient Method and is formulated as

$$\alpha_i = \frac{r_i^T r_i}{d_i^T A d_i} \quad (4.17)$$

$$x_{i+1} = x_i + \alpha_i d_i \quad (4.18)$$

$$r_{i+1} = r_i - \alpha_i A d_i \quad (4.19)$$

$$\beta_{i+1} = \frac{r_{i+1}^T r_{i+1}}{r_i^T r_i} \quad (4.20)$$

$$d_{i+1} = r_{i+1} + \beta_{i+1} d_i \quad (4.21)$$

where iterations are truncated if the norm of the residual r_i is smaller than a certain threshold. It can be proven that the orthonormalization of the new search direction, with respect to all the previous ones, can be done only through the rescaling factor β_{i+1} [55]. CG converges to the exact solution in n steps, moreover, it represents a great method for sparse matrices, because it can be proven to be of complexity $O(m)$, where m is the number of non-zero elements in A [55]. In figure 4.1, a visual representation of the conjugation of search directions and the subsequent exact solution is shown for a two-dimensional problem.

Complex matrices Algorithm (4.17) and the CG method in general can be used for complex matrices, under the condition that A is Hermitian and positive-definite when using the complex inner product, meaning that

$$A = A^\dagger \text{ and } x^\dagger A x > 0. \quad (4.22)$$

Preconditioning

The CG method convergence is known to be limited by the modulus of the condition number of A , $\kappa(A)$, given by [55]

$$\kappa(A) = \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)} \quad (4.23)$$




Figure 4.1: Comparison between the steepest descent method, on the left, and the conjugate gradient method, on the right, for a two-dimensional matrix. Ellipses represent contour lines of the quadratic form $f(x) = x^T A x - b^T x$, in the ‘stretched’ space Ax on the right. As shown in the figure, the conjugation of the search directions eliminates components of the error e_i , until the exact convergence in n (2) steps.

where λ_{\max} and λ_{\min} are respectively the largest and smallest eigenvalues of A in magnitude. If we were able to find a good *preconditioner* M , symmetric and positive-definite, such that $\kappa(M^{-1}A) \ll \kappa(A)$, and M^{-1} is easy to compute, then the algorithm would converge much faster, by solving $M^{-1}Ax = M^{-1}b$, since x is also the solution of $Ax = b$.

$$x = (M^{-1}A)^{-1}M^{-1}b = A^{-1}MM^{-1}b = A^{-1}b. \quad (4.24)$$

Without delving into the details of the preconditioner implementation, detailed in [55], note that, in general, $M^{-1}A$ is neither positive-definite nor symmetric, which requires a Cholesky decomposition [34] $M = EE^T$ to be used, so that the problem may be restated with a symmetric positive-definite matrix $E^{-1}AE^{-T}$.

The catch with preconditioning is that M has no unique recipe. Preconditioners are widely spread across numerical analysis, so many methods have been explored and implemented [45].

Rayleigh-Ritz procedure

A common denominator of all these algorithms is the search of good approximations for the correct eigenvectors in a certain subspace. The method is called Rayleigh-Ritz (RR) procedure [53], and is here outlined.

Let us suppose to have a matrix A of size $n \times n$, with entries in \mathbf{C} and a collection of vectors k organized in a matrix K , where K is of size $n \times k$. Generally speaking, n is large, while k is much smaller.

The best approximation of the true eigenvectors of A in the subspace \mathcal{K} spanned by the vectors in K can be computed by solving the small scale eigenvalue problem

$$K^\dagger A K C = C \Lambda. \quad (4.25)$$

Here matrices $K^\dagger A K$ and C are of size $k \times k$. Computing $K C$ gives a matrix of size $n \times k$, whose column vectors are the best approximations of the true eigenvectors of A in the subspace \mathcal{K} , with their corresponding eigenvalues in the entries of the diagonal matrix Λ .

Shift and Invert

The power iteration is the technique on which Krylov subspace search methods are based [24]. By repeatedly applying matrix A to a vector x , x gets skewed towards the eigenvector whose eigenvalue is of largest magnitude λ_n .

Let us assume A is a hermitian matrix, thus diagonalizable. This means we can write an arbitrary vector $x^{(0)}$ as a linear combination of the eigenvectors $\{v_i\}$ of A .

$$x^{(0)} = \sum_i^n \alpha_i v_i \quad (4.26)$$

If we apply A to $x^{(0)}$ k times, we get

$$x^{(k)} = A^k x^{(0)} = \sum_i^n \alpha_i A^k v_i = \sum_i^n \alpha_i \lambda_i^k v_i \quad (4.27)$$

It can be proven that the ratio of the j -th component of $x_j^{(k)}$ and $x_j^{(k-1)}$ converges to λ_n

$$\lim_{k \rightarrow \infty} \frac{x_j^{(k)}}{x_j^{(k-1)}} = \lambda_n \quad (4.28)$$

which means, that for large enough k , we have the relation

$$A x^{(k)} \approx \lambda_n x^{(k)} \quad (4.29)$$

So $x^{(k)}$ is an approximation of the eigenvector v_n of A whose eigenvalue is λ_n .

Smallest eigenvalue If instead of the largest eigenvalue, we were interested in the smallest one – in magnitude – λ_0 , then we would need to apply the inverse matrix A^{-1} to $x^{(k)}$, which would change the ratio (4.28) to

$$\lim_{k \rightarrow \infty} \frac{x_j^{(k)}}{x_j^{(k-1)}} = \lambda_0 \quad (4.30)$$

Let us assume for a moment that we're solving a nuclear single-particle Hamiltonian, where we have a certain number of bound states of negative energy and a much larger number of unbound states with positive energy. In this case, the inverse power iteration would converge to the states whose energy is closer to zero, avoiding the interesting ones on the bottom of the spectrum.

The solution is, before inverting, to shift the matrix by a quantity σ that is very close to the lowest eigenvalue we want to compute, call it λ_σ (eigenvector v_σ). Now, the eigenvalue of lowest magnitude of $(A - \sigma I)$ is $\lambda_\sigma - \sigma$ and by applying $(A - \sigma I)^{-1}$ to $x^{(k)}$, we will get the approximation to the eigenvector v_σ . This is the procedure implemented in step 8 of algorithm 4.3.

4.2.2. Iterative eigensolvers

Now that the main techniques used by iterative eigensolvers have been laid out, we can look at three general methods, which are the most commonly used ones.

Jacobi-Davidson

The Jacobi-Davidson method is a type of algorithm where at each iteration, the approximation to an eigenpair of matrix A , is improved by correcting the eigenvector through the solution of a certain linear system, as we shall see shortly.

Given an approximation (u, θ) of an eigenpair of matrix A , where u is the approximate eigenvector and θ is the approximate eigenvalue, if the residual

$$r = Au - \theta u \quad (4.31)$$

is ≈ 0 , then the eigenpair converged. Otherwise, we want to find a correction t such that

$$r = A(u + t) - (\theta + \delta\theta)(u + t) = 0 \quad (4.32)$$

Linearizing this equation in t gives

$$(A - \theta I)t = -r \quad (4.33)$$

To avoid singularity of the equation near convergence, since u approximately spans a subspace of the system's kernel $\ker(A - \theta I)$, and enrich the subspace search with a useful orthogonal correction, we project the problem onto the orthogonal subspace of u , which finally gives

$$(I - uu^\dagger)(A - \theta I)(I - uu^\dagger)t = -r \quad (4.34)$$

Although simple, this method is computationally efficient only by using preconditioning,

Algorithm 4.1 Jacobi-Davidson method

- 1: Choose normalized initial vectors $\{u_k\}$, set $V = [u_1, \dots, u_k]$
 - 2: **repeat**
 - 3: Compute eigenpair: $T = V^\dagger AV$, solve $Ty = \theta y$
 - 4: Set $u = Vy$, residual $r = Au - \theta u$
 - 5: **if** $\|r_k\| < \varepsilon \ \forall k$ **then**
 - 6: **return** (θ, u)
 - 7: **end if**
 - 8: Solve approximately $(I - u_k u_k^\dagger)(A - \theta I)(I - u_k u_k^\dagger)t_k = -r_k$ using preconditioned iterative solver, ensuring $t_k \perp u_k$
 - 9: Normalize: $v_k = t_k / \|t_k\|$
 - 10: Expand subspace, setting $V = [V, v]$
 - 11: **until** convergence for $k = 1, \dots, \text{nev}$
-

which is known to be unstable in many cases [53].

Lanczos

Lanczos algorithm [35] is probably the most used iterative eigensolver for hermitian matrices. It's a Krylov subspace search method, meaning the Rayleigh-Ritz procedure is done on a subspace formed as

$$\mathcal{K} = \{v_1, Av_1, A^2v_1, \dots, A^{k-1}v_1\} \quad (4.35)$$

which exploits the power iteration. After orthogonalizing the new approximation to the previous one and diagonalizing the small scale problem, we end up with the new best approximations to the eigenvectors of A . Lanczos is extremely efficient, memory- and CPU-wise for extremal eigenvalues, but this limits its applicability, as one may be interested in the inner portion of the eigenvalue spectrum, such in the case of Hartree-Fock-Bogoliubov

Algorithm 4.2 Lanczos Method

```

1: Choose normalized initial vector  $v_1$ , set  $\beta_0 = 0$ ,  $m =$  subspace size.
2: repeat
3:   for  $j = 1, 2, \dots, m$  do
4:      $w \leftarrow Av_j - \beta_{j-1}v_{j-1}$ 
5:      $\alpha_j \leftarrow v_j^* w$ 
6:      $w \leftarrow w - \alpha_j v_j$ 
7:      $\beta_j \leftarrow \|w\|$ 
8:     if  $\beta_j = 0$  then
9:       break
10:    end if
11:     $v_{j+1} \leftarrow w/\beta_j$ 
12:  end for
13:  Form tridiagonal matrix  $T_m = \text{tridiag}(\beta_{1:m-1}, \alpha_{1:m}, \beta_{1:m-1})$ 
14:  Compute eigen-decomposition  $T_m y_k = \theta_k y_k$ , for  $k = 1, \dots, \text{nev}$ 
15:  Form Ritz approximations  $x_k = V_m y_k$ , where  $V_m = [v_1, \dots, v_m]$ 
16:  Compute residual norms  $r_k = \|Ax_k - \theta_k x_k\|$  for all  $k$ 
17: until convergence for  $k = 1, \dots, \text{nev}$ 

```

(HFB).

A shift-and-invert strategy would be unfeasible in the case of large scale problems, since all Lanczos steps need to be performed exactly to avoid instabilities, a well known problem in the Arnoldi generalization [53].

LOBPCG

The last algorithm of this short list is LOBPCG, it's the newest and most sophisticated one of the three.

Introduced by A. V. Knyazev in 1991 [31], it's a block, preconditioned conjugate gradient method, explicitly targeted at solving large-scale eigenvalue problems, and it has been used in modern solutions of the Schrödinger/KS equation in recent years [36, 38, 44, 64]. We won't go into the details of LOBPCG, since GCG shares with it many aspects, like blocking and search directions calculation.

LOBPCG works very well for large scale problems, but it has limitations. On the one hand, it's not possible to arbitrarily select the portion of the matrix spectrum to calculate, which is required for problems where variational collapse happens, like in HFB or the Dirac equation, which manifests particle/antiparticle solutions [36]. To solve this, an additional filtering step is required [36, 38], which introduces a computational cost in the algorithm. Lastly, LOBPCG may fail when poor conditioning is present or when high precision on the eigenvalues is required [37].

4.2.3. General Conjugate Gradient

The General Conjugate Gradient is an iterative eigensolver designed with the aim of improving LOBPCG, it is a blocked algorithm, which uses the inverse power method and previous search directions to generate the search subspace. GCG is proven to be faster and more stable than LOBPCG [37].

The search subspace is built as

$$V = [X, P, W], \quad (4.36)$$

where X , of dimensions $n \times k$ is the matrix containing the approximations, to the eigenvectors of matrix A , P , of dimensions $n \times k$ is the matrix containing the previous search directions, and W , of dimension $n \times a$ is the matrix containing the eigenvectors on which the inverse power method is applied approximately using the CG.

A slightly different implementation of the algorithm is employed in the present work, detailed in algorithm 4.3, to improve applicability to HF calculations and reduce the computational cost.

Eigenvalue problem The original algorithm aims at solving the general eigenvalue problem $AX = \lambda BX$. Since in our case $B = I$, it is omitted from the procedure, reducing the computational cost of the algorithm, in particular, the one of the search direction block P . After orthonormalization of V , the columns of X are orthonormal as well and the calculation of P is given by

$$P = X_{\text{new}} - X(X^\dagger X_{\text{new}}) \quad (4.37)$$

which is the projection of X_{new} onto the orthogonal complement of X , used in step 12 of algorithm 4.3.

Complex matrix The algorithm has been generalized to the complex case, where the matrix is complex Hermitian and, as such, the transposition operation is replaced by the conjugate transpose.

Blocking The algorithm is designed to allow blocking of the eigenvectors, such that $X = [X_c, X_a, X_r]$, where X_c are the converged eigenvectors, X_a are the active eigenvectors on which we perform the inverse power iteration, with $\text{col}(X_a)$ being a fixed number, and X_r are the remaining eigenvectors, which are to be inserted in X_a as soon as some of its columns converge. This allows to save some computations by avoiding the expensive inverse power on pairs that have already converged. Since in a self-consistent calculation

the matrix changes rapidly at each HF iteration, it is the case that the maximum number of iterations is reached before convergence of all eigenpairs, so we must work at all times on the remaining unconverged eigenvectors. For this reason, we only implement the $X = [X_c, X_a]$ scheme, where the only distinction we make is between converged eigenvectors X_c and unconverged ones X_a .

Orthogonalization The original paper [37] suggests an improved orthogonalization procedure; being beyond the scope of this work, the simpler Gram-Schmidt [7] orthogonalization is used in the present work.

Shift update The shift update is either fixed, in case of known spectrum, eg for HFB $\text{shift} = 0$, or adaptive [37], so that the inverse power step can find the lowest eigenvalues, using the update formula

$$\text{shift} = (\lambda_{\text{nev}} - 100\lambda_1)/99 \quad (4.38)$$

where λ_{nev} is the biggest eigenvalue of the RR procedure and λ_1 is the smallest of the active eigenpairs.

Preconditioning The use of a preconditioner is beyond the scope of this work, a simple diagonal preconditioner is used.

Algorithm 4.3 General Conjugate Gradient algorithm

- 1: **Input:** Matrix A of dimensions $n \times n$, number of desired eigenpairs **nev**, X_{guess} initial guess of dimensions $n \times k$, with $k \geq \text{nev}$, **max_iter** maximum iterations
 - 2: Initialize block $X = [X_a] \leftarrow X_{\text{guess}}$ and X_c as an empty block
 - 3: Initialize blocks P and W as empty blocks
 - 4: Solve the Rayleigh Ritz problem $X^\dagger A X C = C \Lambda$
 - 5: Update $X = X C$
 - 6: Initialize **shift**, Initialize **iter** = 0
 - 7: **while** $\text{col}(X_c) < \text{nev}$ and **iter** < **max_iter** **do**
 - 8: Solve approximately $(A + \text{shift} \cdot I)W = X_a \Lambda$ with some CG steps, initial value X_a to generate W as an approximation to $W = (A + \text{shift} \cdot I)^{-1} X_a \Lambda$
 - 9: Orthogonalize $V = [X, P, W]$, of dimensions $n \times (k + k + a)$
 - 10: Solve the Rayleigh Ritz problem $V^\dagger (A + \text{shift} \cdot I) V C = C \Lambda$
 - 11: Update X_{new} with the first **nev** columns of $V C$ and $\Lambda_{\text{new}} = \Lambda - \text{shift} \cdot I$
 - 12: Compute $P = X_{\text{new}} - X(X^\dagger X_{\text{new}})$
 - 13: Compute the residual $R = A X_{\text{new}} - \Lambda X_{\text{new}}$
 - 14: Check convergence on k -th column norm of R , if $\|R_k\| < \text{tol}$, move $X_{a,k}$ to X_c .
 - 15: Update **shift** using formula (4.38) and **iter** \leftarrow **iter** + 1
 - 16: **end while**
 - 17: **Output:** Approximate eigenpairs (Λ, X)
-

4.3. Code implementation details

In this last section regarding numerical methods, some important features about the actual code implementation of the HF method are discussed. Mainly, the implementation of the Augmented Lagrangian Method to enforce spatial constraints on the HF solution, the pseudocode of the entire self-consistent procedure, and the choice of optimal parameters for the functional minimization.

4.3.1. Constraints

The purpose of spatial constraints is to find the minimum of the energy functional under the condition that the expectation value of a given operator \mathcal{Q} equals a prescribed target value q_0 . Constrained calculations are a fundamental tool to assess the stability of the ground-state minimum and to investigate dynamical properties of the nucleus, such as fission barriers [8].

Constraints can be formulated as an equality-constrained optimization problem (ECP), formulated as

$$\min_{|\Psi\rangle} E \quad (4.39)$$

$$\text{constrained to } \langle \Psi | \mathcal{Q} | \Psi \rangle = \langle \mathcal{Q} \rangle = q_0 \quad (4.40)$$

Which yields the Lagrangian

$$E' = E + \lambda(\langle \mathcal{Q} \rangle - q_0) \quad (4.41)$$

where λ is a Lagrange multiplier determined by the condition $\langle \mathcal{Q} \rangle = q_0$. After finding the minimum of E' , it's trivial to show that for a given λ , we get [22]

$$\frac{dE}{d\langle \mathcal{Q} \rangle} = -\lambda. \quad (4.42)$$

From a numerical standpoint, λ needs to be tuned at each iteration to reach the desired value of q_0 . This method was the one used in early constrained Hartree-Fock calculations [17].

Although this method is simple, it often fails. Moreover, for the same value of λ , many, possibly infinite values of $\langle \mathcal{Q} \rangle$ can be obtained, for which we are only allowed to get the one with the most stable solution.

A different method is provided by the Quadratic Penalty Method (QPM). Briefly speaking, instead of a Lagrange multiplier, we add a quadratic contribution to the functional, such that

$$E' = E + \frac{c}{2}(\langle \mathcal{Q} \rangle - q_0)^2. \quad (4.43)$$

This is a straightforward method; intuitively one penalizes (hence the name), any solution for which $\langle \mathcal{Q} \rangle \neq q_0$ by increasing its energy. However, the success of such procedure is heavily influenced by the choice of c , often leading to instabilities for large values.

What happens is that for small values of c , the penalty may be insufficient to reach the target q_0 , while for bigger values, the penalty may be so strong that the self-consistent calculation oscillates and fails.

Augmented Lagrangian Method

A modern, robust approach, used by HF/HFB codes [12, 50] is given in the form of the Augmented Lagrangian Method (ALM) [57]. Its main idea is to combine the precision of the ECP with the accuracy of the QPM.

Without delving into cumbersome mathematical details, we'll see how the algorithm is practically implemented in the code.

Given the functional

$$E' = E + \lambda(q - q_0) + \frac{c}{2}(q - q_0)^2 \quad (4.44)$$

where $q = \langle \mathcal{Q} \rangle$, the resulting mean field potential will be given by

$$U' = U + \lambda \mathcal{Q} + c(q - q_0) \mathcal{Q} \quad (4.45)$$

$$= U + c(q - q_0(\lambda)) \mathcal{Q} \quad (4.46)$$

where $q_0(\lambda)$ is updated at each iteration with the formula

$$q_0(\lambda) = q_0 - \frac{\lambda}{c} \quad (4.47)$$

$$\lambda^{(i+1)} = \lambda^{(i)} + \mu c(q - q_0) \quad (4.48)$$

Here, a slight deviation from the original ALM is present. Since the original work [57] doesn't provide guidance regarding what is considered an *iteration*, we employ the strategy [12] of using a damping factor $\mu \in [0, 1]$, so λ can be updated at each HF iteration for fast convergence, without large oscillations or instabilities of any kind.

This method is what powers the deformation curves that are shown in chapter 6, allowing to explore the energy surface with arbitrary precision in reaching the value of q_0 at convergence, provided that enough HF iterations are performed.

Note that, since $\lambda^{(0)} = 0$, for $\mu = 0$ ALM reduces to the standard QPM.

4.3.2. Details on the implementation of the code

The whole Hartree-Fock framework presented up to this point, has been implemented using the C++ language [60] and the Eigen linear algebra library [26], which implements linear algebra operations through low level routines such as LAPACK and BLAS. In figure 4.2, the schematics of the program structure is reported.

4.3.3. Optimal parameters choice

Inside the ‘Diagonalize h ’ step in figure 4.2, the execution of the GCG algorithm is performed, using the current iteration’s single-particle Hamiltonian as the matrix to diagonalize and the previous iteration’s orbitals as the initial guess. This is the main computational bottleneck of the code, where a correct choice of the execution parameters can drastically reduce execution times. The parameters that need to be chosen carefully are essentially the inverse power step tolerance and the number of maximum GCG iterations.

Inverse power step tolerance

The first parameter to be tuned is the tolerance on the CG step which approximately solves the system

$$(A + \text{shift} \cdot I)W = X_a \Lambda \quad (4.49)$$

in algorithm 4.3, where A is actually the single-particle Hamiltonian h . When the CG residual $(A + \text{shift} \cdot I)W - X_a \Lambda$ is smaller than the tolerance, the procedure stops and outputs the W block.

In figure 4.3, the relative absolute error of the total energy is calculated against a reference benchmark value (details in the results chapter 5), for different values of the CG tolerance. It’s clear that at least a tolerance of 10^{-3} is needed for good convergence, while tolerances $\geq 10^{-4}$ stop offering increasing returns, rendering a choice between 10^{-4} and 10^{-5} an optimal one.

Inner GCG iterations

The number of inner GCG maximum iterations, here named ‘inverse power steps’ to avoid confusion, is slightly more nuanced than the CG tolerance. The algorithm converges to the true eigenpairs as the power steps are performed, so one could think that a higher number of steps would bring to HF convergence faster, since the precision on the eigenvalues increases, but this is not the case.

In figures 4.4 and 4.5, the convergence of the HF calculation is plotted for different number




Figure 4.2: Pseudocode of the Hartree-Fock program.

of steps, respectively, for the spherical nucleus ^{16}O and the deformed nucleus ^{24}Mg . It's evident that in both cases, a steps number greater than 3 leads to oscillating behaviour near convergence, without accelerating it, while in the case of the spherical nucleus, just one step is enough to quickly, and reliably reach convergence. In any case, it's clear that delaying the inverse power steps to later HF iterations is safer in terms of stability.




Figure 4.3: HF calculation convergence with varying CG tolerance for ^{16}O , box $[-9, 9]$ fm, step size 0.3 fm.

This counter intuitive behavior is likely due to the fact that at each HF iteration the hamiltonian changes and a great number of steps leads to solutions too biased towards the current matrix eigenpairs, at the expense of the next iteration; however, in the case of deformed nuclei, due to sharp shape changes at the start of the calculation, just one step may not be enough to sustain the pace at which the Hamiltonian changes, hence the quicker convergence with more steps.

4.3.4. Numerical stability

As a final remark, the numerical stability of the solver is reported in figure 4.6. The map is produced for a spherical calculation of ^{16}O , with varying box and mesh sizes.

It's possible to observe that for a box whose side is at least ≈ 2.5 times the nuclear radius, the solver numerical stability is loosely dependent on the box extension, but rather on the step size. This is not surprising, as the points separation in space h dictates the precision of the discretized derivatives, as mentioned in section 4.1.




Figure 4.4: HF calculation convergence with varying number of inverse power steps for ^{16}O , box $[-9, 9]$ fm, step size 0.3 fm.




Figure 4.5: HF calculation convergence with varying number of inverse power steps for the deformed nucleus ^{24}Mg , box $[-10, 10]$ fm, step size 0.33 fm.




Figure 4.6: Numerical stability map of the HF solver for ^{16}O for different box and step sizes. Relative error is taken against a benchmark reference value.

5 | Results for spherical nuclei

In this chapter, results for spherical nuclei are presented. These are mostly calculated for double magic nuclei, with the exception of ^{90}Zr . The reason behind choosing spherical nuclei as an initial benchmark, is that numerous spherical HF codes are available and they have the advantage of being one-dimensional, which allows the use of very fine meshes, with a step size that can go down to the physical scale of the problem, which is roughly 0.1 fm in this case, without any major computational limit. We can use these codes as a reference ideal value for the different quantities produced by our code. The choice for benchmarking spherical results in the present work has been the `hfbcs_qrpa` code [15]. This chapter is structured as follows. In section ??, some physical properties of the system are presented, such as the mean square radii and the deformation parameters, which we can use to numerically validate the code. In section ??, benchmark results for ^{16}O are presented, while in section ?? results for the heavier nuclei ^{48}Ca , ^{56}Ni , and ^{90}Zr are shown.

5.1. Physical quantities

After finding a nuclide's ground state, we are able to compute different physical properties of the system. We can use these values as a numerical reference when comparing our results with other codes.

5.1.1. Mean square radii

An important set of quantities characterizing the nuclear density is certainly the one of mean square radii. The individual nuclear species' mean square radius is defined as

$$\langle r_q^2 \rangle = \frac{\int \rho_q(\mathbf{r}) r^2 d\mathbf{r}}{\int \rho_q(\mathbf{r}) d\mathbf{r}}. \quad (5.1)$$

While the charge mean square radius formula is derived from the convolution of the neutron and proton particle densities with their respective internal charge distribution

[5], resulting in equation (5.2).

$$\langle r_{ch}^2 \rangle = \langle r_p^2 \rangle + \langle r^2 \rangle_P + \frac{N}{Z} \langle r^2 \rangle_N + \frac{2}{Z} \left(\frac{\hbar}{mc} \right)^2 \sum_{\alpha q} \mu_q \langle \boldsymbol{\sigma} \cdot \boldsymbol{\ell} \rangle_{\alpha q} \quad (5.2)$$

where q runs over the nuclear species and α runs over all single particle states of species q . $\boldsymbol{\sigma}$ is the vector operator of Pauli matrices, while $\boldsymbol{\ell}$ is the angular momentum operator $-i(\mathbf{r} \times \nabla)$. $\langle r^2 \rangle_P$ and $\langle r^2 \rangle_N$ refer to the square charge radii of the proton and the neutron, while μ_q to their respective magnetic dipole moment in units of nuclear magneton.

All square charge radii computed in this work use the set of parameters in table 5.1, taken to be equal to the spherical benchmark code `hfbcs_qrpa`.

Parameter	Value	Units
$\langle r^2 \rangle_P$	0.64	fm ²
$\langle r^2 \rangle_N$	-0.11	fm ²
μ_p	2.792847	-
μ_n	-1.913043	-

Table 5.1: Parameters used to compute the charge mean square radius.

5.1.2. Deformation parameters

When dealing with deformed nuclei, mean square radii are not sufficient to characterize the nuclear density. The main parameter used is the quadrupole deformation parameter β_2 , similar to the one encountered in section 1.3, it can be computed through the actual mean square radius with formula (5.3)

$$\beta_2 = \frac{4\pi \langle Y_{20} \rangle}{3A \langle r^2 \rangle} \quad (5.3)$$

where $\langle r^2 \rangle$ is the total mean square radius of the nucleus

$$\langle r^2 \rangle = \frac{\int (\rho_n + \rho_p) r^2 d\mathbf{r}}{\int (\rho_n + \rho_p) d\mathbf{r}} = \langle x^2 + y^2 + z^2 \rangle. \quad (5.4)$$

For spherical nuclei, $\beta_2 = 0$, while for deformed ones, thanks to the normalization with respect to the total radius and mass, the β_2 parameter can be used to compare different nuclei across the nuclide chart.

5.2. Parameters and mesh choice

All `hfbcs_qrpa` calculations were performed using a mesh size of 0.1 fm, no pairing interaction, and a radial mesh size whose radius is equal to the side of the box in our computation. The lattice of our code depends on the extension of the nucleus, which is directly determined by its mass A ; since the number of subdivisions that allows reasonable CPU times on a laptop caps around 60 – 70, step sizes vary across different calculations. In the results shown here, for ^{16}O , we are able to reach a 0.3 fm step size, while for the heaviest, ^{90}Zr , we are only able to reach 0.42 fm. The reason behind this choice is that as the nucleus size increases, a bigger box is needed to ensure that all relevant states are able to decay to zero at the boundary. All the data reported in this chapter is computed with the SLy5 parametrization [11].

5.3. Results for ^{16}O

The first results we will take a look at are the ones for ^{16}O . It's the best candidate for gauging the solver's performance, as it is a very light, double magic nucleus, meaning it has no pairing interaction and a spherical shape.

All calculations are performed on a box of size $[-9, 9]$ fm in all three directions and a step size of 0.3 fm, corresponding to $2 \cdot 60^3$ mesh points.

5.3.1. Results neglecting Coulomb interaction

Since the Skyrme functional is complex and nuanced, results are shown for more and more terms in expression (3.101). We start by including only C_0^p , C_1^p , C_0^τ , C_1^τ and neglecting the others and the Coulomb interaction; results are reported in table 5.2. Without further terms, the spin-orbit field $\mathbf{B}(\mathbf{r})$ vanishes, hence the $1p_{3/2}$ and $1p_{1/2}$ levels show degeneration in energy.

Since $N = Z$, assuming equal masses the single-particle equations will be exactly equal between the two species, therefore only neutron results are reported. Note that C_1 terms reduce to being null in this case, that is until we either break the $N = Z$ equality or introduce the Coulomb interaction. In table 5.3 the $C_0^{\nabla \cdot \mathbf{J}}$ and $C_1^{\nabla \cdot \mathbf{J}}$ terms are included just for the spin-orbit field $\mathbf{B}(\mathbf{r})$, but not for the mean field $U(\mathbf{r})$; from an interaction point of view, it's as if we were neglecting the spin-gradient coupling term [11]. As expected, the $1p_{3/2}$ and $1p_{1/2}$ degeneration is removed, displaying the spin-orbit splitting, which lowers the total angular momentum $j = 3/2$ level and raises the $j = 1/2$ level.

Lastly, the $C_0^{\nabla \cdot \mathbf{J}}$ and $C_1^{\nabla \cdot \mathbf{J}}$ terms are also included in the calculation of the mean-field,

Physical quantities					
		GCG	hfbcq_qrpa	Δ	$\Delta\%$
E_{TOT}	[MeV]	-141.582	-141.582	-	-
$\langle r_n^2 \rangle^{1/2}$	[fm]	2.6504	2.6510	0.0006	2.26×10^{-2}
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	2.7486	2.7491	0.0005	1.82×10^{-2}

Neutron energy levels					
		GCG	hfbcq_qrpa	Δ	$\Delta\%$
$1s_{1/2}$	[MeV]	-36.142	-36.139	0.003	8.30×10^{-3}
$1p_{3/2}$	[MeV]	-18.573	-18.572	0.001	5.38×10^{-3}
$1p_{1/2}$	[MeV]	-18.573	-18.572	0.001	5.38×10^{-3}

Table 5.2: ^{16}O including C_0^ρ , C_1^ρ , C_0^τ , C_1^τ terms, neglecting Coulomb interaction.

resulting in the full implementation of the Skyrme functional. As shown in table 5.4, the effect of this addition on the ground state is little, as the spin current $J_{\mu\nu}$ is small in light, closed shell nuclei.

Results including Coulomb interaction

As the final addition to get a complete and accurate description of ^{16}O , the Coulomb interaction is included as detailed in section 3.4. Results are shown in table 5.5.

As shown in tables 5.2, 5.3, 5.4, and 5.5 results for ^{16}O are in great agreement with the output of the `hfbcq_qrpa` code for all the terms in the Skyrme functional.

5.4. Results for heavier nuclei

In the following section, results for some spherical nuclei heavier than ^{16}O are presented in tables 5.6, 5.7 and 5.8. Our code still shows good agreement with the `hfbcq_qrpa` one. A slight increase of the numerical error can be observed as the step size increases, which is compatible with the polynomial error in the finite difference method.

5.4.1. Comparison with experimental binding energies

The Skyrme functional is highly successful at producing theoretical values in great accordance with experimental data, just by fitting a small set of parameters [3]. In table 5.9, binding energies of some of the nuclei studied in this work are compared with experimental values, taken from the Atomic Mass Data Center [1].

Physical quantities					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
E_{TOT}	[MeV]	-142.080	-142.080	-	-
$\langle r_n^2 \rangle^{1/2}$	[fm]	2.6516	2.6516	-	-
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	2.7497	2.7497	-	-
Neutron energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
$1s_{1/2}$	[MeV]	-36.314	-36.312	0.002	5.5×10^{-3}
$1p_{3/2}$	[MeV]	-20.696	-20.696	-	-
$1p_{1/2}$	[MeV]	-14.335	-14.335	-	-

Table 5.3: ^{16}O including C_0^ρ , C_1^ρ , C_0^τ , C_1^τ , $C_0^{\nabla \cdot J}$, $C_1^{\nabla \cdot J}$ terms, neglecting Coulomb interaction and J^2 terms.

Physical quantities					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
E_{TOT}	[MeV]	-142.074	-142.074	-	-
$\langle r_n^2 \rangle^{1/2}$	[fm]	2.6515	2.6516	0.0001	3.77×10^{-3}
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	2.7497	2.7497	-	-
Neutron energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
$1s_{1/2}$	[MeV]	-36.309	-36.308	0.001	2.75×10^{-3}
$1p_{3/2}$	[MeV]	-20.684	-20.685	0.001	4.83×10^{-3}
$1p_{1/2}$	[MeV]	-14.361	-14.361	-	-

Table 5.4: ^{16}O neglecting Coulomb interaction.

Physical quantities					
		GCG	hfbc _s _qrpa	Δ	$\Delta\%$
E_{TOT}	[MeV]	-128.402	-128.400	0.002	1.56×10^{-3}
$\langle r_n^2 \rangle^{1/2}$	[fm]	2.6584	2.6585	0.0001	3.76×10^{-3}
$\langle r_p^2 \rangle^{1/2}$	[fm]	2.6835	2.6836	0.0001	3.73×10^{-3}
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	2.7805	2.7803	0.0002	7.19×10^{-3}
Neutron energy levels					
		GCG	hfbc _s _qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-36.140	-36.137	0.003	8.30×10^{-3}
1p _{3/2}	[MeV]	-20.611	-20.611	-	-
1p _{1/2}	[MeV]	-14.427	-14.428	0.001	6.93×10^{-3}
Proton energy levels					
		GCG	hfbc _s _qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-32.349	-32.345	0.004	1.24×10^{-2}
1p _{3/2}	[MeV]	-17.137	-17.137	-	-
1p _{1/2}	[MeV]	-11.081	-11.082	0.001	9.02×10^{-3}

Table 5.5: ^{16}O complete of the Skyrme functional and Coulomb interaction.

Physical quantities					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
E_{TOT}	[MeV]	-415.955	-415.931	0.024	5.77×10^{-3}
$\langle r_n^2 \rangle^{1/2}$	[fm]	3.6106	3.6110	0.0004	1.11×10^{-2}
$\langle r_p^2 \rangle^{1/2}$	[fm]	3.4502	3.4507	0.0005	1.45×10^{-2}
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	3.5274	3.5060	0.0214	0.610

Neutron energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-49.758	-49.752	0.006	1.21×10^{-2}
1p _{3/2}	[MeV]	-35.952	-35.949	0.003	8.34×10^{-3}
1p _{1/2}	[MeV]	-33.891	-33.891	-	-
1d _{5/2}	[MeV]	-22.170	-22.169	0.001	4.51×10^{-3}
2s _{1/2}	[MeV]	-17.720	-17.720	-	-
1d _{3/2}	[MeV]	-17.431	-17.434	0.003	1.72×10^{-2}
1f _{7/2}	[MeV]	-9.262	-9.261	0.001	1.08×10^{-2}

Proton energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-45.936	-45.930	0.006	1.31×10^{-2}
1p _{3/2}	[MeV]	-34.314	-34.311	0.003	8.74×10^{-3}
1p _{1/2}	[MeV]	-30.482	-30.483	0.001	3.28×10^{-3}
1d _{5/2}	[MeV]	-22.455	-22.454	0.001	4.45×10^{-3}
2s _{1/2}	[MeV]	-16.753	-16.751	0.002	1.19×10^{-2}
1d _{3/2}	[MeV]	-15.337	-15.340	0.003	1.96×10^{-2}

Table 5.6: ^{48}Ca , box size [-12, 12] fm, step size 0.34 fm

Physical quantities					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
E_{TOT}	[MeV]	-482.805	-482.700	0.105	2.18×10^{-2}
$\langle r_n^2 \rangle^{1/2}$	[fm]	3.6422	3.6433	0.0011	3.02×10^{-2}
$\langle r_p^2 \rangle^{1/2}$	[fm]	3.6968	3.6979	0.0011	2.97×10^{-2}
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	3.7722	3.7682	0.0040	0.106

Neutron energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-54.277	-54.260	0.017	3.13×10^{-2}
1p _{3/2}	[MeV]	-41.571	-41.562	0.009	2.16×10^{-2}
1p _{1/2}	[MeV]	-39.613	-39.611	0.002	5.05×10^{-3}
1d _{5/2}	[MeV]	-28.536	-28.530	0.006	2.10×10^{-2}
2s _{1/2}	[MeV]	-23.539	-23.545	0.006	2.55×10^{-2}
1d _{3/2}	[MeV]	-23.367	-23.361	0.006	2.57×10^{-2}
1f _{7/2}	[MeV]	-16.019	-16.018	0.001	6.24×10^{-3}

Proton energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-43.754	-43.740	0.014	3.20×10^{-2}
1p _{3/2}	[MeV]	-31.561	-31.555	0.006	1.90×10^{-2}
1p _{1/2}	[MeV]	-29.545	-29.545	-	-
1d _{5/2}	[MeV]	-19.017	-19.016	0.001	5.26×10^{-3}
2s _{1/2}	[MeV]	-14.004	-14.012	0.008	5.71×10^{-2}
1d _{3/2}	[MeV]	-13.891	-13.887	0.004	2.88×10^{-2}
1f _{7/2}	[MeV]	-6.934	-6.935	0.001	1.44×10^{-2}

Table 5.7: ^{56}Ni , box size [-13, 13] fm, step size 0.37 fm

Physical quantities					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
E_{TOT}	[MeV]	-783.587	-783.325	0.262	3.34×10^{-2}
$\langle r_n^2 \rangle^{1/2}$	[fm]	4.2854	4.2872	0.0018	4.20×10^{-2}
$\langle r_p^2 \rangle^{1/2}$	[fm]	4.2196	4.2212	0.0016	3.79×10^{-2}
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	4.2767	4.2704	0.0063	0.148

Neutron energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-55.636	-55.615	0.021	3.78×10^{-2}
1p _{3/2}	[MeV]	-45.324	-45.309	0.015	3.31×10^{-2}
1p _{1/2}	[MeV]	-44.172	-44.160	0.012	2.72×10^{-2}
1d _{5/2}	[MeV]	-34.148	-34.137	0.011	3.22×10^{-2}
2s _{1/2}	[MeV]	-31.393	-31.391	0.002	6.37×10^{-3}
1d _{3/2}	[MeV]	-29.802	-29.797	0.005	1.68×10^{-2}
1f _{7/2}	[MeV]	-22.755	-22.748	0.007	3.08×10^{-2}
2p _{3/2}	[MeV]	-17.837	-17.840	0.003	1.68×10^{-2}
1f _{5/2}	[MeV]	-17.568	-17.563	0.005	2.85×10^{-2}
2p _{1/2}	[MeV]	-15.729	-15.723	0.006	3.82×10^{-2}
1g _{9/2}	[MeV]	-11.586	-11.580	0.006	5.18×10^{-2}

Proton energy levels					
		GCG	hfbcs_qrpa	Δ	$\Delta\%$
1s _{1/2}	[MeV]	-44.973	-44.956	0.017	3.78×10^{-2}
1p _{3/2}	[MeV]	-36.347	-36.336	0.011	3.03×10^{-2}
1p _{1/2}	[MeV]	-34.121	-34.115	0.006	1.76×10^{-2}
1d _{5/2}	[MeV]	-26.766	-26.759	0.007	2.62×10^{-2}
2s _{1/2}	[MeV]	-22.175	-22.178	0.003	1.35×10^{-2}
1d _{3/2}	[MeV]	-21.216	-21.214	0.002	9.43×10^{-3}
1f _{7/2}	[MeV]	-16.722	-16.718	0.004	2.39×10^{-2}
2p _{3/2}	[MeV]	-10.239	-10.236	0.003	2.93×10^{-2}
1f _{5/2}	[MeV]	-9.613	-9.618	0.005	5.20×10^{-2}
2p _{1/2}	[MeV]	-8.108	-8.104	0.004	4.94×10^{-2}

Table 5.8: ^{90}Zr , box size $[-15, 15]$ fm, step size 0.43 fm

	^{16}O	^{48}Ca	^{56}Ni	^{90}Zr
E_{th}	128.40	415.95	482.80	783.59
E_{exp}	127.62	414.33	483.99	783.89

Table 5.9: Comparison of experimental binding energies in MeV with theoretical calculated values using the SLy5 functional.

6 | Results for deformed nuclei

Having established that the code works well for spherical nuclei, we can start treading in deformation territory.

6.1. ^{24}Mg

In the following section, results for ^{24}Mg are presented, it's a natural choice to study how well deformations are represented by our framework, since it's light, very deformed and shows no pairing interaction in its ground state.

6.1.1. HFBTHO code and calculation details

HFBTHO

To benchmark the code in the case of nuclear deformation, the **HFBTHO** code was used [39], it's a HFB code which minimizes the energy functional on a (Transformed) harmonic oscillator basis. Since ^{24}Mg is a light nucleus, it still works well in this case. All calculations were performed using 12 oscillator shells and assuming a zero pairing interaction. Default parameters were adopted for the quadrupole constraints. Since the version of **HFBTHO** used in this work has been compiled with the J^2 terms disabled, we present the results from our code both with and without them. The results obtained without them serve as a benchmark for the code, while those including the J^2 contribution illustrate their impact on the calculated observables.

Code parameters and axial constraint

As for our code, calculations are performed on a box $[-10, 10]$ fm. In the case of the ground state calculation, a step size of 0.33 fm is used, with a starting guess of a deformed Woods-Saxon with $\beta_2 = 0.4$.

The calculation in the case of the deformation curve is carried out imposing the following

constraints

$$\langle \text{Re } Q_{22} \rangle = 0 \quad (6.1)$$

$$\langle \text{Im } Q_{22} \rangle = 0 \quad (6.2)$$

$$\langle Y_{20} \rangle = q_{20}. \quad (6.3)$$

These constraints altogether impose an axial deformation on the system. This is done because on a full mesh like in our case, the nucleus may deform on a different axis from the chosen one (z), resulting in spurious contributions to the real deformation curve; moreover, the axial symmetry of HFBTHO doesn't allow broken axial symmetry configurations.

Regarding the stiffness c and damping parameter μ of ALM in section 4.3.1, $c = 0.001$ and $\mu = 0.1$ were used. As for convergence criteria, a tolerance of 0.001 on the value of $\beta_2 - \beta_{2,\text{target}}$ was used.

Ground state

Table 6.1 reports data of the comparison for the ground state of ^{24}Mg , while figure 6.1 shows the middle section of the total particle density. Charge radii for the two codes are displayed but not compared, due to different formulas used for their computation. $\langle x^2 \rangle$, $\langle y^2 \rangle$ and $\langle z^2 \rangle$ is reported for our code but not for HFBTHO since it doesn't compute them.

		GCG	GCG no J^2	HFBTHO	Δ	$\Delta\%$
E_{TOT}	[MeV]	-195.854	-197.219	-197.030	0.189	9.52×10^{-2}
$\langle r_n^2 \rangle^{1/2}$	[fm]	3.0124	2.9998	2.9996	0.0002	6.67×10^{-3}
$\langle r_p^2 \rangle^{1/2}$	[fm]	3.0475	3.0346	3.0326	0.0020	6.59×10^{-2}
$\langle r_{ch}^2 \rangle^{1/2}$	[fm]	3.1364	3.1240	3.4614	-	-
$\langle z^2 \rangle^{1/2}$	[fm]	2.145	2.128	-	-	-
$\langle x^2 \rangle^{1/2}$	[fm]	1.511	1.511	-	-	-
$\langle y^2 \rangle^{1/2}$	[fm]	1.514	1.514	-	-	-
β_2	[-]	0.399	0.390	0.390	-	-

Table 6.1: Results for ^{24}Mg ground state, no pairing interaction, box $[-10, 10]$ fm, step size 0.33 fm, SkM* parametrization.

The comparison shows good agreement between the two codes, with the same β_2 minimum and similar ground state properties.




Figure 6.1: Magnesium ground state density $\rho(x, y, 0)$, calculation done on a box $[-10, 10]$ fm, step size 0.33 fm, SkM* parametrization

Deformation curve

In figure 6.2, the deformation curve is shown for ^{24}Mg , without pairing. To counteract the sharp rise in CPU time, due to the high number of points in the curve, a coarser lattice than the one in the ground state calculation is used, hence the shift in energy of the curve.

Figure 6.2 shows the same trend for both codes, with a minimum of the energy in $\beta_2 = 0.390$, albeit a difference in the energies due to the coarse mesh, a gap which is shown in table 6.1 to shrink when increasing the accuracy of the step size.




Figure 6.2: Magnesium deformation curve, no pairing interaction, calculation done on a box $[-10, 10]$ fm, step size 0.66 fm, SkM* parametrization, neglecting J^2 terms.

Bibliography

- [1] Atomic Mass Data Center. Atomic Mass Data Center. URL <https://www-nds.iaea.org/amdc/>. Accessed: 19 October 2025.
- [2] J. Bartel, P. Quentin, M. Brack, C. Guet, and H.-B. Håkansson. Fission barriers of heavy and superheavy nuclei in a self-consistent hartree-fock approach. *Nuclear Physics A*, 386(1):79–100, 1982. doi: 10.1016/0375-9474(82)90302-6.
- [3] M. Bender, P.-H. Heenen, and P.-G. Reinhard. Self-consistent mean-field models for nuclear structure. *Reviews of Modern Physics*, 75(1):121–180, 2003. doi: 10.1103/RevModPhys.75.121.
- [4] D. Benzaid, S. Bentriddi, A. Kerraci, and N. Amrani. Bethe–Weizsäcker semiempirical mass formula coefficients 2019 update based on AME2016. *Nuclear Science and Techniques*, 31(1):5, 2020. doi: 10.1007/s41365-019-0718-8.
- [5] W. Bertozzi, J. Friar, J. Heisenberg, and J. Negele. Contributions of neutrons to elastic electron scattering from nuclei. *Physics Letters B*, 41(4):408–414, 1972. ISSN 0370-2693. doi: [https://doi.org/10.1016/0370-2693\(72\)90662-4](https://doi.org/10.1016/0370-2693(72)90662-4). URL <https://www.sciencedirect.com/science/article/pii/0370269372906624>.
- [6] C. A. Bertulani. Nuclear physics in a nutshell. 2007.
- [7] Å. Björck. Numerics of gram-schmidt orthogonalization. *Linear Algebra and Its Applications*, 197:297–316, 1994.
- [8] P. Bonche, H. Flocard, and P.-H. Heenen. Self-consistent calculation of nuclear rotations: The complete yrast line of ^{24}Mg . *Nuclear Physics A*, 443(1):39–63, 1985. doi: 10.1016/0375-9474(85)90224-7.
- [9] P. Bonche, H. Flocard, and P. H. Heenen. Solution of the skyrme HF + BCS equation on a 3d mesh. *Computer Physics Communications*, 171(1):49–62, 2005. doi: 10.1016/j.cpc.2005.03.001.
- [10] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer. A skyrme parametrization from subnuclear to neutron star densities. *Nuclear*

- Physics A*, 627(4):710–746, 1997. ISSN 0375-9474. doi: [https://doi.org/10.1016/S0375-9474\(97\)00596-4](https://doi.org/10.1016/S0375-9474(97)00596-4). URL <https://www.sciencedirect.com/science/article/pii/S0375947497005964>.
- [11] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, and R. Schaeffer. A skyrme parametrization from subnuclear to neutron star densities part ii. nuclei far from stabilities. *Nuclear Physics A*, 635(1):231–256, 1998. ISSN 0375-9474. doi: [https://doi.org/10.1016/S0375-9474\(98\)00180-8](https://doi.org/10.1016/S0375-9474(98)00180-8). URL <https://www.sciencedirect.com/science/article/pii/S0375947498001808>.
- [12] M. Chen, T. Li, B. Schuetrumpf, P.-G. Reinhard, and W. Nazarewicz. Three-dimensional skyrme hartree-fock-bogoliubov solver in coordinate-space representation. *Computer Physics Communications*, 276:108344, 2022. ISSN 0010-4655. doi: <https://doi.org/10.1016/j.cpc.2022.108344>. URL <https://www.sciencedirect.com/science/article/pii/S0010465522000625>.
- [13] M. Chen, T. Li, B. Schuetrumpf, P.-G. Reinhard, and W. Nazarewicz. Three-dimensional skyrme hartree-fock-bogoliubov solver in coordinate-space representation. *Computer Physics Communications*, 275:108344, 2022. doi: 10.1016/j.cpc.2022.108344. This paper describes the new 3D, symmetry-unrestricted coordinate-space solver HFBFFT, which is based on the Sky3D code.
- [14] M. Chen, T. Li, B. Schüttrumpf, P.-G. Reinhard, and W. Nazarewicz. Three-dimensional skyrme hartree-fock-bogoliubov solver in coordinate-space representation. *Computer Physics Communications*, 284:108578, 2022. doi: 10.1016/j.cpc.2022.108578.
- [15] G. Colo’ and X. Roca-Maza. User guide for the hfbc-sqrpa(v1) code.
- [16] G. Colò. Nuclear density functional theory. *Advances in Physics: X*, 5(1):1740061, 2020. doi: 10.1080/23746149.2020.1740061. URL <https://doi.org/10.1080/23746149.2020.1740061>.
- [17] R. Y. Cusson, P. G. Reinhard, M. R. Strayer, J. A. Maruhn, and W. Greiner. Density as a constraint and the separation of internal excitation energy in tdhf. *Zeitschrift für Physik A Atoms and Nuclei*, 320:475–482, 1985. doi: 10.1007/BF01415725. URL <https://doi.org/10.1007/BF01415725>.
- [18] J. Dobaczewski. Generalization of the bloch-messiah-zumino theorem. *Phys. Rev. C*, 62:017301, May 2000. doi: 10.1103/PhysRevC.62.017301. URL <https://link.aps.org/doi/10.1103/PhysRevC.62.017301>.

- [19] J. Dobaczewski, W. Nazarewicz, T. R. Werner, J. F. Berger, C. R. Chinn, and J. Decharge. Mean-field description of ground-state properties of drip-line nuclei: Pairing and continuum effects. *Phys. Rev. C*, 53(1):2809, 1996. doi: 10.1103/PhysRevC.53.2809.
- [20] J. Dudek and T. Werner. New parameters of the deformed woods-saxon potential for a=110-210 nuclei. *Journal of Physics G: Nuclear Physics*, 4(10):1543, oct 1978. doi: 10.1088/0305-4616/4/10/006. URL <https://doi.org/10.1088/0305-4616/4/10/006>.
- [21] J. Erler, P. Klüpfel, and P.-G. Reinhard. Misfits in skyrme-hartree-fock. *Journal of Physics G: Nuclear and Particle Physics*, 37(6):064001, mar 2010. doi: 10.1088/0954-3899/37/6/064001. URL <https://doi.org/10.1088/0954-3899/37/6/064001>.
- [22] H. Flocard, P. Quentin, A. Kerman, and D. Vautherin. Nuclear deformation energy curves with the constrained hartree-fock method. *Nuclear Physics A*, 203(3):433–472, 1973. ISSN 0375-9474. doi: [https://doi.org/10.1016/0375-9474\(73\)90357-6](https://doi.org/10.1016/0375-9474(73)90357-6). URL <https://www.sciencedirect.com/science/article/pii/0375947473903576>.
- [23] D. FR. Liquid drop model.svg: Illustration of the terms of the semi-empirical mass formula in the liquid drop model of the atomic nucleus. Wikimedia Commons, the free media repository, 2012. URL https://commons.wikimedia.org/wiki/File:Liquid_drop_model.svg. Original work by Honina, derivative by Daniel FR. Licensed under CC BY-SA 3.0.
- [24] G. H. Golub and C. F. van Loan. *Matrix Computations*. JHU Press, fourth edition, 2013. ISBN 1421407949 9781421407944. URL <http://www.cs.cornell.edu/cv/GVL4/golubandvanloan.htm>.
- [25] H.-Q. Gu, H. Liang, W. H. Long, N. Van Giai, and J. Meng. Slater approximation for coulomb exchange effects in nuclear covariant density functional theory. *Phys. Rev. C*, 87:041301, Apr 2013. doi: 10.1103/PhysRevC.87.041301. URL <https://link.aps.org/doi/10.1103/PhysRevC.87.041301>.
- [26] G. Guennebaud, B. Jacob, et al. Eigen. <http://eigen.tuxfamily.org>, 2010.
- [27] R. Hofstadter. Electron scattering and nuclear structure. *Reviews of Modern Physics*, 28:214–254, 1956. doi: 10.1103/RevModPhys.28.214.
- [28] P. Hohenberg and W. Kohn. Inhomogeneous electron gas. *Phys. Rev.*, 136:B864–

- B871, Nov 1964. doi: 10.1103/PhysRev.136.B864. URL <https://link.aps.org/doi/10.1103/PhysRev.136.B864>.
- [29] P. Jachimowicz, M. Kowal, and J. Skalski. Adiabatic fission barriers in superheavy nuclei. *Phys. Rev. C*, 95:014303, Jan 2017. doi: 10.1103/PhysRevC.95.014303. URL <https://link.aps.org/doi/10.1103/PhysRevC.95.014303>.
- [30] J. D. Jackson. *Classical Electrodynamics*. John Wiley & Sons, New York, 3rd edition, 1998. ISBN 978-0471309321.
- [31] A. V. Knyazev. *A Preconditioned Conjugate Gradient Method for Eigenvalue Problems and its Implementation in a Subspace*, pages 143–154. Birkhäuser Basel, Basel, 1991. ISBN 978-3-0348-6332-2. doi: 10.1007/978-3-0348-6332-2_11. URL https://doi.org/10.1007/978-3-0348-6332-2_11.
- [32] W. Kohn and L. J. Sham. Self-consistent equations including exchange and correlation effects. *Phys. Rev.*, 140:A1133–A1138, Nov 1965. doi: 10.1103/PhysRev.140.A1133. URL <https://link.aps.org/doi/10.1103/PhysRev.140.A1133>.
- [33] J. A. Koponen, K. Pellikka, S. Tanimoto, K. Dendo, T. Lesinski, and K. Terasaki. The axial hartree-fock + BCS code SkyAx. *Computer Physics Communications*, 267: 108034, 2021. doi: 10.1016/j.cpc.2021.108034.
- [34] A. Krishnamoorthy and D. Menon. Matrix inversion using cholesky decomposition. In *2013 Signal Processing: Algorithms, Architectures, Arrangements, and Applications (SPA)*, pages 70–72, 2013.
- [35] C. Lanczos. Solution of systems of linear equations by minimized iterations. *J. Res. Nat. Bur. Standards*, 49(1):33–53, 1952.
- [36] B. Li, Z. Ren, and P. Zhao. Efficient solution for the dirac equation in 3d lattice space with the conjugate gradient method. *Physical Review C*, 102(4):044307, 2020.
- [37] Y. Li, H. Xie, R. Xu, C. You, and N. Zhang. A parallel generalized conjugate gradient method for large scale eigenvalue problems. *CCF Transactions on High Performance Computing*, 2(2):111–122, jun 2020. ISSN 2524-4930. doi: 10.1007/s42514-020-00029-6. URL <https://doi.org/10.1007/s42514-020-00029-6>. The corresponding computing package can be downloaded from the web site: <https://github.com/pase2017/GCGE-1.0>.
- [38] L. Lin, S. Shao, and W. E. Efficient iterative method for solving the dirac-kohn-sham density functional theory. *Journal of Computational Physics*, 245:205–217,

2013. ISSN 0021-9991. doi: <https://doi.org/10.1016/j.jcp.2013.03.030>. URL <https://www.sciencedirect.com/science/article/pii/S0021999113002088>.
- [39] P. Marević, N. Schunck, E. Ney, R. Navarro Pérez, M. Verriere, and J. O’Neal. Axially-deformed solution of the skyrme-hartree-fock-bogoliubov equations using the transformed harmonic oscillator basis (iv) hfbtho (v4.0): A new version of the program. *Computer Physics Communications*, 276:108367, 2022. ISSN 0010-4655. doi: <https://doi.org/10.1016/j.cpc.2022.108367>. URL <https://www.sciencedirect.com/science/article/pii/S0010465522000868>.
- [40] P. Marević, N. Schunck, and K. Sato. Axially-deformed solution of the skyrme-hartree-fock-bogoliubov problem in the harmonic-oscillator basis. *Computer Physics Communications*, 283:108640, 2022. doi: 10.1016/j.cpc.2022.108640.
- [41] R. M. Martin. *Electronic Structure: Basic Theory and Practical Methods*. Cambridge University Press, Cambridge, UK, 2004. ISBN 9780521534406.
- [42] J. A. Maruhn, P.-G. Reinhard, P. D. Stevenson, and A. S. Umar. The tdhf code sky3d. *Computer Physics Communications*, 185:2195–2216, 2014. doi: 10.1016/j.cpc.2014.05.005.
- [43] B. Mottelson. The nilsson model and sven gösta nilsson. *Physica Scripta*, 125, 2006.
- [44] T. Nottoli, I. Gianni, A. Levitt, and F. Lipparini. A robust, open-source implementation of the locally optimal block preconditioned conjugate gradient for large eigenvalue problems in quantum chemistry. *Theoretical Chemistry Accounts*, 142(8):69, 2023. ISSN 1432-2234. doi: 10.1007/s00214-023-03010-y. URL <https://doi.org/10.1007/s00214-023-03010-y>.
- [45] J. W. Pearson and J. Pestana. Preconditioners for krylov subspace methods: An overview. *GAMM-Mitteilungen*, 43(4):e202000015, 2020.
- [46] J. C. Pei, M. V. Stoitsov, G. I. Fann, W. Nazarewicz, N. Schunck, and F. R. Xu. Deformed coordinate-space hartree-fock-bogoliubov approach to weakly bound nuclei and large deformations. *Physical Review C*, 78(6):064306, 2008. doi: 10.1103/PhysRevC.78.064306.
- [47] J. C. Pei, A. T. Kruppa, and W. Nazarewicz. Quasi-particle continuum and resonances in the hartree-fock-bogoliubov theory. *Physical Review C*, 84(2):024311, 2011. doi: 10.1103/PhysRevC.84.024311.
- [48] P.-G. Reinhard and H. Flocard. Nuclear effective forces and isotope shifts. *Nuclear Physics A*, 584(3):467–488, 1995. ISSN 0375-9474. doi: <https://doi.org/>

- 10.1016/0375-9474(94)00770-N. URL <https://www.sciencedirect.com/science/article/pii/037594749400770N>.
- [49] P. Ring and P. Schuck. *The nuclear many-body problem*. Springer Science & Business Media, 2004.
- [50] W. Ryssens, V. Hellemans, M. Bender, and P.-H. Heenen. Solution of the skyrme-hf+bcs equation on a 3d mesh, ii: A new version of the ev8 code. *Computer Physics Communications*, 187:175–194, 2015. ISSN 0010-4655. doi: <https://doi.org/10.1016/j.cpc.2014.10.001>. URL <https://www.sciencedirect.com/science/article/pii/S0010465514003361>.
- [51] W. Ryssens, V. Hellemans, M. Bender, and P.-H. Heenen. Solution of the skyrme hf+bcs equation on a 3d mesh. ii. a new version of the ev8 code. *Computer Physics Communications*, 187:175–194, 2015. doi: 10.1016/j.cpc.2014.10.001.
- [52] W. Ryssens, P.-H. Heenen, and M. Bender. Symmetry-unrestricted skyrme mean-field study of heavy nuclei: the mocca code. *arXiv preprint*, 2016. arXiv:1611.01300 [nucl-th].
- [53] Y. Saad. *Numerical Methods for Large Eigenvalue Problems*. Manchester University Press, Manchester, UK, 1992.
- [54] N. Schunck, J. Dobaczewski, J. McDonnell, J. Sarich, A. Staszczak, and M. V. Stoitsov. Solution of the HFB equation in the oscillatory basis: the HFODD code. *Computer Physics Communications*, 183(1):166–191, 2012. doi: 10.1016/j.cpc.2011.09.018.
- [55] J. R. Shewchuk. An introduction to the conjugate gradient method without the agonizing pain. 1994. URL <http://www.cs.cmu.edu/~quake-papers/painless-conjugate-gradient.pdf>.
- [56] T. Skyrme. The effective nuclear potential. *Nuclear Physics*, 9(4):615–634, 1958. ISSN 0029-5582. doi: [https://doi.org/10.1016/0029-5582\(58\)90345-6](https://doi.org/10.1016/0029-5582(58)90345-6). URL <https://www.sciencedirect.com/science/article/pii/0029558258903456>.
- [57] A. Staszczak, M. Stoitsov, A. Baran, and W. Nazarewicz. Augmented lagrangian method for constrained nuclear density functional theory. *The European Physical Journal A*, 46:85–90, 2010. doi: 10.1140/epja/i2010-11018-9. URL <https://doi.org/10.1140/epja/i2010-11018-9>.
- [58] P. Stevenson and M. Barton. Low-energy heavy-ion reactions and the skyrme effective interaction. *Progress in Particle and Nuclear Physics*, 104:142–164, 2019.

- [59] M. V. Stoitsov, J. Dobaczewski, W. Nazarewicz, S. Pittel, and D. J. Dean. Systematic study of deformed nuclei at the drip lines and beyond. *Phys. Rev. C*, 68(5):054312, 2003. doi: 10.1103/PhysRevC.68.054312.
- [60] B. Stroustrup. An overview of c++. In *Proceedings of the 1986 SIGPLAN workshop on Object-oriented programming*, pages 7–18, 1986.
- [61] D. Vautherin and D. M. Brink. Hartree-fock calculations with skyrme’s interaction. i. spherical nuclei. *Phys. Rev. C*, 5:626–647, Mar 1972. doi: 10.1103/PhysRevC.5.626. URL <https://link.aps.org/doi/10.1103/PhysRevC.5.626>.
- [62] C. F. von Weizsäcker. Zur Theorie der Kernmassen. *Zeitschrift für Physik*, 96(7-8): 431–458, 1935. doi: 10.1007/BF01337700.
- [63] Wikipedia contributors. Nilsson model graphical representation – wikipedia. <https://en.wikipedia.org/w/index.php?curid=60892381>, 2025.
- [64] J. Woo, S. Kim, and W. Y. Kim. Gaussian-approximated poisson preconditioner for iterative diagonalization in real-space density functional theory. *The Journal of Physical Chemistry A*, 127(17):3883–3893, 2023. doi: 10.1021/acs.jpca.2c09111. URL <https://doi.org/10.1021/acs.jpca.2c09111>. PMID: 37094552.
- [65] P.-b. Zhou. *Finite Difference Method*, pages 63–94. Springer Berlin Heidelberg, Berlin, Heidelberg, 1993. ISBN 978-3-642-50319-1. doi: 10.1007/978-3-642-50319-1_3. URL https://doi.org/10.1007/978-3-642-50319-1_3.

A | Appendix

A.1. Spherical harmonics

Spherical harmonics, of order λ, μ , are defined as

$$Y_{\lambda\mu}(\theta, \phi) = (-1)^\mu \sqrt{\frac{2\lambda+1}{4\pi} \frac{(\lambda-\mu)!}{(\lambda+\mu)!}} P_\lambda^\mu(\cos \theta) e^{i\mu\phi}. \quad (\text{A.1})$$

Being able to provide the expression for arbitrary μ, λ through an algorithm is important in the current framework, to solve the Poisson equation and investigate nuclear properties. The major challenge is to generate the associated Legendre polynomials P_λ^μ . They can be expressed in the form (for positive μ)

$$P_\lambda^\mu(x) = (1-x^2)^{\mu/2} \frac{d^\mu P_\lambda(x)}{dx^\mu}, \quad (\text{A.2})$$

where $x = \cos \theta$ and

$$P_\lambda(x) = \frac{1}{2^\lambda \lambda!} \frac{d^\lambda (x^2-1)^\lambda}{dx^\lambda}. \quad (\text{A.3})$$

To compute the arbitrary λ, μ associated Legendre polynomial we can employ a recursive approach, setting $\lambda = \mu$

$$P_\mu^\mu(x) = (2\mu-1)!! (1-x^2)^{\mu/2}, \quad (\text{A.4})$$

where $(2\mu-1)!! = 1 \cdot 3 \cdot 5 \dots (2\mu-1)$ denotes the double factorial. Once $P_\mu^\mu(x)$ is known, the next element with $\lambda = \mu + 1$ reads

$$P_{\mu+1}^\mu(x) = x(2\mu+1)P_\mu^\mu(x). \quad (\text{A.5})$$

All higher orders are then generated using the standard upward recurrence relation in λ :

$$(\lambda - \mu + 1) P_{\lambda+1}^\mu(x) = (2\lambda + 1) x P_\lambda^\mu(x) - (\lambda + \mu) P_{\lambda-1}^\mu(x), \quad (\text{A.6})$$

valid for all $\lambda \geq \mu + 1$.

A.1.1. Algorithm

1. Compute the base case P_μ^μ from the closed-form formula.
2. If $\mu = \lambda$ the procedure ends, otherwise
3. Evaluate $P_{\mu+1}^\mu$, if $\lambda = \mu + 1$ the procedure ends, otherwise
4. Apply the recurrence relation $P_{\lambda+1}^\mu$ until the desired degree is reached

This ought to be applied only for $\mu \geq 0$. For $\mu < 0$ the procedure is carried out using $-\mu$ and in the end using the relation

$$Y_{\lambda-\mu} = (-1)^\mu Y_{\lambda\mu}^* \quad (\text{A.7})$$

A.2. 5-point derivatives

The first and second derivatives of a function $\psi(x)$ in $x = x_i$, using 5-points formulae, read

$$\psi'(x_i) = \frac{\psi_{i-2} - 8\psi_{i-1} + 8\psi_{i+1} - \psi_{i+2}}{12h} \quad (\text{A.8})$$

$$\psi''(x_i) = \frac{-\psi_{i-2} + 16\psi_{i-1} - 30\psi_i + 16\psi_{i+1} - \psi_{i+2}}{12h^2} \quad (\text{A.9})$$

A.3. Functional derivatives

Given a functional $\mathcal{F}[\rho]$, the functional derivative is defined as the variation of \mathcal{F} with respect to a small change in the density ρ , formally

$$\frac{\delta \mathcal{F}[\rho]}{\delta \rho} = \lim_{\delta \rho \rightarrow 0} \frac{\mathcal{F}[\rho + \delta \rho] - \mathcal{F}[\rho]}{\delta \rho}.$$

Power dependence Let us suppose to have $\mathcal{F}[\rho] = A\rho^\sigma$, where A is a constant. A variation $\delta\rho$ of ρ yields

$$\mathcal{F}[\rho + \delta\rho] - \mathcal{F}[\rho] = A[(\rho + \delta\rho)^\sigma - \rho^\sigma].$$

Expanding to first order in $\delta\rho$ (Taylor expansion) we obtain

$$(\rho + \delta\rho)^\sigma = \rho^\sigma + \sigma\rho^{\sigma-1}\delta\rho + \mathcal{O}((\delta\rho)^2),$$

hence

$$\mathcal{F}[\rho + \delta\rho] - \mathcal{F}[\rho] = A \sigma \rho^{\sigma-1} \delta\rho + \mathcal{O}((\delta\rho)^2).$$

Dividing by $\delta\rho$ and taking the limit $\delta\rho \rightarrow 0$ gives the usual result for the functional derivative

$$\frac{\delta\mathcal{F}[\rho]}{\delta\rho} = A \sigma \rho^{\sigma-1}. \quad (\text{A.10})$$

Divergence of a vector field Consider the functional

$$\mathcal{G}[\mathbf{J}](\mathbf{r}) = \nabla_{\mathbf{r}} \cdot \mathbf{J}(\mathbf{r}).$$

A small variation $\delta\mathbf{J}$ induces

$$\delta\mathcal{G}(\mathbf{r}) = \nabla_{\mathbf{r}} \cdot \delta\mathbf{J}(\mathbf{r}).$$

We can express the variation $\delta\mathbf{J}(\mathbf{r})$ in terms of its values at all points \mathbf{r}' as

$$\delta\mathbf{J}(\mathbf{r}) = \int d^3r' \delta\mathbf{J}(\mathbf{r}') \delta(\mathbf{r} - \mathbf{r}').$$

Substituting this into the expression for $\delta\mathcal{G}(\mathbf{r})$ yields

$$\delta\mathcal{G}(\mathbf{r}) = \int d^3r' \nabla_{\mathbf{r}} \cdot [\delta\mathbf{J}(\mathbf{r}') \delta(\mathbf{r} - \mathbf{r}')].$$

Since \mathbf{r} and \mathbf{r}' are independent variables, the derivative acts only on the delta function:

$$\delta\mathcal{G}(\mathbf{r}) = \int d^3r' (\nabla_{\mathbf{r}} \delta(\mathbf{r} - \mathbf{r}')) \cdot \delta\mathbf{J}(\mathbf{r}').$$

By the definition of the functional derivative as the kernel relating $\delta\mathcal{G}(\mathbf{r})$ to $\delta\mathbf{J}(\mathbf{r}')$, we can read off

$$\frac{\delta\mathcal{G}[\mathbf{J}](\mathbf{r})}{\delta\mathbf{J}(\mathbf{r}')} = \nabla_{\mathbf{r}} \delta(\mathbf{r} - \mathbf{r}'). \quad (\text{A.11})$$

In compact vector form [30],

$$\frac{\delta(\nabla_{\mathbf{r}} \cdot \mathbf{J}(\mathbf{r}))}{\delta\mathbf{J}(\mathbf{r}')} = \nabla_{\mathbf{r}} \delta(\mathbf{r} - \mathbf{r}').$$

A.4. \overline{U} , \overline{V} matrix structure

The blocked form of the \overline{U} and \overline{V} matrices in section 3.2.2 is given by

$$\bar{U} = \begin{pmatrix} 0 & & & & & & & & & & \\ & \ddots & & & & & & & & & \\ & & 0 & & & & & & & & \\ & & & u_1 & 0 & & & & & & \\ & & & 0 & u_1 & & & & & & \\ & & & & & \ddots & & & & & \\ & & & & & & u_n & 0 & & & \\ & & & & & & 0 & u_n & & & \\ & & & & & & & & 1 & & \\ & & & & & & & & & \ddots & \\ & & & & & & & & & & 1 \end{pmatrix}$$

$$\bar{V} = \begin{pmatrix} 1 & & & & & & & & & & \\ & \ddots & & & & & & & & & \\ & & 1 & & & & & & & & \\ & & & 0 & v_1 & & & & & & \\ & & & -v_1 & 0 & & & & & & \\ & & & & & \ddots & & & & & \\ & & & & & & 0 & v_n & & & \\ & & & & & & -v_n & 0 & & & \\ & & & & & & & & 0 & & \\ & & & & & & & & & \ddots & \\ & & & & & & & & & & 0 \end{pmatrix}$$

List of Figures

1.1	Binding energy per nucleon as a function of A . Due to the short range of the strong force, this value saturates around 7 MeV, with a steady, dim decrease after ^{56}Fe	4
1.2	Visual representation of the liquid drop model from [23]	6
1.3	Graphical representation of a harmonic oscillator shells, together with the spin-orbit coupling. Shells whose total angular momentum is higher are lowered in energy, viceversa for lower total angular momentum.	8
1.4	Nilsson model energy levels trends, as a function of ε	13
1.5	Graphical representation of possible octupole deformations. On the left, the axially symmetric Y_{30} deformation, on the right, the non-axial octupole deformation Y_{32}	14
4.1	Comparison between the steepest descent method, on the left, and the conjugate gradient method, on the right, for a two-dimensional matrix. Ellipses represent contour lines of the quadratic form $f(x) = x^T Ax - b^T x$, in the ‘stretched’ space Ax on the right. As shown in the figure, the conjugation of the search directions eliminates components of the error e_i , until the exact convergence in n (2) steps.	52
4.2	Pseudocode of the Hartree-Fock program.	62
4.3	HF calculation convergence with varying CG tolerance for ^{16}O , box $[-9, 9]$ fm, step size 0.3 fm.	63
4.4	HF calculation convergence with varying number of inverse power steps for ^{16}O , box $[-9, 9]$ fm, step size 0.3 fm.	64
4.5	HF calculation convergence with varying number of inverse power steps for the deformed nucleus ^{24}Mg , box $[-10, 10]$ fm, step size 0.33 fm.	64
4.6	Numerical stability map of the HF solver for ^{16}O for different box and step sizes. Relative error is taken against a benchmark reference value.	65
6.1	Magnesium ground state density $\rho(x, y, 0)$, calculation done on a box $[-10, 10]$ fm, step size 0.33 fm, SkM* parametrization	79

6.2	Magnesium deformation curve, no pairing interaction, calculation done on a box $[-10, 10]$ fm, step size 0.66 fm, SkM* parametrization, neglecting J^2 terms.	80
-----	---	----

List of Tables

5.1	Parameters used to compute the charge mean square radius.	68
5.2	^{16}O including C_0^ρ , C_1^ρ , C_0^τ , C_1^τ terms, neglecting Coulomb interaction. . . .	70
5.3	^{16}O including C_0^ρ , C_1^ρ , C_0^τ , C_1^τ , $C_0^{\nabla\cdot\mathbf{J}}$, $C_1^{\nabla\cdot\mathbf{J}}$ terms, neglecting Coulomb interaction and J^2 terms.	71
5.4	^{16}O neglecting Coulomb interaction.	71
5.5	^{16}O complete of the Skyrme functional and Coulomb interaction.	72
5.6	^{48}Ca , box size $[-12, 12]$ fm, step size 0.34 fm	73
5.7	^{56}Ni , box size $[-13, 13]$ fm, step size 0.37 fm	74
5.8	^{90}Zr , box size $[-15, 15]$ fm, step size 0.43 fm	75
5.9	Comparison of experimental binding energies in MeV with theoretical calculated values using the SLy5 functional.	76
6.1	Results for ^{24}Mg ground state, no pairing interaction, box $[-10, 10]$ fm, step size 0.33 fm, SkM* parametrization.	78

List of Symbols

Variable	Description	SI unit
\boldsymbol{u}	solid displacement	m
\boldsymbol{u}_f	fluid displacement	m

Acknowledgements

Here you might want to acknowledge someone.

