

Recap:

- Introduced notation for discretization: $X \rightarrow x_i$ x_0, x_1, \dots, x_n
 $u(x) \rightarrow u_i$ u_0, u_1, \dots, u_n

- Derivatives

$$\left. \frac{du}{dx} \right|_{x_i} = u'_i = \begin{cases} \frac{u_{i+1} - u_i}{h} + \mathcal{O}(h) \\ \frac{u_i - u_{i-1}}{h} + \mathcal{O}(h) \\ \frac{u_{i+1} - u_{i-1}}{2h} + \mathcal{O}(h^2) \end{cases}$$

$$\left. \frac{d^2u}{dx^2} \right|_{x_i} = u''_i = \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} + \mathcal{O}(h^2)$$

Next: [Good news and bad news:
 - Bad: Lots of tedious writing
 - Good: Will help you a lot with Proj. 7]

- Boundary value problem (example in Pr. 7)
- Formulation as a matrix problem
- Solving matrix eqs., with special focus on tridiag. matrices
- Floating point operations (FLOPs)
- Errors

Background material: Sec 6.4 (6.4.5)

Note: Differences in notation!
 Also, some index typos, I think...

Boundary value problem

- Our case in Pr. 1 : $-\frac{d^2 u}{dx^2} = f(x)$ $x \in [0, 1]$
- $u(x)$ unknown
- $f(x)$ known
- $x \in [0, 1]$
- Boundary values : $u(0) = 0$, $u(1) = 0$ (Dirichlet)

• Special case of :

$$\alpha \frac{d^2 u}{dx^2} + \beta \frac{du}{dx} + \gamma u(x) = f(x)$$

- Ordinary diff. eq. (only one indep. variable x)
- Linear diff. eq. (each term has max. one power of u, u', u'', \dots)
- Second order (highest-order derivative is u'')
- Inhomogeneous ($f(x) \neq 0$)
- Most diff. eqs. in physics are linear. \Rightarrow sum of two solutions is a new valid solution!

• Many approaches :

- Shooting methods (quickly disc.)
- Finite diff. methods (Pr. 1)
- Finite elem. methods (not covered)


Most famous example

Schr. eq. in QM is linear

\hookrightarrow Superposition of quantum states!

Shooting method intuition

- Turn BVP into initial value problem (know $u(x_0)$ and $u'(x_0)$)
- Know $u(x_0)$
- Guess $u'(x_0)$ and solve forward ("shoot")
- \rightarrow one attempt $u_{(1)}(x)$
- Guess another $u'(x_0)$ and solve forward
- \rightarrow second attempt $u_{(2)}(x)$
- Sum of solutions is a new solution (linearity)
- $u_c(x) = c u_{(1)}(x) + (1-c) u_{(2)}(x)$
- Require that $u_c(x_1) = u(x_1)$ (second bound. cond.) \Rightarrow Determines $c \Rightarrow$ solution $u(x) = u_c(x)$!



Finite diff. method

- We have: $-\frac{d^2 u}{dx^2} = f(x)$
- Goal: Find $u(x)$
- Know: $x \in [0, 1]$, $u(0)$, $u(1)$, $f(x)$
- Strategy:
 - ① Express as matrix eq.
 - ② Solve matrix eq.

① Express as matrix eq.

- Discretize

$$-\left[\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} + \mathcal{O}(h) \right] = f_i \quad f_i \equiv f(x_i)$$

- Approximate

- Change notation: $v_i \approx u_i$

$$-\left[\frac{v_{i+1} - 2v_i + v_{i-1}}{h^2} \right] = f_i$$

- Arrange terms

$$(*) \quad \boxed{-v_{i-1} + 2v_i - v_{i+1} = h^2 f_i}$$

Goal: Determine v_1, v_2, \dots, v_{n-1}
 Know: v_0, v_n , all f_i


- Here: $n = 5$ steps

$v_0, v_1, v_2, v_3, v_4, v_5$: 6 pts

v_0, v_5 are boundary pts.

4 unknown: v_1, \dots, v_4

Pr 1: You will generalize this!



$h = 0.2$ (very large...)

• (*) represents a set of eqs. Let's write them out in a suggestive way!

4

$$\begin{aligned}
 (i=1) \quad & -\cancel{v_0} + 2v_1 - v_2 = h^2 f_1 \\
 (i=2) \quad & -v_1 + 2v_2 - v_3 = h^2 f_2 \\
 (i=3) \quad & -v_2 + 2v_3 - v_4 = h^2 f_3 \\
 (i=4) \quad & -v_3 + 2v_4 - \cancel{v_5} = h^2 f_4
 \end{aligned}$$

• v_0, v_5 are known

define
notation
↓

$$\begin{aligned}
 2v_1 - v_2 &= h^2 f_1 + v_0 & \equiv g_1 \\
 -v_1 + 2v_2 - v_3 &= h^2 f_2 & \equiv g_2 \\
 -v_2 + 2v_3 - v_4 &= h^2 f_3 & \equiv g_3 \\
 -v_3 + 2v_4 &= h^2 f_4 + v_5 & \equiv g_4
 \end{aligned}$$

• Can be written as

$$\begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \begin{bmatrix} g_1 \\ g_2 \\ g_3 \\ g_4 \end{bmatrix}$$

$$\boxed{A \bar{v} = \bar{g}}$$

• A, \bar{g} known
• want to solve for \bar{v}

② Solve matrix eq.

• Overview of things we'll discuss

1) Solving a general matrix eq. $A\bar{v} = \bar{g}$

(Gauss elim., LU decomp., ...)

Now \rightarrow 2) Solving $A\bar{v} = \bar{g}$ when A is a general tridiagonal matrix

Gauss elim. \Rightarrow Thomas algorithm

3) Solving $A\bar{v} = \bar{g}$ when A is the special tridiag. matrix

$$A = \begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & -1 & 2 & -1 & \\ & & \ddots & \ddots & \ddots \\ & & & -1 & 2 & -1 \\ & & & & -1 & 2 \end{bmatrix}$$

(You'll think about this in Pr. 7)

• First some words about matrix eq. ...

About solving matrix eqs.

$$\boxed{A \bar{x} = \bar{b}}$$

A, \bar{b} known

\bar{x} unknown

- Comes from set of linear equations

$$m \text{ eqs. } \left\{ \begin{array}{l} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \end{array} \right.$$

n terms, one per unknown variable (x_1, x_2, \dots, x_n)

- In general:

$$\begin{matrix} A & \bar{x} & = & \bar{b} \\ (m \times n) & (n \times 1) & & (n \times 1) \end{matrix}$$

e.g. $\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}$
 $(4 \times 3) \quad (3 \times 1) \quad (4 \times 1)$

- We'll focus on the case $m=n \Leftrightarrow A$ is square
 $\Leftrightarrow n$ eqs. and n unknowns

- If all eqs. are lin. indep. (each eq. represent info not contained in the other eqs.)
 we should be able to solve for all unknowns (x_1, \dots, x_n), i.e. for \bar{x} .

If you have more eqs. than unknowns
 \rightarrow no exact solution, but can fit unknowns.
 Typical case in science!

- All eqs. are lin. indep. $\Leftrightarrow A$ is not singular $\Leftrightarrow \det(A) \neq 0$
 (all eigenvalues of A are $\neq 0$)

• Gaussian elimination

$$A \quad \bar{v} = \bar{g}$$

$$\begin{bmatrix} \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{bmatrix} \begin{bmatrix} \cdot \\ \cdot \\ \cdot \\ \cdot \end{bmatrix} = \begin{bmatrix} \cdot \\ \cdot \\ \cdot \\ \cdot \end{bmatrix}$$

- step 1: Forward subst./elim.

(Turn into upper triang. form)

$$\Rightarrow \begin{bmatrix} \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{bmatrix} \begin{bmatrix} \cdot \\ \cdot \\ \cdot \\ \cdot \end{bmatrix} = \begin{bmatrix} \cdot \\ \cdot \\ \cdot \\ \cdot \end{bmatrix}$$

upper triangular

trivial to solve for last v_i

- step 2: Back subst.

(Use solution for v_i to find v_{i-1})

$$\Rightarrow \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & 0 & & 1 \end{bmatrix} \begin{bmatrix} \cdot \\ \cdot \\ \cdot \\ \cdot \end{bmatrix} = \begin{bmatrix} \cdot \\ \cdot \\ \cdot \\ \cdot \end{bmatrix}$$

solved for all v_i

• Our case: General tridiagonal A

$$\begin{bmatrix} b_1 & c_1 & 0 & 0 \\ a_2 & b_2 & c_2 & 0 \\ 0 & a_3 & b_3 & c_3 \\ 0 & 0 & a_4 & b_4 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \begin{bmatrix} g_1 \\ g_2 \\ g_3 \\ g_4 \end{bmatrix}$$

subdiag: $\bar{a} = [a_2, a_3, a_4]$

main diag: $\bar{b} = [b_1, b_2, b_3, b_4]$

superdiag: $\bar{c} = [c_1, c_2, c_3]$

$$\begin{array}{lcl} (R_1) & b_1 & c_1 \quad 0 \quad 0 \quad | \quad g_1 \\ (R_2) & a_2 & b_2 \quad c_2 \quad 0 \quad | \quad g_2 \\ (R_3) & 0 & a_3 \quad b_3 \quad c_3 \quad | \quad g_3 \\ (R_4) & 0 & 0 \quad a_4 \quad b_4 \quad | \quad g_4 \end{array}$$

$$\boxed{R_2 \rightarrow R_2 - \frac{a_2}{b_1} R_1}$$

$$\begin{array}{lcl} (a_2 - \frac{a_2}{b_1} b_1) & b_1 & c_1 \quad 0 \quad 0 \quad | \quad g_1 \\ \rightarrow 0 & (b_2 - \frac{a_2}{b_1} c_1) & c_2 \quad 0 \quad | \quad (g_2 - \frac{a_2}{b_1} g_1) \\ 0 & a_3 & b_3 \quad c_3 \quad | \quad g_3 \\ 0 & 0 & a_4 \quad b_4 \quad | \quad g_4 \end{array}$$

• New notation :

$$\bullet \tilde{b}_2 = b_2 - \frac{a_2}{\tilde{b}_1} c_1$$

Also: $\tilde{b}_1 = b_1$

$$\bullet \tilde{g}_2 = g_2 - \frac{a_2}{\tilde{b}_1} \tilde{g}_1$$

$$\tilde{g}_1 = g_1$$

$$\begin{array}{cccc|c} \tilde{b}_1 & c_1 & 0 & 0 & \tilde{g}_1 \\ 0 & \tilde{b}_2 & c_2 & 0 & \tilde{g}_2 \\ 0 & \textcircled{a_3} & b_3 & c_3 & g_3 \\ 0 & 0 & a_4 & b_4 & g_4 \end{array}$$

• Cont. like this :

Row op: $R_3 \rightarrow R_3 - \frac{a_3}{b_2} R_2$

Define: $\bullet \tilde{b}_3 = b_3 - \frac{a_3}{\tilde{b}_2} c_2$

$$\bullet \tilde{g}_3 = g_3 - \frac{a_3}{\tilde{b}_2} \tilde{g}_2$$

$$\begin{array}{cccc|c} \tilde{b}_1 & c_1 & 0 & 0 & \tilde{g}_1 \\ 0 & \tilde{b}_2 & c_2 & 0 & \tilde{g}_2 \\ 0 & 0 & \tilde{b}_3 & c_3 & \tilde{g}_3 \\ 0 & 0 & a_4 & b_4 & g_4 \end{array}$$

• Last step

$$R_4 \rightarrow R_4 - \frac{a_4}{\tilde{b}_3} R_3$$

Define: $\bullet \tilde{b}_4 = b_4 - \frac{a_4}{\tilde{b}_3} c_3$

$$\bullet \tilde{g}_4 = g_4 - \frac{a_4}{\tilde{b}_3} \tilde{g}_3$$

$$\Rightarrow \begin{array}{cccc|c} \tilde{b}_1 & c_1 & 0 & 0 & \tilde{g}_1 \\ 0 & \tilde{b}_2 & c_2 & 0 & \tilde{g}_2 \\ 0 & 0 & \tilde{b}_3 & c_3 & \tilde{g}_3 \\ 0 & 0 & 0 & \tilde{b}_4 & \tilde{g}_4 \end{array}$$

Forward subst. done!

For coding: do need for matrix,
but arrays for
 $\bar{a}, \bar{b}, \bar{c}, \bar{g}, \bar{b}, \bar{g}, \bar{v}$

Forward subst. :

$$\left[\begin{array}{l} \tilde{b}_1 = b_1 \\ \tilde{b}_i = b_i - \frac{a_i}{\tilde{b}_{i-1}} c_{i-1} \quad \text{for } i=2,3,4 \\ \tilde{g}_1 = g_1 \\ \tilde{g}_i = g_i - \frac{a_i}{\tilde{b}_{i-1}} \tilde{g}_{i-1} \quad \text{for } i=2,3,4 \end{array} \right]$$

o Back subst.

starting point:

$$\begin{bmatrix} \tilde{b}_1 & c_1 & 0 & 0 \\ 0 & \tilde{b}_2 & c_2 & 0 \\ 0 & 0 & \tilde{b}_3 & c_3 \\ 0 & 0 & 0 & \tilde{b}_4 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = \begin{bmatrix} \tilde{g}_1 \\ \tilde{g}_2 \\ \tilde{g}_3 \\ \tilde{g}_4 \end{bmatrix} \Rightarrow \tilde{b}_4 v_4 = \tilde{g}_4$$

$$\Rightarrow \boxed{v_4 = \frac{\tilde{g}_4}{\tilde{b}_4}}$$

$$R_4 \rightarrow \frac{R_4}{\tilde{b}_4}$$

$$\begin{array}{cccc|c} \tilde{b}_1 & c_1 & 0 & 0 & \tilde{g}_1 \\ 0 & \tilde{b}_2 & c_2 & 0 & \tilde{g}_2 \\ 0 & 0 & \tilde{b}_3 & c_3 & \tilde{g}_3 \\ 0 & 0 & 0 & 1 & v_4 \quad (= \frac{\tilde{g}_4}{\tilde{b}_4}) \end{array}$$

$$R_3 \rightarrow (R_3 - c_3 R_4) / \tilde{b}_3$$

$$\begin{array}{cccc|c} \tilde{b}_1 & c_1 & 0 & 0 & \tilde{g}_1 \\ 0 & \tilde{b}_2 & c_2 & 0 & \tilde{g}_2 \\ 0 & 0 & 1 & 0 & (\tilde{g}_3 - c_3 v_4) / \tilde{b}_3 = v_3 \quad ! \\ 0 & 0 & 0 & 1 & v_4 \end{array}$$

$$\Rightarrow V_3 = \frac{\tilde{g}_3 - c_3 V_4}{\tilde{b}_3}$$

• Can continue upwards like this. In the end

$$\left[\begin{array}{l} V_4 = \frac{\tilde{g}_4}{\tilde{b}_4} \\ V_i = \frac{\tilde{g}_i - c_i V_{i+1}}{\tilde{b}_i} \quad i = 3, 2, 1 \end{array} \right]$$

Back subst. done!

$$\begin{array}{cccc|c} 1 & 0 & 0 & 0 & v_1 \\ 0 & 1 & 0 & 0 & v_2 \\ 0 & 0 & 1 & 0 & v_3 \\ 0 & 0 & 0 & 1 & v_4 \end{array}$$

• Using Gaussian elim. on a general tridiagonal matrix (4×4)

We have solved

$$A \bar{v} = \bar{g}$$

$$\text{for } \bar{v} = [v_1, v_2, v_3, v_4]$$

Two parts to procedure:

- Forward subst. / elim.
- Back subst.