

# Weierstrass sets on finite graphs

Alessio Borzi

University of Warwick

Goal: tropical analogues of Weierstrass semigroups

$X$  smooth projective curve of genus  $g$

$X$  smooth projective curve of genus  $g$ , fix a point  $P \in X$

$X$  smooth projective curve of genus  $g$ , fix a point  $P \in X$

$$H(P) = \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\}$$

$X$  smooth projective curve of genus  $g$ , fix a point  $P \in X$

$$\begin{aligned} H(P) &= \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\} \\ &= \{n \in \mathbb{N} : r(nP) > r((n-1)P)\} \end{aligned}$$

$X$  smooth projective curve of genus  $g$ , fix a point  $P \in X$

$$\begin{aligned} H(P) &= \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\} \\ &= \{n \in \mathbb{N} : r(nP) > r((n-1)P)\} \end{aligned}$$

**Weierstrass semigroup** of  $X$  at  $P$

$X$  smooth projective curve of genus  $g$ , fix a point  $P \in X$

$$\begin{aligned} H(P) &= \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\} \\ &= \{n \in \mathbb{N} : r(nP) > r((n-1)P)\} \end{aligned}$$

**Weierstrass semigroup** of  $X$  at  $P$   
( $\text{ord}_P(f_1 f_2) = \text{ord}_P(f_1) + \text{ord}_P(f_2)$ )



$X$  smooth projective curve of genus  $g$ , fix a point  $P \in X$

$$\begin{aligned} H(P) &= \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\} \\ &= \{n \in \mathbb{N} : r(nP) > r((n-1)P)\} \end{aligned}$$

**Weierstrass semigroup** of  $X$  at  $P$   
( $\text{ord}_P(f_1 f_2) = \text{ord}_P(f_1) + \text{ord}_P(f_2)$ )

Theorem (Weierstrass gap theorem)

$$|\mathbb{N} \setminus H(P)| = g$$

**numerical semigroup** = cofinite submonoid of  $\mathbb{N}$

Question (Hurwitz 1893)

*Which numerical semigroups are Weierstrass semigroups?*

Question (Hurwitz 1893)

*Which numerical semigroups are Weierstrass semigroups?*

Example (Buchweitz 1980)

The semigroup  $S = \langle 13, 14, 15, 16, 17, 18, 20, 22, 23 \rangle$  is not a Weierstrass semigroup

Question (Hurwitz 1893)

*Which numerical semigroups are Weierstrass semigroups?*

Example (Buchweitz 1980)

The semigroup  $S = \langle 13, 14, 15, 16, 17, 18, 20, 22, 23 \rangle$  is not a Weierstrass semigroup

Recent work of Cotterill, Pflueger, Zhang (2022) certifies Weierstrass-realizability of some numerical semigroups.

Baker and Norine (2007)

divisors on **graphs**

Baker and Norine (2007)

divisors on [graphs](#)

Gathmann and Kerber (2008)

[metric graphs](#)

Mikhalkin and Zharkov (2008) (i.e. abstract tropical curves)

Baker and Norine (2007)

divisors on **graphs**

Gathmann and Kerber (2008)

**metric graphs**

Mikhalkin and Zharkov (2008) (i.e. abstract tropical curves)

### Question

*What is the tropical analogue of a Weierstrass semigroup?*

Baker and Norine (2007)

divisors on **graphs**

Gathmann and Kerber (2008)

**metric graphs**

Mikhalkin and Zharkov (2008) (i.e. abstract tropical curves)

### Question

*What is the tropical analogue of a Weierstrass semigroup?*

**graph** := finite connected multigraph with no loops



Baker and Norine (2007)

divisors on **graphs**

Gathmann and Kerber (2008)

**metric graphs**

Mikhalkin and Zharkov (2008) (i.e. abstract tropical curves)

### Question

*What is the tropical analogue of a Weierstrass semigroup?*

**graph** := finite connected multigraph with no loops

**simple graph** := graph with no multiple edges

# Divisor theory on graphs

Let  $G$  be a graph with  $n$  vertices,

# Divisor theory on graphs

Let  $G$  be a graph with  $n$  vertices, a **divisor** is a formal sum of the vertices  $V(G)$

$$D = a_1 P_1 + a_2 P_2 + \cdots + a_n P_n$$

where  $a_i \in \mathbb{Z}$ .

# Divisor theory on graphs

Let  $G$  be a graph with  $n$  vertices, a **divisor** is a formal sum of the vertices  $V(G)$

$$D = a_1 P_1 + a_2 P_2 + \cdots + a_n P_n$$

where  $a_i \in \mathbb{Z}$ . The group of divisors  $\text{Div}(G)$  is the free abelian group on  $V(G)$ .

# Divisor theory on graphs


Let  $G$  be a graph with  $n$  vertices, a **divisor** is a formal sum of the vertices  $V(G)$

$$D = a_1 P_1 + a_2 P_2 + \cdots + a_n P_n$$

where  $a_i \in \mathbb{Z}$ . The group of divisors  $\text{Div}(G)$  is the free abelian group on  $V(G)$ .

## Example

$$D = 3P_1 + P_2 + P_3$$



# Divisor theory on graphs


Let  $G$  be a graph with  $n$  vertices, a **divisor** is a formal sum of the vertices  $V(G)$

$$D = a_1 P_1 + a_2 P_2 + \cdots + a_n P_n$$

where  $a_i \in \mathbb{Z}$ . The group of divisors  $\text{Div}(G)$  is the free abelian group on  $V(G)$ .

## Example

$$D = 3P_1 + P_2 + P_3$$




# Linear equivalence

Linear equivalence can be thought in terms of **chip firing**:

# Linear equivalence


Linear equivalence can be thought in terms of **chip firing**:






# Linear equivalence

Linear equivalence can be thought in terms of **chip firing**:




# Linear equivalence

Linear equivalence can be thought in terms of **chip firing**:




# Linear equivalence

Linear equivalence can be thought in terms of **chip firing**:




# Linear equivalence

Linear equivalence can be thought in terms of **chip firing**:




# Linear equivalence

Linear equivalence can be thought in terms of **chip firing**:





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$



# Principal divisors



Let  $f : V(G) \rightarrow \mathbb{Z}$







# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$



# Principal divisors



Let  $f : V(G) \rightarrow \mathbb{Z}$







# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$





# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$



# Principal divisors

Let  $f : V(G) \rightarrow \mathbb{Z}$



Two divisors  $D, D' \in \text{Div}(G)$  are **linearly equivalent** if

$$D - D' = \Delta f \quad \text{for some } f : V(G) \rightarrow \mathbb{Z}.$$

Notation:  $D(P)$  = coefficient of  $D$  at  $P$ .

Notation:  $D(P)$  = coefficient of  $D$  at  $P$ .

$D \geq D'$  if and only if  $D(P) \geq D'(P)$  for every  $P \in V(G)$ .

Notation:  $D(P)$  = coefficient of  $D$  at  $P$ .

$D \geq D'$  if and only if  $D(P) \geq D'(P)$  for every  $P \in V(G)$ .

$D$  is **effective** if  $D \geq 0$ .

Notation:  $D(P)$  = coefficient of  $D$  at  $P$ .

$D \geq D'$  if and only if  $D(P) \geq D'(P)$  for every  $P \in V(G)$ .

$D$  is **effective** if  $D \geq 0$ .

The **degree** of  $D$  is  $\deg(D) = \sum_{P \in V(G)} D(P)$ .

Notation:  $D(P)$  = coefficient of  $D$  at  $P$ .

$D \geq D'$  if and only if  $D(P) \geq D'(P)$  for every  $P \in V(G)$ .

$D$  is **effective** if  $D \geq 0$ .

The **degree** of  $D$  is  $\deg(D) = \sum_{P \in V(G)} D(P)$ .

Denote by  $\text{Div}_+^d(G)$  the set of effective of divisors of degree  $d$ .



The **linear system** of a divisor  $D \in \text{Div}(G)$  is

$$|D| = \{E \in \text{Div}(G) : D \sim E, E \geq 0\}.$$

The **linear system** of a divisor  $D \in \text{Div}(G)$  is

$$|D| = \{E \in \text{Div}(G) : D \sim E, E \geq 0\}.$$

The **rank** of  $D$  is  $-1$  if  $|D| = \emptyset$ , otherwise

$$r(D) = \max\{d \in \mathbb{N} : |D - E| \neq \emptyset, \forall E \in \text{Div}_+^d(G)\}.$$

# Weierstrass sets

Recall (for curves):

$$\begin{aligned} H(P) &= \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\} \\ &= \{n \in \mathbb{N} : r(nP) > r((n-1)P)\} \end{aligned}$$

# Weierstrass sets

Recall (for curves):

$$\begin{aligned} H(P) &= \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\} \\ &= \{n \in \mathbb{N} : r(nP) > r((n-1)P)\} \end{aligned}$$

Definition (Kang, Matthews, Peachey 2020)

Let  $G$  be a graph and let  $P \in V(G)$ .

**Rank Weierstrass set:**

$$H_r(P) = \{n \in \mathbb{N} : r(nP) > r((n-1)P)\}$$

**Functional Weierstrass set:**

$$H_f(P) = \{n \in \mathbb{N} : \exists f \text{ such that } \Delta f + nP \geq 0, \Delta f(P) = -n\}$$

# Weierstrass sets

Recall (for curves):

$$\begin{aligned} H(P) &= \{n \in \mathbb{N} : \exists f \in K(X) \text{ regular on } X \setminus \{P\}, \text{ord}_P(f) = -n\} \\ &= \{n \in \mathbb{N} : r(nP) > r((n-1)P)\} \end{aligned}$$

Definition (Kang, Matthews, Peachey 2020)

Let  $G$  be a graph and let  $P \in V(G)$ .

**Rank Weierstrass set:**

$$H_r(P) = \{n \in \mathbb{N} : r(nP) > r((n-1)P)\}$$

**Functional Weierstrass set:**

$$H_f(P) = \{n \in \mathbb{N} : \exists f \text{ such that } \Delta f + nP \geq 0, \Delta f(P) = -n\}$$

For curves:  $H_r(P) = H_f(P) = H(P)$ ,

For graphs:  $H_f(P) \setminus H_r(P)$  can be arbitrarily large!

# Which one is the best?

# Which one is the best?

The **genus** of a graph  $G$  is  $g = |E(G)| - |V(G)| + 1$ .

Lemma (Tropical Weierstrass Gap Theorem)

$$|\mathbb{N} \setminus H_r(P)| = g$$

*Not true for  $H_f(P)$ .*

# Which one is the best?

The **genus** of a graph  $G$  is  $g = |E(G)| - |V(G)| + 1$ .

Lemma (Tropical Weierstrass Gap Theorem)

$$|\mathbb{N} \setminus H_r(P)| = g$$


*Not true for  $H_f(P)$ .*

$H_f(P)$  is a semigroup,  $H_r(P)$  is *not*.



## Example

Consider the following graph  $G$



It is the vertex gluing of  $K_{2,3}$  and two copies of  $K_{2,2}$ .  
Let  $P \in V(G)$  be the vertex of degree 7. Then

$$H_r(P) = \{0, 3, 5, 7\} \cup (8 + \mathbb{N}).$$

Note that  $H_r(P)$  is not a semigroup  $6 = 3 + 3 \notin H_r(P)$ .

This result was conjectured by Kang, Matthews and Peachey:

Theorem (B. 2022)

Let  $G$  be a *simple* graph. For every  $P \in V(G)$

$$H_r(P) \subseteq H_f(P)$$

This result was conjectured by Kang, Matthews and Peachey:

Theorem (B. 2022)

Let  $G$  be a *simple* graph. For every  $P \in V(G)$

$$H_r(P) \subseteq H_f(P)$$

Let  $K_{n+1}$  be the complete graph on  $n + 1$  vertices.

Lemma (Kang, Matthews, Peachey 2020)

For every  $P \in V(K_{n+1})$   $H_f(P) = \langle n, n + 1 \rangle$ .

This result was conjectured by Kang, Matthews and Peachey:

Theorem (B. 2022)

Let  $G$  be a *simple* graph. For every  $P \in V(G)$

$$H_r(P) \subseteq H_f(P)$$

Let  $K_{n+1}$  be the complete graph on  $n + 1$  vertices.

Lemma (Kang, Matthews, Peachey 2020)

For every  $P \in V(K_{n+1})$   $H_f(P) = \langle n, n + 1 \rangle$ .

$H_r(P) \subseteq H_f(P)$  and  $|\mathbb{N} \setminus H_r(P)| = g(K_{n+1})$  imply:

Corollary

For every  $P \in V(K_{n+1})$   $H_r(P) = H_f(P) = \langle n, n + 1 \rangle$ .

Let  $K_{m,n}$  be the complete bipartite graph.

### Proposition

For every  $P \in V(K_{m,n})$

$$H_r(P) = H_f(P) = n\mathbb{N} \cup (n(m-1) + \mathbb{N})$$

Let  $K_{m,n}$  be the complete bipartite graph.

### Proposition


For every  $P \in V(K_{m,n})$

$$H_r(P) = H_f(P) = n\mathbb{N} \cup (n(m-1) + \mathbb{N})$$

### Question


*Under which conditions on  $G$  we have  $H_r(P) = H_f(P)$ ?*

# Vertex gluing



**Vertex gluing:** the graph  $G$  obtained by identifying  $v_1$  and  $v_1$

# Vertex gluing




**Vertex gluing:** the graph  $G$  obtained by identifying  $v_1$  and  $v_1$

Proposition

$$H_f^G(v) = H_f^{G_1}(v_1) + H_f^{G_2}(v_2)$$



# Vertex gluing



**Vertex gluing:** the graph  $G$  obtained by identifying  $v_1$  and  $v_1$

## Proposition

$$H_f^G(v) = H_f^{G_1}(v_1) + H_f^{G_2}(v_2)$$

## Theorem (B. 2022)

*Functional Weierstrass sets of graphs*  $\longleftrightarrow$  *submonoids of  $\mathbb{N}$*

*Functional Weierstrass sets of **simple** graphs*  $\longleftrightarrow$  *numerical semigroups*

Fix  $P \in V(G)$ , let  $\lambda_P : \mathbb{N} \rightarrow \mathbb{N}$  defined by

$$\lambda_P(k) = \min\{n \in \mathbb{N} : r(nP) = k\}.$$

Note that  $\lambda_P$  completely determines  $H_r(P)$  and vice versa.

Fix  $P \in V(G)$ , let  $\lambda_P : \mathbb{N} \rightarrow \mathbb{N}$  defined by

$$\lambda_P(k) = \min\{n \in \mathbb{N} : r(nP) = k\}.$$

Note that  $\lambda_P$  completely determines  $H_r(P)$  and vice versa.

### Proposition

$$\lambda_v^G(k) = \max\{\lambda_{v_1}^{G_1}(k_1) + \lambda_{v_2}^{G_2}(k_2) : k_1 + k_2 = k\}$$

Fix  $P \in V(G)$ , let  $\lambda_P : \mathbb{N} \rightarrow \mathbb{N}$  defined by

$$\lambda_P(k) = \min\{n \in \mathbb{N} : r(nP) = k\}.$$

Note that  $\lambda_P$  completely determines  $H_r(P)$  and vice versa.

### Proposition

$$\lambda_v^G(k) = \max\{\lambda_{v_1}^{G_1}(k_1) + \lambda_{v_2}^{G_2}(k_2) : k_1 + k_2 = k\}$$

### Theorem (B. 2022)

Let  $e_1 \geq e_2 \geq \dots \geq e_n \geq 0$  be integers, and set  $s_i = \sum_{j=1}^i e_j$ .  
There exists a *simple* graph  $G$  with a vertex  $P \in V(G)$  such that

$$H_r(P) = \{0, s_1, \dots, s_{n-1}\} \cup (s_n + \mathbb{N})$$

Thank you very much!