



# One-Parametric Presburger Arithmetic has Quantifier Elimination

**Alessio Mansutti and Mikhail Starchak**

MFCS 2025



## Peano arithmetic

$\mathbb{Z}$  +  
 $\times$  A  
 $\leq$  E





## Peano arithmetic

$\mathbb{Z}$  +  
 $\times$  A  
 $\leq$  E



## Presburger arithmetic

$\mathbb{Z}$   $\leq$   
A +  
E



## Peano arithmetic

$\mathbb{Z}$     +  
          $\times$      $\forall$   
          $\leq$      $\exists$



$\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$

## Presburger arithmetic

$\mathbb{Z}$      $\leq$   
         +  
          $\forall$      $\exists$



## Peano arithmetic

$\mathbb{Z}$     $+$   
 $\times$     $\wedge$   
 $\leq$     $\exists$



$\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$

## Presburger arithmetic

$\mathbb{Z}$     $\leq$   
 $\wedge$     $+$   
 $\exists$



## One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.


# One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

**Twisting squares (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)**

$$|2x + (2t - 2)y| \leq t^2 - 2t + 2 \wedge |(2 - 2t)x + 2y| \leq t^2 - 2t + 2$$





# One-parametric Presburger arithmetic (1PPA)


First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

## Twisting squares (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)

$$|2x + (2t - 2)y| \leq t^2 - 2t + 2 \wedge |(2 - 2t)x + 2y| \leq t^2 - 2t + 2$$

$t = 0:$        $|2x - 2y| \leq 2 \wedge |2x + 2y| \leq 2$       5 solutions



# One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

## Twisting squares (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)


$$|2x + (2t - 2)y| \leq t^2 - 2t + 2 \wedge |(2 - 2t)x + 2y| \leq t^2 - 2t + 2$$

$t = 0:$   $|2x - 2y| \leq 2 \wedge |2x + 2y| \leq 2$

5 solutions

$t = 1:$   $|2x| \leq 1 \wedge |2y| \leq 1$

1 solution



# One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

## Twisting squares (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)

$$|2x + (2t - 2)y| \leq t^2 - 2t + 2 \wedge |(2 - 2t)x + 2y| \leq t^2 - 2t + 2$$

$t = 0:$   $|2x - 2y| \leq 2 \wedge |2x + 2y| \leq 2$


$t = 1:$   $|2x| \leq 1 \wedge |2y| \leq 1$

$t = 2:$   $|2x + 2y| \leq 2 \wedge |-2x + 2y| \leq 2$

5 solutions

1 solution

same as  $t = 0$



# One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

## Twisting squares (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)

$$|2x + (2t - 2)y| \leq t^2 - 2t + 2 \wedge |(2 - 2t)x + 2y| \leq t^2 - 2t + 2$$

$t = 0$ :  $|2x - 2y| \leq 2 \wedge |2x + 2y| \leq 2$

5 solutions

$t = 1$ :  $|2x| \leq 1 \wedge |2y| \leq 1$


1 solution

$t = 2$ :  $|2x + 2y| \leq 2 \wedge |-2x + 2y| \leq 2$

same as  $t = 0$

$t = 3$ :  $|2x + 4y| \leq 5 \wedge |-4x + 2y| \leq 5$

5 solutions





# One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .


In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

## Twisting squares (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)

$$|2x + (2t - 2)y| \leq t^2 - 2t + 2 \wedge |(2 - 2t)x + 2y| \leq t^2 - 2t + 2$$

For a fixed  $t \geq 0$ , this formula:

- has  $t^2 - 2t + 5$  solutions when  $t$  is **even**
- has  $t^2 - 2t + 2$  solutions when  $t$  is **odd**



# One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

## “Chinese Remainder Theorem”

Let  $f, g \in \mathbb{Z}[t]$ . The following formula is valid:

$$\underbrace{(f \geq 1 \wedge g \geq 1 \wedge \exists u, v : f \cdot u + g \cdot v = 1)}_{f(t) \text{ and } g(t) \text{ are positive and coprime}} \implies \forall a \forall b \exists x : \underbrace{\begin{array}{l} 0 \leq x < f \cdot g \\ \wedge f \mid x - a \\ \wedge g \mid x - b \end{array}}_{\text{CRT}}$$

where  $(f \mid \tau) := \exists w (w \cdot f = \tau)$ .

# One-parametric Presburger arithmetic (1PPA)

First-order theory of the structure  $\langle \mathbb{Z}, 0, 1, +, (x \mapsto t \cdot x), \leq \rangle$ .

In the **multiplication function**  $x \mapsto t \cdot x$ , the **parameter**  $t$  is a fixed free variable.

A formula  $\varphi(\mathbf{x})$  of 1PPA defines a **parametric Presburger family**  $\{\llbracket \varphi \rrbracket_k : k \in \mathbb{Z}\}$ , where

$\llbracket \varphi \rrbracket_k$ : set of solutions to  $\varphi$  after replacing  $t$  with  $k$

We can ask several questions about  $\varphi$ :

- **satisfiability**: is  $\llbracket \varphi \rrbracket_k$  non-empty for **some**  $k$ ?
- **universality**: is  $\llbracket \varphi \rrbracket_k$  non-empty for **every**  $k$ ?
- **finiteness**: is  $\llbracket \varphi \rrbracket_k$  non-empty only for **finitely many**  $k$ ?

# Eventual quasi-polynomials and 1PPA

## Theorem (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)

Let  $\varphi$  be a 1PPA formula. The counting function  $f(k) := \#\llbracket \varphi \rrbracket_k$  is an EQP.

A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is an **eventual quasi-polynomial (EQP)** whenever there are

- a **threshold**  $T$  and a **period**  $P$ , and
- a family of univariate polynomials  $f_0, \dots, f_{P-1}$

such that for every  $n \geq T$ ,  $f(n) = f_{(n \bmod P)}(n)$ .



# Eventual quasi-polynomials and 1PPA

## Theorem (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)


Let  $\varphi$  be a 1PPA formula. The counting function  $f(k) := \#\llbracket \varphi \rrbracket_k$  is an EQP.

A function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is an **eventual quasi-polynomial (EQP)** whenever there are

- a **threshold**  $T$  and a **period**  $P$ , and
- a family of univariate polynomials  $f_0, \dots, f_{P-1}$

such that for every  $n \geq T$ ,  $f(n) = f_{(n \bmod P)}(n)$ .

Examples:



$$\left\lfloor \frac{x}{2} \right\rfloor = \begin{cases} \frac{x}{2} & \text{if } x \text{ is even} \\ \frac{x-1}{2} & \text{if } x \text{ is odd} \end{cases}$$

## Eventual quasi-polynomials and 1PPA

**Theorem (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)**

*Let  $\varphi$  be a 1PPA formula. The counting function  $f(k) := \# \llbracket \varphi \rrbracket_k$  is an EQP.*

## Eventual quasi-polynomials and 1PPA

**Theorem (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)**

*Let  $\varphi$  be a 1PPA formula. The counting function  $f(k) := \# \llbracket \varphi \rrbracket_k$  is an EQP.*

*Proof idea:* Show the result for quantifier-free formulae. Then,

$$\varphi = \exists x_1 \forall x_2 \dots : \psi$$

# Eventual quasi-polynomials and 1PPA

## Theorem (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)

Let  $\varphi$  be a 1PPA formula. The counting function  $f(k) := \# \llbracket \varphi \rrbracket_k$  is an EQP.

*Proof idea:* Show the result for quantifier-free formulae. Then,

$$\varphi = \exists x_1 \forall x_2 \dots : \psi$$

*bounded quantifier elimination* (Weispfenning. *ISSAC* 1997)

" $\exists y \leq p(t)$ " constrains  $y$  in  $[0..p(t)]$

$$\varphi \equiv \exists y_1 \leq p_1(t) \forall y_2 \leq p_2(t) \dots : \gamma$$




# Eventual quasi-polynomials and 1PPA

## Theorem (Bogart, Goodrick, Woods. *Discrete Analysis* 2017)

Let  $\varphi$  be a 1PPA formula. The counting function  $f(k) := \# \llbracket \varphi \rrbracket_k$  is an EQP.

*Proof idea:* Show the result for quantifier-free formulae. Then,



# Eventual quasi-polynomials and 1PPA

Theorem (Bogart, Goodrick, Woods, *Discrete Analysis* 2017)

Let  $\varphi$

In *Discrete Analysis* 2017, Bogart, Goodrick and Woods ask whether the **parsimonious transformation** can be replaced with **quantifier elimination**.

Proof

$$\varphi = \exists x_1 \forall x_2 \dots : \psi$$

↓ **bounded quantifier elimination** (Weispfenning. *ISSAC* 1997)

“ $\exists y \leq p(t)$ ” constrains  $y$  in  $[0..p(t)]$

$$\varphi \equiv \exists y_1 \leq p_1(t) \forall y_2 \leq p_2(t) \dots : \gamma$$

↓ **parsimonious transformation** (Chen, Li, Sam. *Trans. Amer. Math. Soc.* 2012)

$\# \llbracket \varphi \rrbracket_k = \# \llbracket \varphi' \rrbracket_k$  for every  $k$

↓  $\varphi'$  quantifier-free

# Eventual quasi-polynomials and 1PPA

Theorem (Bogart, Goodrick, Woods, *Discrete Analysis* 2017)

Let  $\varphi$

In *Discrete Analysis* 2017, Bogart, Goodrick and Woods ask whether the **parsimonious transformation** can be replaced with **quantifier elimination**.

Proof

In *Arch. Math. Logic* 2018, Goodrick conjectures that extending 1PPA with a function  $x \mapsto \left\lfloor \frac{x}{p(t)} \right\rfloor$  for every polynomial  $p$  suffices.

$$\varphi \equiv \exists y_1 \leq p_1(t) \forall y_2 \leq p_2(t) \dots : \gamma$$

**parsimonious transformation** (Chen, Li, Sam. *Trans. Amer. Math. Soc.* 2012)

$$\# \llbracket \varphi \rrbracket_k = \# \llbracket \varphi' \rrbracket_k \text{ for every } k$$

$\varphi'$  quantifier-free

# Eventual quasi-polynomials and 1PPA

Theorem (Bogart, Goodrick, Woods, *Discrete Analysis* 2017)

Let  $\varphi$

In *Discrete Analysis* 2017, Bogart, Goodrick and Woods ask whether the **parsimonious transformation** can be replaced with **quantifier elimination**.

Proof

In *Arch. Math. Logic* 2018, Goodrick conjectures that extending 1PPA with a function  $x \mapsto \left\lfloor \frac{x}{p(t)} \right\rfloor$  for every polynomial  $p$  suffices.

$\varphi \equiv \exists y_1 \leq p_1(t) \forall y_2 \leq p_2(t) \dots : \gamma$

**parsimonious transformation**

We prove Goodrick's conjecture.

(Goodrick, *Trans. Amer. Math. Soc.* 2012)

$\# \llbracket \varphi \rrbracket_{\mathbb{N}^k} = \# \llbracket \varphi' \rrbracket_{\mathbb{N}^k}$

$\varphi'$  quantifier-free

# Our results

## Theorem

*There is a quantifier elimination procedure for the extension of 1PPA with the functions:*

- *integer division:*  $x \mapsto \left\lfloor \frac{x}{t^d} \right\rfloor$                       *one function for each  $d \in \mathbb{N}$ , assuming  $t \neq 0$*
- *integer remainder function:*  $x \mapsto (x \bmod p)$                       *for each  $p \in \mathbb{Z}[t]$*
- *divisibility relation:*  $p \mid x$                       *for each  $p \in \mathbb{Z}[t]$*

(The functions  $x \mapsto \left\lfloor \frac{x}{p(t)} \right\rfloor$  capture all these functions and relations.)

# Our results

## Theorem

There is a quantifier elimination procedure for the extension of 1PPA with the functions:

- *integer division*:  $x \mapsto \left\lfloor \frac{x}{t^d} \right\rfloor$  one function for each  $d \in \mathbb{N}$ , assuming  $t \neq 0$
- *integer remainder function*:  $x \mapsto (x \bmod p)$  for each  $p \in \mathbb{Z}[t]$
- *divisibility relation*:  $p \mid x$  for each  $p \in \mathbb{Z}[t]$

(The functions  $x \mapsto \left\lfloor \frac{x}{p(t)} \right\rfloor$  capture all these functions and relations.)

## Theorem

For the class of all *existential* formulae of 1PPA, the following holds:

<i>Satisfiability:</i> <b>NP-complete</b>	<i>Universality:</i> <b>coNEXP-complete</b>	<i>Finiteness:</i> <b>coNP-complete</b>
--	--	--

## Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi(\mathbf{z})$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

## Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi(\mathbf{z})$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

*Step 1. Preprocessing:* Remove divisions and remainder functions

$$\cdots + \left\lfloor \frac{\tau}{t^d} \right\rfloor + \cdots \leq 0 \quad \rightarrow \quad \exists x \left( \cdots + x + \cdots \leq 0 \wedge (t^d x \leq \tau < t^d(x+1)) \right)$$

$$\cdots + (\tau \bmod p) + \cdots \leq 0 \quad \rightarrow \quad \exists x \left( \cdots + x + \cdots \leq 0 \wedge (0 \leq x < p-1) \wedge (p \mid \tau - x) \right)$$



# Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi(\mathbf{z})$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

*Step I. Preprocessing:* Remove divisions and remainder functions

$$\cdots + \left\lfloor \frac{\tau}{t^d} \right\rfloor + \cdots \leq 0 \quad \rightarrow \quad \exists x \left( \cdots + x + \cdots \leq 0 \wedge (t^d x \leq \tau < t^d(x+1)) \right)$$

$$\cdots + (\tau \bmod p) + \cdots \leq 0 \quad \rightarrow \quad \exists x \left( \cdots + x + \cdots \leq 0 \wedge (0 \leq x < p-1) \wedge (p \mid \tau - x) \right)$$

*Step II. Bounded quantifier elimination:*

$$\exists \mathbf{x}' : \varphi'(\mathbf{x}', \mathbf{z}) \quad \rightarrow \quad \exists \mathbf{w} \leq B \bigvee_i \gamma_i(\mathbf{w}, \mathbf{z})$$

# Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Before 2024:** Quantifier elimination procedures for Presburger arithmetic required **2EXPTIME** / **NEXPTIME** to remove one block of existential quantifiers.

$$\cdots + \left\lfloor \frac{\tau}{t^d} \right\rfloor + \cdots \leq 0 \quad \rightarrow \quad \exists x \left( \cdots + x + \cdots \leq 0 \wedge (t^d x \leq \tau < t^d(x+1)) \right)$$

$$\cdots + (\tau \bmod p) + \cdots \leq 0 \quad \rightarrow \quad \exists x \left( \cdots + x + \cdots \leq 0 \wedge (0 \leq x < p-1) \wedge (p \mid \tau - x) \right)$$

*Step II. Bounded quantifier elimination:*

$$\exists \mathbf{x}' : \varphi'(\mathbf{x}', \mathbf{z}) \quad \rightarrow \quad \exists \mathbf{w} \leq B \bigvee_i \gamma_i(\mathbf{w}, \mathbf{z})$$

# Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(x, z)$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Before 2024:** Quantifier elimination procedures for Presburger arithmetic required **2EXPTIME** / **NEXPTIME** to remove one block of existential quantifiers.

**In ICALP'24:** Two different procedures running in **EXPTIME** / **NP** were found, by [Chistikov, M., Starchak] and [Haase, Krishna, Madnani, Mishra, Zetsche].

*Step II. Bounded quantifier elimination:*

We extend the quantifier elimination procedure from [Chistikov, M., Starchak] from Presburger arithmetic to one-parametric Presburger arithmetic.

## Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

*Step I. Preprocessing:* Remove divisions and remainder functions.

*Step II. Bounded quantifier elimination:* compute  $\exists \mathbf{w} \leq B : \bigvee_i \gamma_i(\mathbf{w}, \mathbf{z})$ .

## Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

*Step I. Preprocessing:* Remove divisions and remainder functions.

*Step II. Bounded quantifier elimination:* compute  $\exists \mathbf{w} \leq B : \bigvee_i \gamma_i(\mathbf{w}, \mathbf{z})$ .

*Step III. Remove the divisibility relations:*

$$f \mid \tau(\mathbf{w}) + \sigma(\mathbf{z}) \quad \rightarrow \quad f \mid \tau(\mathbf{w}) + (\sigma(\mathbf{z}) \bmod f)$$

## Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

*Step I. Preprocessing:* Remove divisions and remainder functions.

*Step II. Bounded quantifier elimination:* compute  $\exists \mathbf{w} \leq B : \bigvee_i \gamma_i(\mathbf{w}, \mathbf{z})$ .

*Step III. Remove the divisibility relations:*

$$f \mid \tau(\mathbf{w}) + \sigma(\mathbf{z}) \quad \rightarrow \quad f \mid \tau(\mathbf{w}) + (\sigma(\mathbf{z}) \bmod f)$$

Bounded!

## Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

*Step I. Preprocessing:* Remove divisions and remainder functions.

*Step II. Bounded quantifier elimination:* compute  $\exists \mathbf{w} \leq B : \bigvee_i \gamma_i(\mathbf{w}, \mathbf{z})$ .

*Step III. Remove the divisibility relations:*

$$\begin{aligned} f \mid \tau(\mathbf{w}) + \sigma(\mathbf{z}) &\rightarrow f \mid \tau(\mathbf{w}) + (\sigma(\mathbf{z}) \bmod f) \\ &\rightarrow \exists w \leq p(t) : f \cdot w = \tau(\mathbf{w}) + (\sigma(\mathbf{z}) \bmod f) \end{aligned}$$

## Overview of our procedure

**Input:** A quantifier-free formula  $\varphi(\mathbf{x}, \mathbf{z})$  from the extended language of 1PPA (1PPA<sup>+</sup>).

**Output:** A quantifier-free formula  $\psi$  from 1PPA<sup>+</sup> that is equivalent to  $\exists \mathbf{x} \varphi$ .

*Step I. Preprocessing:* Remove divisions and remainder functions.

*Step II. Bounded quantifier elimination:* compute  $\exists \mathbf{w} \leq B : \bigvee_i \gamma_i(\mathbf{w}, \mathbf{z})$ .

*Step III. Remove the divisibility relations:*

$$\begin{aligned} f \mid \tau(\mathbf{w}) + \sigma(\mathbf{z}) &\rightarrow f \mid \tau(\mathbf{w}) + (\sigma(\mathbf{z}) \bmod f) \\ &\rightarrow \exists w \leq p(t) : f \cdot w = \tau(\mathbf{w}) + (\sigma(\mathbf{z}) \bmod f) \end{aligned}$$

*Step IV. Elimination of bounded quantifiers by “bit blasting”.*



## Step IV: Elimination of bounded quantifiers

$$\exists x \leq t^2 + t - 1 \ \exists z \leq t + 2 : (t + 1) \cdot z = x + (-b \bmod t + 1)$$

Assume  $t \geq 2$ .

## Step IV: Elimination of bounded quantifiers

$$\exists x \leq t^2 + t - 1 \exists z \leq t + 2 : (t + 1) \cdot z = x + (-b \bmod t + 1)$$

Assume  $t \geq 2$ . Bit blast:

$$\begin{aligned} \exists z \leq t + 2 : \varphi \quad \rightarrow \quad \exists z_0, z_1, z_2 \leq t - 1 : \quad & 0 \leq z_2 \cdot t^2 + z_1 \cdot t + z_0 \leq t + 2 \\ & \wedge \varphi[z_2 \cdot t^2 + z_1 \cdot t + z_0 / z] \end{aligned}$$

## Step IV: Elimination of bounded quantifiers

$$\exists x \leq t^2 + t - 1 \exists z \leq t + 2 : (t + 1) \cdot z = x + (-b \bmod t + 1)$$

Assume  $t \geq 2$ . **Bit blast:**

$$\begin{aligned} \exists z \leq t + 2 : \varphi \quad \rightarrow \quad \exists z_0, z_1, z_2 \leq t - 1 : \quad & 0 \leq z_2 \cdot t^2 + z_1 \cdot t + z_0 \leq t + 2 \\ & \wedge \varphi[z_2 \cdot t^2 + z_1 \cdot t + z_0 / z] \end{aligned}$$

The equality  $(t + 1) \cdot z = x - (b \bmod t + 1)$  becomes:

$$(t + 1) \cdot (z_2 \cdot t^2 + z_1 \cdot t + z_0) = (x_2 \cdot t^2 + x_1 \cdot t + x_0) + (-b \bmod t + 1).$$

## Step IV: Elimination of bounded quantifiers

$$\exists x \leq t^2 + t - 1 \exists z \leq t + 2 : (t + 1) \cdot z = x + (-b \bmod t + 1)$$

Assume  $t \geq 2$ . **Bit blast:**

$$\begin{aligned} \exists z \leq t + 2 : \varphi \quad \rightarrow \quad \exists z_0, z_1, z_2 \leq t - 1 : \quad & 0 \leq z_2 \cdot t^2 + z_1 \cdot t + z_0 \leq t + 2 \\ & \wedge \varphi[z_2 \cdot t^2 + z_1 \cdot t + z_0 / z] \end{aligned}$$

The equality  $(t + 1) \cdot z = x - (b \bmod t + 1)$  becomes:

$$-z_2 \cdot t^3 + (x_2 - z_1 - z_2) \cdot t^2 + (x_1 - z_0 - z_1) \cdot t + (x_0 - z_0) + (-b \bmod t + 1) = 0.$$

## Step IV: Elimination of bounded quantifiers

$$\exists x \leq t^2 + t - 1 \exists z \leq t + 2 : (t + 1) \cdot z = x + (-b \bmod t + 1)$$

Assume  $t \geq 2$ . **Bit blast:**

$$\begin{aligned} \exists z \leq t + 2 : \varphi \quad \rightarrow \quad \exists z_0, z_1, z_2 \leq t - 1 : \quad & 0 \leq z_2 \cdot t^2 + z_1 \cdot t + z_0 \leq t + 2 \\ & \wedge \varphi[z_2 \cdot t^2 + z_1 \cdot t + z_0 / z] \end{aligned}$$

The equality  $(t + 1) \cdot z = x - (b \bmod t + 1)$  becomes:

$$-z_2 \cdot t^3 + (x_2 - z_1 - z_2) \cdot t^2 + (x_1 - z_0 - z_1) \cdot t + (x_0 - z_0) + (-b \bmod t + 1) = 0.$$

**Divide by  $t$**  the maximal subterm with no quantified variables:

$$(-b \bmod t + 1) \quad \rightarrow \quad \left\lfloor \frac{-b \bmod t + 1}{t} \right\rfloor \cdot t + ((-b \bmod t + 1) \bmod t)$$

## Step IV: Elimination of bounded quantifiers

$$\begin{aligned} -z_2 \cdot t^3 + (x_2 - z_1 - z_2) \cdot t^2 + \left( x_1 - z_0 - z_1 + \left\lfloor \frac{-b \bmod t + 1}{t} \right\rfloor \right) \cdot t \\ + (x_0 - z_0) + ((-b \bmod t + 1) \bmod t) = 0 \end{aligned}$$

## Step IV: Elimination of bounded quantifiers

$$\begin{aligned} -z_2 \cdot t^3 + (x_2 - z_1 - z_2) \cdot t^2 + \left( x_1 - z_0 - z_1 + \left\lfloor \frac{-b \bmod t + 1}{t} \right\rfloor \right) \cdot t \\ + (x_0 - z_0) + ((-b \bmod t + 1) \bmod t) = 0 \end{aligned}$$

- $(x_0 - z_0) + ((-b \bmod t + 1) \bmod t)$  belongs to  $[-t..2 \cdot t]$ ...
- ...and must be divisible by  $t$ . (This only applies to equalities.)

## Step IV: Elimination of bounded quantifiers

$$\begin{aligned} -z_2 \cdot t^3 + (x_2 - z_1 - z_2) \cdot t^2 + \left( x_1 - z_0 - z_1 + \left\lfloor \frac{-b \bmod t + 1}{t} \right\rfloor \right) \cdot t \\ + (x_0 - z_0) + ((-b \bmod t + 1) \bmod t) = 0 \end{aligned}$$

- $(x_0 - z_0) + ((-b \bmod t + 1) \bmod t)$  belongs to  $[-t..2 \cdot t]$ ...
- ...and must be divisible by  $t$ . (This only applies to equalities.)

**Guess**  $r_0 \in \{-1, 0, 1, 2\}$  and rewrite the equality as

$$\begin{aligned} -z_2 \cdot t^2 + (x_2 - z_1 - z_2) \cdot t + \left( x_1 - z_0 - z_1 + \left\lfloor \frac{-b \bmod t + 1}{t} \right\rfloor \right) + r_0 = 0 \\ \wedge (x_0 - z_0) + ((-b \bmod t + 1) \bmod t) = r_0 \cdot t \end{aligned}$$

**Important:**  $z_0, x_0$  and  $x_1$  now only have integer coefficients!



## Step IV: Elimination of bounded quantifiers

$$-z_2 \cdot t^3 + (x_2 - z_1 - z_2) \cdot t^2 + \left( x_1 - z_0 - z_1 + \left\lfloor \frac{-b \bmod t+1}{t} \right\rfloor \right) \cdot t$$

**2nd iteration:** Also  $z_1$  and  $x_2$  have integer coefficients.

**3rd iteration:** All variables have integer coefficients.

We can then call a quantifier elimination procedure for Presburger arithmetic!

**Guess**  $r_0 \in \{-1, 0, 1, 2\}$  and rewrite the equality as

$$\begin{aligned} & -z_2 \cdot t^2 + (x_2 - z_1 - z_2) \cdot t + \left( x_1 - z_0 - z_1 + \left\lfloor \frac{-b \bmod t+1}{t} \right\rfloor \right) + r_0 = 0 \\ & \wedge (x_0 - z_0) + ((-b \bmod t + 1) \bmod t) = r_0 \cdot t \end{aligned}$$

**Important:**  $z_0, x_0$  and  $x_1$  now only have integer coefficients!

# Our results

## Theorem

There is a quantifier elimination procedure for the extension of 1PPA with the functions:

- *integer division*:  $x \mapsto \left\lfloor \frac{x}{t^d} \right\rfloor$  one function for each  $d \in \mathbb{N}$ , assuming  $t \neq 0$
- *integer remainder function*:  $x \mapsto (x \bmod p)$  for each  $p \in \mathbb{Z}[t]$
- *divisibility relation*:  $p \mid x$  for each  $p \in \mathbb{Z}[t]$

## Theorem

For the class of all *existential* formulae of 1PPA, the following holds:


<i>Satisfiability:</i> <b>NP</b> -complete	<i>Universality:</i> <b>coNEXP</b> -complete	<i>Finiteness:</i> <b>coNP</b> -complete
---	---	---

# How does the following picture change for 1PPA?

## Quantifier elimination

[Presburger, '29]

$$\exists x : \varphi(x, \mathbf{y})$$




$$\psi(\mathbf{y})$$

3EXPTIME

## Automata


[Büchi, '60]



3EXPTIME

## Geometry

[Ginsburg and Spanier, '66]



3EXPTIME