

Linear arithmetic theories: algorithms and applications

Christoph Haase Alessio Mansutti



ESSLI 2023



This course

Goals:

- Introduction to logical, algorithmic and geometric aspects of arithmetic theories
- Showcase how arithmetic theories can be approached with different techniques (formal logic, automata theory and geometry)

Content:

- Classical algorithms and decision procedures
- Geometric description of the sets definable in arithmetic theories
- Recent research developments

Course overview

Monday	Introduction to linear arithmetic
Tuesday	Linear Programming and Integer Linear Programming
Wednesday	Quantifier elimination procedures
Thursday	Automata-based procedures
Friday	Geometric procedures

Required background

Basic familiarity with following topics is helpful:

- Mathematical logic
- Linear algebra
- Automata theory
- Algorithms and computational complexity

Today's lecture

Introduction to linear arithmetic:


- Applications of arithmetic theories
- Syntax, semantics, normal forms

Convex geometry:


- Basic notions: hyperplanes, (convex) cones and polyhedra, hulls, ...
- Farkas' Lemma
- Minkowski-Weyl Theorem

Applications of arithmetic theories

Maximum flow



Maximum flow



Maximum flow

Directed weighted graph $G = (V, E, w)$ such that $w: E \rightarrow \mathbb{R}$:

- w assigns maximum flow capacity to edges in G
- flow is function $f: E \rightarrow \mathbb{R}$
- value of flow is sum of flow leaving s
- goal: find flow with maximum value

Maximum flow

Directed weighted graph $G = (V, E, w)$ such that $w: E \rightarrow \mathbb{R}$:

- w assigns maximum flow capacity to edges in G
- flow is function $f: E \rightarrow \mathbb{R}$
- value of flow is sum of flow leaving s
- goal: find flow with maximum value

For edge $e \in E$, introduce variables f_e encoding flow conditions:

$$\bigwedge_{e \in E} 0 \leq f_e \leq w(e) \quad \wedge \quad \bigwedge_{v \in V \setminus \{s, t\}} \sum_{(u, v) \in E} f_{u, v} = \sum_{(v, u) \in E} f_{v, u}$$

Maximum flow

Directed weighted graph $G = (V, E, w)$ such that $w: E \rightarrow \mathbb{R}$:

- w assigns maximum flow capacity to edges in G
- flow is function $f: E \rightarrow \mathbb{R}$
- value of flow is sum of flow leaving s
- goal: find flow with maximum value

For edge $e \in E$, introduce variables f_e encoding flow conditions:

$$\begin{aligned} & \text{maximize } \sum_{(s,v) \in E} f_{s,v} \\ & \bigwedge_{e \in E} 0 \leq f_e \leq w(e) \quad \wedge \quad \bigwedge_{v \in V \setminus \{s,t\}} \sum_{(u,v) \in E} f_{u,v} = \sum_{(v,u) \in E} f_{v,u} \end{aligned}$$

The Frobenius problem

Given coins in denominations

$$m_1 < \cdots < m_k \in \mathbb{N},$$

what is the largest value c that cannot be generated? Does such a c exist?



The Frobenius problem

Given coins in denominations

$$m_1 < \cdots < m_k \in \mathbb{N},$$

what is the largest value c that cannot be generated? Does such a c exist?



$$n = m_1 \cdot x_1 + \cdots + m_k \cdot x_k$$

The Frobenius problem

Given coins in denominations

$$m_1 < \cdots < m_k \in \mathbb{N},$$

what is the largest value c that cannot be generated? Does such a c exist?



$$\exists x_1 \exists x_2 \cdots \exists x_k : n = m_1 \cdot x_1 + \cdots + m_k \cdot x_k$$

The Frobenius problem

Given coins in denominations

$$m_1 < \dots < m_k \in \mathbb{N},$$

what is the largest value c that cannot be generated? Does such a c exist?



$$\forall n : c < n \rightarrow \exists x_1 \exists x_2 \dots \exists x_k : n = m_1 \cdot x_1 + \dots + m_k \cdot x_k$$

The Frobenius problem

Given coins in denominations

$$m_1 < \cdots < m_k \in \mathbb{N},$$


what is the largest value c that cannot be generated? Does such a c exist?



$$\exists c : \forall n : c < n \rightarrow \exists x_1 \exists x_2 \cdots \exists x_k : n = m_1 \cdot x_1 + \cdots + m_k \cdot x_k$$


Artificial neural networks

Artificial neuron:



Artificial neural networks

Artificial neuron:




Output:

$$b = \begin{cases} 0 & \text{if } \sum_{i=1}^n w_i \cdot a_i < 0 \\ \sum_{i=1}^n w_i \cdot a_i & \text{otherwise} \end{cases}$$

Artificial neural networks

Artificial neuron:



Output:


$$b = \begin{cases} 0 & \text{if } \sum_{i=1}^n w_i \cdot a_i < 0 \\ \sum_{i=1}^n w_i \cdot a_i & \text{otherwise} \end{cases}$$

In logic for $\mathbf{w} \in \mathbb{R}^n$

$$\Phi_{\mathbf{w}}(\mathbf{x}, y) := (\mathbf{w}^{\top} \cdot \mathbf{x} < 0 \rightarrow y = 0) \wedge (\mathbf{w}^{\top} \cdot \mathbf{x} \geq 0 \rightarrow y = \mathbf{w}^{\top} \cdot \mathbf{x})$$

Artificial neural networks

Artificial neuron:



Output:


$$b = \begin{cases} 0 & \text{if } \sum_{i=1}^n w_i \cdot a_i < 0 \\ \sum_{i=1}^n w_i \cdot a_i & \text{otherwise} \end{cases}$$

In logic for $\mathbf{w} \in \mathbb{R}^n$ and $W \in \mathbb{R}^{m \times n}$:

$$\Phi_{\mathbf{w}}(\mathbf{x}, y) := (\mathbf{w}^\top \cdot \mathbf{x} < 0 \rightarrow y = 0) \wedge (\mathbf{w}^\top \cdot \mathbf{x} \geq 0 \rightarrow y = \mathbf{w}^\top \cdot \mathbf{x})$$


$$\Phi_W(\mathbf{x}, \mathbf{y}) := \bigwedge_{1 \leq i \leq m} \Phi_{\mathbf{w}_i}(\mathbf{x}, y_i) \quad \text{where } W = \begin{pmatrix} \mathbf{w}_1 \\ \vdots \\ \mathbf{w}_m \end{pmatrix}$$

Artificial neural networks



$$\Phi(\mathbf{x}, \mathbf{y}) = \exists z_1 \exists z_2 : \Phi_{W_1}(\mathbf{x}, \mathbf{z}_1) \wedge \Phi_{W_2}(\mathbf{z}_1, \mathbf{z}_2) \wedge \Phi_{W_3}(\mathbf{z}_2, \mathbf{y})$$

Artificial neural networks




$$\Phi(\mathbf{x}, \mathbf{y}) = \exists z_1 \exists z_2 : \Phi_{W_1}(\mathbf{x}, \mathbf{z}_1) \wedge \Phi_{W_2}(\mathbf{z}_1, \mathbf{z}_2) \wedge \Phi_{W_3}(\mathbf{z}_2, \mathbf{y})$$

All inputs giving output (1, 0):

$$\{\mathbf{r} = (r_1, r_2, r_3) \in \mathbb{R}^3 : \Phi[\mathbf{r}/\mathbf{x}, (1, 0)/\mathbf{y}] \text{ is true}\}$$

Artificial neural networks



$$\Phi(\mathbf{x}, \mathbf{y}) = \exists z_1 \exists z_2 : \Phi_{W_1}(\mathbf{x}, \mathbf{z}_1) \wedge \Phi_{W_2}(\mathbf{z}_1, \mathbf{z}_2) \wedge \Phi_{W_3}(\mathbf{z}_2, \mathbf{y})$$

All inputs giving output (1, 0):

$$\{\mathbf{r} = (r_1, r_2, r_3) \in \mathbb{R}^3 : \Phi[\mathbf{r}/\mathbf{x}, (1, 0)/\mathbf{y}] \text{ is true}\}$$

Artificial neuron network outputs probability distribution:

$$\forall x_1 \forall x_2 \forall x_3 \forall y_1 \forall y_2 : \Phi(\mathbf{x}, \mathbf{y}) \rightarrow (y_1 + y_2 = 1 \wedge 0 \leq y_1 \wedge 0 \leq y_2)$$

Twin prime conjecture

There are infinitely many primes p such that $p + 2$ is also prime:

Twin prime conjecture

There are infinitely many primes p such that $p + 2$ is also prime:

$$\forall n \exists p : n < p \wedge \Phi_{\text{prime}}(p) \wedge \Phi_{\text{prime}}(p + 2)$$

Twin prime conjecture

There are infinitely many primes p such that $p + 2$ is also prime:

$$\forall n \exists p : n < p \wedge \Phi_{\text{prime}}(p) \wedge \Phi_{\text{prime}}(p + 2)$$

$$\Phi_{\text{prime}}(x) := x > 1 \wedge \forall y \forall z : x = y \cdot z \rightarrow (y = 1 \vee y = x)$$

What do we want from arithmetic theories?

problem	domain	arithmetic functions	relations	Boolean connectives	quantifiers
max-flow	\mathbb{R}	$+$	$=, \leq$	\wedge	\exists
Frobenius	\mathbb{N}	$+$	$=, <$	\wedge, \rightarrow	$\exists \forall \exists$
ANN	\mathbb{R}	$+$	$=, <$	\wedge, \rightarrow	$\forall \exists$
twin primes	\mathbb{N}	$+, \cdot$	$=, <$	\neg, \vee, \rightarrow	$\forall \exists$

What do we want from arithmetic theories?

problem	domain	arithmetic functions	relations	Boolean connectives	quantifiers
max-flow	\mathbb{R}	$+$	$=, \leq$	\wedge	\exists
Frobenius	\mathbb{N}	$+$	$=, <$	\wedge, \rightarrow	$\exists \forall \exists$
ANN	\mathbb{R}	$+$	$=, <$	\wedge, \rightarrow	$\forall \exists$
twin primes	\mathbb{N}	$+, \cdot$	$=, <$	$\neg, \wedge, \vee, \rightarrow$	$\forall \exists$

Problems of interest:

- **Validity**: Is a given formula true?
- **Satisfiability**: Does a satisfying assignment exist?
- **Optimization**: Maximize an objective function.
- **Geometry**: Properties of sets definable in arithmetic theories.

Syntax and semantics of linear arithmetic theories

Syntax

■ $x, y, z, x_1, \dots, x_n \in X$ are first-order variables

■ Atomic formulas, where $a_1, \dots, a_n, b \in \mathbb{Z}$:

$$a_1 \cdot x_1 + \dots + a_n \cdot x_n = b, \quad \sum_{i=1}^n a_i \cdot x_i \leq b, \quad \mathbf{a}^\top \cdot \mathbf{x} \sim b$$

■ Boolean connectives:

$$\neg \quad \wedge \quad \vee \quad \rightarrow$$

■ Quantifiers:

$$\exists x : \Phi(x) \quad \forall x : \Phi(x)$$

Linear arithmetic theories: semantics

Domain of variables are reals (\mathbb{R}) or integers (\mathbb{Z}), or subsets thereof. Write \mathbb{D} for arbitrary domain.

Linear arithmetic theories: semantics

Domain of variables are reals (\mathbb{R}) or integers (\mathbb{Z}), or subsets thereof. Write \mathbb{D} for arbitrary domain. Assignments are mappings $\mathcal{A}: X \rightarrow \mathbb{D}$.

Linear arithmetic theories: semantics

Domain of variables are reals (\mathbb{R}) or integers (\mathbb{Z}), or subsets thereof. Write \mathbb{D} for arbitrary domain. Assignments are mappings $\mathcal{A}: X \rightarrow \mathbb{D}$. Semantics:

$$\blacksquare \mathcal{A} \models \sum_{i=1}^n a_i \cdot x_i \sim b \iff \sum_{i=1}^n a_i \cdot \mathcal{A}(x_i) \sim b$$

$$\blacksquare \mathcal{A} \models \neg \Phi \iff \mathcal{A} \not\models \Phi$$

$$\blacksquare \mathcal{A} \models \Phi \wedge \Psi \iff \mathcal{A} \models \Phi \text{ and } \mathcal{A} \models \Psi$$

$$\blacksquare \mathcal{A} \models \Phi \vee \Psi \iff \mathcal{A} \models \Phi \text{ or } \mathcal{A} \models \Psi$$

$$\blacksquare \mathcal{A} \models \Phi \rightarrow \Psi \iff \mathcal{A} \models \neg \Phi \text{ or } \mathcal{A} \models \Psi$$

$$\blacksquare \mathcal{A} \models \exists x : \Phi(x) \iff \text{there is } a \in \mathbb{D} \text{ such that } \mathcal{A} \models \Phi[a/x]$$

$$\blacksquare \mathcal{A} \models \forall x : \Phi(x) \iff \text{for all } a \in \mathbb{D}, \mathcal{A} \models \Phi[a/x]$$

Simplifying formulas (1)

- Can assume negation-free formulas:

$$\neg(a = b) \iff a < b \vee b < a$$

$$\neg(a < b) \iff b \leq a$$

$$\neg(a \leq b) \iff b < a$$

- Equality not needed:

$$a = b \iff a \leq b \wedge b \leq a$$

- Over \mathbb{Z} only one of \leq and $<$ needed:

$$a < b \iff a + 1 \leq b$$

Simplifying formulas (2)

Prenex form:

$Q_1x_1 Q_2x_2 \cdots Q_kx_k : \Phi(x_1, \dots, x_k)$ and Φ is quantifier-free

Simplifying formulas (2)

Prenex form:


$Q_1x_1 Q_2x_2 \cdots Q_kx_k : \Phi(x_1, \dots, x_k)$ and Φ is quantifier-free

Can wlog assume formula to be in prenex form:

- push negations in front of atomic formulas
- apply equivalences from previous slide to remove negation
- ensure no two quantifiers refer to the same variable
- pull quantifiers outwards


Convex geometry

Hyperplanes



Two vectors $v, w \in \mathbb{R}^d$ are **orthogonal** whenever $v^\top \cdot w = 0$.


Hyperplanes



Two vectors $v, w \in \mathbb{R}^d$ are **orthogonal** whenever $v^\top \cdot w = 0$.

What is the set of vectors/points orthogonal to a **non-zero** vector $v \in \mathbb{R}^d$?

Hyperplanes




Two vectors $v, w \in \mathbb{R}^d$ are **orthogonal** whenever $v^\top \cdot w = 0$.

What is the set of vectors/points orthogonal to a **non-zero** vector $v \in \mathbb{R}^d$?

$$H := \left\{ x \in \mathbb{R}^d : v^\top \cdot x = 0 \right\} \leftarrow \text{a hyperplane!}$$

Hyperplanes



Two vectors $v, w \in \mathbb{R}^d$ are **orthogonal** whenever $v^\top \cdot w = 0$.

What is the set of vectors/points orthogonal to a **non-zero** vector $v \in \mathbb{R}^d$?


$$H := \{x \in \mathbb{R}^d : v^\top \cdot x = 0\} \leftarrow \text{a hyperplane!}$$

$$H^+ := \{x \in \mathbb{R}^d : v^\top \cdot x \geq 0\} \text{ and}$$

$$H^- := \{x \in \mathbb{R}^d : v^\top \cdot x \leq 0\} \text{ are its half-spaces}$$

(v is always in H^+)

Hyperplanes




$$H := \left\{ \mathbf{x} \in \mathbb{R}^d : \mathbf{v}^\top \cdot \mathbf{x} = 0 \right\} \leftarrow \text{hyperplane}$$

Given $c \in \mathbb{Z} \setminus \{0\}$,

$$H_c := \left\{ \mathbf{x} \in \mathbb{R}^d : \mathbf{v}^\top \cdot \mathbf{x} = c \right\} \leftarrow \text{affine hyperplane!}$$

If $c > 0$ then $H_c \subseteq H^+$ else $H_c \subseteq H^-$


(Polyhedral) cones



$$\mathbf{C} := \left\{ \mathbf{A} \cdot \boldsymbol{\lambda} : \boldsymbol{\lambda} \in \mathbb{R}^d \text{ and } \boldsymbol{\lambda} \geq \mathbf{0} \right\} \leftarrow \text{a cone!}$$

We denote this set by $\text{cone}(\mathbf{A})$.

(Polyhedral) cones



$$\mathbf{C} := \left\{ \mathbf{A} \cdot \boldsymbol{\lambda} : \boldsymbol{\lambda} \in \mathbb{R}^d \text{ and } \boldsymbol{\lambda} \geq \mathbf{0} \right\} \leftarrow \text{a cone!}$$

We denote this set by $\text{cone}(\mathbf{A})$.

Equivalently,

$$\mathbf{C} := \left\{ \mathbf{x} \in \mathbb{R}^d : \mathbf{B} \cdot \mathbf{x} \geq \mathbf{0} \right\} \leftarrow \text{also a cone!}$$

Farkas' lemma

Lemma (Farkas; 1902)

Let $A \in \mathbb{R}^{m \times d}$ and $\mathbf{b} \in \mathbb{R}^m$. Exactly one of the following two assertions holds:

1. There is $\mathbf{x} \in \mathbb{R}^d$ such that $A \cdot \mathbf{x} = \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
2. There is $\mathbf{v} \in \mathbb{R}^m$ such that $\mathbf{v}^\top \cdot A \geq \mathbf{0}$ and $\mathbf{v}^\top \cdot \mathbf{b} < 0$.

Farkas' lemma

Lemma (Farkas; 1902)

Let $A \in \mathbb{R}^{m \times d}$ and $\mathbf{b} \in \mathbb{R}^m$. Exactly one of the following two assertions holds:


1. There is $\mathbf{x} \in \mathbb{R}^d$ such that $A \cdot \mathbf{x} = \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
2. There is $\mathbf{v} \in \mathbb{R}^m$ such that $\mathbf{v}^\top \cdot A \geq \mathbf{0}$ and $\mathbf{v}^\top \cdot \mathbf{b} < 0$.

Proof.

Either \mathbf{b} belongs to $\text{cone}(A) = \{A \cdot \boldsymbol{\lambda} : \boldsymbol{\lambda} \in \mathbb{R}^d \text{ and } \boldsymbol{\lambda} \geq \mathbf{0}\}$ or there is a hyperplane separating $\text{cone}(A)$ from \mathbf{b} . The former case implies (1), the latter implies (2). ■


Convex sets and convex hulls

A set $S \subseteq \mathbb{R}^d$ is **convex** whenever $[x, y] \subseteq S$ for every $x, y \in S$



Convex sets and convex hulls

A set $S \subseteq \mathbb{R}^d$ is **convex** whenever $[x, y] \subseteq S$ for every $x, y \in S$



For $S \subseteq \mathbb{R}^d$, the **convex hull** $\text{conv}(S)$ of S is:


- the (unique) smallest convex set containing S , or
- the intersection of all convex sets containing S , or
- the set of all **convex combinations** of elements of S :

$$\text{conv}(S) := \left\{ \lambda_1 \mathbf{v}_1 + \dots + \lambda_n \mathbf{v}_n : \mathbf{v}_1, \dots, \mathbf{v}_n \in S, \sum_{i=1}^n \lambda_i = 1 \text{ and } \lambda_1, \dots, \lambda_n \geq 0 \right\}.$$

Convex polyhedra

A set S is a **polyhedron** if:

- $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{b}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$ and vector $\mathbf{b} \in \mathbb{Q}^d$, **or**
- $S = \text{conv}(V) + \text{cone}(W)$ for some finite sets $V, W \subseteq \mathbb{Q}^d$.




Convex polyhedra

A set S is a **polyhedron** if:

- $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{b}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$ and vector $\mathbf{b} \in \mathbb{Q}^d$, **or**
- $S = \text{conv}(V) + \text{cone}(W)$ for some finite sets $V, W \subseteq \mathbb{Q}^d$.

H is a **supporting hyperplane** of S whenever $S \cap H \neq \emptyset$ and $S \subseteq H^+$.



Convex polyhedra


A set S is a **polyhedron** if:

- $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{b}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$ and vector $\mathbf{b} \in \mathbb{Q}^d$, or
- $S = \text{conv}(V) + \text{cone}(W)$ for some finite sets $V, W \subseteq \mathbb{Q}^d$.

H is a **supporting hyperplane** of S whenever $S \cap H \neq \emptyset$ and $S \subseteq H^+$.

A set F is a **face** of S if

- either $F = S$, or
- $F = S \cap H$ for some supporting hyperplane H of S .



The geometry of a system of inequalities over the **reals**

Theorem (Minkowski-Weyl; 1897, 1935)

Consider $S \subseteq \mathbb{R}^d$. The two following statements are equivalent:

- (H) $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{b}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$ and vector $\mathbf{b} \in \mathbb{Q}^n$
- (V) $S = \text{conv}(V) + \text{cone}(W)$ for some finite sets $V, W \subseteq \mathbb{Q}^d$.


The geometry of a system of inequalities over the **reals**

Theorem (Minkowski-Weyl; 1897, 1935)

Consider $S \subseteq \mathbb{R}^d$. The two following statements are equivalent:

- (H) $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{b}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$ and vector $\mathbf{b} \in \mathbb{Q}^n$
- (V) $S = \text{conv}(V) + \text{cone}(W)$ for some finite sets $V, W \subseteq \mathbb{Q}^d$.

Proof by authority.

“This classical result is an outstanding example of a fact which is completely obvious to geometric intuition, but which wields important algebraic content and is not trivial to prove.” (R. T. Rockafellar) 



The two definitions of convex cones are equivalent

Minkowski-Weyl theorem for cones

Consider $S \subseteq \mathbb{R}^d$. The two following statements are equivalent:

(H) $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{0}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$

(V) $S = \text{cone}(W)$ for some finite sets $W \subseteq \mathbb{Q}^d$.




The two definitions of convex cones are equivalent

Minkowski-Weyl theorem for cones


Consider $S \subseteq \mathbb{R}^d$. The two following statements are equivalent:

(H) $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{0}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$

(V) $S = \text{cone}(W)$ for some finite sets $W \subseteq \mathbb{Q}^d$.



Smart calls to
Farkas' Lemma




The two definitions of convex cones are equivalent

Minkowski-Weyl theorem for cones

Consider $S \subseteq \mathbb{R}^d$. The two following statements are equivalent:

(H) $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{0}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$

(V) $S = \text{cone}(W)$ for some finite sets $W \subseteq \mathbb{Q}^d$.



Minkowski-Weyl theorem: reducing polyhedra to cones

Minkowski-Weyl theorem for cones

Consider $S \subseteq \mathbb{R}^d$. The two following statements are equivalent:

(H) $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{0}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$

(V) $S = \text{cone}(W)$ for some finite sets $W \subseteq \mathbb{Q}^d$.

Minkowski-Weyl theorem for polyhedra

Consider $S \subseteq \mathbb{R}^d$. The two following statements are equivalent:

(H) $S = \{\mathbf{x} \in \mathbb{R}^d : A \cdot \mathbf{x} \geq \mathbf{b}\}$ for some matrix $A \in \mathbb{Q}^{n \times d}$ and vector $\mathbf{b} \in \mathbb{Q}^d$



(V) $S = \text{conv}(V) + \text{cone}(W)$ for some finite sets $V, W \subseteq \mathbb{Q}^d$.

Minkowski-Weyl theorem: reducing polyhedra to cones

Mi Every polyhedron in dimension d is the slice of a cone in dimension $d+1$


Con
(H)
(V)

Mi
Con
(H)
(V)




Summary of today's lecture

- Applications of arithmetic theories
- Syntax and semantics of linear arithmetic theories
- Convex geometric objects
- Farkas' lemma
- Minkowski—Weyl theorem



Summary of today's lecture

- Applications of arithmetic theories
- Syntax and semantics of linear arithmetic theories
- Convex geometric objects
- Farkas' lemma
- Minkowski—Weyl theorem



Agenda for the rest of the week

Tomorrow Linear programming and Integer linear programming

Wednesday Quantifier elimination procedures

Thursday Automata-based procedures

Friday Geometric procedures