

# Statistics and Probability

Alex Corner

Sheffield Hallam University

# Statistics

## Some terminology

## Some terminology

- ▶ **Population:** *All* relevant data.

## Some terminology

- ▶ **Population:** *All* relevant data.
- ▶ **Sample:** *Some* of the relevant data, hopefully enough to be *representative* of the population.

## Some terminology

- ▶ **Population:** *All* relevant data.
- ▶ **Sample:** *Some* of the relevant data, hopefully enough to be *representative* of the population.
  - ▶ This should be a **random sample** with no **bias** towards a group or individual in the population.

## Some terminology

- ▶ **Population:** *All* relevant data.
- ▶ **Sample:** *Some* of the relevant data, hopefully enough to be *representative* of the population.
  - ▶ This should be a **random sample** with no **bias** towards a group or individual in the population.
  - ▶ This can be difficult to achieve and careful thought needs to be given as to how data is collected.

## Some terminology

- ▶ **Population:** *All* relevant data.
- ▶ **Sample:** *Some* of the relevant data, hopefully enough to be *representative* of the population.
  - ▶ This should be a **random sample** with no **bias** towards a group or individual in the population.
  - ▶ This can be difficult to achieve and careful thought needs to be given as to how data is collected.
- ▶ Sample/population **data** can be:
  - ▶ **Discrete:** E.g., number of attacks on a port in a given hour of the day.



## Some terminology

- ▶ **Population:** *All* relevant data.
- ▶ **Sample:** *Some* of the relevant data, hopefully enough to be *representative* of the population.
  - ▶ This should be a **random sample** with no **bias** towards a group or individual in the population.
  - ▶ This can be difficult to achieve and careful thought needs to be given as to how data is collected.
- ▶ Sample/population **data** can be:
  - ▶ **Discrete:** E.g., number of attacks on a port in a given hour of the day. Usually integer-valued, e.g., 0, 1, 2, etc.

## Some terminology

- ▶ **Population:** *All* relevant data.
- ▶ **Sample:** *Some* of the relevant data, hopefully enough to be *representative* of the population.
  - ▶ This should be a **random sample** with no **bias** towards a group or individual in the population.
  - ▶ This can be difficult to achieve and careful thought needs to be given as to how data is collected.
- ▶ Sample/population **data** can be:
  - ▶ **Discrete:** E.g., number of attacks on a port in a given hour of the day. Usually integer-valued, e.g., 0, 1, 2, etc.
  - ▶ **Continuous:** E.g., amount of time elapsed since previous attack.

## Some terminology

- ▶ **Population:** *All* relevant data.
- ▶ **Sample:** *Some* of the relevant data, hopefully enough to be *representative* of the population.
  - ▶ This should be a **random sample** with no **bias** towards a group or individual in the population.
  - ▶ This can be difficult to achieve and careful thought needs to be given as to how data is collected.
- ▶ Sample/population **data** can be:
  - ▶ **Discrete:** E.g., number of attacks on a port in a given hour of the day. Usually integer-valued, e.g., 0, 1, 2, etc.
  - ▶ **Continuous:** E.g., amount of time elapsed since previous attack. Often classified into groups, e.g., 0 – 5 minutes, 5 – 10 minutes, 10 – 15 minutes, etc.

## Grouping Data

- ▶ Suppose that we have data from a random sample of 100 people, giving their age at their last birthday.

## Grouping Data

- ▶ Suppose that we have data from a random sample of 100 people, giving their age at their last birthday.

66	70	10	54	62	13	11	15	69	26	49	11	3
67	10	54	42	32	56	39	60	79	33	12	47	24
19	47	63	32	7	70	55	46	11	20	15	39	37
28	72	46	64	61	51	56	53	61	11	80	53	28
76	6	5	39	58	29	52	54	47	60	62	51	72
41	57	32	12	33	17	40	20	10	27	47	71	68
44	7	23	17	81	23	12	33	16	46	71	48	58
79	80	43	31	72	68	36	41	11				

- ▶ It's a little hard to get a feel for the data, so we could count the frequency in each bin, order the data numerically, or calculate various **statistics**.

## Grouping Data

- ▶ Suppose that we have data from a random sample of 100 people, giving their age at their last birthday.

66	70	10	54	62	13	11	15	69	26	49	11	3
67	10	54	42	32	56	39	60	79	33	12	47	24
19	47	63	32	7	70	55	46	11	20	15	39	37
28	72	46	64	61	51	56	53	61	11	80	53	28
76	6	5	39	58	29	52	54	47	60	62	51	72
41	57	32	12	33	17	40	20	10	27	47	71	68
44	7	23	17	81	23	12	33	16	46	71	48	58
79	80	43	31	72	68	36	41	11				


- ▶ It's a little hard to get a feel for the data, so we could count the frequency in each bin, order the data numerically, or calculate various **statistics**.
- ▶ In doing so, we lose some of the information but might get a clearer overview.

## Grouping Data

- Here we have placed the data into bins of **class width 5**.

Group Number	Age Range	Tally	Number in Group (Frequency)
1	$0 \leq x < 5$		1
2	$5 \leq x < 10$		4
3	$10 \leq x < 15$		12
4	$15 \leq x < 20$		6
5	$20 \leq x < 25$		5
6	$25 \leq x < 30$		5
7	$30 \leq x < 35$		7
8	$35 \leq x < 40$		5
9	$40 \leq x < 45$		6
10	$45 \leq x < 50$		9
11	$50 \leq x < 55$		8
12	$55 \leq x < 60$		6
13	$60 \leq x < 65$		8
14	$65 \leq x < 70$		5
15	$70 \leq x < 75$		7
16	$75 \leq x < 80$		3
17	$80 \leq x < 85$		3
18	$85 \leq x < 90$		0
19	$90 \leq x < 95$		0
20	$95 \leq x < 100$		0
21	$100 \leq x < 105$		0

## Charts: Bar Chart





# Charts: Histogram



## Charts: Histogram



# Measures of Central Tendency

# Measures of Central Tendency

- ▶ **Arithmetic Mean:** Add up all of the values and divide by how many there are.

## Measures of Central Tendency

- ▶ **Arithmetic Mean:** Add up all of the values and divide by how many there are.
- ▶ **Median:** Order the data and look at the middle value(s).

# Measures of Central Tendency

- ▶ **Arithmetic Mean:** Add up all of the values and divide by how many there are.
- ▶ **Median:** Order the data and look at the middle value(s).
- ▶ **Mode:** Tally the data and select the one with the highest frequency.

## (Arithmetic) Mean

## (Arithmetic) Mean

- In a sample of five values (6, 9, 2, 4, 3) the mean value is:

$$\frac{6 + 9 + 2 + 4 + 3}{5} = \frac{24}{5} = 4.8.$$



## (Arithmetic) Mean

- ▶ In a sample of five values (6, 9, 2, 4, 3) the mean value is:

$$\frac{6 + 9 + 2 + 4 + 3}{5} = \frac{24}{5} = 4.8.$$

- ▶ The one hundred values from the previous histogram example add up to 4165 and so the mean value is  $\frac{4165}{100} = 41.65$ .

## (Arithmetic) Mean

- ▶ In a sample of five values (6, 9, 2, 4, 3) the mean value is:

$$\frac{6 + 9 + 2 + 4 + 3}{5} = \frac{24}{5} = 4.8.$$

- ▶ The one hundred values from the previous histogram example add up to 4165 and so the mean value is  $\frac{4165}{100} = 41.65$ .
- ▶ For  $n$  data values  $x_1, x_2, \dots, x_n$ , the formula for the mean is:

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n}.$$

## (Arithmetic) Mean

- ▶ In a sample of five values (6, 9, 2, 4, 3) the mean value is:

$$\frac{6 + 9 + 2 + 4 + 3}{5} = \frac{24}{5} = 4.8.$$

- ▶ The one hundred values from the previous histogram example add up to 4165 and so the mean value is  $\frac{4165}{100} = 41.65$ .
- ▶ For  $n$  data values  $x_1, x_2, \dots, x_n$ , the formula for the mean is:

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n}.$$

## Median

## Median

- ▶ Here we arrange the data values in ascending (or descending) order. Then the **median** is:
  1. the middle item if there is an odd number of data items;
  2. the average of the middle two items if there is an even number of data items.

## Median

- ▶ Here we arrange the data values in ascending (or descending) order. Then the **median** is:
  1. the middle item if there is an odd number of data items;
  2. the average of the middle two items if there is an even number of data items.
- ▶ E.g., if the ordered data is 2, 3, 4, 6, 9, then the middle item is 4 and so the median is 4.

## Median

- ▶ Here we arrange the data values in ascending (or descending) order. Then the **median** is:
  1. the middle item if there is an odd number of data items;
  2. the average of the middle two items if there is an even number of data items.
- ▶ E.g., if the ordered data is 2, 3, 4, 6, 9, then the middle item is 4 and so the median is 4.
- ▶ From our histogram example, the sample of one hundred data values, the ordered data is:

3	5	6	7	7	10	10	10	11	11	11	11	11
12	12	12	13	15	15	16	17	17	19	20	20	23
23	24	26	27	28	28	29	31	32	32	32	33	33
33	36	37	39	39	39	40	41	41	42	43	44	46
46	46	47	47	47	47	48	49	51	51	52	53	53
54	54	54	55	56	56	57	58	58	60	60	61	61
62	62	63	64	66	67	68	68	69	70	70	71	71
72	72	72	76	79	79	80	80	81				

## Median

- ▶ Here we arrange the data values in ascending (or descending) order. Then the **median** is:
  1. the middle item if there is an odd number of data items;
  2. the average of the middle two items if there is an even number of data items.
- ▶ E.g., if the ordered data is 2, 3, 4, 6, 9, then the middle item is 4 and so the median is 4.
- ▶ From our histogram example, the sample of one hundred data values, the ordered data is:

3	5	6	7	7	10	10	10	11	11	11	11	11
12	12	12	13	15	15	16	17	17	19	20	20	23
23	24	26	27	28	28	29	31	32	32	32	33	33
33	36	37	39	39	39	40	41	41	42	43	44	46
46	46	47	47	47	47	48	49	51	51	52	53	53
54	54	54	55	56	56	57	58	58	60	60	61	61
62	62	63	64	66	67	68	68	69	70	70	71	71
72	72	72	76	79	79	80	80	81				

- ▶ The middle two values are 43 and 44, hence the median is 43.5.



## Mode

## Mode


- ▶ The **mode** is the data value which occurs most in the sample.

## Mode

- ▶ The **mode** is the data value which occurs most in the sample.
- ▶ For grouped data we can talk of a **modal group** or **modal class**, sometimes specifying the mid-point in order to give a single value.

# Mode

- ▶ The **mode** is the data value which occurs most in the sample.
- ▶ For grouped data we can talk of a **modal group** or **modal class**, sometimes specifying the mid-point in order to give a single value.
- ▶ For the one hundred data values in the histogram example the mode is clearly indicated on the bar chart as the tallest bar: 11. We can also tally values to find it.



# Probability

## Definition of Probability

## Definition of Probability

- ▶  $A$  is a statement which is either true or false.

## Definition of Probability

- ▶  $A$  is a statement which is either true or false.
- ▶  $N$  is the number of trials, or observations.



## Definition of Probability

- ▶  $A$  is a statement which is either true or false.
- ▶  $N$  is the number of trials, or observations.
- ▶  $S$  is the number of successes (so  $N - S$  is the number of failures).

## Definition of Probability

- ▶  $A$  is a statement which is either true or false.
- ▶  $N$  is the number of trials, or observations.
- ▶  $S$  is the number of successes (so  $N - S$  is the number of failures).
- ▶ For a single trial: Probability that  $A$  is true, or the **probability of success**, is:

$$P(A) = \frac{S}{N}.$$

## Definition of Probability

- ▶  $A$  is a statement which is either true or false.
- ▶  $N$  is the number of trials, or observations.
- ▶  $S$  is the number of successes (so  $N - S$  is the number of failures).
- ▶ For a single trial: Probability that  $A$  is true, or the **probability of success**, is:

$$P(A) = \frac{S}{N}.$$

- ▶ Since  $0 \leq S \leq N$ , then  $0 \leq P(A) \leq 1$ .

## Definition of Probability

- ▶  $A$  is a statement which is either true or false.
- ▶  $N$  is the number of trials, or observations.
- ▶  $S$  is the number of successes (so  $N - S$  is the number of failures).
- ▶ For a single trial: Probability that  $A$  is true, or the **probability of success**, is:

$$P(A) = \frac{S}{N}.$$

- ▶ Since  $0 \leq S \leq N$ , then  $0 \leq P(A) \leq 1$ .
- ▶ The *probability of failure* is:

$$P(\bar{A}) = \frac{N - S}{N} = 1 - \frac{S}{N} = 1 - P(A).$$

## Probability: Example 1

Tossing a coin:

## Probability: Example 1

Tossing a coin:

- ▶ Toss a coin 4 times: we *expect* 2 heads.

## Probability: Example 1

Tossing a coin:

- ▶ Toss a coin 4 times: we *expect* 2 heads.
- ▶ Toss a coin 100 times: we *expect* 50 heads.

## Probability: Example 1

Tossing a coin:

- ▶ Toss a coin 4 times: we *expect* 2 heads.
- ▶ Toss a coin 100 times: we *expect* 50 heads.
- ▶ Toss a coin once: what do we *expect*?



## Probability: Example 1

Tossing a coin:

- ▶ Toss a coin 4 times: we *expect* 2 heads.
- ▶ Toss a coin 100 times: we *expect* 50 heads.
- ▶ Toss a coin once: what do we *expect*?

$$P(\text{one head in one trial}) = \frac{1}{2}.$$

## Probability: Example 1

Tossing a coin:

- ▶ Toss a coin 4 times: we *expect* 2 heads.
- ▶ Toss a coin 100 times: we *expect* 50 heads.
- ▶ Toss a coin once: what do we *expect*?

$$P(\text{one head in one trial}) = \frac{1}{2}.$$

- ▶  $A = \text{'a head shows when a coin is tossed'}$ , so  $P(A) = \frac{1}{2}$ .

# Probability: Example

Group Number	Age Range	Tally	Number in Group (Frequency)
1	$0 \leq x < 5$		1
2	$5 \leq x < 10$		4
3	$10 \leq x < 15$		12
4	$15 \leq x < 20$		6
5	$20 \leq x < 25$		5
6	$25 \leq x < 30$		5
7	$30 \leq x < 35$		7
8	$35 \leq x < 40$		5
9	$40 \leq x < 45$		6
10	$45 \leq x < 50$		9
11	$50 \leq x < 55$		8
12	$55 \leq x < 60$		6
13	$60 \leq x < 65$		8
14	$65 \leq x < 70$		5
15	$70 \leq x < 75$		7
16	$75 \leq x < 80$		3
17	$80 \leq x < 85$		3
18	$85 \leq x < 90$		0
19	$90 \leq x < 95$		0
20	$95 \leq x < 100$		0
21	$100 \leq x < 105$		0

Age at last birthday:

## Probability: Example

Group Number	Age Range	Tally	Number in Group (Frequency)
1	$0 \leq x < 5$		1
2	$5 \leq x < 10$		4
3	$10 \leq x < 15$		12
4	$15 \leq x < 20$		6
5	$20 \leq x < 25$		5
6	$25 \leq x < 30$		5
7	$30 \leq x < 35$		7
8	$35 \leq x < 40$		5
9	$40 \leq x < 45$		6
10	$45 \leq x < 50$		9
11	$50 \leq x < 55$		8
12	$55 \leq x < 60$		6
13	$60 \leq x < 65$		8
14	$65 \leq x < 70$		5
15	$70 \leq x < 75$		7
16	$75 \leq x < 80$		3
17	$80 \leq x < 85$		3
18	$85 \leq x < 90$		0
19	$90 \leq x < 95$		0
20	$95 \leq x < 100$		0
21	$100 \leq x < 105$		0

Age at last birthday:

- Probability that one of the people selected at random is in the 15-19 age group.

## Probability: Example

Group Number	Age Range	Tally	Number in Group (Frequency)
1	$0 \leq x < 5$		1
2	$5 \leq x < 10$		4
3	$10 \leq x < 15$		12
4	$15 \leq x < 20$		6
5	$20 \leq x < 25$		5
6	$25 \leq x < 30$		5
7	$30 \leq x < 35$		7
8	$35 \leq x < 40$		5
9	$40 \leq x < 45$		6
10	$45 \leq x < 50$		9
11	$50 \leq x < 55$		8
12	$55 \leq x < 60$		6
13	$60 \leq x < 65$		8
14	$65 \leq x < 70$		5
15	$70 \leq x < 75$		7
16	$75 \leq x < 80$		3
17	$80 \leq x < 85$		3
18	$85 \leq x < 90$		0
19	$90 \leq x < 95$		0
20	$95 \leq x < 100$		0
21	$100 \leq x < 105$		0

Age at last birthday:

- ▶ Probability that one of the people selected at random is in the 15-19 age group.
- ▶ The total number of outcomes is 100 and the number of 'successes' is 6.

## Probability: Example

Group Number	Age Range	Tally	Number in Group (Frequency)
1	$0 \leq x < 5$		1
2	$5 \leq x < 10$		4
3	$10 \leq x < 15$		12
4	$15 \leq x < 20$		6
5	$20 \leq x < 25$		5
6	$25 \leq x < 30$		5
7	$30 \leq x < 35$		7
8	$35 \leq x < 40$		5
9	$40 \leq x < 45$		6
10	$45 \leq x < 50$		9
11	$50 \leq x < 55$		8
12	$55 \leq x < 60$		6
13	$60 \leq x < 65$		8
14	$65 \leq x < 70$		5
15	$70 \leq x < 75$		7
16	$75 \leq x < 80$		3
17	$80 \leq x < 85$		3
18	$85 \leq x < 90$		0
19	$90 \leq x < 95$		0
20	$95 \leq x < 100$		0
21	$100 \leq x < 105$		0

Age at last birthday:

- ▶ Probability that one of the people selected at random is in the 15-19 age group.
- ▶ The total number of outcomes is 100 and the number of 'successes' is 6.
- ▶ The probability of success is:  $P(\text{Success}) = \frac{6}{100}$ .

## Probability: Example

Group Number	Age Range	Tally	Number in Group (Frequency)
1	$0 \leq x < 5$		1
2	$5 \leq x < 10$		4
3	$10 \leq x < 15$		12
4	$15 \leq x < 20$		6
5	$20 \leq x < 25$		5
6	$25 \leq x < 30$		5
7	$30 \leq x < 35$		7
8	$35 \leq x < 40$		5
9	$40 \leq x < 45$		6
10	$45 \leq x < 50$		9
11	$50 \leq x < 55$		8
12	$55 \leq x < 60$		6
13	$60 \leq x < 65$		8
14	$65 \leq x < 70$		5
15	$70 \leq x < 75$		7
16	$75 \leq x < 80$		3
17	$80 \leq x < 85$		3
18	$85 \leq x < 90$		0
19	$90 \leq x < 95$		0
20	$95 \leq x < 100$		0
21	$100 \leq x < 105$		0

Age at last birthday:

- ▶ Probability that one of the people selected at random is in the 15-19 age group.
- ▶ The total number of outcomes is 100 and the number of 'successes' is 6.
- ▶ The probability of success is:  $P(\text{Success}) = \frac{6}{100}$ .
- ▶ The probability of failure is:  $P(\text{Failure}) = \frac{94}{100}$ .

## Probability: Example 3

Rolling dice:



## Probability: Example 3

Rolling dice:

- ▶ Two dice are rolled and the sum of the spots is calculated.

## Probability: Example 3

Rolling dice:

- ▶ Two dice are rolled and the sum of the spots is calculated.
- ▶  $A =$  'The sum is 7 when two dice are rolled.'

## Probability: Example 3

Rolling dice:

- ▶ Two dice are rolled and the sum of the spots is calculated.
- ▶  $A =$  'The sum is 7 when two dice are rolled.'

(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	<b>(1, 6)</b>
(2, 1)	(2, 2)	(2, 3)	(2, 4)	<b>(2, 5)</b>	(2, 6)
(3, 1)	(3, 2)	(3, 3)	<b>(3, 4)</b>	(3, 5)	(3, 6)
(4, 1)	(4, 2)	<b>(4, 3)</b>	(4, 4)	(4, 5)	(4, 6)
(5, 1)	<b>(5, 2)</b>	(5, 3)	(5, 4)	(5, 5)	(5, 6)
<b>(6, 1)</b>	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

## Probability: Example 3

Rolling dice:

- ▶ Two dice are rolled and the sum of the spots is calculated.
- ▶  $A =$  'The sum is 7 when two dice are rolled.'

(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	<b>(1, 6)</b>
(2, 1)	(2, 2)	(2, 3)	(2, 4)	<b>(2, 5)</b>	(2, 6)
(3, 1)	(3, 2)	(3, 3)	<b>(3, 4)</b>	(3, 5)	(3, 6)
(4, 1)	(4, 2)	<b>(4, 3)</b>	(4, 4)	(4, 5)	(4, 6)
(5, 1)	<b>(5, 2)</b>	(5, 3)	(5, 4)	(5, 5)	(5, 6)
<b>(6, 1)</b>	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

- ▶ There are 36 possible outcomes when rolling two dice. But only six of these outcomes make  $A$  true.

## Probability: Example 3

Rolling dice:

- ▶ Two dice are rolled and the sum of the spots is calculated.
- ▶  $A =$  'The sum is 7 when two dice are rolled.'

(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	<b>(1, 6)</b>
(2, 1)	(2, 2)	(2, 3)	(2, 4)	<b>(2, 5)</b>	(2, 6)
(3, 1)	(3, 2)	(3, 3)	<b>(3, 4)</b>	(3, 5)	(3, 6)
(4, 1)	(4, 2)	<b>(4, 3)</b>	(4, 4)	(4, 5)	(4, 6)
(5, 1)	<b>(5, 2)</b>	(5, 3)	(5, 4)	(5, 5)	(5, 6)
<b>(6, 1)</b>	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

- ▶ There are 36 possible outcomes when rolling two dice. But only six of these outcomes make  $A$  true.
- ▶ Number of trials  $N = 36$ , number of successes  $S = 6$ .

$$P(A) = \frac{6}{36} = \frac{1}{6}.$$



## And, Or: Intersection, Union

- ▶ Last semester you used  $A \cdot B$  to mean  $A$  AND  $B$ . We will sometimes use  $A \wedge B$  instead.

## And, Or: Intersection, Union

- ▶ Last semester you used  $A \cdot B$  to mean  $A$  AND  $B$ . We will sometimes use  $A \wedge B$  instead.
- ▶ Its analogue in set theory is **intersection**:  $A \cap B$ .



## And, Or: Intersection, Union

- ▶ Last semester you used  $A \cdot B$  to mean  $A$  AND  $B$ . We will sometimes use  $A \wedge B$  instead.
- ▶ Its analogue in set theory is **intersection**:  $A \cap B$ .
- ▶ Last semester you used  $A + B$  to mean  $A$  OR  $B$ . We will sometimes use  $A \vee B$  instead.

## And, Or: Intersection, Union

- ▶ Last semester you used  $A \cdot B$  to mean  $A$  AND  $B$ . We will sometimes use  $A \wedge B$  instead.
- ▶ Its analogue in set theory is **intersection**:  $A \cap B$ .
- ▶ Last semester you used  $A + B$  to mean  $A$  OR  $B$ . We will sometimes use  $A \vee B$  instead.
- ▶ Its analogue in set theory is **union**:  $A \cup B$ .

## Law of multiplication

## Law of multiplication


- ▶ Consider two events  $A$  and  $B$ .

## Law of multiplication

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.

## Law of multiplication

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$



- ▶ The intersection of the sets is  $P(A \cap B)$ : this is the probability  $A$  and  $B$  are both true.

## Law of multiplication


- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$



- ▶ The intersection of the sets is  $P(A \cap B)$ : this is the probability  $A$  and  $B$  are both true.
- ▶ If  $A$  and  $B$  are mutually exclusive (i.e., can't both be true at the same time), then  $P(A \cap B) = 0$ .

## Law of multiplication

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$



- ▶ The intersection of the sets is  $P(A \wedge B)$ : this is the probability  $A$  and  $B$  are both true.
- ▶ If  $A$  and  $B$  are mutually exclusive (i.e., can't both be true at the same time), then  $P(A \wedge B) = 0$ .
- ▶ If  $A$  and  $B$  are **independent** (i.e., the outcome of one does not affect the other), then  $P(A \wedge B) = P(A) \times P(B)$ .



## Probability: Example 4

## Probability: Example 4

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.

## Probability: Example 4

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then replaced in the pack.

## Probability: Example 4

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then replaced in the pack.
- ▶ Then a second card is drawn from the pack.

## Probability: Example 4

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then replaced in the pack.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.

## Probability: Example 4

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then replaced in the pack.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.
- ▶ Let  $A =$  'The first card is red' and  $B =$  'The second card is red'.

## Probability: Example 4

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then replaced in the pack.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.
- ▶ Let  $A =$  'The first card is red' and  $B =$  'The second card is red'.
- ▶  $P(A) = \frac{26}{52} = \frac{1}{2}$  and  $P(B) = \frac{26}{52} = \frac{1}{2}$ . The two events are independent, so:

$$P(A \wedge B) = P(A) \times P(B) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}.$$

# Law of Addition




## Law of Addition

- ▶ Consider two events  $A$  and  $B$ .


## Law of Addition

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$



## Law of Addition

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$



- ▶ The union of the sets is  $P(A \cup B)$ : this is the probability that  $A$  or  $B$  are true.

## Law of Addition

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$



- ▶ The union of the sets is  $P(A \cup B)$ : this is the probability that  $A$  or  $B$  are true.
- ▶ If  $A$  and  $B$  are mutually exclusive, then  $P(A \cup B) = P(A) + P(B)$ .

## Law of Addition

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$




- ▶ The union of the sets is  $P(A \vee B)$ : this is the probability that  $A$  or  $B$  are true.
- ▶ If  $A$  and  $B$  are mutually exclusive, then  $P(A \vee B) = P(A) + P(B)$ .
- ▶ In general,

$$P(A \vee B) = P(A) + P(B)$$

## Law of Addition

- ▶ Consider two events  $A$  and  $B$ .
- ▶  $P(A)$  and  $P(B)$  are denoted by the sets in the Venn diagram.  $U$



- ▶ The union of the sets is  $P(A \vee B)$ : this is the probability that  $A$  or  $B$  are true.
- ▶ If  $A$  and  $B$  are mutually exclusive, then  $P(A \vee B) = P(A) + P(B)$ .
- ▶ In general,

$$P(A \vee B) = P(A) + P(B) - P(A \wedge B).$$

## Probability: Example 4 continued

## Probability: Example 4 continued

- ▶ Same setup as our previous example, but now we want the probability that at least one card is red. ( $P(A) = P(B) = \frac{1}{2}$ )



## Probability: Example 4 continued

- ▶ Same setup as our previous example, but now we want the probability that at least one card is red. ( $P(A) = P(B) = \frac{1}{2}$ )
- ▶ We want to find  $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$ . Since  $A$  and  $B$  are independent, then  $P(A \wedge B) = P(A) \times P(B)$ .


## Probability: Example 4 continued

- ▶ Same setup as our previous example, but now we want the probability that at least one card is red. ( $P(A) = P(B) = \frac{1}{2}$ )
- ▶ We want to find  $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$ . Since  $A$  and  $B$  are independent, then  $P(A \wedge B) = P(A) \times P(B)$ . So:

$$P(A) + P(B) - P(A) \times P(B) = \frac{1}{2} + \frac{1}{2} - \frac{1}{2} \times \frac{1}{2} = \frac{3}{4}.$$

## Example: Network 1

- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in series as shown below.



## Example: Network 1

- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in series as shown below.



- ▶ The link functions only if both ( $a$ ) and ( $b$ ) both function.

## Example: Network 1

- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in series as shown below.



- ▶ The link functions only if both ( $a$ ) and ( $b$ ) both function.
- ▶ The two sections are independent. If one fails, the other is not affected.

## Example: Network 1

- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in series as shown below.



- ▶ The link functions only if both ( $a$ ) and ( $b$ ) both function.
- ▶ The two sections are independent. If one fails, the other is not affected.
- ▶ The probability that ( $a$ ) functions is  $P(A) = 0.8$ , while for ( $b$ ) the probability is  $P(B) = 0.9$ .

## Example: Network 1

- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in series as shown below.




- ▶ The link functions only if both ( $a$ ) and ( $b$ ) both function.
- ▶ The two sections are independent. If one fails, the other is not affected.
- ▶ The probability that ( $a$ ) functions is  $P(A) = 0.8$ , while for ( $b$ ) the probability is  $P(B) = 0.9$ .
- ▶ The probability that the network functions (both sections function) is:

$$P(A \wedge B) = P(A) \times P(B) = 0.8 \times 0.9 = 0.72.$$

## Example: Network 2


- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in parallel as shown below.





## Example: Network 2


- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in parallel as shown below.



- ▶ The link functions if either of ( $a$ ) or ( $b$ ) function (or both).

## Example: Network 2

- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in parallel as shown below.



- ▶ The link functions if either of ( $a$ ) or ( $b$ ) function (or both).
- ▶ The two sections are independent. If one fails, the other is not affected.

## Example: Network 2


- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in parallel as shown below.



- ▶ The link functions if either of ( $a$ ) or ( $b$ ) function (or both).
- ▶ The two sections are independent. If one fails, the other is not affected.
- ▶ The probability that ( $a$ ) functions is  $P(A) = 0.8$ , while for ( $b$ ) the probability is  $P(B) = 0.9$ .

## Example: Network 2

- ▶ A short network link consists of two sections ( $a$ ) and ( $b$ ), in parallel as shown below.



- ▶ The link functions if either of ( $a$ ) or ( $b$ ) function (or both).
- ▶ The two sections are independent. If one fails, the other is not affected.
- ▶ The probability that ( $a$ ) functions is  $P(A) = 0.8$ , while for ( $b$ ) the probability is  $P(B) = 0.9$ .
- ▶ The probability that the network functions is:

$$P(A \vee B) = P(A) + P(B) - P(A \wedge B) = 0.8 + 0.9 - 0.72 = 0.98.$$

# Conditional Probability

- ▶ When two events are *not independent*, then the process is not as simple.

# Conditional Probability

- ▶ When two events are *not independent*, then the process is not as simple.
- ▶ Consider a single trial consisting of two events  $A$  and  $B$ .

# Conditional Probability

- ▶ When two events are *not independent*, then the process is not as simple.
- ▶ Consider a single trial consisting of two events  $A$  and  $B$ .
- ▶ The **conditional probability**  $P(B|A)$  means 'the probability that  $B$  will be true *given that*  $A$  is already true'.

## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.



## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then *kept*.

## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then *kept*.
- ▶ Then a second card is drawn from the pack.

## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then *kept*.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.

## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then *kept*.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.
- ▶ Let  $A =$  'The first card is red' and  $B =$  'The second card is red'.

## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then *kept*.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.
- ▶ Let  $A =$  'The first card is red' and  $B =$  'The second card is red'.
- ▶  $P(A) = \frac{26}{52} = \frac{1}{2}$  as before but  $P(B)$  is a conditional probability because it depends on the result of event  $A$ .

## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then *kept*.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.
- ▶ Let  $A =$  'The first card is red' and  $B =$  'The second card is red'.
- ▶  $P(A) = \frac{26}{52} = \frac{1}{2}$  as before but  $P(B)$  is a conditional probability because it depends on the result of event  $A$ .
- ▶ If the first card was red, then there are now only 25 cards left in the remaining 51 cards, so:

$$P(B|A) = \frac{25}{51}.$$

## Probability: Example 5

- ▶ A pack of 52 playing cards has 26 red cards and 26 black cards.
- ▶ One card is drawn from the full pack at random, then *kept*.
- ▶ Then a second card is drawn from the pack.
- ▶ Calculate the probability that both cards are red.
- ▶ Let  $A =$  'The first card is red' and  $B =$  'The second card is red'.
- ▶  $P(A) = \frac{26}{52} = \frac{1}{2}$  as before but  $P(B)$  is a conditional probability because it depends on the result of event  $A$ .
- ▶ If the first card was red, then there are now only 25 cards left in the remaining 51 cards, so:

$$P(B|A) = \frac{25}{51}.$$

- ▶ This means the probability that the second card is red given the first card was red is:

$$P(A \wedge B) = P(B|A) \times P(A) = \frac{25}{51} \times \frac{1}{2} = \frac{25}{102}.$$

## Conditional Probability

- ▶ The **conditional probability** can be expressed as

$$P(B|A) = \frac{P(A \wedge B)}{P(A)}.$$



## Conditional Probability

- ▶ The **conditional probability** can be expressed as

$$P(B|A) = \frac{P(A \wedge B)}{P(A)}.$$

- ▶ This can be rearranged as

$$P(A \wedge B) = P(B|A) \times P(A).$$

## Conditional Probability

- ▶ The **conditional probability** can be expressed as

$$P(B|A) = \frac{P(A \wedge B)}{P(A)}.$$

- ▶ This can be rearranged as

$$P(A \wedge B) = P(B|A) \times P(A).$$

- ▶ If  $A$  and  $B$  are independent, then

$$P(B|A) = P(B).$$

## Probability: Example 6

- Calculate the probability that the first card is black and the second card is red.

## Probability: Example 6

- ▶ Calculate the probability that the first card is black and the second card is red.
- ▶ Both cards are red:

$$P(A \wedge B) = P(B|A) \times P(A) = \frac{25}{51} \times \frac{1}{2} = \frac{25}{102}.$$

## Probability: Example 6

- ▶ Calculate the probability that the first card is black and the second card is red.
- ▶ Both cards are red:

$$P(A \wedge B) = P(B|A) \times P(A) = \frac{25}{51} \times \frac{1}{2} = \frac{25}{102}.$$

- ▶ First card is black, second card is red:

$$P(\bar{A} \wedge B) = P(B|\bar{A}) \times P(\bar{A}) = \frac{26}{51} \times \frac{1}{2} = \frac{26}{102}.$$

# Probability Trees

- ▶ For small examples, we can often visualise things in the form of a **probability tree**.

# Probability Trees

- For small examples, we can often visualise things in the form of a **probability tree**.

