

Treewidth of Products of Graphs with High Treewidth

Raj Kaul

Abstract

Treewidth is the standard measure for how “tree-like” a graph is. This paper studies how the treewidth of a product graph depends on the treewidth of its factors. Kozawa, Otachi, and Yamazaki [2014] and Hickingbotham and Wood [2025] independently showed that $\text{tw}(G \boxtimes H) \geq (\text{tw}(G) + 1)\text{had}(H) - 1$ for all graphs G and H , where $\text{had}(H)$ is the Hadwiger number of H . We improve this bound to $\text{tw}(G \boxtimes H) \geq (\text{tw}(G) + 1)(\text{tw}(H) + 1) - 1$, thereby solving an open problem of Hickingbotham and Wood. We also prove analogous product inequalities for pathwidth, Cartesian products, and strict bramble number, which is a parameter that is tied to treewidth. As an application of our results, we show that products of expanders have large subgraphs that are expanders.

1 Introduction

Treewidth¹ is a foundational parameter in structural and algorithmic graph theory [4, 9, 17] that measures the structural complexity of a graph. Informally, graphs with small treewidth are more “tree-like”. Grids are the canonical graphs with large treewidth, since the $n \times n$ -grid has treewidth n and, by the Grid-Minor Theorem of Robertson and Seymour [18], every graph with large enough treewidth contains the $n \times n$ -grid as a minor.² Inspired by the fact that grids are Cartesian products of paths, this paper studies the treewidth of various products of general graphs.

Date: July 21, 2026.

School of Mathematics, Monash University, Melbourne, Australia (raj.kaul@monash.edu). Supported by an Australian Government Research Training Program Scholarship.

¹We consider simple, finite, and undirected graphs. A set $A \subseteq V(G)$ is *connected* in G if $G[A]$ is connected. If T is a tree and $x, y \in V(T)$, then xTy denotes the unique path in T whose set of endpoints is $\{x, y\}$. A *tree-decomposition* of a graph G is a pair (T, β) consisting of a tree T and a function $\beta : V(T) \rightarrow 2^{V(G)}$ with the following two properties: (*vertex-property*) for every vertex $v \in V(G)$, the set $\{t \in V(T) : v \in \beta(t)\}$ is non-empty and connected in T ; and (*edge-property*) for every edge $uv \in E(G)$, there exists $t \in V(T)$ such that $\{u, v\} \subseteq \beta(t)$. The vertices of T are called *nodes*, and the sets $\beta(t)$ are called *bags*. If the underlying tree T is a path, then (T, β) may be called a *path-decomposition*. The *width* of (T, β) is $\max_{t \in V(T)} |\beta(t)| - 1$, and the *treewidth* of G , denoted by $\text{tw}(G)$, is the minimum width of a tree-decomposition of G . The pathwidth of G , denoted by $\text{pw}(G)$, is the minimum width of a path-decomposition of G .

²A graph H is a *minor* of a graph G if H can be obtained from G by a sequence of vertex deletions, edge deletions, and edge contractions.

Our first contribution, proven in [Section 2](#), is a lower bound on the treewidth of a strong product. The *strong product* of graphs G and H , denoted by $G \boxtimes H$, is the graph with vertex set $V(G) \times V(H)$, where $(g_1, h_1)(g_2, h_2) \in E(G \boxtimes H)$ if and only if $g_1 = g_2$ and $h_1 h_2 \in E(H)$; or $g_1 g_2 \in E(G)$ and $h_1 = h_2$; or $g_1 g_2 \in E(G)$ and $h_1 h_2 \in E(H)$.

Theorem 1. *For all graphs G and H , $\text{tw}(G \boxtimes H) \geq (\text{tw}(G) + 1)(\text{tw}(H) + 1) - 1$.*

[Theorem 1](#) improves upon the bound $\text{tw}(G \boxtimes H) \geq (\text{tw}(G) + 1)\text{had}(H) - 1$, where $\text{had}(H)$ denotes the Hadwiger number of H ,³ which was shown independently by Kozawa, Otachi, and Yamazaki [12] and Hickingbotham and Wood [11]. Furthermore, [Theorem 1](#) solves an open problem of Hickingbotham and Wood [11], who asked whether there exists a constant $c > 0$ such that $\text{tw}(G \boxtimes H) \geq c \text{tw}(G) \text{tw}(H)$ for all graphs G and H . [Theorem 1](#) says the answer is ‘yes’ with $c = 1$.

Our second contribution, proven in [Section 3](#), is that [Theorem 1](#) holds when treewidth is replaced by pathwidth:

Theorem 2. *For all graphs G and H , $\text{pw}(G \boxtimes H) \geq (\text{pw}(G) + 1)(\text{pw}(H) + 1) - 1$.*

[Theorems 1](#) and [2](#) are optimal since $\text{tw}(K_n) = \text{pw}(K_n) = n - 1$ and $\text{tw}(K_m \boxtimes K_n) = \text{tw}(K_{mn}) = mn - 1 = (\text{tw}(K_m) + 1)(\text{tw}(K_n) + 1) - 1$.

In addition to the strong product, this paper also studies Cartesian products and lexicographic products, which we now introduce. The *Cartesian product* of graphs G and H , denoted by $G \square H$, is the graph with vertex set $V(G) \times V(H)$, where $(g_1, h_1)(g_2, h_2) \in E(G \square H)$ if and only if $g_1 = g_2$ and $h_1 h_2 \in E(H)$; or $g_1 g_2 \in E(G)$ and $h_1 = h_2$. The *lexicographic product* of graphs G with H , denoted by $G \circ H$, is the graph with vertex set $V(G) \times V(H)$, where $(g_1, h_1)(g_2, h_2) \in E(G \circ H)$ if and only if $g_1 g_2 \in E(G)$; or $g_1 = g_2$ and $h_1 h_2 \in E(H)$.

Both $\square$ and $\boxtimes$ are commutative (through isomorphism). On the other hand, $\circ$ is not commutative. As illustrated in [Figure 1](#), the basic relationship between the different products is $G \square H \subseteq G \boxtimes H \subseteq G \circ H$.




Figure 1: Examples of P_4 times C_4 using different products. Notice that $P_4 \boxtimes C_4$ is $P_4 \square C_4$ plus the red edges, and $P_4 \circ C_4$ is $P_4 \boxtimes C_4$ plus the blue edges.

Our next result, proven in [Section 2](#), is a lower bound on the treewidth of a Cartesian product:

³The *Hadwiger number* of a graph H is the maximum integer t such that K_t is a minor of H . Note that since treewidth is minor-monotone, $\text{tw}(H) \geq \text{tw}(K_{\text{had}(H)}) = \text{had}(H) - 1$.

Theorem 3. For all graphs G and H , $\text{tw}(G \square H) \geq \frac{1}{2}(\text{tw}(G) + 1)(\text{tw}(H) + 1) - 1$.

The constant $\frac{1}{2}$ in Theorem 3 is best possible since Harvey and Wood [10] showed that $\text{pw}(K_m \square K_n) = (\frac{1}{2} + o(1))mn$ for $m \geq n$ (see [14] for the $m = n$ case).

A key tool in the proof of Theorem 1 is the following dual notion to treewidth introduced by Seymour and Thomas [20] called the *bramble number*. Two connected subsets A and B of $V(G)$ *touch* if $G[A \cup B]$ is connected. A *bramble* in G is a collection $\mathcal{B}$ of non-empty connected subsets of $V(G)$ that pairwise touch. A *hitting set* for $\mathcal{B}$ is a set $S \subseteq V(G)$ that intersects every element of $\mathcal{B}$. The *order* of $\mathcal{B}$, denoted by $\text{ord}(\mathcal{B})$, is the minimum size of a hitting set for $\mathcal{B}$. The *bramble number* of G , denoted by $\text{bn}(G)$, is the maximum order of a bramble in G .

Theorem 4 ([20]). For every graph G , $\text{tw}(G) + 1 = \text{bn}(G)$.

As an aside, Bellenbaum and Diestel [2] and Mazoit [15] each gave a short proof of Theorem 4.

Brambles are useful for lower bounding the treewidth. For example, recall that the $n \times n$ -grid has treewidth n . The lower bound of n follows from Theorem 4 by considering the bramble consisting of the top row, the last column minus the top vertex, and the crosses in the bottom-left $(n-1) \times (n-1)$ -grid. A well-known generalisation of this lower bound says that $\text{tw}(G \square H) \geq n$ for all connected graphs G and H with at least $n \geq 2$ vertices (see [7] for a proof). Wood [21] improved this bound by taking into account the connectivity of G and H .⁴

Theorem 5 ([21]). For all k -connected graphs G and H with at least $n \geq 2$ vertices,

$$\text{tw}(G \square H) \geq k(n - 2k + 2).$$

Qualitatively, the bound in Theorem 5 is of a very different character from the bounds in Theorems 1 and 3, which are solely in terms of treewidth. This is because treewidth is a global structural parameter, unlike connectivity, and the number of vertices in a graph can be much larger than its treewidth. Quantitatively, the bound in Theorem 5 is better when the factors have treewidth comparable to their connectivity. On the other hand, Theorems 1 and 3 are superior when the factors have large treewidth and small connectivity, such as random d -regular graphs. An important consequence of Theorems 1 and 3, which is made precise in Section 4, is that products of expanders have large subgraphs that are expanders.

Since the treewidth of a graph is at least its connectivity [6, Exercise 12.26], Theorem 5 inspires the question: Can the bound $\text{tw}(G \square H) \in \Omega(\text{tw}(G)\text{tw}(H))$ in Theorem 3 be improved to $\text{tw}(G \square H) \in \Omega(\text{tw}(G)|V(H)|)$? The answer is ‘no’ since Hickingbotham and Wood [11, Proposition 21] showed that for all $n \geq k+1$, there exists an n -vertex treewidth k graph $G_{k,n}$ such that $\text{tw}(G_{k,n} \boxtimes G_{k,n}) \in \mathcal{O}(n + k^2)$.

On a related note, Hickingbotham and Wood [11] asked whether $\text{tw}(G \boxtimes H) \in \mathcal{O}(\text{tw}(G \square H))$? Our next theorem shows that the analogous relationship fails for $\circ$ and $\boxtimes$, that is $\text{tw}(G \circ H) \notin \mathcal{O}(\text{tw}(G \boxtimes H))$. More strongly, in Section 5 we show that $\text{tw}(G \boxtimes H)$ and $\text{tw}(G \circ H)$ are not tied.

⁴Theorem 5 is stated in [21] as $\text{tw}(G \square H) \geq k(n - 2k + 2) - 1$, however it is implicit in the proof that the strict bramble number of $G \square H$, which we define later, is at least $k(n - 2k + 2)$. Then [1, Lemma 2.6] implies $\text{tw}(G \square H) \geq k(n - 2k + 2)$.

Theorem 6. *There is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\text{tw}(G \circ H) \leq f(\text{tw}(G \boxtimes H))$ for all graphs G and H .*

Guided by the role of brambles in our proof of [Theorem 1](#), we next consider the strict bramble number. A bramble in a graph G is *strict* if its elements are pairwise intersecting. The *strict bramble number* of G , denoted by $\text{sbn}(G)$, is the maximum order of a strict bramble in G .

The next two theorems give strict bramble analogues of our earlier product inequalities. The first is proven in [Section 2](#):

Theorem 7. *For all graphs G and H , $\text{tw}(G \square H) \geq \text{sbn}(G)(\text{tw}(H) + 1) - 1$.*

[Theorem 7](#) strengthens the bound $\text{tw}(G \square H) \geq \text{sbn}(G)(\text{had}(H) + 1) - 1$ by Kozawa, Otachi, and Yamazaki [12], who introduced the notion of strict brambles under the name “PI family”.⁵ They also showed that strict bramble number is tied to treewidth, in particular, $\text{sbn}(G) \geq \frac{1}{2}(\text{tw}(G) + 1)$. Therefore [Theorem 7](#) implies [Theorem 3](#).

A duality result ([Theorem 19](#) below) for strict brambles analogous to [Theorem 4](#) was proved by Lardas, Protopapas, Thilikos, and Zoros [13]. In [Section 6](#), we use their duality result to obtain the following lower bound on the strict bramble number of Cartesian products:

Theorem 8. *For all graphs G and H , $\text{sbn}(G \square H) \geq \text{sbn}(G)\text{sbn}(H)$.*

We now highlight parallels between [Theorems 1](#) and [8](#). By [Theorem 4](#), we may rewrite [Theorem 1](#) as $\text{bn}(G \boxtimes H) \geq \text{bn}(G)\text{bn}(H)$. Therefore [Theorems 1](#) and [8](#) motivate viewing $(\boxtimes, \text{bn})$ and $(\square, \text{sbn})$ as natural ‘product-parameter pairings’, in the sense that the parameter respects the product via a multiplicative lower bound. [Theorem 8](#) pertains to the more restrictive product, which necessitates a more restrictive parameter. Indeed, the proofs of [Theorems 1](#) and [8](#) are nearly identical, differing only in the modifications required to accommodate the more restrictive setting. The pairing is also supported by the optimality of [Theorem 3](#), specifically, if bn is used in place of sbn in [Theorem 8](#), then a factor of $\frac{1}{2}$ must be introduced.

2 Treewidth of Product Graphs

This section proves [Theorems 1](#), [3](#) and [7](#) (restated below).

Recall that in [Section 1](#) it was explained that [Theorem 3](#) is a consequence of [Theorem 7](#). As such, we only prove [Theorems 1](#) and [7](#).

Let G be a graph and let (T, β) be a tree-decomposition of G . For each set $X \subseteq V(G)$, let $(T, \beta)\langle X \rangle := \{t \in V(T) : X \cap \beta(t) \neq \emptyset\}$.

The following are well-known facts about tree-decompositions and brambles (see [16] for proofs):

Fact 9. *For every graph G and every tree-decomposition (T, β) of G , if $X \subseteq V(G)$ is connected in G , then $(T, \beta)\langle X \rangle$ is connected in T .*

⁵The name “strict bramble” is by Aidun, Dean, Morrison, Yu, and Yuan [1]

Fact 10. For every bramble $\mathcal{B}$ and every tree-decomposition of a graph G , some bag is a hitting set for $\mathcal{B}$.

We now prove [Theorem 1](#), which states that for all graphs G and H

$$\text{tw}(G \boxtimes H) \geq (\text{tw}(G) + 1)(\text{tw}(H) + 1) - 1.$$

Proof of Theorem 1. Let (T, β) be a minimum-width tree-decomposition of $G \boxtimes H$ and let $\mathcal{B}$ be a maximum-order bramble in G . For each vertex $v \in V(H)$, let $\mathcal{B}_v := \{A \times \{v\} : A \in \mathcal{B}\}$. Then each $\mathcal{B}_v$ is a bramble in $G \boxtimes H$ and $\text{ord}(\mathcal{B}_v) = \text{ord}(\mathcal{B}) = \text{bn}(G)$. For each node $t \in V(T)$, let

$$\beta'(t) := \{v \in V(H) : \beta(t) \text{ is a hitting set for } \mathcal{B}_v\}.$$

We show that (T, β') is a tree-decomposition of H . Consider any vertex $v \in V(H)$. Since $\mathcal{B}_v$ is a bramble in $G \boxtimes H$ and (T, β) is a tree-decomposition of $G \boxtimes H$, [Fact 10](#) implies that there exists a node $x \in V(T)$ such that $\beta(x)$ is a hitting set for $\mathcal{B}_v$, which implies $v \in \beta'(x)$. Suppose there exists a node $y \in V(T - x)$ such that $v \in \beta'(y)$. Then for all $A \times \{v\} \in \mathcal{B}_v$, since $A \times \{v\}$ is connected and intersects both $\beta(x)$ and $\beta(y)$, [Fact 9](#) implies that $A \times \{v\}$ intersects $\beta(t)$ for every node $t \in V(xy)$. Consequently, $\beta(t)$ is a hitting set for $\mathcal{B}_v$ for all $t \in V(xy)$. Thus $\{t \in V(T) : v \in \beta'(t)\}$ is non-empty and connected in T , and (T, β') has the vertex-property. Next, consider any edge $uv \in E(H)$. The following shows that $\mathcal{B}_u \cup \mathcal{B}_v$ is a bramble in $G \boxtimes H$: Since $\mathcal{B}_u$ and $\mathcal{B}_v$ are brambles in $G \boxtimes H$, it suffices to show that every $U \times \{u\} \in \mathcal{B}_u$ and every $V \times \{v\} \in \mathcal{B}_v$ touch in $G \boxtimes H$. Since $\mathcal{B}$ is a bramble in G , U and V touch in G . If $U \cap V = \emptyset$, then there exists an edge $pq \in E(G)$ with $p \in U$ and $q \in V$, therefore $(p, u)(q, v)$ is an edge between $U \times \{u\}$ and $V \times \{v\}$. On the other hand, there exists $w \in U \cap V$. Thus $(w, u)(w, v)$ is an edge between $U \times \{u\}$ and $V \times \{v\}$. In either case, $U \times \{u\} \in \mathcal{B}_u$ and $V \times \{v\} \in \mathcal{B}_v$ touch in $G \boxtimes H$, hence $\mathcal{B}_u \cup \mathcal{B}_v$ is a bramble in $G \boxtimes H$. Then by [Fact 10](#), there exists a node $t \in V(T)$ such that $\beta(t)$ is a hitting set for $\mathcal{B}_u \cup \mathcal{B}_v$. Therefore $\{u, v\} \subseteq \beta'(t)$, and (T, β') has the edge-property. Hence (T, β') is a tree-decomposition of H as claimed. Fix $t \in V(T)$ such that $|\beta'(t)|$ is maximum. So $|\beta'(t)| \geq \text{tw}(H) + 1$. Then

$$\text{tw}(G \boxtimes H) + 1 \geq |\beta(t)| \geq \sum_{v \in \beta'(t)} |\beta(t) \cap (V(G) \times \{v\})| \geq \sum_{v \in \beta'(t)} \text{ord}(\mathcal{B}_v) \geq \text{bn}(G)(\text{tw}(H) + 1).$$

Therefore $\text{tw}(G \boxtimes H) + 1 \geq (\text{tw}(G) + 1)(\text{tw}(H) + 1)$ by [Theorem 4](#). $\square$

Next we prove [Theorem 7](#), which states that for all graphs G and H

$$\text{tw}(G \square H) \geq \text{sbn}(G)(\text{tw}(H) + 1) - 1.$$

We remark that our proof of [Theorem 7](#) follows the same method as [Theorem 1](#), except we use strict brambles.

Proof of Theorem 7. Let (T, β) be a minimum-width tree-decomposition of $G \square H$ and let $\mathcal{S}$ be a maximum-order strict bramble in G . For each vertex $v \in V(H)$, let $\mathcal{S}_v := \{A \times \{v\} : A \in \mathcal{S}\}$. Then each $\mathcal{S}_v$ is a strict bramble in $G \square H$ and $\text{ord}(\mathcal{S}_v) = \text{ord}(\mathcal{S}) = \text{sbn}(G)$. For each node $t \in V(T)$, let

$$\beta'(t) := \{v \in V(H) : \beta(t) \text{ is a hitting set for } \mathcal{S}_v\}.$$

We show that (T, β') is a tree-decomposition of H . Consider any vertex $v \in V(H)$. Since $\mathcal{S}_v$ is a bramble in $G \square H$ and (T, β) is a tree-decomposition of $G \square H$, [Fact 10](#) implies that there exists a node $x \in V(T)$ such that $\beta(x)$ is a hitting set for $\mathcal{S}_v$, which implies $v \in \beta'(x)$. Suppose there exists a node $y \in V(T - x)$ such that $v \in \beta'(y)$. Then for all $A \times \{v\} \in \mathcal{S}_v$, since $A \times \{v\}$ is connected and intersects both $\beta(x)$ and $\beta(y)$, [Fact 9](#) implies that $A \times \{v\}$ intersects $\beta(t)$ for every node $t \in V(xy)$. Consequently, $\beta(t)$ is a hitting set for $\mathcal{S}_v$ for all $t \in V(xy)$. Thus $\{t \in V(T) : v \in \beta'(t)\}$ is non-empty and connected in T , and (T, β') has the vertex-property. Next, consider any edge $uv \in E(H)$. The following shows that $\mathcal{S}_u \cup \mathcal{S}_v$ is a bramble in $G \square H$: Since $\mathcal{S}_u$ and $\mathcal{S}_v$ are brambles in $G \square H$, it suffices to show that every $U \times \{u\} \in \mathcal{S}_u$ and every $V \times \{v\} \in \mathcal{S}_v$ touch in $G \square H$. Since $\mathcal{S}$ is a strict bramble in G , there exists $w \in U \cap V$, which implies that $(w, u)(w, v)$ is an edge between $U \times \{u\}$ and $V \times \{v\}$. Therefore $U \times \{u\}$ and $V \times \{v\}$ touch in $G \square H$, and $\mathcal{S}_u \cup \mathcal{S}_v$ is a bramble in $G \square H$. Then by [Fact 10](#), there exists a node $t \in V(T)$ such that $\beta(t)$ is a hitting set for $\mathcal{S}_u \cup \mathcal{S}_v$, so $\{u, v\} \subseteq \beta'(t)$, and (T, β') has the edge-property. Hence (T, β') is a tree-decomposition of H as claimed. Fix $t \in V(T)$ such that $|\beta'(t)|$ is maximum. So $|\beta'(t)| \geq \text{tw}(H) + 1$. Then

$$\text{tw}(G \square H) + 1 \geq |\beta(t)| \geq \sum_{v \in \beta'(t)} |\beta(t) \cap (V(G) \times \{v\})| \geq \sum_{v \in \beta'(t)} \text{ord}(\mathcal{S}_v) \geq \text{sbn}(G)(\text{tw}(H) + 1),$$

as required. $\square$

In our proof of [Theorem 7](#), the underlying tree T in the tree-decomposition of $G \square H$ also indexes the final tree-decomposition of H . Hence if we instead take (T, β) to be a minimum-width path-decomposition of $G \square H$, we would conclude $\text{pw}(G \square H) \geq \text{sbn}(G)(\text{pw}(H) + 1) - 1$. Since $\text{sbn}(G) \geq \frac{1}{2}(\text{tw}(G) + 1)$, we deduce

Corollary 11. *For all graphs G and H , $\text{pw}(G \square H) \geq \frac{1}{2}(\text{tw}(G) + 1)(\text{pw}(H) + 1) - 1$.*

3 Pathwidth of a Strong Product

This section proves [Theorem 2](#), which states that for all graphs G and H

$$\text{pw}(G \boxtimes H) \geq (\text{pw}(G) + 1)(\text{pw}(H) + 1) - 1.$$

In their proof of the Tree-Minor Theorem,⁶ Bienstock, Robertson, Seymour, and Thomas [3] introduced a dual parameter for pathwidth, called blockages. The first step in their proof is a duality result ([Theorem 12](#) below) that is analogous to [Theorem 4](#). We use their duality result in our proof of [Theorem 2](#).

Let G be a graph and let k be a non-negative integer. For each set $X \subseteq V(G)$, let ∂X be the set of vertices in X that have a neighbour in $V(G) \setminus X$. A *k -cut* in G is a pair (A, B) of subsets of $V(G)$ such that $|A \cap B| \leq k$ and $G[A] \cup G[B] = G$. A *cut* is a k -cut for some k . A *blockage of order k* in G is a set $\mathcal{B} \subseteq 2^{V(G)}$ that satisfies the following *blockage axioms*:

- if $X \in \mathcal{B}$, then $|\partial X| \leq k$;

⁶The Tree-Minor Theorem states that if a forest F is not a minor of G , then $\text{pw}(G) \leq |V(F)| - 2$.

- if $Y \subseteq X \in \mathcal{B}$ and $|\partial Y| \leq k$, then $Y \in \mathcal{B}$;
- if (X, Y) is a k -cut in G , then $\mathcal{B}$ contains exactly one of X or Y .

Theorem 12 ([3]). *G has a blockage of order k if and only if $\text{pw}(G) \geq k$.*

Subsequent proofs of the Tree-Minor Theorem by Diestel [5] and Seymour [19] avoid pathwidth-blockage duality by employing a shorter and more direct approach.

For convenience, our proof of Theorem 2 uses the notion of stoppages, which were also introduced by Bienstock et al. [3, §4] and are essentially equivalent to blockages. Let G be a graph and let k be a non-negative integer. A *stoppage of order k* in G is a set $\mathcal{Y}$ of k -cuts in G that satisfies the following *stoppage axioms*:

- if (A, B) is a k -cut in G , then $(A, B) \in \mathcal{Y}$ or $(B, A) \in \mathcal{Y}$;
- if $(A, B) \in \mathcal{Y}$ and $(C, D) \in \mathcal{Y}$, then $G[A] \cup G[C] \neq G$.

Bienstock et al. [3] stated the relationship between blockages and stoppages but left the details to the reader. We provide the details for completeness.

Lemma 13 ([3, 4.1 (reworded)]). *G has a blockage of order k if and only if G has a stoppage of order k .*

Proof. ($\implies$) Suppose $\mathcal{B}$ is a blockage of order k in G . Let $\mathcal{Y}$ be the set of all k -cuts (A, B) with $A \in \mathcal{B}$. We show that $\mathcal{Y}$ is a stoppage of order k in G . The first stoppage axiom follows from the third blockage axiom. For the second stoppage axiom, suppose there exists $(A, B) \in \mathcal{Y}$ and $(C, D) \in \mathcal{Y}$ with $G[A] \cup G[C] = G$. Let X be the set of vertices in $A \cap B$ that have no neighbour in $B \setminus A$, and let $B' := B \setminus X$. Then (A, B') is a k -cut. Since $A \in \mathcal{B}$, the third blockage axiom implies $B' \notin \mathcal{B}$. Note that $B' \setminus A \subseteq C$ because $G[A] \cup G[C] = G$. Moreover, each vertex $u \in A \cap B'$ has a neighbour $v \in B' \setminus A$. Then since $uv \notin E(G[A])$, $uv \in E(G[C])$, thus $u \in C$. Hence $B' \subseteq C$, so the second blockage axiom implies $B' \in \mathcal{B}$, contradiction.

($\impliedby$) Suppose $\mathcal{Y}$ is a stoppage of order k in G . Let $\mathcal{B} := \{A : (A, B) \in \mathcal{Y}\}$. We show that $\mathcal{B}$ is a blockage of order k in G . Since every element of $\mathcal{Y}$ is a k -cut, the first blockage axiom holds. For the second blockage axiom, suppose $Y \subseteq X \in \mathcal{B}$ and $|\partial Y| \leq k$. Then there exists $W \subseteq V(G)$ such that $(X, W) \in \mathcal{Y}$. Furthermore, by letting $Z := (V(G) \setminus Y) \cup \partial Y$, we have that (Y, Z) is a k -cut in G . Since $Y \subseteq X$, it follows that $(V(G) \setminus X) \cup \partial X \subseteq (V(G) \setminus Y) \cup \partial Y$, thus $G[X] \cup G[Z] = G$. Hence by the second stoppage axiom, $(Z, Y) \notin \mathcal{Y}$, thus $(Y, Z) \in \mathcal{Y}$ by the first stoppage axiom, so $Y \in \mathcal{B}$. Next we prove the third blockage axiom. Suppose that (X, Y) is a k -cut in G . Then the first stoppage axiom implies $X \in \mathcal{B}$ or $Y \in \mathcal{B}$. If $X \in \mathcal{B}$ and $Y \in \mathcal{B}$, then $(X, W) \in \mathcal{Y}$ and $(Y, Z) \in \mathcal{Y}$ for some $W, Z \subseteq V(G)$. However since (X, Y) is a cut, $G[X] \cup G[Y] = G$, which contradicts the second stoppage axiom. $\square$

We now prove Theorem 2.

Proof of Theorem 2. Let $k := (\text{pw}(G)+1)(\text{pw}(H)+1)-1$. Let $\mathcal{G}$ be a stoppage of order $\text{pw}(G)$ in G , and let $\mathcal{H}$ be a stoppage of order $\text{pw}(H)$ in H . For each $u \in V(H)$, let

$$G_u := (G \boxtimes H)[V(G) \times \{u\}] \quad \text{and} \quad \mathcal{G}_u := \{(A \times \{u\}, B \times \{u\}) : (A, B) \in \mathcal{G}\}.$$

Then $\mathcal{G}_u$ is a stoppage of order $\text{pw}(G)$ in G_u . For each $u \in V(H)$ and each subset $S \subseteq V(G \boxtimes H)$, let $S_u := V(G_u) \cap S$ and let $\pi(S) := \{v \in V(G) : (v, w) \in S\}$.

Consider each k -cut (A, B) in $G \boxtimes H$. Note that (A_u, B_u) is a cut in G_u for each $u \in V(H)$. Hence we may define

$$X_{(A,B)} := \{u \in V(H) : |A_u \cap B_u| > \text{pw}(G) \text{ or } (A_u, B_u) \in \mathcal{G}_u\}.$$

Note that the first stoppage axiom for $\mathcal{G}_u$ implies:

$$(A_u, B_u) \in \mathcal{G}_u \text{ for all } u \in X_{(A,B)} \setminus X_{(B,A)}. \quad (\star)$$

Claim 1. *If (A, B) is a k -cut in $G \boxtimes H$, then $(X_{(A,B)}, X_{(B,A)})$ is a $\text{pw}(H)$ -cut in H .*

Subproof. We first show that $X_{(A,B)} \cup X_{(B,A)} = V(H)$. Consider any $u \in V(H) \setminus X_{(A,B)}$. Then $|A_u \cap B_u| \leq \text{pw}(G)$ and $(A_u, B_u) \notin \mathcal{G}_u$, therefore by the first stoppage axiom for $\mathcal{G}_u$, $(B_u, A_u) \in \mathcal{G}_u$, implying that $u \in X_{(B,A)}$. Hence $X_{(A,B)} \cup X_{(B,A)} = V(H)$.

By the stoppage axioms, $X_{(A,B)} \cap X_{(B,A)} = \{u \in V(H) : |A_u \cap B_u| > \text{pw}(G)\}$, thus

$$(\text{pw}(G) + 1)|X_{(A,B)} \cap X_{(B,A)}| \leq \sum_{u \in X_{(A,B)} \cap X_{(B,A)}} |A_u \cap B_u| \leq |A \cap B| \leq k,$$

which implies $|X_{(A,B)} \cap X_{(B,A)}| \leq \text{pw}(H)$.

Finally, we show that there is no edge $uv \in E(H)$ such that $u \in X_{(A,B)} \setminus X_{(B,A)}$ and $v \in X_{(B,A)} \setminus X_{(A,B)}$. Assume for a contradiction there is such an edge $uv \in E(H)$. Then $(\star)$ implies that $(A_u, B_u) \in \mathcal{G}_u$ and $(B_v, A_v) \in \mathcal{G}_v$. Therefore $(\pi(A_u), \pi(B_u)) \in \mathcal{G}$ and $(\pi(B_v), \pi(A_v)) \in \mathcal{G}$. Then by the second stoppage axiom for $\mathcal{G}$, $G[\pi(A_u)] \cup G[\pi(B_v)] \neq G$. We now show that $(\pi(A_u), \pi(B_v))$ is a cut in G , which will contradict the previous sentence. As a first step, we show that $\pi(A_u) \cup \pi(B_v) = V(G)$. Consider any $b \in V(G) \setminus \pi(A_u)$. Thus $(b, u) \notin A_u$, implying that $(b, u) \notin A$. Therefore since (A, B) is a cut in $G \boxtimes H$, $(b, u) \in B \setminus A$. Then $(b, v) \in N_{G \boxtimes H}(b, u) \subseteq B$, which implies $b \in \pi(B_v)$. Hence $\pi(A_u) \cup \pi(B_v) = V(G)$. Next, assume for a contradiction that there exists an edge $ab \in E(G)$ such that $a \in \pi(A_u) \setminus \pi(B_v)$ and $b \in \pi(B_v) \setminus \pi(A_u)$. Since $a \notin \pi(B_v)$, $(a, v) \notin B_v$, implying that $(a, v) \notin B$. Therefore since (A, B) is a cut in $G \boxtimes H$, $(a, v) \in A \setminus B$. Similarly, from $b \notin \pi(A_u)$ we deduce that $(b, u) \in B \setminus A$. Now since $ab \in E(G)$ and $uv \in E(H)$, $(a, v)(b, u) \in E(G \boxtimes H)$, which along with $(a, v) \in A \setminus B$ and $(b, u) \in B \setminus A$ contradicts (A, B) being a cut in $G \boxtimes H$. Hence we have proved that $(\pi(A_u), \pi(B_v))$ is a cut in G , so $G[\pi(A_u)] \cup G[\pi(B_v)] = G$, the desired contradiction. $\blacksquare$

Let $\mathcal{Y}$ be the set of all k -cuts (A, B) in $G \boxtimes H$ such that $(X_{(A,B)}, X_{(B,A)}) \notin \mathcal{H}$.

Claim 2. *$\mathcal{Y}$ is a stoppage of order k in $G \boxtimes H$.*

Subproof. We begin by proving the first stoppage axiom for $\mathcal{Y}$. Suppose there exists a k -cut (A, B) in $G \boxtimes H$ such that $(A, B) \notin \mathcal{Y}$ and $(B, A) \notin \mathcal{Y}$. Then $(X_{(A,B)}, X_{(B,A)}) \in \mathcal{H}$ and $(X_{(B,A)}, X_{(A,B)}) \in \mathcal{H}$. Therefore $(X_{(A,B)}, X_{(B,A)})$ is a cut in H , and $H[X_{(A,B)}] \cup H[X_{(B,A)}] \neq H$ by the second stoppage axiom for $\mathcal{H}$, a contradiction. Hence $(A, B) \in \mathcal{Y}$ or $(B, A) \in \mathcal{Y}$, as required.

We now prove the second stoppage axiom for $\mathcal{Y}$. Consider any $(A, B) \in \mathcal{Y}$ and any $(C, D) \in \mathcal{Y}$. By definition of $\mathcal{Y}$, $(X_{(A,B)}, X_{(B,A)}) \notin \mathcal{H}$ and $(X_{(C,D)}, X_{(D,C)}) \notin \mathcal{H}$. Note that by **Claim 1**, $(X_{(A,B)}, X_{(B,A)})$ and $(X_{(C,D)}, X_{(D,C)})$ are $\text{pw}(H)$ -cuts in H , thus the first stoppage axiom for $\mathcal{H}$ implies that $(X_{(B,A)}, X_{(A,B)}) \in \mathcal{H}$ and $(X_{(D,C)}, X_{(C,D)}) \in \mathcal{H}$.

Then the second stoppage axiom for $\mathcal{H}$ implies $H[X_{(B,A)}] \cup H[X_{(D,C)}] \neq H$. Hence there exists $u \in X_{(A,B)} \setminus X_{(B,A)}$ and $v \in X_{(C,D)} \setminus X_{(D,C)}$ such that $\{u\}$ and $\{v\}$ touch in H . Moreover, $(\star)$ implies that $(A_u, B_u) \in \mathcal{G}_u$ and $(C_v, D_v) \in \mathcal{G}_v$. Then $(\pi(A_u), \pi(B_u)) \in \mathcal{G}$ and $(\pi(C_v), \pi(D_v)) \in \mathcal{G}$, so $G[\pi(A_u)] \cup G[\pi(C_v)] \neq G$ by the second stoppage axiom for $\mathcal{G}$. Hence there exists $b \in \pi(B_u) \setminus \pi(A_u)$ and $d \in \pi(D_v) \setminus \pi(C_v)$ such that $\{b\}$ and $\{d\}$ touch in G . Consequently, $\{(b, u)\}$ and $\{(d, v)\}$ touch in $G \boxtimes H$. To complete the argument, we show that $(b, u) \in B \setminus A$ and $(d, v) \in D \setminus C$, which will imply that $(G \boxtimes H)[A] \cup (G \boxtimes H)[C] \neq G \boxtimes H$. Indeed $(b, u) \in B_u \subseteq B$, and if $(b, u) \in A$, then $b \in \pi(A_u)$, contrary to the definition of b . Similarly we deduce that $(d, v) \in D \setminus C$, as desired. $\blacksquare$

Therefore [Theorem 12](#), [Lemma 13](#), and [Claim 2](#) imply that $\text{pw}(G \boxtimes H) \geq k$. $\square$

4 Application to Vertex Expansion

This section shows that products of graph classes with positive vertex expansion contain linearly-sized graphs with positive vertex expansion ([Theorem 16](#) below).

Let G be a graph. For a set $X \subseteq V(G)$, let $N_G(X) := \bigcup_{x \in X} N_G(x) \setminus X$. Define the *vertex expansion* of G to be

$$\varphi(G) := \min_{X \subseteq V(G): 0 < |X| \leq \frac{1}{2}|V(G)|} \frac{|N_G(X)|}{|X|}.$$

Define the *vertex expansion* of a graph class $\mathcal{G}$ to be $\varphi(\mathcal{G}) := \inf_{G \in \mathcal{G}} \varphi(G)$.

Grohe and Marx [\[8\]](#) established the following relationships between treewidth and vertex expansion:

Theorem 14 ([\[8, Proposition 1\]](#)). *For every graph G , $\text{tw}(G) \geq \lfloor \frac{1}{4}\varphi(G)|V(G)| \rfloor$.*

Theorem 15 ([\[8, Proposition 2\]](#)). *Let n be a positive integer and let $\epsilon > 0$. For every n -vertex graph G with $\text{tw}(G) \geq \epsilon n$, there exists a subgraph $S \subseteq G$ such that $\text{tw}(S) \geq \frac{1}{2}\epsilon n$ and $\varphi(S) \geq \frac{1}{2}\epsilon$.*

For $*$ in $\{\square, \boxtimes, \circ\}$, define $\mathcal{G} * \mathcal{H} := \{G * H : G \in \mathcal{G}, H \in \mathcal{H}\}$.

Theorem 16. *For all $*$ in $\{\square, \boxtimes, \circ\}$, if $\mathcal{G}$ and $\mathcal{H}$ are graph classes with positive vertex expansion, then there exists a constant $c > 0$ such that every graph $G * H \in \mathcal{G} * \mathcal{H}$ has a subgraph S with $\text{tw}(S) \geq c|V(G * H)|$, hence $|V(S)| \geq c|V(G * H)| - 1$, and $\varphi(S) \geq c$.*

Proof. Let $\epsilon_0 := \min\{\varphi(\mathcal{G}), \varphi(\mathcal{H})\}$ and let $\mathcal{F} := \{F \in \mathcal{G} * \mathcal{H} : \frac{1}{64}\epsilon_0^2|V(F)| < 1\}$. Then $\mathcal{F}$ is finite. Since $\varphi(\mathcal{G}) > 0$, every graph in $\mathcal{G}$ has an edge, thus $\epsilon_1 := \min\{\text{tw}(F)/|V(F)| : F \in \mathcal{F}\}$ is positive. Let $\epsilon := \min\{\frac{1}{64}\epsilon_0^2, \epsilon_1\}$. Now consider each $G * H \in \mathcal{G} * \mathcal{H}$. By [Theorem 3](#) and [Theorem 14](#),

$$\begin{aligned} \text{tw}(G * H) &\geq \text{tw}(G \square H) \geq \frac{1}{2}(\text{tw}(G) + 1)(\text{tw}(H) + 1) - 1 \geq \frac{1}{2}(\frac{1}{4}\epsilon_0|V(G)|)(\frac{1}{4}\epsilon_0|V(H)|) - 1 \\ &= \frac{1}{32}\epsilon_0^2|V(G * H)| - 1. \end{aligned}$$

If $G * H \notin \mathcal{F}$, then $\text{tw}(G * H) \geq \frac{1}{32}\epsilon_0^2|V(G * H)| - 1 \geq \frac{1}{64}\epsilon_0^2|V(G * H)|$. On the other hand if $G * H \in \mathcal{F}$, then $\text{tw}(G * H) \geq \epsilon_1|V(G * H)|$. Hence $\text{tw}(G * H) \geq \epsilon|V(G * H)|$ for all $G * H \in \mathcal{G} * \mathcal{H}$. Therefore [Theorem 15](#) implies the result with $c := \frac{1}{2}\epsilon$. $\square$

5 Strong Product Versus Lexicographic Product

This section proves [Theorem 6](#), which states that there is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\text{tw}(G \circ H) \leq f(\text{tw}(G \boxtimes H))$ for all graphs G and H .

We begin with the following lemma:

Lemma 17. *For all graphs G and H , if $\mathcal{B}$ is a bramble in G such that $|A| \geq 2$ for every $A \in \mathcal{B}$, then $\text{tw}(G \circ H) \geq \text{ord}(\mathcal{B})|V(H)| - 1$.*

Proof. For each $v \in V(G)$, let $H_v := (G \circ H)[\{v\} \times V(H)]$. For every subset $X \subseteq V(G)$ and every function $\sigma : X \rightarrow V(G \circ H)$, if $\sigma(v) \in V(H_v)$ for each $v \in X$, then we say that σ *represents* X . Observe that if uv is an edge in $G[X]$, then $\sigma(u)\sigma(v)$ is an edge in $G \circ H$. Therefore if X is connected in G , then $\sigma(X)$ is connected in $G \circ H$. For each $X \in \mathcal{B}$, let $\mathcal{R}_X := \{\sigma(X) : \sigma \text{ represents } X\}$. Then $\mathcal{B}' := \bigcup_{X \in \mathcal{B}} \mathcal{R}_X$ is a family of connected sets of vertices in $G \circ H$. We claim that $\mathcal{B}'$ is a bramble in $G \circ H$. To see this, consider any $X', Y' \in \mathcal{B}'$. Then there exists $X, Y \in \mathcal{B}$ and there exists σ and τ that represent X and Y respectively such that $X' = \sigma(X)$ and $Y' = \tau(Y)$. Since $\mathcal{B}$ is a bramble in G and $|X| \geq 2$ and $|Y| \geq 2$, there exists an edge $xy \in E(G)$ such that $x \in X$ and $y \in Y$. Since $\sigma(x) \in V(H_x)$ and $\tau(y) \in V(H_y)$, $\sigma(x)\tau(y)$ is an edge in $G \circ H$ between X' and Y' . Hence X' and Y' touch, and $\mathcal{B}'$ is a bramble in $G \circ H$ as claimed.

Let S be a minimum-sized hitting set for $\mathcal{B}'$, and let $F := \{v \in V(G) : V(H_v) \cap S \neq \emptyset\}$. We claim that if $v \in F$, then $V(H_v) \subseteq S$. Indeed for if not, then there exists $(v, u) \in V(H_v) \cap S$ and $(v, w) \in V(H_v) \setminus S$. By choice of S , there exists $X' \in \mathcal{B}'$ such that $S \cap X' = \{(v, u)\}$. Then there exists $X \subseteq V(G)$ and σ that represents X such that $X' = \sigma(X)$. Define $\tau : X \rightarrow V(G \circ H)$ by $\tau(v) = (v, w)$ and $\tau(x) = \sigma(x)$ for all $x \neq v$. Then τ represents X , so $\tau(X) \in \mathcal{B}'$. However since $(v, w) \notin S$, $\tau(X) \cap S = \emptyset$, which implies that S is not a hitting set for $\mathcal{B}'$, a contradiction. It follows that $\bigcup_{v \in F} V(H_v) = S$. Next, assume for a contradiction that $|S| < \text{ord}(\mathcal{B})|V(H)|$. Then,

$$|F||V(H)| = \sum_{v \in F} |V(H_v)| = |S| < \text{ord}(\mathcal{B})|V(H)|,$$

implying that $|F| \leq \text{ord}(\mathcal{B}) - 1$. Then F is not a hitting set for $\mathcal{B}$, so there exists $X \in \mathcal{B}$ such that $X \cap F = \emptyset$, which implies that every element of $\mathcal{R}_X$ is disjoint from S , a contradiction. Hence $\text{bn}(G \circ H) \geq \text{ord}(\mathcal{B}') = |S| \geq \text{ord}(\mathcal{B})|V(H)|$. Then by [Theorem 4](#), $\text{tw}(G \circ H) \geq \text{ord}(\mathcal{B})|V(H)| - 1$. $\square$

As an aside, we point out the following corollary of [Lemma 17](#) that may be of independent interest.

Corollary 18. *For all graphs G and H , if G has a maximum-order bramble with no singleton sets, then $\text{tw}(G \circ H) = (\text{tw}(G) + 1)|V(H)| - 1$.*

We now prove [Theorem 6](#).

Proof of Theorem 6. The edges of K_3 form a strict bramble of order 2. Therefore $\text{tw}(K_3 \circ P_n) \geq 2n - 1$ by [Lemma 17](#). On the other hand for each $n \geq 2$, $\text{tw}(K_3 \boxtimes P_n) \leq (\text{tw}(P_n) + 1)|V(K_3)| - 1 = 5$. Consequently, there is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\text{tw}(K_3 \circ P_n) \leq f(\text{tw}(K_3 \boxtimes P_n))$. $\square$

6 Strict Bramble Number of Cartesian Products

This section proves [Theorem 8](#), which states that for all graphs G and H

$$\text{sbn}(G \square H) \geq \text{sbn}(G) \text{sbn}(H).$$

The proof of [Theorem 8](#) is essentially the same as the proof of [Theorem 1](#). The main difference is that we replace [Theorem 4](#) with the following duality for strict bramble number by Lardas, Protopapas, Thilikos, and Zoros [13]. A *lenient tree-decomposition* of a graph G is a pair (T, ℓ) consisting of a tree T and a function $\ell : V(T) \rightarrow 2^{V(G)}$ with the following two properties: (*vertex-property*) for every vertex $v \in V(G)$, the set $\{t \in V(T) : v \in \ell(t)\}$ is non-empty and connected in T ; and (*lenient-property*) for every edge $uv \in E(G)$, the sets $\{t \in V(T) : u \in \ell(t)\}$ and $\{t \in V(T) : v \in \ell(t)\}$ touch in T . The vertices of T are called *nodes*, and the sets $\ell(t)$ are called *bags*. The *width* of (T, ℓ) is $\max_{t \in V(T)} |\ell(t)|$. Note that a lenient tree-decomposition may not be a tree-decomposition.

Theorem 19 ([13]). *For every graph G , the minimum width of a lenient tree-decomposition of G equals $\text{sbn}(G)$.*

We define analogous notation from [Section 2](#) for lenient tree-decompositions. Let G be a graph and let (T, ℓ) be a lenient tree-decomposition of G . For each set $X \subseteq V(G)$, let $(T, \ell)\langle X \rangle := \{t \in V(T) : X \cap \ell(t) \neq \emptyset\}$.

The following basic facts about lenient tree-decompositions and strict brambles have proofs that are similar to [Facts 9](#) and [10](#):

Fact 20. *For every graph G and every lenient tree-decomposition (T, ℓ) of G , if $X \subseteq V(G)$ is connected in G , then $(T, \ell)\langle X \rangle$ is connected in T .*

Fact 21. *For every strict bramble $\mathcal{S}$ and every lenient tree-decomposition of a graph G , some bag is a hitting set for $\mathcal{S}$.*

We now prove [Theorem 8](#).

Proof of Theorem 8. Let (T, ℓ) be a minimum-width lenient tree-decomposition of $G \square H$. Then [Theorem 19](#) implies $\text{sbn}(G \square H) \geq |\ell(t)|$ for all $t \in V(T)$. Let $\mathcal{S}$ be a maximum-order strict bramble in G . For each vertex $v \in V(H)$, let $\mathcal{S}_v := \{A \times \{v\} : A \in \mathcal{S}\}$. Then each $\mathcal{S}_v$ is a strict bramble in $G \square H$ and $\text{ord}(\mathcal{S}_v) = \text{ord}(\mathcal{S}) = \text{sbn}(G)$. For each node $t \in V(T)$, let

$$\ell'(t) := \{v \in V(H) : \ell(t) \text{ is a hitting set for } \mathcal{S}_v\}.$$

We show that (T, ℓ') is a lenient tree-decomposition of H . Consider any vertex $v \in V(H)$. Since $\mathcal{S}_v$ is a strict bramble in $G \square H$ and (T, ℓ) is a lenient tree-decomposition of $G \square H$, [Fact 21](#) implies that there exists a node $x \in V(T)$ such that $\ell(x)$ is a hitting set for $\mathcal{S}_v$, which implies $v \in \ell'(x)$. Suppose there exists a node $y \in V(T - x)$ such that $v \in \ell'(y)$. Then for all $A \times \{v\} \in \mathcal{S}_v$, since $A \times \{v\}$ is connected and intersects both $\ell(x)$ and $\ell(y)$, [Fact 20](#) implies that $A \times \{v\}$ intersects $\ell(t)$ for every node $t \in V(xy)$. Consequently, $\ell(t)$ is a hitting set for $\mathcal{S}_v$ for all $t \in V(xy)$. Thus $\{t \in V(T) : v \in \ell'(t)\}$ is non-empty and connected in T , and

(T, ℓ') has the vertex-property. Next, assume for a contradiction that (T, ℓ') does not have the lenient-property. Then there exists an edge $uv \in E(H)$ such that the sets of nodes $U := \{t \in V(T) : u \in \ell'(t)\}$ and $V := \{t \in V(T) : v \in \ell'(t)\}$ do not touch. As argued above, U and V are non-empty and connected in T . Then there exists $t \in V(T) \setminus (U \cup V)$ that separates U and V in T . Since $t \notin U \cup V$, $\ell(t)$ is neither a hitting set for $\mathcal{S}_u$ nor for $\mathcal{S}_v$. Hence there exists $X \times \{u\} \in \mathcal{S}_u$ and $Y \times \{v\} \in \mathcal{S}_v$ that are disjoint from $\ell(t)$. Since U and V are non-empty, $(T, \ell) \langle X \times \{u\} \rangle \cap U \neq \emptyset$ and $(T, \ell) \langle Y \times \{v\} \rangle \cap V \neq \emptyset$, therefore $(T, \ell) \langle X \times \{u\} \rangle$ and $(T, \ell) \langle Y \times \{v\} \rangle$ lie in different components of $T - t$. Then the lenient-property of (T, ℓ) implies that $X \times \{u\}$ and $Y \times \{v\}$ do not touch in $G \square H$. However since $\mathcal{S}$ is strict, $X \cap Y \neq \emptyset$, thus $uv \in E(H)$ implies that $X \times \{u\}$ and $Y \times \{v\}$ touch in $G \square H$, a contradiction. Therefore (T, ℓ') has the lenient-property, and (T, ℓ') is a lenient tree-decomposition of H as claimed. Fix $t \in V(T)$ such that $|\ell'(t)|$ is maximum. So $|\ell'(t)| \geq \text{sbn}(H)$ by [Theorem 19](#). Then

$$\text{sbn}(G \square H) \geq |\ell(t)| \geq \sum_{v \in \ell'(t)} |\ell(t) \cap (V(G) \times \{v\})| \geq \sum_{v \in \ell'(t)} \text{ord}(\mathcal{S}_v) \geq \text{sbn}(G) \text{sbn}(H). \quad \square$$

We end with two open problems. The first is about the pathwidth of a Cartesian product:

Question 22. Does [Corollary 11](#) hold when $\text{tw}(G)$ is replaced by $\text{pw}(G)$?

Since Harvey and Wood [10] showed that $\text{pw}(K_m \square K_n) = (\frac{1}{2} + o(1))mn$ for $m \geq n$, a positive answer to [Question 22](#) would be optimal. The second is the question of Hickingbotham and Wood [11] that inspired [Theorem 6](#):

Question 23 (Hickingbotham and Wood [11]). Is $\text{tw}(G \boxtimes H) \in \mathcal{O}(\text{tw}(G \square H))$?

The following argument due to Neel Kaul (private communication, 2026) shows that $\text{tw}(G \boxtimes H) \leq (\text{degen}(G) + 1)(\text{tw}(G \square H) + 1) - 1$, where $\text{degen}(G)$ is the degeneracy of G : Let (T, β) be a minimum-width tree-decomposition of $G \square H$ and consider an orientation of the edges of G such that $\Delta^+(G) = \text{degen}(G)$. For each node $t \in V(T)$, let $\beta'(t) := \bigcup_{(g,h) \in \beta(t)} (\{g\} \cup N_G^+(g)) \times \{h\}$. Then (T, β') is a tree-decomposition $G \square H$. Now consider any edge $(g_1, h_1)(g_2, h_2) \in E(G \boxtimes H) \setminus E(G \square H)$. Then $g_1 g_2 \in E(G)$ and $h_1 h_2 \in E(H)$. Without loss of generality $g_2 \in N_G^+(g_1)$. Since $(g_1, h_1)(g_1, h_2) \in E(G \square H)$, there exists $t \in V(T)$ such that $\{(g_1, h_1), (g_1, h_2)\} \subseteq \beta(t)$. Hence $\{(g_1, h_1), (g_2, h_2)\} \subseteq \beta'(t)$. Therefore (T, β') is a tree-decomposition of $G \boxtimes H$. Finally, since $|\beta'(t)| \leq (\Delta^+(G) + 1)|\beta(t)|$ for all $t \in V(T)$, we have $\text{tw}(G \boxtimes H) \leq (\text{degen}(G) + 1)(\text{tw}(G \square H) + 1) - 1$. A consequence of this argument is that $\text{tw}(G \square H)$ and $\text{tw}(G \boxtimes H)$ are polynomially tied.

Acknowledgements

Thanks to David Wood and Neel Kaul for fruitful discussions and for their feedback on the paper. Thanks to Jung Hon Yip for feedback on the paper.

References

- [1] Ivan Aidun, Frances Dean, Ralph Morrison, Teresa Yu, and Julie Yuan. [Treewidth and gonality of glued grid graphs](#). *Discrete Appl. Math.*, 279:1–11, 2020.
- [2] Patrick Bellenbaum and Reinhard Diestel. [Two short proofs concerning tree-decompositions](#). *Combin. Probab. Comput.*, 11(6):541–547, 2002.
- [3] Dan Bienstock, Neil Robertson, Paul Seymour, and Robin Thomas. [Quickly excluding a forest](#). *J. Combin. Theory Ser. B*, 52(2):274–283, 1991.
- [4] Hans L. Bodlaender. [A partial \$k\$ -arboretum of graphs with bounded treewidth](#). *Theoret. Comput. Sci.*, 209(1):1–45, 1998.
- [5] Reinhard Diestel. [Graph minors I: A short proof of the path-width theorem](#). *Combin. Probab. Comput.*, 4(1):27–30, 1995.
- [6] Reinhard Diestel. *Graph theory*, volume 173 of *Graduate Texts in Mathematics*. Springer, 6th edition, 2025.
- [7] Vida Dujmović, Pat Morin, David R. Wood, and David Worley. [Grid minors and products](#), 2025. arXiv:2402.14181.
- [8] Martin Grohe and Dániel Marx. [On tree width, bramble size, and expansion](#). *J. Combin. Theory Ser. B*, 99(1):218–228, 2009.
- [9] Daniel J. Harvey and David R. Wood. [Parameters tied to treewidth](#). *J. Graph Theory*, 84(4):364–385, 2017.
- [10] Daniel J. Harvey and David R. Wood. [The treewidth of line graphs](#). *J. Combin. Theory Ser. B*, 132:157–179, 2018.
- [11] Robert Hickingbotham and David R. Wood. [Structural properties of graph products](#). *J. Graph Theory*, 109(2):107–136, 2025.
- [12] Kyohei Kozawa, Yota Otachi, and Koichi Yamazaki. [Lower bounds for treewidth of product graphs](#). *Discrete Appl. Math.*, 162:251–258, 2014.
- [13] Emmanouil Lardas, Evangelos Protopapas, Dimitrios M. Thilikos, and Dimitris Zoros. [On strict brambles](#). *Graphs Combin.*, 39(2):24, 2023.
- [14] Brian Lucena. [Achievable sets, brambles, and sparse treewidth obstructions](#). *Discrete Appl. Math.*, 155(8):1055–1065, 2007.
- [15] Frédéric Mazoit. [A simple proof of the tree-width duality theorem](#), 2013. arXiv:1309.2266.
- [16] Bruce A. Reed. [Tree width and tangles: a new connectivity measure and some applications](#). In R. A. Bailey, editor, *Surveys in Combinatorics*, volume 241 of *London Math. Soc. Lecture Note Ser.*, pages 87–162. Cambridge University Press, 1997.
- [17] Bruce A. Reed. [Algorithmic aspects of tree width](#). In *Recent advances in algorithms and combinatorics*, volume 11, pages 85–107. Springer, 2003.
- [18] Neil Robertson and Paul Seymour. [Graph minors. V. Excluding a planar graph](#). *J. Combin. Theory Ser. B*, 41(1):92–114, 1986.
- [19] Paul Seymour. [A shorter proof of the path-width theorem](#), 2023. arXiv:2309.05100.
- [20] Paul Seymour and Robin Thomas. [Graph searching and a min-max theorem for tree-width](#). *J. Combin. Theory Ser. B*, 58(1):22–33, 1993.
- [21] David R. Wood. [Treewidth of Cartesian products of highly connected graphs](#). *J. Graph Theory*, 73(3):318–321, 2013.