

PIVCATA LEARNER'S MATERIAL

QUARTER 2

Mathematics

G7





DepEd CALABARZON
Curriculum and Jearning Management Division

COVERNMENT PROPE

Republic Act 8293, section 176 states that: No copyright shall subsist in any work of the Government of the Philippines. However, prior approval of the government agency or office wherein the work is created shall be necessary for exploitation of such work for profit. Such agency or office may, among other things, impose as a condition the payment of royalties.

Borrowed materials (songs, stories, poems, pictures, photos, brand names, trademarks, etc.) included in this module are owned by their respective copyright holders. Every effort has been exerted to locate and seek permission to use these materials from their respective copyright owners. The publisher and the authors do not represent nor claim ownership over them.

This module was carefully examined and revised in accordance with the standards prescribed by the DepEd Regional Office 4A and CLMD CALABARZON. All parts and sections of the module are assured not to have violated any rules stated in the Intellectual Property Rights for learning standards.

The Editors

PIVOT 4A Learner's Material Quarter 2 Second Edition, 2021

Mathematics Grade 7

Job S. Zape, Jr.

PIVOT 4A Instructional Design & Development Lead

Jisela N. Ulpina

Content Creator & Writer

Jhonathan S. Cadavido & Angelo D. Uy Internal Reviewer & Editors

Lhovie A. Cauilan & Jael Faith T. Ledesma
Layout Artist & Illustrator

Jhucel A. del Rosario & Melanie Mae N. Moreno Graphic Artist & Cover Designer

Ephraim L. Gibas IT & Logistics

Published by: Department of Education Region IV-A CALABARZON

Regional Director: Francis Cesar B. Bringas

Guide in Using PIVOT 4A Learner's Material

For the Parents/Guardians

This module aims to assist you, dear parents, guardians, or siblings of the learners, to understand how materials and activities are used in the new normal. It is designed to provide information, activities, and new learning that learners need to work on.

Activities presented in this module are based on the Most Essential Learning Competencies (MELCs) in **Mathematics** as prescribed by the Department of Education.

Further, this learning resource hopes to engage the learners in guided and independent learning activities at their own pace. Furthermore, this also aims to help learners acquire the essential 21st century skills while taking into consideration their needs and circumstances.

You are expected to assist the children in the tasks and ensure the learner's mastery of the subject matter. Be reminded that learners have to answer all the activities in their own notebook.

For the Learners

The module is designed to suit your needs and interests using the IDEA instructional process. This will help you attain the prescribed grade-level knowledge, skills, attitude, and values at your own pace outside the normal classroom setting.

The module is composed of different types of activities that are arranged according to graduated levels of difficulty—from simple to complex. You are expected to:

- a. answer all activities on separate sheets of paper;
- b. accomplish the **PIVOT Assessment Card for Learners on page 38** by providing the appropriate symbols that correspond to your personal assessment of your performance; and
- c. submit the outputs to your respective teachers on the time and date agreed upon.

Parts of PIVOT 4A Learner's Material

	K to 12 Learning Delivery Process	Descriptions	
Introduction	What I need to know	This part presents the MELC/s and the desired learning outcomes for the day or week, purpose of the lesson, core content and relevant samples.	
Intro	What is new	This maximizes awareness of his/her own knowledge as regards content and skills required for the lesson.	
ent	What I know	This part presents activities, tasks and contents of value and interest to learner. This exposes	
Development	What is in	him/her on what he/she knew, what he/she does not know and what he/she wants to know and learn. Most of the activities and tasks simply and	
ď	What is it	directly revolve around the concepts of developing mastery of the target skills or MELC/s.	
	What is more	In this part, the learner engages in various tasks and opportunities in building his/her knowledge skills and attitude/values (KSAVs) to meaningfully connect his/her concepts after	
Engagement	What I can do	doing the tasks in the D part. This also exposes him/her to real life situations/tasks that shall: ignite his/ her interests to meet the expectation; make his/her performance satisfactory; and/or	
, E	What else I can do	produce a product or performance which will help him/her fully understand the target skills and concepts.	
ation	What I have learned	This part brings the learner to a process where he/she shall demonstrate ideas, interpretation, mindset or values and create pieces of information that will form part of his/her	
Assimilation	What I can achieve	knowledge in reflecting, relating or using them effectively in any situation or context. Also, this part encourages him/her in creating conceptual structures giving him/her the avenue to integrate new and old learnings.	

This module is a guide and a resource of information in understanding the Most Essential Learning Competencies (MELCs). Understanding the target contents and skills can be further enriched thru the K to 12 Learning Materials and other supplementary materials such as Worktexts and Textbooks provided by schools and/or Schools Division Offices, and thru other learning delivery modalities, including radio-based instruction (RBI) and TV-based instruction (TVI).

Measuring Quantities

Lesson

Measurement is a number that characterized an object or event, which can be compared with other objects or events. You use basic unit of measure for length, time, weight/mass, electric current, volume, angles, temperature and rate. There are two systems of measures, the Metric System or the International System of Units (SI) and the English System. Most countries, including the Philippines, use the SI Unit of measure.

The table below shows the commonly used units of measure for different quantities.

	Units of Measure					
Quantity	Quantity Metric English					
Length	millimeter (mm) centimeter (cm) meter (m) kilometer (km)	inch (in) foot (ft) yard (yd) mile (mi)				
Volume	cubic centimeter (cm³) cubic meter (m³)	cubic inch (in³) cubic foot (ft³) cubic yard (yd³)				
Mass	gram (g) kilogram (kg)	ounce (oz) ton (t) pound (lb)				
Temperature	Kelvin (K) degree Celsius (°C)	degree Fahrenheit (°F)				
Time	second (s) minute (min) hour (h)	second (s) minute (min) hour (h)				
Capacity	milliliter (mL) liter (L)	fluid ounce (fl. oz) pint gallon				
Angle	degree radian	degree radian				



Learning Task 1. Identify the appropriate unit of measure that can be used to the following quantities.

Physical Quantity	Unit of measure	Physical Quantity	Unit of measure
1. height of the table		5. distance of your place to Manila	
2. length of a handkerchief		6. oven temperature while baking cake	
3. volume of an ice cream		7. mass of a balikbayan box	
4. length of a ball pen		8. size of the shoes	

Each unit of measure has its own use in any physical quantities.

1. **Length** describes how long a physical quantity is. It includes distance, height, depth and the like. In the previous page you can see the different units of measures used in measuring length. The smallest unit of measure for length is millimeter (mm) and the biggest unit of measure is kilometer (km). Length is use to determine perimeter and area of other geometric figures. The basic instrument used in measuring length is a ruler. It has four units of measure, millimeter (mm), centimeter (cm), inches (in), and foot. Meter stick is used to measure longer length like measur-



Instruments Used to Measure Length


ing the floor of your house, the distance between the floor and the ceiling and many more. In English System, yard stick is used. Distance from one town to another is measured by kilometers in metric system or miles in English System. The instrument use is odometer.

One unit of measure can be expressed in another unit of measure. It can be from smaller to bigger unit and vice versa or from Metric unit to English unit of measure and vice versa.

The table below shows the equivalent measure of one unit to meter since it is the basic unit of measure in metric system.

Unit	Meaning	
Millimeter (mm)	$\frac{1}{1000} = 0.001$	meter
Centimeter (cm)	$\frac{1}{100} = 0.01$	meter
Decimeter (dm)	$\frac{1}{10} = 0.1$	meter
Dekameter (dam)	10	meters
Hectometer (hm)	$100 = 10^2$	meters
Kilometer (km)	$1000 = 10^3$	meters

In converting smaller unit to bigger unit, divide the given measure by the corresponding powers of 10 while from bigger to smaller multiply the given measure by powers of 10.



1. Convert 3 km to a) m b) cm

Solution: Given: 3 km to be converted to smaller unit. Hence you multiply by powers of 10.

- a. Since meter is 3 steps from kilometer, then $3 \text{ km} = 3 \text{ x } 10^3 = 3,000 \text{ meters}$
- b. Centimeter is 5 steps from kilometer, then $3 \text{ km} = 3 \text{ x } 10^5 = 300\ 000 \text{ cm}$
- 2. Convert 560,000 cm to a) dam b) km

Solution: Given: 560,000 cm to be converted to bigger unit. Hence you divide by powers of 10.

- a. Since dam is 3 steps from cm, then $560,000 \text{ cm} = 560,000 \div 10^3 = 560 \text{ dam}$
- b. Kilometer is 5 steps from centimeter, then 560 000cm = $560,000 \div 10^5 = 5.6 \text{ km}$
- 3. The length of the rectangular lot is 12 m and the width is 550 cm. Find the perimeter and the area.

Solution: Given l = 12m w = 550 cm

Before solving for the perimeter and the area of the rectangular lot, make sure that the length and the width have the same unit of measure. Hence you can convert one unit in terms of the other unit. In this case centimeter is converted to meters. $550 \text{ cm} = 550 \div 10^2 = 5.5 \text{ m}$. The width now is w = 5.5 m

$$P = 2l + 2w$$
 (formula for the perimeter) $A = lw$ (formula for the area)
= $2(12) + 2(5.5)$ = $12(5.5)$
= $24 + 11 = 35$ meters = 66 m^2 (square meter)

The table below shows how to convert the English unit to metric unit of measure and vice versa.

Converting	to	Multiply by	Result
Inch (in)	cm	2.54	1 in = 2.54 cm
Foot (ft)	cm	30.5	1 ft = 30.5 cm
Yard (yd)	m	0.9	1 yd = 0.914 m
	cm	90	= 91.4 cm
Mile (mi)	km	1.61	1 mi = 1.61 km
Centimeter (cm)	in	0.4	1 cm = 0.4 in
Meter (m)	in	39.4	1m = 39.4 in
	ft	3.28	= 3.28 ft
	yd	1.1	= 1.1 yd
Kilometer (km)	mi	0.62	I km = 0.62 mi

1. What is 4.2 inches in centimeter?

Solution: Since 1 in = 2.54 cm, then 4.2 in = 4.2(2.54) = 10.67 cm

2. What is 10.5 m in yards?

Solution: Since 1 m = 1.1 yd, then 10.5m = 10.5(1.1) = 11.55 yds.

3. Three yards of ribbon will be cut into 7 equal pieces. How long in centimeters is each piece?

Solution: Convert 3 yards to centimeters.


Since I yd = 91.4 cm, then 3 yd = 3(91.4) = 274.2 cm

To get the length of each piece : $274.2 \div 7 = 39.17$ cm per piece.

2. Volume and Capacity

Volume is the measure of how much amount of space is occupied by a 3-dimensional figure. **Capacity** is the amount contained in a 3-dimensional figure. Volume can be measured using SI units like cubic meter (m³) for solid, which is the basic unit for volume. Other unit of measure is Liter for liquid which is equivalent to 1000 cubic centimeter. For English System, gallon, pints or fluid ounce are used to measure the capacity of liquid contained in a 3-dimensional figures.

The illustration at the right is a cube whose edge measures 10 cm. The volume of the cube is 1000 cm³ or cc (cubic centimeter). This means that it can hold a capacity of 1000cc or of liquid equivalent to 1 Liter (L). Your water consumption is measured in cubic centimeter. You can verify it through your water bill. For larger capacity , cubic meter is used. Smaller capacity is measured using milliliter (mL) , which is commonly used for liquid medicine prescriptions.



Below is the conversion table for volume.

Unit	Equal to	Unit	Equal to
Cubic kilometer (km³)	$10^9 \; {\rm m}^3$	Cubic meter (m³)	1000 L
Cubic hectometer (hm³)	$10^6\mathrm{m}^3$	Hectoliter (hL)	100 L
Cubic dekameter (dam³)	10^3 m^3	dekaLiter (daL)	10 L
Cubic meter (m³)	$10^0 \text{ m}^3 = 1$	Cubic decimeter (dm³)	1 L
Cubic decimeter (dm³)	10 ⁻³ m ³	Deciliter (dL)	0.1 L
Cubic centimeter (cm³)	10 ⁻⁶ m ³	Centiliter (cL)	0.01 L
Cubic millimeter (mm³)	10 ⁻⁹ m ³	Milliliter (mL) = cc	0.001 L

1. Convert 8 m³ to mm³.

Solution: Since the conversion is from bigger unit to smaller unit, you have to multiply every step by 10^3 . Cubic millimeter (mm³) is 3 steps from m³, therefore $8\text{m}^3 = 8(10^9) = 8 \times 10^9$ cubic millimeter.

2. Convert 340,000 cm³ to km³

Solution: Since the conversion is from smaller to bigger unit, you have to divide every step by 10^3 . Cubic kilometer (km³) is 5 steps from cm³, therefore $430000~\text{cm}^3$ = $430~000 \div 10^{15}$ = $430~00~\text{x}~10^{-15}$ or $4.3~\text{x}~10^{-19}~\text{km}^3$.

3. The rectangular prism measures 0.054 m in length, the height is 5 cm and the width is 42 mm, what is the volume of the prism in cubic centimeter? How many liters of liquid can it hold?

Solution. All units of measure must be in centimeter. Convert 0.054 m and 42 mm to cm. 0.054 m = 0.054 x 10^2 = 5.4 cm; 42 mm = $42 \div 10$ = 4.2 cm

The volume of the prism is $V = lwh = (5.4 cm) (4.2 cm)(5cm) = 113.4 cm^3$

Since 1 cc = 0.001 L, then $113.4 \text{ cm}^3 \text{ or cc} = 113.4 \text{ x} \cdot 0.001 = 0.1134 \text{ L}$

Other measures for capacity that are very useful in your home are shown in the table below:

Convert	to	Multiply by	Result
Teaspoon (tsp)	mL	5	1 tsp = 5 mL
Tablespoon (tbsp)	mL	15	1 tbsp = 15 ml
Fluid ounce (fl.oz)	mL	30	1 fl.oz = 30 mL
Cup (c)	L	0.28	1 c = 0.28 L
Pint (pt)	L	0.47	1 pt = 0.47 L
	С	2	= 2 c
Quart (qt)	L	0.95	1qt = 0.95
	С	4	= 4 c
	pt	2	= 2 pt
Gallon (gal)	L	3.8	1gal = 3.8 L
	qt	4	= 4 qt

Examples:

1. How many liters is 16 cups of milk?

Solution: Since 1 cup is 0.28 L, then 16 cups = 16×0.28 = 4.48 L.

2. A bottle of cough syrup contains 5 fl. oz . If you are going to drink the medicine and the doctor prescribed 5 mL three times a day, how many days you're going to drink the one bottle of medicine?

Solution: Convert 5 fl.oz to mL. 5 fl.oz = $5 \times 30 = 150$ mL. If you are going to drink 5 mL, 3 times a day, in one day you will consume 15 mL.


The number of days you can consume the medicine is $150 \div 15 = 10$ days

3. **Mass** is a measure of the amount of matter in a sub stance or an object. The basic SI unit for mass is the kilogram (kg), but smaller masses may be measured in grams (g), for great amount of mass, the unit of measure is ton. To measure mass, you would use a balance or weighing scale. Weighing scales can be in different forms according to its use. There are weighing scales intended for kitchen use, for markets and some for heavy weights. At the right are samples of weighing scales.



The table below shows the metric unit of measure and its equivalent in grams.

Unit	Equivalent
Milligrams (mg)	1 mg = 0.001g
Centigram (cg)	1 cg = 0.01 g
Decigram (dg)	1dg = 0.1 g
Gram (g)	1g = 1 g
Dekagram (dag)	1 dag = 10 g
Hectogram (hg)	1 hg = 100 g
Kilogram (kg)	1 kg = 1000 g
Metric ton (t)	1 t = 1 000 000 g
	= 1000 kg



Examples:

1. Every cup cake needs 2250 mg of flour . How many grams of flour is needed for 12 cup cakes?

Solution: Convert first 2250mg to g. Since converting smaller unit to bigger divide every step by 10. Gram is 3 steps from mg, then $2250\text{mg} = 2250 \div 10^3 = 2.25 \text{ g}$

12 cup cakes need 2. 25 g x 12 = 27 g of flour

2. In your village, the garbage collector collects 980 kg of garbage everyday. How many metric ton of garbage in 30 days can they collect?

Solution: 1 metric ton = 1000 kg , therefore $980 \text{ kg} = 980 \div 1000 = 0.98 \text{ t}$ In 30 days they can collect $0.98 \times 30 = 29.4 \text{ t}$ of garbage.

4. **Temperature** refers to the measure of the hotness or coldness of an object or substance with reference to some standard value. The instrument used to determine temperature is a thermometer. It is measured by Kelvin (K), degree Celsius (°C), and degree Fahrenheit (°F) The commonly used units of measure for temperature are degree Celsius and degree Fahrenheit.

Comparing the three units of measure as you can see in the figure the boiling point of water, body temperature, room temperature and freezing point of water are different from each other. You can convert one unit of measure to the other using these formulas:

Fahrenheit to Celsius


$${}^{0}\text{C} = \frac{5}{9} ({}^{0}F - 32)$$
 or ${}^{0}\text{C} = 0.56 ({}^{0}\text{F} - 32)$

Celsius to Fahrenheit

$${}^{0}F = \frac{9}{5}({}^{0}C) + 32$$
 or ${}^{0}F = 1.8({}^{0}C) + 32$

Celsius to Kelvin

$$K = {}^{0}C + 273$$



Solution: Find how much higher the melting point of candle than butter:

$$55^{\circ}$$
C - 31° C = 24° C. Convert 24° C to ${}^{\circ}$ F. ${}^{\circ}$ F = $1.8(24) + 32 = 75.2$

310.15

273.15

-273.15

212 Water Boils

98.6 Body temperature

Water Freezes

459.67 Absolute Zero

The melting point of candle is 75.2 °F higher than the butter.

2. The recipe for a certain cake to be baked in an oven calls for a 475°F tempera ture. If your oven is set in degree Celsius, what should be your temperature setting?

Solution: Change ${}^{0}F$ to ${}^{0}C$: ${}^{0}C = 0.56(475 - 32) = 0.56(443) = 248.08 {}^{0}C$ or you may round off to 250 °C in your oven setting.

5. Time is very important and the fundamental quantities of physical world. As the saying goes "Time is gold". Everything you do is bounded with time. Time is a period during which an action or event occurs.

Earth revolves around the sun for 365 days, 5 hours, 48 minutes and 46 seconds. However, the 365 days is adapted as equivalent to one year. Nevertheless, the 5 hours, 48 minute and 46 seconds are still considered that is why we have leap year, which is 366 days every 4 years.

The table shows the unit of time:

60 seconds (s)	= 1 minute	366 days	= 1 leap year
60 minute (min)	= 1 hour	10 years	= 1 decade
24 hours (h)	= 1 day	20 years	= 1 score
12 months (mo)	= 1 year (yr)	100 years	= 1 century
365 days	= 1 year	1000 years	= 1 millennium

Examples:

1. Joy recorded a song for 140 minutes. How many hours did she spend in recording the song?

Solution: Since 1 hour = 60 minutes, then $140 \text{ min} = 140 \div 60 = 2.33 \text{ hours}$


2. Approximately it takes 1575.5 minutes to travel by sea from Manila to Leyte. How many days, hours, minute and seconds does it take to travel from Manila to Leyte?

Solution: Convert 1575.5 min to hour \rightarrow 1575.5 min = 1575.5 \div 60 = 26.26 hours Convert 26.26 hours to days. \rightarrow 26.26 h = 26.26 \div 24 = 1. 09 days Convert 0.09 days to hour \rightarrow 0.09 day = 0.09(24) = 3.06 hours Convert 0.06 hour to min. \rightarrow 0.06 hr = 0.06(60) = 3.6 min Convert 0.6 min to second. \rightarrow 0.6min = 0.6(60) = 36 seconds

Thus, it takes 1 day, 3 hours, 3 minutes and 36 seconds to travel from Manila to Leyte.

6. Angle is formed by two rays intersecting at a point called the **vertex** of the angle. The two rays are called the **sides of the angle**. The amount of the opening of the two sides determines the measure of the angle. The unit of measure is de-

grees and the instrument used to measure the angle is a protractor. The protractor has inner and outer scale. The inner scale is read in a counter clockwise direction while the outer scale is read in a clock wise direction. The angle can be named using the letter assigned to the vertex. However if there are several angles with the same vertex you use the 3 letters



in naming the angle. In the figure at the right , you can name several angles like $\angle BAC$ or $\angle CAB$ (the symbol \angle is read as angle) The measure of this angle is 45° , reading the inner scale. Make sure that one side of the angle lies on the zero line. This angle is an **acute angle**. Any angle that measures less than 90° is an acute angle. $\angle BAD = 90^{\circ}$. This angle is a right angle. **Right angle** measures 90° . An angle whose measure is more than 90° is an **obtuse angle** like $\angle FAC = 135^{\circ}$, the measure is read using the outer scale.

Examples:

Using the figure above determine the measure and kind of angles are the following:

- a. ∠*FAE*
- b $\angle BAE$
- c ∠*DAF*
- $d \angle EAC$

Solution: a. 40°, acute angle

- c. 900 right angle
- b 140°, obtuse angle
- d. $\angle EAC = \angle FAC \angle FAE = 95^{\circ}$ obtuse angle

7. Rate is a ratio of two different quantities. Among these ratios are speed, which is the ratio between distance and time, water consumption is the ratio of the price per cubic meter and many more.

1. What is the speed of the car if it travels 100 kilometer in 2 hours?

Solution: The ratio between the distance and the time is the speed.

 $\frac{100}{2}$ = 55 The speed is 55 km per hour or 55kph



Learning Task 2

A. Convert to the indicated unit of measure

6.
$$3,569 \text{ sec} = ___days$$

3.
$$108 \text{ cc} = \underline{\qquad} \text{mL}$$

4.
$$8 \text{ tbsp} = ___ \text{mL}$$

9.
$$200 \, {}^{\circ}\text{F} = \underline{\hspace{1cm}}^{\circ}\text{C}$$

B. Identify the kind of angle being represented by the following angle measures.

- C. Solve the following;
- 1. Elisse has 20 meters of ribbon. How many 15 cm of ribbon can be cut from it?
- 2. Mr. JB has a rectangular lot that measures 550 m by 0.75 km. What is the area of the land in square meter. He is selling the whole lot for 206, 250,000 pesos, how much is the price per square meter?
- 3. Drinking 8 glasses of water a day is good for your health. If each glass of water is equivalent to 148 mL, How many liters of water you need to consume everyday?
- 4. Your baby brother has a body temperature of 100 °F. What is his body temperature in °C? Does he has a normal body temperature? How do you say so?
- 5. Your mother asked you to buy 2.5 kg of sugar. Each pack of sugar in the store weighs 250 grams. How many packs of sugar will you buy?
- 6. The 120 grams powdered milk make 12 cups of milk. How many kilograms of powdered milk can produce 60 cups of milk. How many liters is 60 cups of

7. You have 8 modules to study in a day. Your teacher gave you this schedule for 5 days from Monday to Friday.

Subject	Number of minutes per day	Subject	Number of minutes per day
Math	45	MAPEH	30
Science	40	EPP	30
English	40	ESP	25
Filipino	35		


- A. How many hours in a day you spend studying all these subjects?
- B. If you will add the number of minutes you spend in each subject in one week, how many hours you spend in one week for each subject?
- C. Your mother gives you 2700 seconds playtime everyday, how many minutes is your playtime?



Refer to the figure at the right. Find the measure and identify the kind of angle.

1.

4. *∠NAP*



B. Below is a recipe of a cake. Find the equivalent amount for 3 cakes.

Ingredients	Measure for one cake	Measure for 3 cakes
Cake flour	12,000 mg	g
Milk	2 cups	mL
Brown sugar	1 pint	cups
butter	5 mg	g
water	0.006 L	ml
Chocolate chips	100 mg	g

- 1. If the cost of all the ingredients for 3 cakes is P 360.00, how much is the cost of the ingredients per cake?
- 2. If a cake was sold for P220.00, how much is the profit for the three cakes?

Algebraic Expressions

Lesson

In the English subject you learn about phrases and sentences. Mathematics has also mathematical phrases and sentences. You have to translate verbal phrases to mathematical symbol to form mathematical phrases. Mathematical phrase consists of operational symbols like +, -, () or x, \div or /. There is/are corresponding word(s) for these symbols. Aside from operational symbols, it also includes variables and numbers.



Learning Task 1: Identify the mathematical operation that corresponds to the following words. Do this in a separate sheet of paper.

Mathematical term	Mathematical symbol/operations	Mathematical term	Mathematical symbol/ operations
1. sum		6. product	
2. increased by		7. quotient	
3. difference		8. times	
4. decreased by	W 11	9. divided by	
5. less than		10. more than	

In mathematics you cannot do away with symbols. Mathematical word problems cannot be solved unless you translate it into symbols. Symbols used for operations are called **operational symbols** while symbols used to determine relation between quantities are called **relational symbols**.

Operational symbols	Words associated to symbols	Operational symbols	Words related to symbols
Plus sign (+)	Plus, add, increased by, more than, sum of	Division sign ÷, / or bar	Divide, quotient, ratio
Minus sign (-)	Minus, subtract from, decreases by, dimin- ished, difference of, less, less than	Involution (exponents)	Raised to the power of, squared, cube,
Multiplication sign {(), •}	Multiply, multiplied by, the product of, times (no symbol be- tween variables means multiplication	Evolution Radical sign	Roots, where the integer

In grammar, a phrase is a group of words that does not express complete thoughts. Mathematical phrase does not express a complete thought also unless it becomes an equation.

Examples: Translate the following to mathematical symbols:

1. Thrice the sum of 5 and a number.

Solution: let m be the number

Sum of 5 and number (5 + m)

Thrice means 3 times

The mathematical phrase is 3(5 + m)

2. Subtract two thirds of the number from thirty

Solution: let x be the number

Two thirds of the number - the word "of" means multiply. $\frac{2}{3}x$

The mathematical phrase is $30 - \frac{2}{3}x$

3. One half the square root of twice the square of the number.

Solution: Let y be the number. Twice the square of the number: 2y2

Square root of $2y^2$: $\sqrt{2y^2}$

The mathematical phrase is $\frac{\sqrt{2y^2}}{2}$

4. Translate 2x - 5 to verbal phrase.

Solution: 2x means twice a number

Possible verbal phrase: Twice a number decreased by 5 or subtract 5 from twice a number.

In the above examples the mathematical phrases are called **algebraic expressions.** Algebraic expression can be a number, a single variable or a combination of letters, numbers and operational symbols.

Examples:

- A. 3, m, -5x, 3y, 7xy are algebraic expressions with one term
- B. $4x^2 8xy$, xy + 5 are algebraic expressions with two terms.

Terms in algebraic expression are separated by plus (+) or minus (-) signs. When the operations between variables or variable and number is multiplication or division, it is considered as one term only. 4xyz is a single

term algebraic expression so as $\frac{24x}{ab}$ since the operations involve are multiplication and division

Algebraic expressions are named according to number of terms.

- A. Monomial is an algebraic expression with one term
- B. Binomial is an algebraic expression with two terms
- C. Trinomial is an algebraic expression wit 3 terms
- D. Multinomial or polynomial is an algebraic expression with more than 3 terms.

The **degree** of the algebraic expressions is the highest exponent of an expression with one variable or the highest sum of the exponents of the variables in a term of the expressions. The **constant** of the algebraic expressions is a number with fixed value. A **variable** is a letter which represents a number.

- 1. $5x^4$ 3x + 12. The expression is trinomial, the degree is 4 which is the highest exponent, the variable is x and the constant is 12.
- 2. $3x^2yz + x^3yz^2 2xyz + 4xy^2z$. This is a multinomial with variables xyz. The highest sum of the exponent is 6 which is on the second term, hence the degree is 6 and there is no constant in this expression.



Learning Task 2

A. Match the verbal phrase in column A with mathematical phrase in column B. Write the letter of the correct answer.

COLUMN A	COLUMN B
1. The difference between a number and 5	A. $(m + 2)^2 + 5$
2. Five times the sum of a number and 6	B. 4(p - 6)
3. Divide the sum of the squares of \boldsymbol{a} and \boldsymbol{b} by the square of c	C. m - 5
4. The square of the sum of a number and 2 increased by 5.	D. 5(a + 6)
5. The difference of a number and 6 multiplied by 4	$E. \frac{a^2 + b^2}{c^2}$

B. Identify the kind of algebraic expression and determine the degree, variables and constant .

Algebraic Expression	Type/kind	Variables	Degree	Constant
1. $7x + 4x^3 - 17$				
2. $3abc^2 + a^2bc^2 - abc + 2$				
3. $x + 2x^2 - 6x^3 + 9x^4 + 1$				
4. 3xyz ² + 12				
5. 14				



Express the following verbal phrases into mathematical phrases. Write your answers in your answer sheet.

- 1. Zab is x years old now. What is his age 7 years from now?
- 2. Joan is twice as old as her sister now. What was her age 6 years ago?
- 3. A square has a side 3x 2, what is the perimeter?
- 4. Zab's income is 7x + 6 every day. What is his total income in x days?

Addition and Subtraction of Polynomials



Lesson

Algebraic expressions are said to be similar if the expressions has the same literal coefficients or variables, like 3x and x are similar while x and x^2 are not.

In general, algebraic expressions are called polynomials. You can only perform addition and subtraction to polynomials that are similar.

In the previous lesson you define variables as a letter that represents a number. An expression can have a value if you replace the variable with a number and perform the operations involved.

The terms in the expression 3x + 5 are not similar, hence you cannot add them. However if you give value to x, then you can perform the operations.

Example:

1. Find the value of 3x + 5, if x = 2

Solution: 3x + 5 = 3(2) + 5 = 6 + 5 = 11.

The value of the expression 3x + 5 is 11 when x = 2.

2. Find the value of $\frac{6ab-c}{abc}$ if a = 5; b = 3; c = 10

Solution: $\frac{6ab-c}{abc} = \frac{6(5)(3)-10}{5(3)(10)} = \frac{80}{150} = \frac{8}{15}$

Rules in Adding or Subtracting Polynomials

- Step 1. Arrange the polynomial in descending or ascending order if possible.
- Step 2. Group the terms which are similar.
- Step 3. Add/subtract the numerical coefficients, applying the rules in adding and subtracting integers and copy the literal coefficients.

Examples:

1. Add: $4x^2 - 2x + 5$ and $7 + 5x^3 + 4x - x^2$ $5x^3 + (4 - 1)x^2 + (-2 + 4)x + (5 + 7) = 5x^3 + 3x^2 + 4x + 12$



Learning Task 1: Evaluating Algebraic Expression

Find the value of the expression given the value of the variables

1.
$$3x + 5$$
; $x = 2$

4.
$$3(xy + 6)$$
; $x = -6$ and $y - 1$

2.
$$5xy+x-4$$
, $x = -3$ and $y = 5$

5.
$$\frac{2x+5}{3x-y}$$
 x = -3 and y = -8

3.
$$\frac{6ab-c}{abc}$$
; $a = 5$; $b = 3$; $c = 10$

Solution:

Step 1.
$$4x^2 - 2x + 5$$
 and $5x^3 - x^2 + 4x + 7$

Step 2.
$$5x^3 + (4x^2 - x^2) + (-2x + 4x) + (5 + 7)$$

Step 3.
$$5x^3 + (4 - 1)x^2 + (-2 + 4)x + 12$$

$$5x^3 + 3x^2 + 2x + 12$$

You can also solve it using the vertical method.

All similar terms must be in one column.

$$4x^{2} - 2x + 5$$

$$+ 5x^{3} - x^{2} + 4x + 7$$

$$5x^{3} + 3x^{2} + 2x + 12$$

2. Subtract 4xy + 5x - 7y + 3 from xy - 6x + y + 6

Grouping Similar Terms

$$(xy - 6x + y + 6) - (4xy + 5x - 7y + 3)$$

Change the sign of the subtrahend

$$xy - 6x + y + 6 - 4xy - 5x + 7y - 3$$

Group similar terms and perform the operation.

$$(xy - 4xy) + (-6x - 5x) + (y + 7y) + (6 - 3)$$

-3xy + (-11x) + 8y + 3
-3xy - 11x + 8y + 3

Vertical Method

Similar terms must be placed in one column, change the sign of the subtrahend then add.

$$xy - 6x + y + 6$$

$$(-)4xy^{(+)}5x^{(+)}7y^{(+)}3$$

$$- 3xy - 11x + 8y + 3$$

Note: Terms without numerical coefficient written beside the variable, it is understood the numerical coefficient is 1, like xy, x, etc.

3. Add:
$$5a^5 - 6a^2 + 8$$
; $3a^3 + a^2 - 8a$; $4a^4 + a^3 - 6a^2 + a - 10$

Align similar terms vertically, you may replace the term with the missing exponent with zero as its numerical coefficient or just provide a space.

 $5a^5 + 0a^4 + 0a^3 - 6a^2 + 0a + 8$ (the terms with missing exponents are 4,3 and 1) + $3a^3 + a^2 - 8a$

$$4a^4 + a^3 - 6a^2 + a - 10$$

$$5a^5 + 4a^4 + 4a^3 - 11a^2 - 7a - 2$$

4. Subtract $3x^3 + 8x - 5$ from the sum of x + 4 and $x^2 - 3$

Add first
$$(x + 4) + (x^2 - 3)$$
 Subtract: $(x^2 + x - 7) - (3x^3 + 8x - 5)$
 $(x^2 + x - 4 - 3)$ $x^2 + x - 7 - 3x^3 - 8x + 5$
 $(x^2 + x - 7)$ $-3x^3 + x^2 + (x - 8x) + (-7 - 5)$
 $-3x^3 + x^2 - 7x - 12$



Learning Task 2:

A. Evaluate the algebraic expressions

Let $a = 4$, $b = -5$, $c = 0.6$, $x = -3$, $y = 0.4$; $z = \frac{1}{2}$			
Algebraic Expression	Value	Algebraic Expression	Value
1. 3abc + bc - a		6. $3x^2 + 2xy - 3(ab)^2$	_%
2. bxy + 5ab - ay		7. $25 - 2x^2y + 3a^2b - az^2$	
3. abz - 8z + b		8. $(az)^2 + 5x^2y - 4bc$	
4. ax + by - cz	33	9. 2x ² y ^{2 -} 12abz - 16	
5. 4ax ÷ cz		10. $8z^3 - 4x + y^2$	

B. Perform the indicated operations

Add	Subtract:
1. $(5x^4 - 3x^2 + 4) + (6x^3 - 4x^2 - 7)$	6. $(5x^3 - 7x^2 + 3x - 4) - (8x^3 + 2x^2 + 3x - 7)$
2. $-7x^3y + 4x^2y^2 - 2$ and $4x^3y + 1 - 8x^2y^2$	7. $(2x^2y - 5xy + 3y^2) - (7xy - 6y^2 + 5x^2y)$
3. $(5 + 24y^3 - 7y^2) + (-6y^3 + 7y^2 + 5)$	8. $(9x^5 - 6x^3 + 7x^2) - (7x^3 - 6x^5 + 2x^2)$
4. $(2x^5 - 6x^3 - 12x^2 - 4) + (-11x^5 + 8x + 2x^2 + 6)$	9. Subtract $4x^3 - 5x - 8$ from $6x^2 - 3x + 8$
5. $(3y^5 - 2y + y^4 + 2y^3 + 5)$ and $(2y^5 + 3y^3 + 2 + 7)$	10. $(1.5y^3 + 4.8y^2 + 12) - (y^3 - 1.7y^2 + 2y)$



Solve the following. Do this in a separate sheet of paper.

- 1. A box is (2x-3) by (x + 5) by (3x + 1), what is the volume of the box if x = 3 cm?
- 2. The formula for the area of a triangle is $A = \frac{bh}{2}$ If the base (b) = 10 cm and the height (h) = 6 cm, what is the area of the triangle?
- 3. The length (l) of the rectangle is $x^2 + 2x 3$ and the width (w) is 5x + 4, what is the perimeter of the rectangle?
- 4. From the sum of $3x^3 + 7x^2 5$ and $2x^2 + 3x + 8$ take away $5x^2 + x 5$.
- 5. What should be added to $3x^3 + 4x^2 7$ to have a sum of $4x^3 + x^2 + 5$?

Laws of Exponents and Its Application

Lesson

In the process of multiplication and division of polynomials you need to apply the laws of exponents. Remember that exponent tells how many times the base will be multiplied by itself. Any number $\boldsymbol{a^n}$, a is the base and n is the exponent. 5^3 means $5 \cdot 5 \cdot 5 = 125$ In $x \cdot x \cdot x \cdot x \cdot x$, the variable x is multiplied by itself 5 times, hence you can write it as x^5 . Here, the base is x and the exponent is 5



Simplify by writing in exponential form	Write in expanded form	
1. $2(2)(2)(2)(2) = 2^6$	$(2x)^4 = (2x)(2x)(2x)(2x)$	
2. (3a)(3a)(3a) =	$(xy)^7 =$	
3. (-4)(-4)(-4)(-4) =	(-ab) ⁶ =	

Exponents have its own rules in performing mathematical operations.

The Laws of Exponents

1. Product Law of Exponent.

Any numbers m and integers x and y,

The base must be the same before you nents.

$$m^x \cdot m^y = m^{x+y}$$
 can add the expo-

Example:

Similarly: $b^2 \cdot b^3 = (b \cdot b)(b \cdot b \cdot b) = b^5$

2. Quotient $b^2 \cdot b^3 = b^{2+3} = b^5$ Law of Exponents

Any numbers m and integers x and y,

a. b
$$\frac{m^x}{m^y} = m^{x-y}$$
, if $x > y$ $\frac{m^x}{m^y} = m^{x-y} = 1$, if $x = y$ $\frac{m^x}{m^y} = \frac{1}{m^{y-x}}$, if $x < y$ Examples:

$$a. \quad \frac{a^5}{a^2} = \frac{a \cdot a \cdot a \cdot a \cdot a}{a \cdot a} = a \cdot a \cdot a = a^3 \text{ similarly } \frac{a^5}{a^2} = a^{5-2} = a^3$$

b.
$$\frac{a^3}{a^3} = \frac{a \cdot a \cdot a}{a \cdot a \cdot a} = 1$$
 similarly $\frac{a^3}{a^3} = a^{3-3} = a^0 = 1$

c. $\frac{a^2}{a^4} = \frac{a \cdot a}{a \cdot a \cdot a \cdot a} = \frac{1}{a^2} \quad similarly \quad \frac{a^2}{a^4} = \frac{1}{a^{4-2}} = \frac{1}{a^2}$

Any number raised to zero is always equal to 1 except 0.

3. Power of a Power Rule

Any numbers m and with exponents of integers x and y, $(m^x)^y = m^{xy}$

Example: a.
$$(a^2)^3 = a^2 \cdot a^2 \cdot a^2 = a^{2+2+2} = a^6$$
 or $(a^2)^3 = a^{2(3)} = a^6$
b $\left(\frac{a^2}{b^3}\right)^2 = \left(\frac{a^{2(2)}}{b^{3(2)}}\right) = \frac{a^4}{b^6}$

4. Power of a Product

Any numbers m and n integers x, $(mn)^x = m^x n^x$

Example:
$$(2ab)^3 = 2^3a^3b^3 = 8a^3b^3$$

5. Power of a Quotient

Any numbers m and n integers x, $\left(\frac{m}{n}\right)^x = \frac{m^x}{n^x}$

Example:
$$\left(\frac{3}{2}\right)^3 = \frac{3^3}{2^3} = \frac{27}{8}$$

Multiplication and Division of Polynomials

You can apply the laws of exponents in multiplying or dividing polynomials.

Examples:

1. Multiply 3xy and 4xz.

Solution: $3xy(4xz) = 3(4)(x)(x)(y)(z) = 12x^2yz$, multiply the numerical coefficients, add only the exponents of the same variable and copy the rest of the variables

2. Find the product of 3abc and $(4a^2 + 3a - 5)$.

Solution: Use Distributive Property

$$3abc(4a^2 + 3a - 5) = 3abc(4a^2) + 3abc(3a) + 3abc(-5)$$

= $12a^3bc + 9a^2bc - 15abc$

3. What is the product of $(m - 4)(m^2 + 3m + 7)$?

Solution:

Distributive Method (Horizontal Form)	Vertical Form	
(m - 4)(2m ² + 2m - 3)	2m ² + 2m - 3	
$m(2m^2 + 2m - 3) - 4(2m^2 + 2m - 3))$	m - 4	
$2m^3 + 2m^2 - 3m - 8m^2 - 8m + 12$	$2m^3 + 2m^2 - 3m$ Partial	
$2m^3 + (2m^2 - 8m^2) + (-3m - 8m) + 12$	+ $-8m^2 - 8m + 12$ products	
2m ³ - 6m ² - 11m + 12	2m ³ - 6m ² - 11m +12	

4. Find the quotient : $25x^5y^2z^3 \div 5xy^3z^2$

Solution:

$$\frac{25x^5y^2z^3}{5xy^3z^2} = \frac{5x^{5-1}z^{3-2}}{y^{3-2}}$$
 Divide the numerical coefficients, then apply the quotient law of exponents

5. What is the quotient : $\frac{4x^8 + 6x^6 - 2x^4 - 10x^2}{2x^2}$?

Solution:
$$\frac{4x^8 + 6x^6 - 2x^4 - 10x^2}{2x^2} = \frac{4x^8}{2x^2} + \frac{6x^6}{2x^2} - \frac{2x^4}{2x^2} - \frac{10x^2}{2x^2} = 2x^6 + 3x^4 - x^2 - 5$$

In dividing polynomial by another polynomial you have to apply the following steps:

- 1. Arrange both dividend and divisor in descending or ascending powers of common variables, leaving a space or replacing the missing term with zero.
- 2. Divide the first term of the dividend with the first term of the divisor to get the first term of the quotient.
- 3. Multiply the entire divisor by term of the quotient and write the product under the dividend of similar terms.
- 4. Subtract similar terms. Apply rules in subtracting polynomials
- 5. Bring down the next terms of the dividend.
- 6. Divide the first term of the difference by the first term of the divisor to get the second term of the quotient.
- 7. Repeat steps 3 6 until the result of subtraction is zero or the degree of the term of the difference is less than the degree of the divisor.

Example: Divide $15n^2 - 2n - 24$ by 3n - 4.

The quotient is 5n + 6.

To check if the answer is correct, multiply the divisor and the quotient to get the dividend. $15n^2 - 2n - 24 = (3n - 4)(5n + 6)$

$$= 3n(5n + 6) - 4(5n + 6) = 15n^{2} + 18n - 20n - 24$$

$$= 15n^{2} - 2n - 24$$

which is 6, then

subtract.

2. Divide $(6x^3 - 3x^2 + 2z + 3) \div (x - 2)$

$$6x^2 + 9x + 20$$

$$(x-2)6x^3-3x^2+2x+3$$
 Divide $6x^3$ by x, to get the quotient $6x^2$

$$6x^3 - 12x^2$$

The product of multiplying (x-2) by $6x^2$

$$9x^{2} + 2x + 3
9x^{2} - 18x
20x + 3$$

Subtract then divide $9x^2$ by x to get the quotient 9xMultiply (x—2) by 9x

 $\frac{20x + 3}{20x + 3}$ Subtract then divide 20x by x to obtain the quotient 20 Multiply x-2 by 20

, Subtract. 43 cannot be divided by x. Hence 43 is the re-

The quotient is $6x^2 + 9x + 20$ R. 43 or $6x^2 + 9x + 20 + \frac{43}{x-2}$

Learning Task 1:

- A. Simplify
- B. Perform the indicated operations. Simplify your answer
- 1. $(-m^4)^3$

E

- 1. $2ab(4a^2 + 3ab 7b^2)$
- 7. $\frac{16a^3b^2c^5 24a^5bc^4 + 44a^7b^6c^6}{4a^2bc^3}$

- $2. \quad \left(\frac{10c^2}{2d^3}\right)^2$
- 2. $-3xy(x^3y^3 3x^2y + 7xy^2)$
- 8. $\frac{30c^2 + 19ac 63a^2}{6c 7a}$

- 3. $(3a^2bc^3)^3$ 4. $(2x-1)(2x^2+4x+3)$

3. $(x + 2)(x^2 - 3x + 2)$

- 4. $(2x^3y^2z)(3x^2yz^2)$ 5. $(12y^2-9y+16)(8y^3-14y+5)$ 9. $(a^6+b^6) \div (a^2+b^2)$

- 5. $\left(\frac{3m^2n^3}{xy^2}\right)^2 \left(\frac{x^3yz^2}{2mn^2}\right)^3$ 6. $\frac{25a^5b^3c}{5ab^4c}$

 $10. \ \frac{6w^3 + 7w^2 - 12w + 15}{2w^2 + 3w - 5}$

Solve the following problems:

- 1. What is the area of the rectangle whose length is (x + 5) and width (x 5)?
- 2. What is the area of the square whose sides measure (3x + 4)?
- 3. The area of the rectangle is $3x^2 + 7x 6$, what is the length if the width is (x + 3)?
- 4. What is the average speed of the car that covers a distance of $(2y^3-7y^2 + 5y - 1)$ km in (2y-1) hour?
- 5. Multiply $(m^2 + 2m-2)$ by the sum of (m + 3) and (2m 3).

Special Products



Lesson

There are binomials or trinomials that when you multiply the products form a pattern. Such are called **special products.** Using the laws of exponents you will be able to find the product of a) square of a binomial b) sum and difference of binomials, c) cube of a binomial d) square of a trinomial.

Did you find a pattern in the product when you multiply the binomials or trinomials in the Learning Task Number 1? These are special types of polynomials.

A. Square of a Binomial

$$(a + b)^2 = a^2 + 2ab + b^2$$

1st 2nd Square of Twice the prod-
term term the 1st uct of the first 2nd term
term and the 2nd

Using the pattern above for $(a - b)^2 = (a)^2 + 2(a)(-b) + (-b)^2 = a^2 - 2ab + b^2$

Square of Binomial
$$(a + b)^2 = a^2 + 2ab + b^2$$
 $(a - b)^2 = a^2 - 2ab + b^2$

Examples:

1.
$$(2x + 3)^2 = (2x)^2 + 2(2x)(3) + 3^2 = 4x^2 + 12x + 9$$

2.
$$(x - 2y)^2 = (x)^2 + 2(x)(-2y) + (-2y)^2 = x^2 - 4xy + 4y^2$$

B. Square of a Trinomial

$$(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$$

$$1st term 2nd term 3rd term Square of Square of Square of Square of Twice the Twice the the 1st the 2nd the 3rd 1st and 1st and 2nd and term term term 2nd term 3rd term 3rd term 3rd term 3rd term 3rd term$$

Square of Trinomial:
$$(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$$

Example:
$$(2x + y - 3z)^2 = (2x)^2 + y^2 + (-3z)^2 + 2(2x)(y) + 2(2x)(-3z) + 2(y)(-3z)$$

= $4x^2 + y^2 + 9z^2 + 4xy - 12xz - 6yz$

C. Sum and Difference of two Binomials

$$(a + b) (a - b) = a(a) + a (-b) + a(b) + (b)(-b)$$

Product Product Product Product of inner the 2nd

1st terms terms terms terms

Sum and Difference of two Binomials:

The product of the sum and difference of binomial is the product of the first terms plus product of the second terms.

$$(a + b)(a - b) = (a)(a) + (b)(-b) = a^2 - b^2$$

Examples:
$$(x + 2y)(x - 2y) = (x)^2 - (2y)^2 = x^2 - 4y^2$$

 $(2a^2 - 3b)(2a^2 + 3b) = (2a^2)^2 - (3b)^2 = 4a^4 - 9b^2$
 $[x + (3a - 2)][x - (3a - 2)] = (x)^2 - (3a - 2)^2$
 $= x^2 - [(3a)^2 - 2(3a)(2) + 2^2]$
 $= x^2 - (9a^2 - 12a + 4)$
 $= x^2 - 9a^2 + 12a - 4$

D. Cube of a Binomial

(a - b)
3
 = (a) 3 + 3(a) 2 (-b) + 3(a)(-b) 2 + (-b) 3 = a^3 - $3a^2b$ + $3ab^2$ - b^3 Examples : (2x + 3y) 3 = (2x) 3 + 3(2x) 2 (3y) + 3(2x)(3y) 2 + (3y) 3 = $8x^3$ + $36x^2y$ + $54xy^2$ + $27y^3$ (3a - 2) = (3a) 3 - 3(3a) 2 (2) + 3(3a)(2) 2 - (2) 3 = $27a^3$ - $54a^2$ + $36a$ - 8

Cube of a Binomial:
$$(a + b)^3 = a^3 + 3a^2 b + 3ab^2 + b^3$$

 $(a - b)^3 = a^3 - 3a^2 b + 3ab^2 - b^3$



Learning Task 1: Find the product. Do this in your separate sheet of paper.

1. $(a + b)^2$

4. $(a + b)^3$

2. $(a - b)^2$

5. $(a - b)^3$

3. (a + b)(a - b)

6. $(a + b + c)^2$



Learning Task 2: Find the product. Do this in a separate sheet of paper.

- 1. $(a + 5)^2$
- 6. (4 2x)(4 + 2x)

11. $(2 + 2x)^3$

- 2. $(3xy 7)^2$
- 7. (3xy abc)(3xy + abc)
- 12. $(x^2 3)^3$

- 3. $[x + (y 2)]^2$
- 8. $(x^2 + 4y^2)(x^2 4y^2)$
- 13. $(2x + 5)^3$


- 4. $(2x + 3y + 5)^2$
- 9. [2 + (x-1)][2 (x-1)]
- 14. $[(x + 1) y]^3$

- 5. $(3a b + 2c)^2$
- 10. [(x+2) (y+1)][(x+2) + (y+1)]
- 15. $[x + (y 2)]^3$



Given the square figure at the right, find the following:

- 1. Area of Fig. 1
- 2. Area of Fig. 2
- 3. Area of Fig. 3
- 4. Area of Fig. 4
- 5. Area of the whole figure



WEEK 7

Linear Equations and Inequalities in One Variable

Lesson

In the previous lesson, you translate verbal phrases to mathematical symbols which become an algebraic expression. In grammar, a phrase does not express a complete thought only sentence does. Similarly, in mathematics, a mathematical phrase or expression does not express a complete thought, only mathematical sentence does.

Mathematical sentence makes use of relational symbol like equal (=), less than (<), greater than (>), less than or equal to (\leq) , and greater than or equal to (\geq) . If the mathematical sentence uses equal sign (=), then it is an equation, other wise it is an inequality.

Word problems can be solved if you will translate it into an equation or inequality. An **equation** is a mathematical sentence that shows two equal expressions. 3x + 5 = 20 is an equation. The left side of equation 3x + 5 must be equal to the right side of the equation which is 20. 3x + 5 = 20 is an example of a linear equation in one variable. A **linear equation** is in the form of ax + b = c, where a, b and c are real numbers and $a \ne 0$.

An equation may either be true or false.

8 + 2 = 10 is a true equation since 8 + 2 is really 10.

2x + 5 = 15 is an equation that can either be true or false, depending on the value of x. There is only one value of x that will make the equation true.

An inequality is a mathematical sentence that shows two unequal quantities. One might be greater or lesser than the other.

5 > 1 is a true inequality.

2x + 1 < 10, may be true or false, depending on the values of x. More than one value of x can make the inequality true.

You have to interpret or translate word problem correctly to be able to solve equation or inequality.

Examples.

1. The sum of three consecutive integers is 57.

Consecutive means that the next number must 1 greater than the previous.

If x represent the number, the next number is x + 1, this is one greater than x, the third number is (x + 1) + 1, also 1 greater than the previous. When simplified it is x + 2.

Hence the 3 consecutive integers are represented by: x, x + 1 and x + 2

The equation is: x + (x + 1) + (x + 2) = 57

2. Angel's age is twice the age of Gideon 5 years ago. The sum of their ages now is 35

You can represent the ages in tabular form.

	Age now	Age 5 year ago
Gideon	x	(x - 5)
Angel	2(x - 5) = 2x - 10	2x - 15

Equation: x + 2x - 10 = 35

3. EJ drives her car 50kph. After 30 minutes her father followed her driving an other car at 60 kph. How long will it take for the father to be at the side of her daughter?

Remember that the formula for distance is d = r t, where r is the rate or speed and t is the time.

	Rate (r)	Time (t)	Distance (d)
EJ	50	t	50t
Father	60	$t-\frac{1}{2}$	60($t-\frac{1}{2}$)

Note: 30 minutes is one half hour.

There is a point where EJ and the father will be beside each other before he can overtake her. Therefore they cover the same distance.

Equation: $d_1 = d_2$

50t
$$(t-\frac{1}{2})$$
 =60

4. The difference between six times the number and eight is greater than 15.

Let x be the number

Six times the number: 6x

The difference between 6 times the number and 8: 6x - 8

Greater than means that the problem is an inequality problem

Inequality: 6x - 8 > 15

5. Ten divided by twice the number is less than one-fourth.

Let x be the number; twice the number: 2x

Ten divided by twice the number: $\frac{10}{2x}$

Inequality: $\frac{10}{2x} < \frac{1}{4}$

6. Choi's monthly commission plus monthly salary of 15,000.00 is at most 20,000.00

Choi's commission is x; at most means it can be equal to or less than

Inequality: $x + 15,000 \le 20,000.00$

7. Elisse's score in 3 of the 4 quizzes in math are 80, 78, 90. What score must she get on the 4th quiz to have an average of at least 85.

Let x be the 4th score; average score is to add the 4 scores divided by 4 At least means equal to or greater than.

Inequality:

$$\frac{80 + 78 + 90 + x}{4} \ge 85$$



Learning Task 1: Translate the verbal sentence to mathematical sentence. Use variable \mathbf{x} to represent number. Do this in a separate sheet of paper.

Verbal Sentence	Mathematical sentence
1. The sum of a number and 5 is 12.	
2. Twice a number is 54	
3. Thrice a number increase by 2 is less than 50.	
4. One half the sum of a number and 4 is less than 28.	
5. A number increased by one-third of the number is 5.	

Learning Task 2:

A. Translate the following sentences into equation. Use the x to represent the unknown number.

- 1. The perimeter of the rectangle is 96 when the length of a rectangle is twice the width.
- 2. The perimeter of an equilateral (equal sides) triangle is 24.
- 3. Two-thirds of the number is 72.
- 4. The sum of two consecutive integers is 29.
- 5. Four times a number increased by 8 is 54.
- B. Translate the following sentence to inequality. Use y as the variable to represent a number.
- 1. The discount is not less than 100.00.
- 2. The costs of the book is at most 300.00.
- 3. His monthly income is at least 20,000.00.
- 4. Twice the number is at least 80.
- 5. Twice the measure of an acute angle is not more than 90 degrees.

Learning Task 3: Write the equation or inequality described in the following problems.

1. JP's age is twice the age of Reyna. The sum of their ages does not exceed 51.

2. Twice the sum of a number and 10 is 55.

3. Cut a 60 cm ribbon into two such that one part is one-third of the other.

4. The sum of two consecutive even integers is 162.

5. The sum of three consecutive integers is not more than 57.

6. Nine subtracted from the quotient of a number and three is twenty-one.

7. If seven is subtracted from six times a number, the result is at least 10.

8. The sum of 6 times a number and fifteen is no more than forty-two.

9. Five times the measure of an angle is an acute angle.

10. The circumference of a circle with radius y is at most 25.



A. Find the value of x that will make the equality true or correct.

1. x + 2 = 8

2. 3x = 12

3. 2x - 3 = 1

4. 3(x + 5) = 18

5. 5x + 8 = -7

B. Find 3 values of x to make the inequality true or correct

1. x < -1

2. 2x > 0

3. x + 2 < 7

4. x - 4 > -1

5. 3x + 5 < -1

Solving Linear Equations and Inequality in One Variable



Lesson

Solving equation or inequality is finding the value or values of the variable that will satisfy the equation or inequality.

The equation x + 5 = 12 is a mathematical sentence that is a conditional equation because it can either be true or false. It can only be true if x = 7. By substituting 7 to x, you have 7 + 5 = 12. Both sides of the equation name the same number which is 12. Thus 7 is called the solution of the equation or sometimes it is called the root of the equation.

The inequality x + 5 < 12 can either be true or false too. The value of x that will make the inequality true are numbers that are less than 7. If you replace x with 6 which is less than 7, then 6 + 5 < 12. 11 is really less than 12. You can substitute any number to x as long as it is less than 7.

You have to consider properties of equality or inequality before you can solve an equation or inequality.

For any real numbers a, b and c

Properties of Equality

Addition Property of Equality (APE)

If a = b, then a + c = b + c

Multiplication Property of Equality

(MPE).

If a = b, then ac = bc

Reflexive Property of Equality

Any number or expression is equal to it self. 3 = 3; m = m; x = x

Symmetric Property of Equality (SPE)

The sides of the equation can be interchanged. If ab = cd, then cd = ab.

Transitive Property of Equality (TPE)

If two equations are equal to a third quantity, then they are all equal to each other. If a = b and b = c, then a = c

Properties of Inequality

Axiom of comparison

a < b a = b a > b

Transitive Property of Inequality

If a < b and b < c, then a < c

Addition property of Inequality (API)

If a < b then a = c < b + c

Multiplication Property of Inequality

(MPI)

If a < b and c > 0, then ac < bc and $\frac{a}{c} < \frac{b}{c}$

If a < b and c < 0, then ac > bc and $\frac{a}{c} > \frac{b}{c}$

If a > b and c > 0, then ac > bc and $\frac{a}{c} > \frac{b}{c}$

If a > b and c < 0, then ac < bc and

APE	If $x + 5 = 4$, then $x + 5 + (-2) = 4 + (-2)$ or $x + 3 = 2$
MPE	If $x = 10$, then $2(x) = 2(10)$ or $2x = 20$
API	If $x - 2 < 6$, then $x - 2 + 2 = 6 + 2$ or $x < 8$
MPI	If $-x < 3$, then $(-1)(-x) > (-1)(3)$ or $x > -3$

Linear Equation in One Variable

To solve equations, you apply the properties of equality and follow some steps.

- 1. Simplify both sides of the equation. This includes applying Distributive Property of Multiplication over Addition (DPMA) and/ or combining similar terms.
- 2. If the equation has a fraction clear the denominators by multiplying both sides of the equation by the LCD of all the denominators.
- 3. If equation has decimal, clear the decimal by multiplying every term of the equation by powers of 10 depending on the greatest number of decimal places in the term has.
- 4. If a constant is added to a term with a variable, add its opposite to both sides of the equation or apply APE.
- 5. If a variable has a numerical coefficient other than 1, multiply both sides by the reciprocal of the numerical coefficient or divide both sides of the equation by the numerical coefficient or by MPE
- 6. Substitute the solved value to the variable in the equation to check if you get the right answer.

Examples: Solve for the value of x.

1. 2x + 5 = 21

Solution. Since the equation is already in simplified form, the go to step 4.

By APE: 2x + 5 + (-5) = 21 + (-5) (the opposite of 5 is negative 5)

2x = 16 (simplify)

By MPE: $\frac{1}{2}(2x) = \frac{1}{2}(16)$ (the reciprocal of 2 is 1/2.

Simplify: x = 8 the solution of the equation.

Check: $2x + 5 = 21 \longrightarrow 2(8) + 5 = 21 \longrightarrow 16 + 5 = 21 \longrightarrow 21 = 21$

2. The sum of three consecutive odd integers is 81. Find the numbers.

Solution: The difference between any two consecutive odd integers is 2.

If **x** is the first odd integer, then **x** + 2 is the second odd integer and the third is (x + 2) + 2 or x + 4

Since the sum of these numbers is 82, then the equation is:

$$x + (x + 2) + (x + 4) = 81$$

 $3x + 6 = 81$ (combine similar terms)
 $3x + 6 - 6 = 81 - 6$ (by APE)
 $3x = 75$
 $x = 25$ (by MPE, multiplying both sides by 1/3 or dividing both sides by 3)

The first number is x = 25

Second number is x + 2 therefore 25 + 2 = 27

Third number is x + 4, therefore 25 + 4 = 29

Check if the sum of these odd numbers is 81: 25 + 27 + 29 = 81

3. Anton is 2 more than three times older than her daughter, Zab. In 16 years , he will be twice as old as his daughter. What are their present ages?

Solution.

	Present Age	Age 16 from now
Zab	X	x + 16
Anton	3x + 2	(3x +2) + 16
151		2(x + 16)

In forming an equation, consider equating the same quantity expressed in different ways. In this case, you can equate Anton's age 16 years from now.

$$(3x + 2) + 16 = 2(x + 16)$$

 $3x + 18 = 2x + 32$ simplify both sides
 $3x + 18 - 18 = 2x + 32 - 18$ by APE
 $3x = 2x + 14$ simplify
 $3x - 2x = 2x - 2x + 14$ by APE - to combine 3x and 2x
 $x = 14$ the age of Zab now
 $3(14) + 2 = 44$ the age of Anton now

To check: If Zab's age now is 14, her age in 16 years is 14 + 16 = 30.

If Anton's age now is 44, his age in 16 years is 44 + 16 = 60

The age of Anton is twice as the age of Zab's 16 yrs from now, which is the condition in the problem.

Linear Inequality in One Variable

To solve inequality you have to use the properties of inequality and follow the same steps in solving for the equation.

Examples: Solve for x

1.
$$\frac{x}{3} + 4 > 5$$

Solution: Clear fraction by multiplying each term in the inequality by the de nominator which is 3.

$$\frac{x}{3} + 4 > 5 \longrightarrow (3)\frac{x}{3} + 4(3) > 5(3) \longrightarrow x + 12 > 15$$

By API
$$x + 12 - 12 > 15 - 12$$

 $x > 3$

The solutions of the inequality are all numbers that are greater than 3.

The solution of the inequality can be written in different ways:

- (a) set notation: $\{x/x > 3\}$ read as set of all x such that x is greater than 3
- (b) interval notation $(3,+\infty)$ the symbol ∞ means infinity.

Take note that the point that corresponds to 3 is not solid, this means that 3 is not included in the solution set of the inequality.

The point must be solid if the number corresponding to that point is included as one of the solutions of the inequality. For interval notation, a bracket is used if the number is a part of the solution set.

Suppose the inequality is $\frac{x}{3} + 4 \ge 5$, the inequality is equal or greater than.

The solution is $x \ge 3$

- (a) by set notation $\{x/x \ge 3\}$
- (b) By interval notation: $[3, +\infty)$, the bracket is used to indicate that 3 is included.
- (c) Graphical method: -1 0 1 2 3 4 5 6

The point that corresponds to 3 is solid, which means that 3 is included.

2. Find the value of x that satisfy both inequalities -3 < 2x + 1 and $2x + 1 \le 9$.

Solution: You can write the inequality as

$$-3 < 2x + 1 \le 9$$

 $-3 - 1 < 2x + 1 - 1 \le 9 - 1$ by API
 $-4 < 2x \le 8$ Simplify
 $-2 < x \le 4$ by MPI

The solution set are numbers between - 2 and 4 including 4.

Set notation: $\{x/-2 < x \le 4\}$

Interval notation: (-2, 4], parenthesis means that -2 is not included in the solution set while bracket means that 4 is included in the solution set.

Graph: -4 -3 -2 -1 0 1 2 3 4 5

The point that corresponds to -2 is not solid while the point that corresponds to 4 is solid.

3. The length of the rectangle is 4 more than twice the width. If the perimeter is not more than 80 cm, find the possible dimensions of the rectangle.

Solution: Let x be the width

2x + 4 is the length (4 more than twice the width)

P = 21 + 2w, however in the problem it must not be more than 80.

 $21 + 2w \le 80$, the perimeter can be 80 but not more than

$$2(2x + 4) + 2(x) \le 80$$

$$4x + 8 + 2x \le 80$$

$$6x + 8 \le 80$$
 simplify

$$6x + 8 - 8 \le 80 - 8$$
 by API

$$6x \le 72$$

$$x \le 12$$
 by MPI

The possible width is $0 < width \le 12$, since there is no negative measurement.

Since 1 = 2x + 4, if x = 0 then the length is 4, however the width must be greater than 0, therefore the length is greater than 4. The highest possible width is 12, then the length is 2(12) + 4 = 28, therefore the length can be equal or less than 28.

The possible length is $4 < length \le 28$.

Possible dimensions of rectangle are: w = 12, 1 = 28; w = 11, 1 = 26, etc.



Learning Task 1: What value or values of x that will make the equation or inequality true?

1.
$$2x = 6$$

6.
$$5x > 5$$

2.
$$2x + 2 = 8$$

7.
$$3x - 4 < 5$$

3.
$$\frac{x}{6} = 3$$

8.
$$\frac{2x}{3} > 4$$

4.
$$3(x - 2) = 0$$

9.
$$2(x + 3) < 0$$

5.
$$3 + x = 2x - 1$$

10.
$$5 + x < 2x - 1$$

Learning Task 2: Do this in a separate sheet of paper.

A. Solve for x in the equation.

1.

1.
$$5x + 12 = 2x - 3$$

2.
$$3x + 6 = -2x + 1$$

3.
$$\frac{x}{2} - 4x = 7$$

4.
$$1.4x - 3.8 = 0.4x + 6.2$$

5.
$$5(x-2) + 2x = 7(x+4) - 38$$

B. Solve for x in the inequality. Write your answer in a) set notation, b) interval notation and c) graph

1.
$$2x \le 4$$

$$2. -3 < 2x + 3 < 15$$

3.
$$5x - 4 > 6$$

4.
$$\frac{2x}{3} + \frac{1}{2} > x - 5$$

$$5. \ 0.8 + 0.6x \ge 4.2$$



Solve the word problems:

- 1. The sides of the quadrilateral are consecutive numbers. If the perimeter is 160, how long is each side?
- 2. Inzo is 10 years older than Migz. Five years ago Migs was one-third as old as Inzo. What are their present ages?
- 3. An express train travels 100 kph from station A to station B. A local train travelling at 45 kph, takes 45 minutes longer for the same trip. How far apart is station A from station B?
- 4. The sum of two consecutive even integers is less than 60 but greater than 24. Find at least 3 pairs of integers.
- 5. The perimeter of the square is less than 152 m. Find the possible length of the sides of the square.

PIVOT Assessment Card for Learners

Personal Assessment on Learner's Level of Performance

Using the symbols below, choose one which best describes your experience in working on each given task. Draw it in the column for Level of Performance (LP). Be guided by the descriptions below.





- I was able to do/perform the task without any difficulty. The task helped me in understanding the target content/lesson.



- I was able to do/perform the task. It was quite challenging but it still helped me in understanding the target content/lesson.



I was not able to do/perform the task. It was extremely difficult. I need additional enrichment activities to be able to do/perform this task.

Distribution of Learning Tasks Per Week for Quarter 2

Week 1	LP	Week 2	LP	Week 3	LP	Week 4	LP
Learning Task 1	= 11	Learning Task 1	- 1	Learning Task 1		Learning Task 1	
Learning Task 2		Learning Task 2	A	Learning Task 2		Learning Task 2	
Learning Task 3		Learning Task 3	1,2%	Learning Task 3		Learning Task 3	
Learning Task 4		Learning Task 4		Learning Task 4		Learning Task 4	
Learning Task 5	ļ	Learning Task 5		Learning Task 5		Learning Task 5	
Learning Task 6	6	Learning Task 6	- N=13 -	Learning Task 6		Learning Task 6	
Learning Task 7		Learning Task 7		Learning Task 7		Learning Task 7	
Learning Task 8		Learning Task 8		Learning Task 8		Learning Task 8	
Week 5	LP	Week 6	LP	Week 7	LP	Week 8	LP
Learning Task 1		Learning Task 1		Learning Task 1		Learning Task 1	
Learning Task 1 Learning Task 2		Learning Task 1 Learning Task 2		Learning Task 1 Learning Task 2			
		_		_		Learning Task 1	
Learning Task 2		Learning Task 2		Learning Task 2		Learning Task 1 Learning Task 2	
Learning Task 2 Learning Task 3		Learning Task 2 Learning Task 3		Learning Task 2 Learning Task 3		Learning Task 1 Learning Task 2 Learning Task 3	
Learning Task 2 Learning Task 3 Learning Task 4		Learning Task 2 Learning Task 3 Learning Task 4		Learning Task 2 Learning Task 3 Learning Task 4		Learning Task 1 Learning Task 2 Learning Task 3 Learning Task 4	
Learning Task 2 Learning Task 3 Learning Task 4 Learning Task 5		Learning Task 2 Learning Task 3 Learning Task 4 Learning Task 5		Learning Task 2 Learning Task 3 Learning Task 4 Learning Task 5		Learning Task 1 Learning Task 2 Learning Task 3 Learning Task 4 Learning Task 5	

Note: If the lesson is designed for two or more weeks as shown in the eartag, just copy your personal evaluation indicated in the first Level of Performance found in the second column up to the succeeding columns, ie. if the lesson is designed for weeks 4-6, just copy your personal evaluation indicated in the LP column for week 4, week 5 and week 6. Thank you.

Answer Key

B. Cake flour - 36 g Milk - 1680 ml Sugar - 6 cups Bulter - 0.015 g Water - 18 ml Chocolate chips -0.3g 1.120 pesos/ cake 2.300 pesos	WEELK1-2: Learning Task 1 1. meter or yard 2. Centimeter or inches 3. Gallon, pint or quarts 4. Centimeter or miles 5. Kilometer or miles 6. °O Celsius or °Pshrenheit 7. Kilogram or Pound 8. Inches or centimeter	5. 5y5+y ⁺ +5y ² -y+12 63x ³ -9x ² + 3 73x ² y -12xy +9y ² 8. 15x ² -13xy +5y ² 94x ³ +6x ² +2x +16 10. 0.5y ³ +6.5y ² - 12y + 12 Logarning Task 3 1. 240 cm ² 2. 30 cm ² 3. 2x ² +14x + 2 4. 3x ³ +4x ² + 2x + 8 5. x ³ -3x ² + 12	2, 2x 3, 2x 4, 4 5, x ² +4x + 4 B. 1, x ² -2x 3, 2x + 3, 2x + 4 5, 2x + 4	9, 5x < 90 10. 2Hy < 25 2, 4 3, 2 4, 1 5, 3 4, 1 7, 1, 2, 3 1, 2, -3, -4 2, 1, 2, 3 3, 5, 4, 3 4, 4, 5, 6 5, 4, 3 4, 4, 5, 6 5, -4, -5
A.1. 70°, acute 2. 45°, acute 3. 45°, acute 4. 45°, acute 5. 75°, acute	7 + x . 1 22 2x - 6 3. 12x - 8 4. 7x² + 6x	1, 5x ⁴ -6x ³ - 7x ² -3 2, -3x ³ y -4x ² y ² -1 3, 18y ³ + 10 4, -9x ⁵ -6x ³ -10x ² +2	8-821-5yx8 8-821-5yx8- sirty yrT A 2x .1	$1z = e - \frac{x}{\epsilon} \cdot \delta$ $01 \le 7 - x\delta \cdot 7$ $04 \ge \delta 1 + x\delta \cdot \delta$
C. 133 2. 412,500 m² 2. 412,500 m² 5. 412,500 m² 5. 412,500 m² 6. 0.6 kg of pow-dered milk 7. A. 4.08 hours B. math—3.75 Sci & Eng. –3.33 Fil. –2.92 RAPEH & EPP—2.5 Sci & Eng. –3.33 Fil. –2.92 C. 4.5 minutes	71- 6 2 2 - 17. milai	1. x2-25 2. 9x2+24x+16 3. 3x-2 4. y2-3y+1 5. 2m3+4m3-4m Learning Task 1 1. 11 3. 5. 10 143 6 1. 2x4, 6 295, 6 7223.2 319 8. 10 414.3 9. 106.88 5160 1013.4 8160 1013.4 9160 1013.4	2. 924 + 10a + 25 2. 9x2y 2 - 42xy - 49 3. x2 + y2 + 2xy - 4x 4. 4x + 9y 2 + 12xy - 4x 4. 4x + 9y 2 + 12xy - 4x 5. 9a ² + b ² + 4c ² - 6a 6ab + 12ac - 4bc 6. 16 - 4x ² 7. 9x2y 2 - 8x2bc ² 8. x ⁴ - 16y ⁴ 9x ² + 2x + 3 10. (x ² + 2x + 2x + 3 11. 8+2,4x+2,4x+8x ³ 12. x ² + 2x + 3 11. 8+2,4x+2,4x+8x ³ 11. 8+2,4x+2,4x+8x ³ 11. 8+2,4x+2,4x+8x ³ 11. 8+3,4x+2,4x+8x ³ 12. x ³ + 150x ³ 13. 8x ³ + 60x ³ 14. x ³ + 3x ³ + 3x ³ 15. x ³ + 3x ³ + 3x ³ 17. x ³ + 3x ³ + 3x ³ 18. x ³ + 3x ³ + 3x ³ 19. x ³ + 3x ³ + 3x ³ 19. x ³ + 3x ³ + 3x ³ 10. x ³ + 3x ³ + 3x ³ 11. x ³ + 3x ³ + 3x ³ 12. x ³ + 3x ³ + 3x ³ 13. x ³ + 3x ³ + 3x ³ 14. x ³ + 3x ³ + 3x ³ 15. x ³ + 3x ³ + 3x ³ 17. x ³ + 3x ³ + 3x ³ 18. x ³ + 3x ³ + 3x ³ 19. x ³ + 3x ³ + 3x ³ 10. x ³ + 3x ³ + 3x ³ 10. x ³ + 3x ³ + 3x ³ 11. x ³ + 3x ³ + 3x ³ 12. x ³ + 3x ³ + 3x ³ 13. x ³ + 3x ³ + 3x ³ 14. x ³ + 3x ³ + 3x ³ 15. x ³ + 3x ³ + 3x ³ 17. x ³ + 3x ³ + 3x ³ 18. x ³ + 3x ³ + 3x ³ 19. x ³ + 3x ³ + 3x ³ 19. x ³ + 3x ³ + 3x ³ 19. x ³ + 3x ³ + 3x ³ + 3x ³ 19. x ³ + 3x ³ + 3x ³ + 3x ³	1.36 & 38 4.12&14 2. Migz-15, Inzo-25 3. 60.75km 5. x < 38 WEEK 7 Learning Task 2 B. I. y > 20,000 A. y > 20,000 A. y > 20,000 A. y > 20,000 A. x > 43 x = 60 S. x < 51 S. x <
2. 1560 dam 3. 108ml 4. 120ml 5. 4.34gal 6. 2.48 days 7. 960 mo. 8. 26°C 9. 94.08°C 10. 0.010866 kg B. 26°C 11. acute 2. Obtuse 3. Acute 4. Acute 5. Obtuse 5. Obtuse 5. Obtuse 7. Obtuse 8. Acute 8. Acute 7. Obtuse 8. Acute 8. Acute 9. Acute 1. Acute 8. Acute 9. Obtuse 1. Acute 1. Acute 1. Acute 1. Acute 1. Acute 2. Obtuse 3. Acute 5. Obtuse 5. Obtuse 5. Obtuse 7. Obtuse 8. Acute 8. Acute 8. Acute 8. Acute 9. Acute 1. Acute 8. Acute 9. Acute 1. Acute	3. minus (-) 8. mulitiply (x,·) 4. minus (-) 9. divide (+,·/) 5. mins (-) 10. plus (+) 1. C 2. D 3. E 4. A 5. B 5. B 5. B 5. Wind Varia- Degree Conbe stnat	21χ ² γ ³ 5. χ ³ – χ ² – 4χ + 4 4. 4χ ³ + 6χ ² – 4χ + 2 5. 96γ ⁵ – 72γ ⁴ + 296γ ³ 6. 6. 68γ ² – 269γ+80 7. 48θσ ² – 68β ² + 7 118β ² β ² σ ² 8. 5σ + 9β 9. 8 ⁴ –82λ ² + b ⁴ 1 + w3 · 1 · 1 · w2 · 1 1 + w3 · 1 · w3 · 1 1 - w3 · 1 · w3 · 1 1 - w3 · 1 · w3 · 1 1 · w3 · 1 · w3 · 1	4, x +(x+1) = 29 5, 4x + 8 = 54 Learning Task I 1, at - 2ab + bz 2, at - 2ab + bz 3, at - bz 4, at - 3a2b + bz 4, at - 3a2b + bz 6, at - 4ab2 - bz 7, at - 2ab + bz 7, at - 2ab + bz 8, at - bz 4, at - 3a2b - bz 7, at - 2ab + bz	8 · 0 · 2 · 8 · 8 · 8 · 8 · 8 · 8 · 9 · 8 · 9 · 9
WEEK 1-2 Learning Task 2 A. 1. 0.708m	WEEK 3 Learning Task I L. plus (+) 6. multiply (x,·) 2. plus (+) 7. divide (+, /) 2.	1m ¹² 2. 2. \frac{25e^4}{ds} 3. 27a6b3c9 4. \text{6x5y3z3} 5. \frac{3mnx}{2y} 2. \frac{3mnx}{2y} 2. \frac{3mnx}{2} 3. \frac{14ab3}{2} 3. \frac{14ab3}{2} 4. \text{8a3b} + 6a2b2 - \frac{14ab3}{2} 3. \frac{23x^4y^4 + 9x^3y^2 - \frac{14ab3}{2} 3. \frac{23x^4y^4 + 9x^3y^2 - \frac{14ab3}{2} 3. \frac{23x^4y^4 + 9x^3y^2 - \frac{1}{2} 3. \	2. $2x = 54$ 3. $3x + 2 < 50$ 4. $\frac{1}{2}(x+4) < 28$ 4. $\frac{1}{2}x + 4$ 5. $x + \frac{1}{3}x = 5$ Learning Task 2 A. I. $4x + 2x = 96$ 2. $3x = 24$ 2. $3x = 24$ 3. $\frac{2}{3}x = 72$	8 < x .01 E > x .7 81 .8 4 4 4 5 4 4 5 4 6 4 6 4 6 4 6 4 6 4 6 4 6 4 6
Refe	erence	WEEK 2 Vestuing Task 1	WEEK 7 Learning Task 1 1. x + 5 = 12	WEEK 8 Learning Task 1 1.3 5.4 8.x > 6 2.3 6.x > 1 9.x < -3

Department of Education. (2020). *K to 12 Most Essential Learning Competencies with Corresponding CG Codes*. Pasig City: Department of Education Curriculum and Instruction Strand.

Department of Education Region 4A CALABARZON. (2020). PIVOT 4A Budget of Work in all Learning Areas in Key Stages 1-4: Version 2.0. Cainta, Rizal: Department of Education Region 4A CALABARZON.

For inquiries or feedback, please write or call:

Department of Education Region 4A CALABARZON

Office Address: Gate 2, Karangalan Village, Cainta, Rizal

Landline: 02-8682-5773, locals 420/421

https://tinyurl.com/Concerns-on-PIVOT4A-SLMs

