

## Fasting Increases Aggression and Differentially Modulates Local And Systemic Steroid Levels in Male Zebra Finches

H. Bobby Fokidis, Nora H. Prior, and Kiran K. Soma

Department of Psychology (H.B.F., K.K.S.), Department of Zoology (N.H.P., K.K.S.), Graduate Program in Neuroscience (K.K.S.), and Brain Research Centre (K.K.S.), University of British Columbia, Vancouver, British Columbia, V6T-1Z4, Canada

Aggression enables individuals to obtain and retain limited resources. Studies of the neuroendocrine regulation of aggression have focused on territorial and reproductive contexts. By contrast, little is understood concerning the neuroendocrine regulation of aggression over other resources, such as food. Here, we developed a paradigm to examine the role of steroids in food-related aggression. In groups of male zebra finches, a 6-hour fast decreased body mass and increased aggressive interactions among subjects that competed for a point source feeder. Fasting also dramatically altered circulating steroid levels by decreasing plasma testosterone but not estradiol ( $E_2$ ). By contrast, both plasma corticosterone and dehydroepiandrosterone (DHEA) concentrations were elevated with fasting. Interestingly, short-term access to food (15 minutes) after fasting normalized circulating steroid levels. Fasting increased corticosterone levels in a wide range of peripheral tissues but increased DHEA levels specifically in adrenal glands and liver; these effects were quickly normalized with refeeding. DHEA can be metabolized within specific brain regions to testosterone and  $E_2$ , which promote the expression of aggression. We measured  $E_2$  in micro-dissected brain regions and found that fasting specifically increased local  $E_2$  levels in 3 regions: the periaqueductal gray, ventral tegmental area, and ventromedial nucleus of the hypothalamus. These regions are part of the vertebrate social behavior network and regulate the expression of aggression. Together, these data suggest that fasting stimulates secretion of DHEA from the adrenals and liver and subsequent conversion of DHEA to  $E_2$  within specific brain regions, to enable individuals to compete for limited food resources. (*Endocrinology* 154: 4328–4339, 2013)

Aggression is an ancient behavior that evolved to enable organisms to cope with limited resources. Aggression is context dependent, has multiple causes, and involves the coordination of peripheral endocrine signals and central neural processes. Sex steroids are well-known modulators of aggression within the context of dyadic encounters, such as resident-intruder interactions for territorial defense (1) or male-male competition for mates (2, 3). Aggression in these contexts is typically attributed to the actions of gonadal T, which can bind to neural androgen receptors (4, 5) or be converted locally to  $17\beta$ -estradiol ( $E_2$ ) by the enzyme aromatase and bind to neural

estrogen receptors (6, 7). Importantly, aggression occurs in other contexts, and its most ubiquitous expression is the defense and monopolization of food resources (8, 9), but little is understood of the physiologic regulation of aggression in this context.

In social species, limited food availability or fasting can spur aggressive interactions within the group or decreased social stability (10, 11). Fasting induces a myriad of effects on steroid profiles, including adrenal secretion of glucocorticoids (eg, corticosterone [CORT]) to mobilize energy reserves (12, 13) and increase food intake (14, 15). Increased CORT secretion also suppresses gonadal sex ste-

ISSN Print 0013-7227 ISSN Online 1945-7170  
Printed in U.S.A.

Copyright © 2013 by The Endocrine Society

Received February 20, 2013. Accepted August 2, 2013.

First Published Online August 12, 2013

Abbreviations: BST, bed nucleus of stria terminalis; CB, cerebellum; CORT, corticosterone; DHEA, dehydroepiandrosterone;  $E_2$ ,  $17\beta$ -estradiol; HP, hippocampus; HSD, hydroxysteroid dehydrogenase; LS, lateral septum; PAG, periaqueductal gray; POA, preoptic area; VMH, ventromedial nucleus of the hypothalamus; VTA, ventral tegmental area.

roid (eg, T and E<sub>2</sub>) secretion (16). Because fasting decreases sex steroid secretion, it is unlikely that circulating T or E<sub>2</sub> levels directly promote the heightened aggression seen in fasted animals.

Alternatively, the androgen precursor dehydroepiandrosterone (DHEA) can maintain aggression, even when circulating sex steroid levels are low (17). Secreted by the adrenal glands, as well as the liver, the prohormone DHEA can be locally converted to active T and/or E<sub>2</sub> within behaviorally relevant neural circuits that possess the necessary steroidogenic enzymes (17, 18). Adrenal DHEA secretion increases with acute stress in humans (19, 20) and with injections of ACTH in red squirrels (21). In contrast, in song sparrows (*Melospiza melodia*), both acute restraint stress and social stress do not affect DHEA levels in the brachial vein, but modulate DHEA levels in the jugular vein (22–24). Fasting increases serum DHEA levels in humans and mice (25) as well as hepatic DHEA secretion in mice (25, 26). Because circulating DHEA levels change with fasting and DHEA promotes agonistic behavior, it is a good candidate for a steroid regulator of food-related aggression.

Here, we examine the effects of fasting on aggression, circulating steroid levels, and local steroid levels in peripheral tissues and specific brain regions in a songbird, the zebra finch (*Taeniopygia guttata*). Songbirds are excellent models for understanding the links between food availability, behavior, and steroids. Compared with laboratory rodents, songbirds have higher circulating DHEA concentrations (17), higher rates of DHEA conversion and neurosteroid synthesis in the brain (18, 27), higher mass-specific metabolic rates (28, 29), and more robust responses to even mild fasts (30, 31). Highly gregarious zebra finches are the most extensively studied songbird. Short-term fasting (ie, 2–6 hours) increases plasma CORT and decreases plasma T levels and alters courtship behavior and singing rates (32, 33). A suite of steroidogenic enzymes is expressed within the zebra finch brain (27, 34–36), thus indicating the capacity for de novo sex steroid synthesis.

If aggression over food is regulated by neurally synthesized sex steroids, then we predict that a fast would increase aggression, decrease circulating sex steroid levels, and increase circulating DHEA and CORT levels and/or local sex steroid levels in behaviorally relevant brain regions such as the bed nucleus of stria terminalis (BST), lateral septum (LS), ventromedial nucleus of the hypothalamus (VMH), periaqueductal gray (PAG), and the ventral tegmental area (VTA). In particular, E<sub>2</sub> is a key regulator of male aggression (17, 37), and we measured local E<sub>2</sub> levels within the social behavior network, a highly conserved circuit of brain regions (including those listed

above) in vertebrates that regulates a variety of social behaviors (38, 39). To our knowledge, this is the first study to investigate the role of steroids in food-related aggression in any species.

## Materials and Methods

All methods complied with the Canadian Council on Animal Care and University of British Columbia guidelines for animal care and use.

### Experiment 1: effects of an acute fast on aggressive behavior

This experiment tested the hypothesis that acute fasting increases aggressive interactions within a group. Adult male zebra finches that had not previously interacted were housed overnight in groups of 4 with ad libitum access to food (50:50 panicum and white millet, ~12% protein, 4.7% lipid; Just for Birds) and water. Subjects were held on a 14-hour light, 10-hour dark photoperiod. Prior to lights on the following day, all food was removed to prevent subjects from feeding and accumulating seeds in their crop, an alimentary structure that stores food prior to digestion. Subjects were transferred as a group to a testing room. Subjects were weighed ( $\pm 0.25$  g), their furcular fat and pectoralis muscle scores (both 0–5) were recorded, and an empty crop was confirmed via palpation. All subjects were marked on their breast feathers with a blue or green washable marker for individual identification. Zebra finch plumage lacks these colors, and thus they are unlikely to influence social interactions.

On either side of the testing cage (38 ½" × 19 ¾" × 19", Corner's Ltd) were 2 wooden perches, a single food dish for general feeding, a water tower, and a point source feeder that allows only one subject to access food at a time. Groups of 4 subjects were randomly assigned to either 1) ad libitum food (control) with a full food dish or 2) a 6-hour fasting treatment (fasted) with an empty food dish ( $n = 15$  groups per treatment, 60 subjects per treatment, 120 subjects total). All cages began with an empty point source feeder, which was replaced with full one at the onset of the behavioral trial. Each testing room housed 2 groups (one control, one fasted). Subjects were left undisturbed for 6 hours, after which food dishes (empty or full) were removed from all cages, and subjects were video-recorded for 15 minutes. Then empty point source feeders were replaced with full ones, and subjects were video recorded for an additional 30 minutes. Birds were then captured, reweighed, rescored for fat and muscle, and returned to the colony. Several subjects ( $n = 19$ ) were tested multiple times, but with at least 6 days between tests.

Videos were scored by a single observer (HBF) that was blind with respect to treatment, and various behaviors were assessed within 2 separate time blocks: 1) from 0–15 minutes and 2) from 15–30 minutes after introduction of the full point source feeder. Behaviors were categorized into those in which aggression focused on accessing the feeder and those not directly related to the feeder. The former category includes displacements of one subject by another from the point source feeder, the total amount of time each subject spent at the point source feeder, and the order in which subjects accessed the point source feeder. The latter category includes aggression away from the point source feeder, such as chases, displacements from a perch, and bill fencing

(rapid exchanges of facial pecks). We quantified the total number of behaviors from both categories that were initiated as well as received by an individual.

### Experiment 2: effects of an acute fast on circulating steroid levels

This experiment tested the hypothesis that acute fasting decreases circulating sex steroid levels and increases circulating CORT and DHEA levels. Adult male zebra finches were treated as described above (Experiment 1), but here subjects were tested in pairs. Pair housing allowed blood samples to be collected from all subjects within a short time frame to avoid steroid changes resulting from handling stress, while simultaneously avoiding isolation stress. Subjects were again weighed and scored for fat and muscle.

Subjects were then exposed to 1 of 3 treatments: 1) ad libitum food (control); 2) a 6-hour fast (fasted); or 3) a 6-hour fast followed by 15 minutes of access to a full point source feeder (refed). The 2 former groups were provided with an empty point source feeder as a control. All subjects ( $n = 12$  subjects per vein per treatment) were captured and bled from either the brachial vein (with heparinized capillary tubes) or the jugular vein (with a 1-cc heparinized insulin syringe). For each subject, only a single blood sample was collected from only one vein. Steroid levels in plasma from the brachial vein reflect systemic levels, whereas steroid levels in plasma from the jugular vein indirectly reflect the neural steroidal milieu, including the synthesis and metabolism of steroids within the brain (24). Thus steroid profiles may differ between brachial plasma and jugular plasma, providing information on systemic vs central steroid levels. Bleed times averaged  $4.7 \pm 1.6$  minutes, and blood was stored on ice until plasma was separated via centrifugation. Plasma was stored at  $-80^\circ\text{C}$ .

### Experiment 3: effects of an acute fast on tissue steroid levels

This experiment tested the hypothesis that acute fasting modulates local steroid levels within peripheral and central tissues. We repeated the above study (Experiment 2) in adult male finches with the same 3 treatment groups. Because handling stress can alter steroid concentrations, only one subject from each pair was used for this study, and the other subject was only a “companion” to avoid isolation stress ( $n = 8$  subjects per treatment). After treatment, 1 subject from each cage was captured and euthanized via rapid decapitation within  $1.4 \pm 0.3$  minutes of entering the testing room. Trunk blood was collected into heparinized capillary tubes. The brain was rapidly removed and separated into 4 parts: cerebellum (CB), diencephalon, rostral telencephalon, and caudal telencephalon (40). Peripheral tissues were collected in the following order: pectoralis muscle, furcular fat, liver, pancreas, small intestine, testes, and adrenal glands. All tissues were snap frozen on dry ice and time of collection was recorded. Trunk blood was centrifuged and plasma was collected. All samples were stored at  $-80^\circ\text{C}$ .

### Experiment 4: effects of an acute fast on $E_2$ levels in microdissected brain regions

This experiment tested the hypothesis that acute fasting increases neural  $E_2$  levels within the social behavior network, using the Palkovits punch technique to sample tissue from specific regions (41–43). We repeated the above study (Experiment 3) in

adult male zebra finches, and after treatment, 1 subject per cage was euthanized, with blood and other tissues collected as above, but here whole brains were snap frozen on powdered dry ice. Time of collection was again recorded and all samples were stored at  $-80^\circ\text{C}$ .

Brains were coronally sectioned at  $300\ \mu\text{m}$  using a cryostat set at  $-10^\circ\text{C}$  in a rostral to caudal direction and oriented to match the plane of sectioning in the zebra finch brain atlas (44). Brain regions were located relative to prominent neuroanatomical landmarks and punched using stainless steel cannulae (Brain Punch Set, catalog no. 57401, Stoelting Co). Punches for each brain region were taken from a single section and bilateral punches were pooled for subsequent analysis. Punches were either 1 or 2 mm in diameter (Table 1), depending on the region of interest, and were collected into microcentrifuge tubes and stored at  $-80^\circ\text{C}$ . The brain regions sampled are area X (X), the magnocellular nucleus of anterior nidopallium, hippocampus (HP; 3 subregions: rostral, rHP; intermediate, iHP; caudal, cHP), preoptic area (3 subregions: rostral, rPOA; intermediate, iPOA; caudal, cPOA), nucleus rotundus, BST, LS, ventromedial nucleus of the hypothalamus (VMH), optic tectum, ventral tegmental area (VTA), periaqueductal gray (PAG), nucleus taeniae of the amygdala, caudomedial nidopallium, robust nucleus of the arcopallium, CB, locus coeruleus, and the nucleus of the solitary tract. These regions were selected because they are involved in the regulation of social behavior, energy balance, or the stress response, or because they served as negative controls (ie, little change in steroid levels was expected in optic tectum, CB, and nucleus rotundus).

### Steroid extraction

For Experiment 2,  $5\ \mu\text{L}$  of plasma diluted 1:50 with assay buffer were used to measure CORT with a double-antibody  $^{125}\text{I}$  RIA (MP Biomedicals; catalog no. 07120103; see below for assay details). Remaining plasma and all tissue samples had steroids first extracted using solid-phase extraction with C-18 columns (Varian, Bond-Elut; Catalog no. 12113045) as previously described (40, 41, 45). For soft tissues, samples were first homogenized in a bead homogenizer (settings: 8 m/s for 30 seconds  $\times$  2, Omni Bead Ruptor 24, Omni) in ice-cold 80% HPLC-grade methanol (MeOH) and left overnight at  $4^\circ\text{C}$ . Samples were centrifuged the following day and the supernatant was collected (up to 1 mL) to which was added 10 mL of  $\text{dH}_2\text{O}$ , prior to loading on C-18 columns. Columns had been primed with 3 mL 100% MeOH and then equilibrated with 10 mL  $\text{dH}_2\text{O}$ . Samples were then loaded on the column. This was followed by an interference elution with 10 mL 40% MeOH; this elution aimed to remove interfering lipids and glucuronidated and sulfated steroids. Finally, steroids were eluted with 5 mL 90% MeOH. Samples were then dried at  $40^\circ\text{C}$  in a vacuum centrifuge (Thermo Electron SPD111V Speedvac; Thermo Scientific) and stored at  $-20^\circ\text{C}$ .

### RIAs

Dried extracts from Experiment 2 were resuspended with  $6\ \mu\text{L}$  of absolute ethanol and  $1194\ \mu\text{L}$  of phosphate buffer containing sodium chloride, gelatin, and sodium azide (PBSG). This provided sufficient volume to measure T and DHEA in duplicate and  $E_2$  in singleton. For  $E_2$ , samples were run as singletons to maximize the chance of detecting  $E_2$  (41). Thus for Experiment

**Table 1.** Concentrations of E<sub>2</sub> (in pg/g) in Brain Regions of Adult Male Zebra Finches Exposed to ad Libitum Food (Control), a 6 Hour Fast (Fasted), or a 6-Hour Fast Followed by 15 Minutes of Access to a Point Source Feeder (Refed)

Brain Region	Punch Number and Size	Control	Fasted	Refed
Social Behavior Network				
POA				
Rostral	2 × 2 mm	434.3 ± 15.5 (100)	456.9 ± 18.3 (100)	466.1 ± 11.7 (100)
Intermediate	2 × 2 mm	425.4 ± 21.2 (100)	435.0 ± 25.5 (100)	393.6 ± 17.2 (100)
Caudal	2 × 2 mm	545.4 ± 29.8 (100)	503.7 ± 48.4 (100)	536.3 ± 56.7 (100)
BST	2 × 1 mm	ND	202.6 ± 39.1 (38)	ND
LS	1 × 2 mm	390.9 ± 71.5 (88)	509.9 ± 75.8 (86)	466.6 ± 81.5 (88)
PAG	1 × 2 mm	198.2 ± 81.5 (29)	612.6 ± 119.2 (80)	363.6 ± 108.6 (63)
VTA	2 × 1 mm	1083 ± 57.7 (100)	1388.5 ± 64.4 (100)	1189.6 ± 70.9 (88)
VMH	2 × 1 mm	583.6 ± 58.3 (100)	847.2 ± 78.2 (100)	782 ± 84.2 (100)
Nucleus taeniae of the amygdala	2 × 1 mm	72.3 (14)	ND	ND
Other brain areas				
Area X	2 × 2 mm	289 ± 69.6 (38)	19.0 (14)	300.8 ± 58.8 (38)
Magnocellular nucleus of the anterior nidopallium	2 × 2 mm	113.5 ± 44.3 (25)	86.1 ± 86.1 (13)	68.8 ± 68.8 (25)
Robust nucleus of the arcopallium	2 × 1 mm	1462.5 ± 186.1 (100)	1418 ± 164.0 (100)	1475.2 ± 178.7 (100)
Caudomedial nidopallium	2 × 2 mm	3251.4 ± 241.8 (100)	3451.7 ± 308.1 (100)	3027.9 ± 116.4 (100)
Nucleus rotundus	2 × 1 mm	231.5 ± 231.5 (13)	56.9 ± 56.9 (13)	ND
Nucleus tractus solitarius	1 × 2 mm	235.1 ± 9.5 (100)	ND	102.8 ± 28.6 (50)
Locus coeruleus	2 × 1 mm	ND	121.6 ± 21.6 (25)	187.6 ± 55.1 (40)
Rostral hippocampus	2 × 2 mm	381.7 ± 117.8 (100)	388.5 ± 140.4 (100)	464.1 ± 104.6 (100)
Intermediate hippocampus	2 × 2 mm	387.8 ± 132.6 (100)	367.8 ± 124.3 (100)	380.1 ± 126.7 (100)
Caudal hippocampus	2 × 2 mm	261.5 ± 20.1 (100)	290.5 ± 16.0 (100)	280.7 ± 7.5 (100)
Optic tectum	2 × 2 mm	ND	ND	ND
CB	1 × 2 mm	ND	ND	ND

All data are mean ± SEM and numbers in parentheses indicate percentage of detectable samples out of 8 birds within each group. ND, not detectable.

2, the T, DHEA, and E<sub>2</sub> assays used 28%, 36%, and 36% of the original sample, respectively. For Experiment 3, samples were resuspended using 4.5 µL of absolute ethanol and 645.5 µL PBSG. Assays for CORT, T, DHEA, and E<sub>2</sub> were all run in singleton to maximize the chance of detecting the analytes and used 8%, 28%, 16%, and 46% of the original sample, respectively. For Experiment 4, the small size of punches necessitated the entire sample be used to measure E<sub>2</sub> in singleton.

We used specific double-antibody <sup>125</sup>I RIAs to measure 4 steroids: CORT (see above), T (MP Biomedicals; catalog no. 07189102), DHEA (Beckman-Coulter; Immunotech DSL 8900), and 17β-E<sub>2</sub> (Beckman-Coulter; Immunotech DSL 4800). These were run according to manufacturer's recommendations; however, assays were modified to improve sensitivity and were previously validated for songbird samples (24, 40, 46). The detectable ranges for the assays were as follows: CORT = 1.5 to 1000 pg/tube; DHEA = 2–600 pg/tube; T = 0.3125 to 500 pg/tube; and E<sub>2</sub> = 0.192 to 19.2 pg/tube. Additional validations presented here include demonstrating parallelism between standard curves and serially diluted plasma and tissue (Supplemental Figure 1 published on The Endocrine Society's Journals Online web site at <http://endo.endojournals.org>) and estimates of recovery of known steroid quantities within several tissues, prior to solid-phase extraction and subsequent comparison with “unspiked” samples (Supplemental Table 1). All data were corrected for recovery. Intrassay and interassay variation are reported in Supplemental Table 2.

## Statistical analysis

All data were tested for normality of distribution (Kolmogorov-Smirnov test), homoscedasticity (Levene's test of equal variance), and for the presence of influential outlying data (Chauvenet's test). When required, data were arcsine square root or log transformed. Outliers were removed if they altered the results of the data analysis. We used repeated measures ANOVA to assess individual changes in body mass and fat and muscle scores.

To assess changes in behaviors with treatment (Experiment 1), we used a mixed model ANOVA approach. The repeated measure was the behavior before, immediately after, and 15 minutes after introduction of the point source feeder (ie, pre, post, 15-minutes post). To account for variation at the group level, we tested individual behaviors nested within groups (ie, cage identification). This enabled us to detect changes in behavior with food introduction, while controlling for the effects of specific groupings of subjects. Treatment was the between-subjects factor, and initial body mass, and the change in body mass were included as random factors in the model. Two behaviors (the occurrence of chase events and bill fencing) had a Poisson distribution and thus were analyzed using a repeated-measures logistic regression providing a Wald's  $\chi^2$  statistic, with treatment as the independent variable and food introduction (pre, post, 15 minutes post) as the repeated measure. To assess whether body mass or change in body mass influenced the order in which the point source feeder was accessed (ordinal data), we used a non-parametric Kruskal-Wallis one-way ANOVA (*H* test statistic) coupled to Tamhane's *T*<sub>2</sub> post hoc tests for unequal variances.



Changes in plasma steroid levels in response to treatment (Experiment 2) were assessed using multivariate analysis of covariance with treatment, steroid, vein where blood was sampled (brachial vs. jugular), and their interactions as the between-subjects factors, and with change in body mass as a covariate. Tissue steroid levels (Experiment 3) were analyzed using multivariate analysis of covariance with treatment and tissue type as between-subjects factors, and initial body mass as a covariate. Levels of  $E_2$  in brain punches (Experiment 4) were analyzed using ANOVA with treatment and brain region as between- and within-subjects factors, respectively. Due to large number of brain regions sampled in this study ( $n = 21$ ) that reduces statistical power, we split our analyses into those regions that make up the social behavior network (PAG, POA, BST, LS, VMH, VTA, and nucleus taeniae of the amygdala) (39, 47) and the other regions. Relevant interactions between independent variables were also included, and differences between tissues (Experiment 3) or brain regions (Experiment 4) were assessed using Tukey's honestly significant difference post hoc tests. The critical  $\alpha$  level was set at  $P \leq .05$  and statistics were performed using Windows versions of Statistica 9.1 (StatSoft Inc), SPSS 13.0 (SPSS), and GraphPad Prism 5 (GraphPad).


## Results

### Experiment 1: effects of fasting on morphology and aggressive behavior

Fasted subjects exhibited significant decreases in body mass ( $F_{1,59} = 19.631$ ,  $P \leq .001$ ; Figure 1A) and muscle score ( $F_{1,59} = 6.893$ ,  $P = .002$ ; Figure 1B), but not fat score ( $F_{1,59} = 0.205$ ,  $P = .653$ ; Figure 1C) compared with controls.

Prior to presentation of a point source feeder, control and fasted subjects behaved similarly and showed little social interaction. Both immediately after (fasted group) and 15 minutes after (refed group) the presentation of a point source feeder, subjects spent more time feeding ( $F_{1,54} = 9.271$ ,  $P \leq .001$ ; Figure 2A) and displayed more point source feeder displacements ( $F_{1,54} = 10.365$ ,  $P \leq .001$ ; Figure 2B) than controls, but the refed subjects displayed fewer overall behavioral acts than fasted subjects. Fasted and refed subjects initiated ( $F_{1,54} = 5.201$ ,  $P \leq .001$ ; Figure 2C) and received ( $F_{1,54} = 3.770$ ,  $P = .008$ ; Figure 2C) more behavioral acts than controls, but in both cases the trend toward fewer behavioral acts in the refed subjects was not significant from the fasted subjects.

Along with point source feeder-centered behaviors, other social behaviors were also more frequent in fasted compared with control and/or refed subjects. The occurrence of chases was more frequent in fasted compared with refed or control subjects (Wald's  $\chi^2 = 6.937$ ,  $df = 1$ ,  $P = .048$ ; Figure 2D). Displacements from a perch were more frequent in fasted and refed subjects compared with controls ( $F_{1,54} = 4.002$ ,  $P = .016$ ; Figure 2E). However, no



**Figure 1.** Effects of an acute fast (6 hour) on (A) body mass, (B) pectoralis muscle score, and (C) furcular fat score in adult male zebra finches ( $n = 60$  per treatment). All data are mean  $\pm$  SEM. \*,  $P \leq .05$ ; \*\*,  $P \leq .01$ ; and \*\*\*,  $P \leq .001$ .


difference in the occurrence of bill-fencing behavior was observed among groups (Wald's  $\chi^2 = 0.844$ ,  $df = 1$ ,  $P = .441$ ; Figure 2F).

Both initial body mass and the change in body mass influenced the order in which fasted subjects accessed the point source feeder, with heavier subjects accessing the feeder prior to lighter subjects (Kruskal-Wallis  $H = 4.731$ ,  $df = 1$ ,  $P = .032$ ). A stronger predictor of access was the proportion of body mass lost during the fast, with subjects that lost more mass accessing the point source feeder prior to those that lost less mass (Kruskal-Wallis  $H = 5.922$ ,  $df = 1$ ,  $P = .016$ ; Figure 3).

### Experiment 2: effects of fasting on circulating steroid levels

Overall, plasma steroid concentrations were affected by treatment but not the vein sampled; however, the treatment  $\times$  vein interaction was often significant (Supplemental Table 3). The change in body mass also affected plasma steroid levels (Supplemental Table 3).

Fasted subjects had significantly higher brachial and jugular CORT concentrations than control and refed subjects (Figure 4A and Supplemental Table 3). Plasma CORT levels also tended to be higher in subjects that lost



**Figure 2.** Aggressive behaviors within zebra finch groups exposed to ad libitum food (Control), a 6-hour fast (Fasted), or a 6-hour fast followed by 15 minutes of access to a point source feeder (Refed). A, The total time group spent at the point source feeder. B, The number of displacements of subjects from a point source feeder. C, The number of behavioral acts initiated by and received by an individual. D, The proportion of behavioral trials with an observed chase. E, The number of displacements of subjects from perches. F, The occurrence of bill fencing.  $n = 15$  groups per treatment. All data are mean  $\pm$  SEM. \*,  $P \leq .05$ ; \*\*,  $P \leq .01$ ; and \*\*\*,  $P \leq .001$ .

more body mass with fasting; however, this correlation was not significant (Pearson's  $r = 0.274$ ;  $P = 0.091$ ).

Plasma DHEA concentrations showed a significant treatment  $\times$  vein interaction. In the brachial plasma, DHEA concentrations were significantly higher in fasted subjects than either control or refed subjects (Figure 4B and Supplemental Table 3). In stark contrast, in the jugular plasma, fasting did not affect DHEA concentrations, although DHEA concentrations in refed subjects were significantly lower than controls (Figure 4B).


Plasma T concentrations also showed a significant treatment  $\times$  vein interaction (Supplemental Table 3). Fasted subjects had lower T levels in the brachial plasma, but not in the jugular plasma, compared with controls (Figure 4C). Refeeding increased T levels in the jugular plasma, but not in the brachial plasma, compared with fasted subjects (Figure 4C).

Plasma  $E_2$  concentrations were not significantly affected by treatment, the vein sampled, or change in body mass (Supplemental Table 3). However, as expected,  $E_2$  concentrations tended to be lower in brachial than jugular plasma for control subjects (Figure 4D).

### Experiment 3: effects of fasting on tissue steroid levels

Overall, tissue steroid concentrations were significantly affected by treatment, tissue type, and often their interaction, but not by body mass (Supplemental Table 4).

CORT was detectable in all tissues and was above the limit of detection in all samples. Fasting increased CORT concentrations in all tissues, except furcular fat (Figure 5A). Notably, tissue CORT concentrations did not differ between controls and refed subjects (Figure 5A). As expected, CORT concentrations were highest in the adrenal gland (Figure 5A).



**Figure 3.** Box plot exhibiting the order of access to a point source feeder by groups of 4 adult male zebra finches in relation to the degree of body mass lost during a 6-hour fast. Different letters indicate significant differences at  $P \leq .05$ .

DHEA was detectable in all tissues, except for 3 samples (muscle-Refed:  $n = 1$ ; rostral telencephalon-Fasted:  $n = 2$ ). Treatment influenced DHEA concentrations in a tissue-specific manner (Supplemental Table 4). Fasting increased DHEA concentrations in the trunk plasma, liver, adrenal glands, caudal telencephalon, and diencephalon only (Figure 5B).

T was detectable in all tissues, but was below the limit of detection in 5 samples (muscle-Control:  $n = 1$ ; muscle-Refed:  $n = 1$ ; Intestine-Fasted:  $n = 1$ ; diencephalon-Fasted:  $n = 2$ ). Tissue T levels were not significantly affected by treatment but did vary across tissues (Supplemental Table 4). As expected, T concentrations were highest in the testes, which exhibited a nonsignificant trend for lower T levels in fasted subjects (Figure 5C).


Levels of  $E_2$  in peripheral tissues were not assessed because no significant effect of treatment was observed on circulating  $E_2$  concentrations.

#### Experiment 4: effects of fasting on $E_2$ levels in microdissected brain regions

In microdissected brain punches ( $\leq 1$  mg), levels of  $E_2$  were detectable (ie, within the range of the standard curve) in 19 of the 21 brain regions examined (Table 1). However, for some regions, only a small proportion of samples had detectable amounts of  $E_2$ , even though we used an ultrasensitive assay that can measure as little as 0.2 pg of  $E_2$  per tube. To avoid testing small sample sizes, only regions with  $>25\%$  of samples detectable in each treatment group were considered for further analysis (Table 1).

For both the HP and preoptic area (POA), we did not observe differences in  $E_2$  concentrations among rostral, intermediate, and caudal levels (HP:  $F_{2,272} = 1.427$ ,  $P = .242$ ; POA:  $F_{2,272} = 1.427$ ,  $P = .242$ ; Table 1). Thus, a mean  $E_2$  concentration was calculated for the HP and POA for subsequent analysis.


In the social behavior network, there was a significant treatment  $\times$  brain region interaction (Supplemental Table 5). In contrast, in the other regions, there was an effect of brain region only, with no effect of treatment and no treatment  $\times$  brain region interaction (Supplemental Table 5). In the social behavior network, post hoc comparisons revealed that 3 regions significantly responded to fasting (Figure 6). Fasted subjects exhibited significantly higher  $E_2$  levels than controls within the PAG, VTA, and VMH (Figure 6). In addition, refeeding significantly reduced  $E_2$  levels in 2 of these regions: PAG and VTA (Figure 6).



**Figure 4.** Plasma concentrations of (A) CORT, (B) DHEA, (C) T, and (D)  $E_2$  in the brachial and jugular veins of adult male zebra finches exposed to ad libitum food (Control), a 6-hour fast (Fasted), or a 6-hour fast followed by 15 minutes of access to a point source feeder (Refed).  $n = 12$  subjects per vein per treatment. All data are mean  $\pm$  SEM. \*,  $P \leq .05$ .


## Discussion

Here, we propose a novel mechanism for regulating aggression over access to food resources, involving adrenal and hepatic DHEA secretion and subsequent local conversion of DHEA to  $E_2$  within specific brain regions. An acute fast increased several aggressive behaviors immediately after food presentation, and these behaviors decreased (or trended toward a decrease) after 15 minutes of access to food presentation. Fasted subjects had reduced systemic T levels compared with controls. Thus systemic T levels cannot account for the aggressive competition over



**Figure 5.** Concentrations of (A) CORT, (B) DHEA, and (C) T in trunk plasma and tissues of adult male zebra finches exposed to ad libitum food (Control), a 6-hour fast (Fasted), or a 6-hour fast followed by 15 minutes of access to a point source feeder (Refed) ( $n = 8$  subjects per treatment). All data are mean  $\pm$  SEM. \*,  $P \leq .05$ .

food. Fasting, however, increased plasma levels of 2 adrenal steroids, CORT (in both brachial and jugular veins) and DHEA (in brachial vein only). The CORT increase was widespread among tissues, consistent with its role in energy mobilization. In contrast, the adrenal glands and liver were the only peripheral tissues that exhibited increases in DHEA content with fasting. Both tissues had higher DHEA levels than plasma, suggesting these organs are the major sites of DHEA secretion during fasting.



**Figure 6.** Concentration of E<sub>2</sub> within 3 brain regions in the social behavior network in adult male zebra finches exposed to ad libitum food (Control), a 6-hour fast (Fasted), or a 6-hour fast followed by 15 minutes of access to a point source feeder (Refed). Dotted line indicates the mean E<sub>2</sub> concentration in trunk plasma.  $n$  ranges from 3 to 8 subjects per treatment; percentage detectability of samples is detailed in Table 1. All data are mean  $\pm$  SEM. \*,  $P \leq .05$ ; \*\*,  $P \leq .01$ .

DHEA can be converted to active sex steroids (T and E<sub>2</sub>) within the brain to modulate social behavior. Consistent with this idea, we observed localized increases in E<sub>2</sub> concentrations with fasting in three areas (PAG, VTA, and VMH) of the social behavior network. Surprisingly, providing access to food for only 15 minutes after the fast was often sufficient to normalize steroid concentrations in plasma, peripheral tissues, and the brain to control levels, which was consistent with the changes in behavior. Taken together, these data support a novel role for steroids as mediators of food-related aggression in male zebra finches, which may involve peripheral DHEA secretion in response to an energetic challenge and subsequent neural conversion of DHEA to E<sub>2</sub> to influence aggression.

### An acute fast has dramatic effects on body mass, muscle, and adrenal steroids

In this study, zebra finches exhibited robust responses to a 6-hour fast, including an 8.5% decline in body mass and a decrease in pectoralis muscle score. These data support other songbird studies, indicating that even short-term food limitation constitutes a major energetic challenge (30, 48). The reduced muscle score likely reflects increased protein catabolism to meet the energetic demands of fasting. One likely mediator of this effect is CORT, which increased with fasting in plasma and across nearly all tissues investigated. During fasting, CORT maintains glucose levels (16) by promoting gluconeogenesis using substrates derived from muscle and fat catabolism (12, 30, 49). Furthermore, CORT encourages locomotor activity in captive birds (30, 31, 50), and this may contribute to the decreases in body mass and muscle score. Although fasting elevated CORT levels in most tissues, it did not significantly increase CORT levels in furcular fat.



We also did not observe a decrease in fat score with fasting, which collectively suggests that CORT does not mobilize energy from furcular fat under the current paradigm.

Refeeding lowered CORT concentrations to control levels, suggesting the rapid ( $\leq 15$  minutes) metabolism or clearance of CORT once food becomes available. Many tissues express  $11\beta$ -hydroxysteroid dehydrogenase type 2 ( $11\beta$ -HSD2) which inactivates CORT to dehydrocorticosterone (51, 52). Earlier studies have reported a half-life for CORT in plasma of 10–25 minutes under baseline conditions (53, 54), but more recent studies suggest shorter times of 5–10 minutes in rats (55–57). The higher metabolic rate of songbirds, compared with similarly-sized mammals (28), suggests an equivalent or faster rate of clearance of CORT in circulation. However, white-crowned sparrows (*Zonotrichia leucophrys*) that ingested mealworms injected with CORT showed a peak in circulating CORT after 7 minutes, and baseline levels were restored at about 1 hour (58). Ingested CORT, however, may be released slowly during digestion, in contrast to an injection of CORT. Fasting and subsequent refeeding may rapidly increase steroid clearance by up-regulating  $11\beta$ -HSD2 activity. Fasting increases  $11\beta$ -HSD2 in the rat kidney (59), and restraint stress elevates  $11\beta$ -HSD2 activity in the rat placenta to buffer the fetus from high maternal CORT levels (60); however, no studies have investigated other tissues or the effects of refeeding.

Fasting also increased brachial, but not jugular, DHEA concentrations, and refeed subjects had plasma DHEA levels that were equivalent to those of controls. This pattern was also observed in the adrenal glands and liver, where DHEA levels were higher than in trunk plasma. Previous studies in songbirds have demonstrated that both restraint and social stress were incapable of altering brachial DHEA levels, but could alter jugular and hepatic DHEA levels (23). The effects of fasting on brachial DHEA levels may involve increased hepatic DHEA secretion, potentially allowing metabolic processes in the liver to affect the brain and social behavior. The decline in plasma, adrenal, and hepatic DHEA levels within 15 minutes of refeeding is consistent with human studies, in which insulin (which increases in response to feeding) increases the metabolic clearance rate for DHEA (61, 62). Early studies suggest a much faster metabolic clearance rate for DHEA in rodents and rabbits as compared to primates (63, 64), with half-lives in circulation ranging from 15–40 minutes (65–67), but studies in birds have not been done.

### **An acute fast increases aggressive interactions but not systemic sex steroid levels**

Fasted subjects displayed more agonistic behaviors than controls, which is consistent with prior studies (8,

68). As expected, this increase in aggression was observed in behaviors aimed at accessing the point source feeder, but also in interactions not directly related to the food, such as chases and perch displacements. Furthermore, these behaviors subsided after 15 minutes of refeeding concurrent with the decreases in circulating and brain steroid profiles. Another interesting observation was the relationship between the change in body mass and the likelihood of accessing the point source feeder first. One possible interpretation is that subjects that lost more body mass (ie, had a lower energy balance) were more motivated to feed, raising the possibility of individual variation in aggression and competitiveness driven by individual differences in energy balance.

CORT increases locomotor activity (30, 31) but is unlikely to solely contribute to fasting effects on aggressive interactions, because fasted and control subjects did not differ in behaviors prior to introduction of the point source feeder. Several lines of evidence suggest that systemic but not neural T levels were reduced by fasting: 1) fasting decreased T levels in brachial plasma but not jugular plasma; 2) refeeding increased T levels in jugular plasma but not brachial plasma; and 3) fasting tended to decrease T levels in the testes but not the brain. Taken together, these data suggest that systemic T is an unlikely candidate for regulating the heightened aggression during food-related contests.

Circulating  $E_2$  levels tended to be higher in jugular plasma than brachial plasma in controls, although this was not significant. Our analysis limited univariate comparisons, but the lack of overlap in error bars (Figure 2D) suggests that, in controls,  $E_2$  levels could be higher in the jugular vein than brachial vein. These data are consistent with previous findings that the brain is the predominant source of circulating  $E_2$  in male zebra finches (69). However, neither brachial nor jugular plasma  $E_2$  concentrations changed in response to fasting or subsequent refeeding. Thus circulating  $E_2$  is also an unlikely regulator of aggression in a food-related context.

### **Neurosteroids, fasting, and aggression**

The brain can synthesize active steroids (ie, neurosteroids) de novo from cholesterol as well as from circulating inactive precursors, such as DHEA (70, 71). Both  $E_2$  and DHEA concentrations are typically higher in brain than plasma (69, 72) and are maintained even after castration, adrenalectomy, or gonadal regression (73). However, the factors that regulate neurosteroid production remain elusive.

Neural  $E_2$  synthesis maintains aggression when systemic sex steroid levels are low, such as during the non-breeding season (17, 37). Here,  $E_2$  was present throughout

the male zebra finch brain, as previously reported (41), with highest levels in the caudomedial nidopallium, an aromatase-rich forebrain area involved in auditory processing of song (74). Statistical considerations limited our analyses to regions with reliably detectable levels of  $E_2$ . Interestingly, fasting specifically increased  $E_2$  levels within 3 regions of the social behavior network: the PAG, VTA, and VMH. Furthermore, in the PAG and VTA,  $E_2$  concentrations were rapidly reduced to baseline levels by re-feeding, suggesting a time course consistent with a rapid nongenomic effect of  $E_2$  on aggression.

The PAG regulates multiple behaviors, including vocalization, mating, fight or flight, and aggressive behaviors (75). The VTA is part of the mesolimbic dopaminergic system involved in motivation and reproductive behavior (38), and the VTA innervates the PAG and other social behavior regions (76). Aromatase is present in cell bodies in VTA and in fibers in PAG (77, 78), and aromatase activity is modulated by dopamine (79). In zebra finches, dopaminergic cells of the VTA respond to conspecific interactions (80), and in rats, VTA neurons are involved in the anticipation of feeding (81). Both the VTA and PAG project to regions regulating aggressive behavior, including the BST and LS (79, 82). The VMH is involved in satiety (83, 84), and although  $E_2$  can suppress food intake (85), this action does not appear to involve the VMH (86). In song sparrows (*M. melodia*),  $E_2$  treatment decreases immunoreactivity for phosphorylated tyrosine hydroxylase, the rate-limiting enzyme in catecholamine synthesis in the VMH (87). Together, these observations suggest higher  $E_2$  levels within the PAG, VTA, and VMH may alter dopaminergic signaling to efferent regions that regulate aggression. Future research manipulating local  $E_2$  levels in these regions is vital to elucidate the role of  $E_2$  in mediating food-related aggression.

## Acknowledgments

We thank Jennifer Ferris, Michael Xu, Mary Shen, Kang Nian Yap, Anne Cheng, and Alice Chan for help with animal care; Dr. Chunqi Ma, Matthew Taves, and Dr. Sarah Heimovics for laboratory assistance and advice; Dr. Joanne Weinberg for the use of her  $\gamma$ -counter; and Matthew Taves and Valerie J. Fokidis for comments on the manuscript.

Address all correspondence and requests for reprints to: H. Bobby Fokidis, Ph.D., Rollins College, 1000 Holt Avenue, Winter Park, Florida 32789–4499, USA E-mail: hfokidis@rollins.edu.

This work was supported by a postdoctoral fellowship (to H.B.F.) from the Natural Sciences and Engineering Research Council of Canada, a predoctoral fellowship (to N.H.P.) from

the National Science Foundation, and an Operating Grant (to K.K.S.) from the Canadian Institutes of Health Research.

Disclosure Summary: The authors have nothing to disclose.

## References

1. Scotti MA, Schmidt KL, Newman AE, Bonu T, Soma KK, Demas GE. Aggressive encounters differentially affect serum dehydroepiandrosterone and testosterone concentrations in male Siberian hamsters (*Phodopus sungorus*). *Horm Behav*. 2009;56:376–381.
2. Enstrom DA, Ketterson ED, Nolan V. Testosterone and mate choice in the dark-eyed junco. *Anim Behav*. 1997;54:1135–1146.
3. Rilling JK, Winslow JT, Kilts CD. The neural correlates of mate competition in dominant male rhesus Macaques. *Biol Psychiatry*. 2004;56:364–375.
4. Lu SF, Mo Q, Hu S, Garipha C, Simon NG. Dehydroepiandrosterone upregulates neural androgen receptor level and transcriptional activity. *J Neurobiol*. 2003;57:163–171.
5. Holmes MM, Wade J. Testosterone regulates androgen receptor immunoreactivity in the copulatory, but not courtship, neuromuscular system in adult male green anoles. *J Neuroendocrinol*. 2005;17:560–569.
6. Balthazart J, Castagna C, Ball GF. Aromatase inhibition blocks the activation and sexual differentiation of appetitive male sexual behavior in Japanese quail. *Behav Neurosci*. 1997;111:381–397.
7. Fusani L, Metzendorf R, Hutchison JB, Gahr M. Aromatase inhibition affects testosterone-induced masculinization of song and the neural song system in female canaries. *J Neurobiol*. 2003;54:370–379.
8. de la Mora AN, Drummond H, Wingfield JC. Hormonal correlates of dominance and starvation-induced aggression in chicks of the blue-footed booby. *Ethology*. 1996;102:748–761.
9. Militzer K. Social dominance and bodily conditions in small groups of male and female laboratory rats of known familiarity. *Int J Mammal Biol*. 1995;60:97–111.
10. Coleman SL, Mellgren RL. Social enhancement and interference of food finding in zebra finches (*Taeniopygia guttata*). *J Comp Psychol*. 1997;111:242–250.
11. Gelezen C, Collier DA, Campbell IC, Oppelaar H, Kas MJ. Behavioral, physiological, and molecular differences in response to dietary restriction in three inbred mouse strains. *Am J Physiol Endocrinol Metab*. 2006;291:E574–E581.
12. Challet E, Lemaho Y, Robin JP, Malan A, Cherel Y. Involvement of corticosterone in the fasting-induced rise in protein-utilization and locomotor activity. *Pharmacol Biochem Behav*. 1995;50:405–412.
13. Dallman MF, Akana SF, Bhatnagar S, et al. Starvation: early signals, sensors, and sequelae. *Endocrinology*. 1999;140:4015–4023.
14. Morimoto Y, Arisue K, Yamamura Y. Relationship between circadian rhythm of food intake and that of plasma corticosterone and effect of food restriction on circadian adrenocortical rhythm in the rat. *Neuroendocrinology*. 1977;23:212–222.
15. Warne JP, Akana SF, Ginsberg AB, Horneman HF, Pecoraro NC, Dallman MF. Disengaging insulin from corticosterone: roles of each on energy intake and disposition. *Am J Physiol Regul Integr Comp Physiol*. 2009;296:R1366–R1375.
16. Sapolsky RM, Romero LM, Munck AU. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev*. 2000;21:55–89.
17. Soma KK, Scotti MA, Newman AE, Charlier TD, Demas GE. Novel mechanisms for neuroendocrine regulation of aggression. *Frontiers in Neuroendocrinology*. 2008;29:476–489.
18. Soma KK, Alday NA, Hau M, Schlinger BA. Dehydroepiandrosterone metabolism by  $3\beta$ -hydroxysteroid dehydrogenase/8 $\beta$ -8 $\alpha$  isomerase in adult zebra finch brain: sex difference and rapid effect of stress. *Endocrinology*. 2004;145:1668–1677.

19. Kroboth PD, Salek FS, Pittenger AL, Fabian TJ, Frye RF. DHEA and DHEA-S: A review. *J Clin Pharmacol*. 1999;39:327–348.
20. Oberbeck R, Benschop RJ, Jacobs R, et al. Endocrine mechanisms of stress-induced DHEA-secretion. *J Endocrinol Invest*. 1998;21:148–153.
21. Boonstra R, Lane JE, Boutin S, et al. Plasma DHEA levels in wild, territorial red squirrels: Seasonal variation and effect of ACTH. *Gen Comp Endocrinol*. 2008;158:61–67.
22. Newman AE, Soma KK. Corticosterone and dehydroepiandrosterone in songbird plasma and brain: effects of season and acute stress. *Eur J Neurosci*. 2009;29:1905–1914.
23. Newman AE, Soma KK. Aggressive interactions differentially modulate local and systemic levels of corticosterone and DHEA in a wild songbird. *Horm Behav*. 2011;60:389–396.
24. Newman AE, Pradhan DS, Soma KK. Dehydroepiandrosterone and corticosterone are regulated by season and acute stress in a wild songbird: Jugular versus Brachial plasma. *Endocrinology*. 2008;149:2537–2545.
25. Grasfeder LL, Gaillard S, Hammes SR, et al. Fasting-Induced hepatic production of DHEA is regulated by PGC-1 $\alpha$ , ERR $\alpha$ , and HNF4 $\alpha$ . *Mol Endocrinol*. 2009;23:1171–1182.
26. Bauer M, Hamm AC, Bonaus M, et al. Starvation response in mouse liver shows strong correlation with life-span-prolonging processes. *Physiol Genomics*. 2004;17:230–244.
27. London SE, Monks DA, Wade J, Schlinger BA. Widespread capacity for steroid synthesis in the avian brain and song system. *Endocrinology*. 2006;147:5975–5987.
28. Braun EJ, Sweazea KL. Glucose regulation in birds. *Comp Biochem Physiol B-Biochem Mol Biol*. 2008;151:1–9.
29. Prinzinger R, Pressmar A, Schleucher E. Body temperature in birds. *Comp Biochem Physiol a-Physiology*. 1991;99:499–506.
30. Fokidis HB, Hurley L, Rogowski C, Sweazea K, Deviche P. Effects of captivity and body condition on plasma corticosterone, locomotor behavior, and plasma metabolites in curve-billed thrashers. *Physiol Biochem Zool*. 2011;84:595–606.
31. Lynn SE, Breuner CW, Wingfield JC. Short-term fasting affects locomotor activity, corticosterone, and corticosterone binding globulin in a migratory songbird. *Horm Behav*. 2003;43:150–157.
32. Lynn SE, Stamps TB, Barrington WT, Weida N, Hudak CA. Food, stress, and reproduction: Short-term fasting alters endocrine physiology and reproductive behavior in the zebra finch. *Horm Behav*. 2010;58:214–222.
33. Ritschard M, Brumm H. Zebra finch song reflects current food availability. *Evol Ecol*. 2012;26:801–812.
34. Balthazart J, Foidart A, Surlémont C, Vockel A, Harada N. Distribution of aromatase in the brain of the Japanese quail, ring dove, and zebra finch: an immunocytochemical study. *J Comp Neurol*. 1990;301:276–288.
35. Shen P, Schlinger BA, Campagnoni AT, Arnold AP. An atlas of aromatase mRNA expression in the zebra finch brain. *J Comp Neurol*. 1995;360:172–184.
36. Metzdorf R, Gahr M, Fusani L. Distribution of aromatase, estrogen receptor, and androgen receptor mRNA in the forebrain of songbirds and nonsongbirds. *J Comp Neurol*. 1999;407:115–129.
37. Soma KK, Tramontin AD, Wingfield JC. Oestrogen regulates male aggression in the non-breeding season. *Proc Soc Biol*. 2000;267:1089–1096.
38. O'Connell LA, Hofmann HA. The Vertebrate mesolimbic reward system and social behavior network: A comparative synthesis. *J Comp Neurol*. 2011;519:3599–3639.
39. Goodson JL, Kabelik D. Dynamic limbic networks and social diversity in vertebrates: From neural context to neuromodulatory patterning. *Front Neuroendocrinol*. 2009;30:429–441.
40. Taves MD, Schmidt KL, Ruhr IM, Kapusta K, Prior NH, Soma KK. Steroid concentrations in plasma, whole blood and brain: effects of saline perfusion to remove blood contamination from brain. *Plos One*. 2010;5:e15727.
41. Charlier TD, Po KW, Newman AE, Shah AH, Saldanha CJ, Soma KK. 17 $\beta$ -Estradiol levels in male zebra finch brain: combining Palkovits punch and an ultrasensitive radioimmunoassay. *Gen Comp Endocrinol*. 2010;167:18–26.
42. Palkovits M. Isolated removal of hypothalamic or other brain nuclei of the rat. *Brain Res*. 1973;59:449–450.
43. Vockel A, Pröve E, Balthazart J. Sex- and age-related differences in the activity of testosterone-metabolizing enzymes in microdissected nuclei of the zebra finch brain. *Brain Res*. 1990;511:291–302.
44. Nixdorf-Bergweiler BE, Bischof HJ. *Stereotaxic Atlas of the Brain of the Zebra Finch, Taeniopygia guttata, with Special Emphasis on Telencephalic Visual and Song System Nuclei in Transverse and Sagittal Sections*. Bethesda, MD: National Library of Medicine NCB; 2007.
45. Taves MD, Ma C, Heimovics SA, Saldanha CJ, Soma KK. 2011 Measurement of steroid concentrations in brain tissue: methodological considerations. *Front Endocrinol (Lausanne)*. 2011;2:39.
46. Newman AE, Chin EH, Schmidt KL, Bond L, Wynne-Edwards KE, Soma KK. Analysis of steroids in songbird plasma and brain by coupling solid phase extraction to radioimmunoassay. *Gen Comp Endocrinol*. 2008;155:503–510.
47. Newman SW. 1999 The medial extended amygdala in male reproductive behavior: a node in the mammalian social behavior network. In: McGinty JF, ed. *Advancing from the Ventral Striatum to the Extended Amygdala: Implications for Neuropsychiatry and Drug Abuse: In Honor of Lennart Heimer*; New York, New York Academy of Sciences. 1999; 242–257.
48. Fokidis HB, des Roziers MB, Sparr R, Rogowski C, Sweazea K, Deviche P. Unpredictable food availability induces metabolic and hormonal changes independent of food intake in a sedentary songbird. *J Exp Biol*. 2012;215:2920–2930.
49. Widmaier EP, Rosen K, Abbott B. Free fatty acids activate the hypothalamic-pituitary-adrenocortical axis in rats. *Endocrinology*. 1992;131:2313–2318.
50. Sashkov VA, Sel'verova NB, Morenkov ED, Ermakova IV, Buraya TI. Relationship between level of neuroactive steroids in the brain and behavior and anxiety of maturing and mature female rats during the estrous cycle. *Neurochem J*. 2009;3:35–43.
51. Klusonová P, Kucka M, Ergas P, Mikšík I, Bryndová J, Pácha J. Cloning of chicken 11 $\beta$ -hydroxysteroid dehydrogenase type 1 and its tissue distribution. *J Steroid Biochem Mol Biol*. 2008;111:217–224.
52. Schmidt KL, Chin EH, Shah AH, Soma KK. Cortisol and corticosterone in immune organs and brain of European starlings: developmental changes, effects of restraint stress, comparison with zebra finches. *Am J Physiol Regul Integr Comp Physiol*. 2009;297:R42–R51.
53. Sainio EL, Lehtola T, Roininen P. Radioimmunoassay of total and free corticosterone in rat plasma: measurement of the effect of different doses of corticosterone. *Steroids*. 1988;51:609–622.
54. Birrenkott GP, Wiggins ME. Determination of dexamethasone and corticosterone half-lives in male broilers. *Poult Sci*. 1984;63:1064–1068.
55. Andrews MH, Wood SA, Windle RJ, Lightman SL, Ingram CD. Acute glucocorticoid administration rapidly suppresses basal and stress-induced hypothalamo-pituitary-adrenal axis activity. *Endocrinology*. 2012;153:200–211.
56. Windle RJ, Wood SA, Shanks N, Lightman SL, Ingram CD. Ultradian rhythm of basal corticosterone release in the female rat: dynamic interaction with the response to acute stress. *Endocrinology*. 1998;139:443–450.
57. Hess GD, Riegler GD. Effects of chronic ACTH stimulation on adrenocortical function in young and aged rats. *Am J Physiol*. 1972;222:1458–1461.
58. Breuner CW, Greenberg AL, Wingfield JC. Noninvasive corticosterone treatment rapidly increases activity in Gambel's white-crowned



- sparrows (*Zonotrichia leucophrys gambelii*). *Gen Comp Endocrinol*. 1998;111:386–394.
59. Riddle MC, McDaniel PA. Renal 11  $\beta$ -hydroxysteroid dehydrogenase activity is enhanced by ramipril and captopril. *J Clin Endocrinol Metab*. 1994;78:830–834.
  60. Welberg LA, Thrivikraman KV, Plotsky PM. Chronic maternal stress inhibits the capacity to up-regulate placental 11 $\beta$ -hydroxysteroid dehydrogenase type 2 activity. *J Endocrinol*. 2005;186:R7–R12.
  61. Lavallée B, Provost PR, Kahwash Z, Nestler JE, Belanger A. Effect of insulin on serum levels of dehydroepiandrosterone metabolites in men. *Clin Endocrinol (Oxf)*. 1997;46:93–100.
  62. Kahwash Z, O'Brien R, Nestler JE. Insulin acutely increases the metabolic-clearance rate (MCR) of dehydroepiandrosterone (DHEA) in men. *Clin Res*. 1993;41:A800–A800.
  63. Leblanc M, Labrie C, Bélanger A, Candas B, Labrie F. Bioavailability and pharmacokinetics of dehydroepiandrosterone in the cynomolgus monkey. *J Clin Endocrinol Metab*. 2003;88:4293–4302.
  64. Wang DY, Bulbrook RD, Sneddon A, Hamilton T. The metabolic clearance rates of dehydroepiandrosterone, testosterone and their sulphate esters in man, rat and rabbit. *J Endocrinol*. 1967;38:307–318.
  65. Longcope C. Dehydroepiandrosterone metabolism. *J Endocrinol*. 1996;150:S125–S127.
  66. Baulieu EE. Dehydroepiandrosterone (DHEA): a fountain of youth? *J Clin Endocrinol Metab*. 1996;81:3147–3151.
  67. Bird CE, Murphy J, Boroomand K, Finnis W, Dressel D, Clark AF. Dehydroepiandrosterone: kinetics of metabolism in normal men and women. *J Clin Endocrinol Metab*. 1978;47:818–822.
  68. David M, Giraldeau LA. Zebra finches in poor condition produce more and consume more food in a producer-scrouter game. *Behav Ecol*. 2012;23:174–180.
  69. Schlinger BA, Arnold AP. Brain is the major site of estrogen synthesis in a male songbird. *Proc Natl Acad Sci USA*. 1991;88:4191–4194.
  70. Baulieu EE, Robel P. Neurosteroids: a new brain-function? *J Steroid Biochem Mol Biol*. 1990;37:395–403.
  71. Mellon SH, Deschepper CF. Neurosteroid biosynthesis: Genes for adrenal steroidogenic enzymes are expressed in the brain. *Brain Res*. 1993;629:283–292.
  72. Boross M, Bodrogi L, Poteczin E, Feher T. Age-Dependent Change in Brain and Blood Content of Neurosteroids Dehydroepiandrosterone (DHEA) and Dehydroepiandrosterone-Sulfate (DHEA-S) in Rat. 40128 Bologna: Monduzzi Editore;1993.
  73. Mensah-Nyagan AG, Do-Rego JL, Beaujean D, Luu-The V, Pelletier G, Vaudry H. Neurosteroids: Expression of steroidogenic enzymes and regulation of steroid biosynthesis in the central nervous system. *Pharmacol Rev*. 1999;51:63–81.
  74. Tremere LA, Pinaud R. Brain-generated estradiol drives long-term optimization of auditory coding to enhance the discrimination of communication signals. *J Neurosci*. 2011;31:3271–3289.
  75. Kingsbury MA, Kelly AM, Schrock SE, Goodson JL. Mammal-like organization of the avian midbrain central gray and a reappraisal of the intercollicular nucleus. *Plos One*. 2011;6:e20720.
  76. Schrott A, Kabai P. ABCD: a functional database for the avian brain. *J Neurosci Methods*. 2008;167:393–395.
  77. Saldanha CJ, Tuerk MJ, Kim YH, Fernandes AO, Arnold AP, Schlinger BA. Distribution and regulation of telencephalic aromatase expression in the zebra finch revealed with a specific antibody. *J Comp Neurol*. 2000;423:619–630.
  78. Absil P, Ritters LV, Balthazart J. Preoptic aromatase cells project to the mesencephalic central gray in the male Japanese quail (*Coturnix japonica*). *Horm Behav*. 2001;40:369–383.
  79. Absil P, Baillien M, Ball GF, Panzica GC, Balthazart J. The control of preoptic aromatase activity by afferent inputs in Japanese quail. *Brain Res Rev*. 2001;37:38–58.
  80. Kabelik D, Schrock SE, Ayres LC, Goodson JL. Estrogenic regulation of dopaminergic neurons in the opportunistically breeding zebra finch. *Gen Comp Endocrinol*. 2011;173:96–104.
  81. Skibicka KP, Hansson C, Alvarez-Crespo M, Friberg PA, Dickson SL. Ghrelin directly targets the ventral tegmental area to increase food motivation. *Neuroscience*. 2011;180:129–137.
  82. Bailhache T, Balthazart J. The catecholaminergic system of the quail brain: immunocytochemical studies of dopamine  $\beta$ -hydroxylase and tyrosine-hydroxylase. *J Comp Neurol*. 1993;329:230–256.
  83. Egawa M, Inoue S, Satoh S, Takamura Y, Nagai K, Nakagawa H. Acute and chronic effects of VMH lesions on circadian rhythms in food intake and metabolites. *Brain Res Bull*. 1993;31:293–299.
  84. Resch JM, Boisvert JP, Hourigan AE, Mueller CR, Yi SS, Choi S. Stimulation of the hypothalamic ventromedial nuclei by pituitary adenylate cyclase-activating polypeptide induces hypophagia and thermogenesis. *Am J Physiol Regul Integr Comp Physiol*. 2011;301:R1625–R1634.
  85. Archer ZA, Findlay PA, McMillen SR, Rhind SM, Adam CL. Effects of nutritional status and gonadal steroids on expression of appetite-regulatory genes in the hypothalamic arcuate nucleus of sheep. *J Endocrinol*. 2004;182:409–419.
  86. Santollo J, Torregrossa AM, Eckel LA. Estradiol acts in the medial preoptic area, arcuate nucleus, and dorsal raphe nucleus to reduce food intake in ovariectomized rats. *Horm Behav*. 2011;60:86–93.
  87. Heimovics SA, Prior NH, Maddison CJ, Soma KK. Rapid and widespread effects of 17 $\beta$ -estradiol on intracellular signaling in the male songbird brain: a seasonal comparison. *Endocrinology*. 2012;153:1364–1376.