

# MATH 240 Formulas and Definitions

<b>Terminology</b>	<b>1</b>
<b>1.1 Systems of Linear Equations</b>	<b>1</b>
<b>1.2 Row Reduction and Echelon Forms</b>	<b>2</b>
<b>1.3 Vector Equations</b>	<b>2</b>
<b>1.4 The Matrix Equation</b>	<b>3</b>
<b>1.5 Solution Sets of Linear Systems</b>	<b>4</b>
<b>1.7 Linear Independence</b>	<b>5</b>

## Terminology

Linear Equation / Linear System:

- Can be written in the form  $a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$

Linear combination:

- Vector  $\mathbf{y}$  defined by:  $\mathbf{y} = c_1\mathbf{v}_1 + \cdots + c_p\mathbf{v}_p$

## 1.1 Systems of Linear Equations

Elementary row operations

### ELEMENTARY ROW OPERATIONS

1. (Replacement) Replace one row by the sum of itself and a multiple of another row.<sup>1</sup>
2. (Interchange) Interchange two rows.
3. (Scaling) Multiply all entries in a row by a nonzero constant.

## 1.2 Row Reduction and Echelon Forms

Criteria for echelon form and reduced echelon form

A rectangular matrix is in **echelon form** (or **row echelon form**) if it has the following three properties:

1. All nonzero rows are above any rows of all zeros.
2. Each leading entry of a row is in a column to the right of the leading entry of the row above it.
3. All entries in a column below a leading entry are zeros.

If a matrix in echelon form satisfies the following additional conditions, then it is in **reduced echelon form** (or **reduced row echelon form**):

4. The leading entry in each nonzero row is 1.
5. Each leading 1 is the only nonzero entry in its column.

Existence and uniqueness theorem

### Existence and Uniqueness Theorem

A linear system is consistent if and only if the rightmost column of the augmented matrix is *not* a pivot column—that is, if and only if an echelon form of the augmented matrix has *no* row of the form

$$[0 \quad \cdots \quad 0 \quad b] \quad \text{with } b \text{ nonzero}$$

If a linear system is consistent, then the solution set contains either (i) a unique solution, when there are no free variables, or (ii) infinitely many solutions, when there is at least one free variable.

## 1.3 Vector Equations

Algebraic properties of vectors

### Algebraic Properties of $\mathbb{R}^n$

For all  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  in  $\mathbb{R}^n$  and all scalars  $c$  and  $d$ :

- |                                                                                                                             |                                                              |
|-----------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------|
| (i) $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$                                                                     | (v) $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$ |
| (ii) $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$                                      | (vi) $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$         |
| (iii) $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$                                                      | (vii) $c(d\mathbf{u}) = (cd)\mathbf{u}$                      |
| (iv) $\mathbf{u} + (-\mathbf{u}) = -\mathbf{u} + \mathbf{u} = \mathbf{0}$ ,<br>where $-\mathbf{u}$ denotes $(-1)\mathbf{u}$ | (viii) $1\mathbf{u} = \mathbf{u}$                            |

## Definition of a Span

If  $\mathbf{v}_1, \dots, \mathbf{v}_p$  are in  $\mathbb{R}^n$ , then the set of all linear combinations of  $\mathbf{v}_1, \dots, \mathbf{v}_p$  is denoted by  $\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  and is called the **subset of  $\mathbb{R}^n$  spanned** (or **generated**) by  $\mathbf{v}_1, \dots, \mathbf{v}_p$ . That is,  $\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is the collection of all vectors that can be written in the form

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p$$

with  $c_1, \dots, c_p$  scalars.

## 1.4 The Matrix Equation

### The matrix equation

If  $A$  is an  $m \times n$  matrix, with columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$ , and if  $\mathbf{x}$  is in  $\mathbb{R}^n$ , then the **product of  $A$  and  $\mathbf{x}$** , denoted by  $A\mathbf{x}$ , is the **linear combination of the columns of  $A$  using the corresponding entries in  $\mathbf{x}$  as weights**; that is,

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n$$

### Equivalence of matrix equation, vector equation, and linear equation

If  $A$  is an  $m \times n$  matrix, with columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$ , and if  $\mathbf{b}$  is in  $\mathbb{R}^m$ , the matrix equation

$$A\mathbf{x} = \mathbf{b} \tag{4}$$

has the same solution set as the vector equation

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n = \mathbf{b} \tag{5}$$

which, in turn, has the same solution set as the system of linear equations whose augmented matrix is

$$\begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n & \mathbf{b} \end{bmatrix} \tag{6}$$

### Equivalence of statements

Let  $A$  be an  $m \times n$  matrix. Then the following statements are logically equivalent. That is, for a particular  $A$ , either they are all true statements or they are all false.

- For each  $\mathbf{b}$  in  $\mathbb{R}^m$ , the equation  $A\mathbf{x} = \mathbf{b}$  has a solution.
- Each  $\mathbf{b}$  in  $\mathbb{R}^m$  is a linear combination of the columns of  $A$ .
- The columns of  $A$  span  $\mathbb{R}^m$ .
- $A$  has a pivot position in every row.

### Properties of the Matrix-Vector Product $A\mathbf{x}$

If  $A$  is an  $m \times n$  matrix,  $\mathbf{u}$  and  $\mathbf{v}$  are vectors in  $\mathbb{R}^n$ , and  $c$  is a scalar, then:

- $A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v}$ ;
- $A(c\mathbf{u}) = c(A\mathbf{u})$ .

### Row-Vector rule for computing $A\mathbf{x}$

#### Row-Vector Rule for Computing $A\mathbf{x}$

If the product  $A\mathbf{x}$  is defined, then the  $i$ th entry in  $A\mathbf{x}$  is the sum of the products of corresponding entries from row  $i$  of  $A$  and from the vector  $\mathbf{x}$ .

## 1.5 Solution Sets of Linear Systems

### Nontrivial solution to the homogeneous equation $A\mathbf{x} = \mathbf{0}$

The homogeneous equation  $A\mathbf{x} = \mathbf{0}$  has a nontrivial solution if and only if the equation has at least one free variable.

### Parametric Vector form

$$\mathbf{x} = s\mathbf{u} + t\mathbf{v} \quad (s, t \text{ in } \mathbb{R})$$

### Translate a solution

Suppose the equation  $A\mathbf{x} = \mathbf{b}$  is consistent for some given  $\mathbf{b}$ , and let  $\mathbf{p}$  be a solution. Then the solution set of  $A\mathbf{x} = \mathbf{b}$  is the set of all vectors of the form  $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ , where  $\mathbf{v}_h$  is any solution of the homogeneous equation  $A\mathbf{x} = \mathbf{0}$ .

## 1.7 Linear Independence

Definition of linear independence

An indexed set of vectors  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  in  $\mathbb{R}^n$  is said to be **linearly independent** if the vector equation

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \dots + x_p\mathbf{v}_p = \mathbf{0}$$

has only the trivial solution. The set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  is said to be **linearly dependent** if there exist weights  $c_1, \dots, c_p$ , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0} \quad (2)$$

Two linearly dependent vectors

A set of two vectors  $\{\mathbf{v}_1, \mathbf{v}_2\}$  is linearly dependent if at least one of the vectors is a multiple of the other. The set is linearly independent if and only if neither of the vectors is a multiple of the other.

Characterization of Linearly Dependent Sets

### Characterization of Linearly Dependent Sets

An indexed set  $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  of two or more vectors is linearly dependent if and only if at least one of the vectors in  $S$  is a linear combination of the others. In fact, if  $S$  is linearly dependent and  $\mathbf{v}_1 \neq \mathbf{0}$ , then some  $\mathbf{v}_j$  (with  $j > 1$ ) is a linear combination of the preceding vectors,  $\mathbf{v}_1, \dots, \mathbf{v}_{j-1}$ .

A Set with more vectors than entries is linearly dependent

If a set contains more vectors than there are entries in each vector, then the set is linearly dependent. That is, any set  $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  in  $\mathbb{R}^n$  is linearly dependent if  $p > n$ .

A set with the zero vector is linearly dependent

If a set  $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  in  $\mathbb{R}^n$  contains the zero vector, then the set is linearly dependent.