

# Topology Note

Alex

June 19, 2022



# Contents

<b>1</b>	<b>Topology Spaces and Continuous Function</b>	<b>5</b>
1.1	Basic Definition of Topology . . . . .	5
1.2	Basis for a Topology . . . . .	6
1.2.1	Exercise . . . . .	7
1.3	The Order Topology . . . . .	7
1.4	The Product Topology . . . . .	8

**Definitions****K**

K-topology on  $\mathbb{R}$ , 6

**L**

larger, 5

strictly larger, 5

lower limit topology on  $\mathbb{R}$ , 6

**B**

basis, 6

**T**

topology, 5

topology generated by basis, 6

topology space, 5

trivial topology, 5

**C**

coarser, 5

strictly coarser, 5

**R**

ray, 8

closed ray, 8

open ray, 8

**O**

open set, 5

open sets, 5

order topology, 7

**I**

interval, 7

closed interval, 7

half-open interval, 7

open interval, 7

**D**

discrete topology, 5

**F**

finer, 5

strictly finer, 5

finite complement topology, 5

**S**

smaller, 5

strictly smaller, 5

standard topology on  $\mathbb{R}$ , 6

subbasis, 7

# Chapter 1

## Topology Spaces and Continuous Function

### 1.1 Basic Definition of Topology

**Definition 1.1.1** (topology). A **topology** on a set  $\mathbb{X}$  is a collection  $\mathbb{T}$  of subsets of  $\mathbb{X}$  having the following properties:

- $\emptyset$  and  $\mathbb{X}$  are in  $\mathbb{T}$
- The union of the elements of any sub collection of  $\mathbb{T}$  is in  $\mathbb{T}$
- The intersection of the elements of any **finite** sub collection of  $\mathbb{T}$  is in  $\mathbb{T}$

**Definition 1.1.2** (topology space). A **topological space** is a set  $\mathbb{X}$  for which a topology  $\mathbb{T}$  has been specified.

**Definition 1.1.3** (open set). A **open set**  $\mathbb{U}$  is a subset of  $\mathbb{X}$  that belongs to a topology  $\mathbb{T}$  of  $\mathbb{X}$ .

**Definition 1.1.4** (open sets). A topology can also be called a **open sets**

**Definition 1.1.5** (discrete topology). The set of all subsets of a set  $\mathbb{X}$  formed a topology called **discrete topology**

**Definition 1.1.6** (trivial topology). The set consisting the set  $\mathbb{X}$  and  $\emptyset$  only formed a topology of  $\mathbb{X}$  called **trivial topology**

**Definition 1.1.7** (finite complement topology). Let  $\mathbb{X}$  be a set. Let  $\mathbb{T}_f$  be the collection of all subsets  $\mathbb{U}$  of  $\mathbb{X}$  such that  $\mathbb{X} - \mathbb{U}$  either if a **finite**<sup>1</sup> of is all of  $\mathbb{X}$ . Then  $\mathbb{T}_f$  is a topology on  $\mathbb{X}$ , called the .

**Definition 1.1.8** (finer, larger, strictly finer, strictly larger, coarser, smaller, strictly coarser, strictly smaller, comparable). Let  $\mathbb{T}$  and  $\mathbb{T}'$  be two topology on a given set  $\mathbb{X}$ . If  $\mathbb{T}$  is a subset of  $\mathbb{T}'$ , we say that  $\mathbb{T}'$  is **finer** or **larger** than  $\mathbb{T}$ . If  $\mathbb{T}$  is a proper subset of  $\mathbb{T}'$ , we say that  $\mathbb{T}'$  is **strictly finer** or **strictly larger** than  $\mathbb{T}$ . We also say that  $\mathbb{T}$  is **coarser** or **smaller** or **strictly coarser** or **strictly smaller** than  $\mathbb{T}'$ . We say that  $\mathbb{T}$  and  $\mathbb{T}'$  is **comparable** if either  $\mathbb{T}$  is a subset of  $\mathbb{T}'$  or  $\mathbb{T}'$  is a subset of  $\mathbb{T}$ .

---

<sup>1</sup>The set  $\mathbb{U}$  can form a topology because of the definition of topology is intersection of finite sub collection. If this can be intersection of infinite sub collection,  $\mathbb{U}$  will not be a topology.

## 1.2 Basis for a Topology

**Definition 1.2.1** (basis). If  $\mathbb{X}$  is a set, a **basis** for a topology on  $\mathbb{X}$  is a collection  $\mathbb{B}$  of subsets of  $\mathbb{X}$  (called **basis elements**) such that:

- For each  $x \in \mathbb{X}$ , there is at least one basis element  $B$  containing  $x$
- If  $x$  belongs to the intersection of two basis elements  $B_1$  and  $B_2$ , then there is another element  $x \in B_3 \in \mathbb{B}$  such that  $B_3 \subseteq B_1 \cap B_2$

**Definition 1.2.2** (topology generated by basis). Let  $\mathbb{B}$  be a basis on  $\mathbb{X}$ . Let  $\mathbb{U}$  be a set containing all subsets  $U$  of  $\mathbb{X}$  such that for each element  $x \in U$ , there is  $B \in \mathbb{B}$  that  $x \in B \subseteq U$ . Such  $\mathbb{U}$  formed a topology on  $\mathbb{X}$ , called **topology  $\mathbb{T}$  generated by  $\mathbb{B}$**

**Lemma 1.2.1.** Let  $\mathbb{X}$  be a set. Let  $\mathbb{B}$  be a basis for a topology  $\mathbb{T}$  on  $\mathbb{X}$ . Then  $\mathbb{T}$  equals to the set of all possible unions of elements of  $\mathbb{B}$ .

*Proof.* Let set  $\mathbb{U}$  be the set of all possible unions of elements of  $\mathbb{B}$ . For any  $U \in \mathbb{U}$ .  $U = \cup B$  <sup>2</sup> for some  $B \in \mathbb{B}$ . Thus, for every  $x \in U$ , there exist a  $B' \in \mathbb{B}$  that  $x \in B' \subseteq U$ . Thus,  $U \in \mathbb{T}$ .

Conversely, for any  $U \in \mathbb{T}$ . For any  $x \in U$ , let  $x \in B_x \in \mathbb{B}$ . Then,  $U = \cup_{x \in U} B_x$ . Thus,  $U \in \mathbb{U}$ .

Therefore,  $\mathbb{U}$  equals to  $\mathbb{T}$ . □

**Lemma 1.2.2.** <sup>3</sup> Let  $\mathbb{X}$  be a topological space. Suppose that  $\mathbb{C}$  is a collection of open sets of  $\mathbb{X}$  such that for each open set  $U$  of  $\mathbb{X}$  and each  $x \in U$ , there is an element  $C \in \mathbb{C}$  such that  $x \in C \subseteq U$ . Then  $\mathbb{C}$  is a basis for the topology of  $\mathbb{X}$ .

**Lemma 1.2.3.** <sup>4</sup> Let  $\mathbb{B}$  and  $\mathbb{B}'$  be basis for the topologies  $\mathbb{T}$  and  $\mathbb{T}'$ , respectively, on  $\mathbb{X}$ . Then the following are equivalent:

- $\mathbb{T}'$  is finer than  $\mathbb{T}$
- For each  $x \in \mathbb{X}$  and each basis element  $B \in \mathbb{B}$  containing  $x$ , there is a basis element  $B' \in \mathbb{B}'$  such that  $x \in B' \subseteq B$ .

**Definition 1.2.3** (standard topology on the real line). Let be  $\mathbb{B} = \{B | B = \{x | a < x < b\}, a < b, a \in \mathbb{R}, b \in \mathbb{R}\}$ .  $\mathbb{B}$  formed a basis on real line. The topology generated by  $\mathbb{B}$  is called the **standard topology on the real line** <sup>5</sup>.

**Definition 1.2.4** (lower limit topology on the real line). Let be  $\mathbb{B} = \{B | B = \{x | a \leq x < b\}, a < b, a \in \mathbb{R}, b \in \mathbb{R}\}$ .  $\mathbb{B}$  formed a basis on real line. The topology generated by  $\mathbb{B}$  is called the **lower limit topology on the real line**. When  $\mathbb{R}$  is given this topology, we denote it by  $\mathbb{R}_l$ .

**Definition 1.2.5** (K-topology on the real line). Let be  $\mathbb{B} = \{B | B = \{x | a < x < b\}, a < b, a \in \mathbb{R}, b \in \mathbb{R}\}$ . Let  $K = \{x | x = \frac{1}{n}, n \in \mathbb{Z}_+\}$ .  $\mathbb{B} \cup \{B - K | B \in \mathbb{B}\}$  formed a basis on real line. The topology generated by  $\mathbb{B}$  is called the **K-topology on the real line**. When  $\mathbb{R}$  is given this topology, we denote it by  $\mathbb{R}_K$ .

**Lemma 1.2.4.** <sup>6</sup> The topologies  $\mathbb{R}_l$  and  $\mathbb{R}_K$  is strictly finer than the standard topology on  $\mathbb{R}$ .

<sup>2</sup>Note that this expression may not be unique.

<sup>3</sup>We omit the proof of this lemma as it is obvious.

<sup>4</sup>We omit the proof of this lemma as it is obvious.

<sup>5</sup>Whenever we consider  $\mathbb{R}$ , we shall suppose it is given this topology unless we specifically state otherwise.

<sup>6</sup>We omit the proof of this lemma as it is obvious.

**Lemma 1.2.5.** *The topologies of  $\mathbb{R}_l$  and  $\mathbb{R}_\mathbb{K}$  is not comparable.*

*Proof.* Let  $\mathbb{T}_l$  and  $\mathbb{T}_\mathbb{K}$  be topologies of  $\mathbb{R}_l$  and  $\mathbb{R}_\mathbb{K}$  respectively. Let  $K = \{x | x = \frac{1}{n}, n \in \mathbb{Z}_+\}$ .

We first proof that  $\mathbb{T}_l$  is not finer than  $\mathbb{T}_\mathbb{K}$ . Let  $U = \{x | -1 < x < 1\} - K, x = 0$ . If there exist  $B = \{x | a \leq x < b\} \in \mathbb{T}_l$  such that  $x \in B \subseteq U$ , then  $0 < b < 1$ . Thus, there exist  $n \in \mathbb{Z}_+$  that  $0 < \frac{1}{n} < b$ . Thus  $B$  is not a subset of  $U$ .

Then we proof that  $\mathbb{T}_\mathbb{K}$  is not finer than  $\mathbb{T}_l$ . Let  $U' = \{x | a' \leq x < b'\}$ . If there exist  $B' = \{x | a'' < x < b''\} \text{ or } \{x | a'' < x < b''\} - K$  such that  $a' \in B \subseteq U$ . Thus  $a'' < a < b''$ . Thus there exist  $c$  that  $a'' < x < a, x \in B, x \notin U'$ . Thus  $B' \not\subseteq U'$ .

Thus the topologies of  $\mathbb{R}_l$  and  $\mathbb{R}_\mathbb{K}$  is not comparable.  $\square$

**Definition 1.2.6** (subbasis). *A **subbasis**  $\mathbb{S}$  for a topology on  $\mathbb{X}$  is a collection of subsets of  $\mathbb{X}$  whose union equals  $\mathbb{X}$ . The **topology generated by the subbasis**  $\mathbb{S}$  is defined to be the collection  $\mathbb{T}$ <sup>7</sup> of all unions of finite intersections of elements of  $\mathbb{S}$ .*

### 1.2.1 Exercise

1. Show that if  $\mathbb{A}$  is a basis for a topology on  $\mathbb{X}$ , then the topology generated by  $\mathbb{A}$  equals the intersection of all topologies on  $\mathbb{X}$  that contain  $\mathbb{A}$ . Prove the same if  $\mathbb{A}$  is a subbasis.

*Proof.* As a subbasis is also a basis, we will directly prove the case of subbasis here.

Let  $\mathbb{S} = \{\mathbb{T}_\alpha\}$  be set contain all the topologies that contain  $\mathbb{A}$ . Let  $\mathbb{T}$  be the topology that  $\mathbb{A}$  generated. Let  $\mathbb{T}' = \cap \mathbb{T}_\alpha$ .<sup>8</sup>

First,  $\mathbb{A} \subseteq \mathbb{T}_\alpha$ . Thus,  $\mathbb{T} \subseteq \mathbb{T}_\alpha$ . Thus,  $\mathbb{T} \subseteq \mathbb{T}'$ .

Also,  $\mathbb{A} \subseteq \mathbb{T}$ . Thus,  $\mathbb{T} \in \mathbb{S}$ . Thus,  $\mathbb{T}' \subseteq \mathbb{T}$ .

Thus,  $\mathbb{T} = \mathbb{T}'$   $\square$

## 1.3 The Order Topology

**Definition 1.3.1** (interval). *Let  $\mathbb{X}$  is a set having a simple order relation  $<$ . Given elements  $a$  and  $b$  of  $\mathbb{X}$  such that  $a < b$ , there are four subsets of  $\mathbb{X}$  that are called **intervals** determined by  $a$  and  $b$ :*

- $(a, b) = \{x | a < x < b\}$
- $(a, b] = \{x | a < x \leq b\}$
- $[a, b) = \{x | a \leq x < b\}$
- $[a, b] = \{x | a \leq x \leq b\}$

$(a, b)$  is called an **open interval** on  $\mathbb{X}$ .  $[a, b]$  is called an **closed interval** on  $\mathbb{X}$ .  $(a, b]$  and  $[a, b)$  is called **half-open intervals**.

**Definition 1.3.2** (order topology).<sup>9</sup> *Let  $\mathbb{X}$  be a set with a simple order relation; assume  $\mathbb{X}$  has more than one element. Let  $\mathbb{B}$  be the collection of all sets of the following types:*

- All open intervals  $(a, b)$  in  $\mathbb{X}$ .

<sup>7</sup>It is obvious that  $\mathbb{T}$  is a topology, we just omit the proof here.

<sup>8</sup>It is obvious that  $\mathbb{T}'$  is also a topology, we just omit the proof here.

<sup>9</sup>The standard topology on  $\mathbb{R}$  is an order topology derived from the usual order on  $\mathbb{R}$ .

- All intervals of the form  $[a_0, b)$ , where  $a_0$  is the smallest element(if exist) of  $\mathbb{X}$ .
- All intervals of the form  $(a, b_0]$ , where  $b_0$  is the largest element(if exist) of  $\mathbb{X}$ .

The collection  $\mathbb{B}$  formed a basis for a topology on  $\mathbb{X}$ , which is called the order topology.

**Definition 1.3.3** (ray).<sup>1011</sup> If  $\mathbb{X}$  is an ordered set, and  $a$  is an element of  $\mathbb{X}$ , there are four subsets of  $\mathbb{X}$  that are called **rays** determined by  $a$ :

- $(a, +\infty) = \{x | x > a\}$
- $(-\infty, a) = \{x | x < a\}$
- $[a, +\infty) = \{x | x \geq a\}$
- $(-\infty, a] = \{x | x \leq a\}$

$(a, +\infty)$  and  $(-\infty, a)$  are called **open rays**.  $[a, +\infty)$  and  $(-\infty, a]$  are called **closed rays**.

## 1.4 The Product Topology

**Definition 1.4.1** (product topology). Let  $\mathbb{X}$  and  $\mathbb{Y}$  be topological spaces. The **product topology** on  $\mathbb{X} \times \mathbb{Y}$  having a basis  $\mathbb{B}$  containing all sets of the form  $U \times V$ , where  $U$  and  $V$  is open sets of  $\mathbb{X}$  and  $\mathbb{Y}$  respectively.

**Theorem 1.4.1.**<sup>12</sup> If  $\mathbb{B}$  and  $\mathbb{C}$  is basis for the topology of  $\mathbb{X}$  and  $\mathbb{Y}$  respectively, then the collection

$$\mathbb{D} = \{B \times C | B \in \mathbb{B} \text{ and } C \in \mathbb{C}\}$$

is a basis for the topology of  $\mathbb{X} \times \mathbb{Y}$

**Definition 1.4.2** (projection). Let  $\pi_1 : \mathbb{X} \times \mathbb{Y} \rightarrow \mathbb{X}$  be defined by the equation:

$$\pi_1(x, y) = x$$

Let  $\pi_2 : \mathbb{X} \times \mathbb{Y} \rightarrow \mathbb{Y}$  be defined by the equation:

$$\pi_2(x, y) = y$$

The maps  $\pi_1$  and  $\pi_2$  are called the **projections** of  $\mathbb{X} \times \mathbb{Y}$  onto its first and second factors, respectively.

**Theorem 1.4.2.**<sup>13</sup> The collection

$$\mathbb{S} = \{\pi_1^{-1}(U) | U \text{ open in } \mathbb{X}\} \cup \{\pi_2^{-1}(V) | V \text{ open in } \mathbb{Y}\}$$

is a subbasis for the product topology on  $\mathbb{X} \times \mathbb{Y}$ .

<sup>10</sup>open rays are always open sets in the order topology

<sup>11</sup>the open rays also formed a subbasis of the order topology

<sup>12</sup>We omit the proof of this lemma as it is obvious.

<sup>13</sup>We omit the proof of this lemma as it is obvious.