

Alessandro Scala



Quantum Abstract Interpretation

Seminar for the **Introduction to Quantum Computing** course

Università di Pisa
Dipartimento di Informatica

Pisa, 24 Luglio 2023

Roadmap



Introduction

As quantum computing advances, we would like to have some means to prove correctness properties on quantum programs, *especially* since quantum programming is counterintuitive.



Reasons

The naive way to check properties of a program is to run it and observe its behaviour.



Reasons

The naive way to check properties of a program is to run it and **observe** its behaviour.

We **cannot observe** the state of a quantum program!



Reasons

The naive way to check properties of a program is to run it and **observe** its behaviour.

We **cannot observe** the state of a quantum program!

Could **simulation** on a classical machine solve this issue?



Reasons

The naive way to check properties of a program is to run it and **observe** its behaviour.

We **cannot observe** the state of a quantum program!

Could **simulation** on a classical machine solve this issue?

No: **exponential** space and time cost.



Example

$$n_{qubits} = 1$$

$$|0\rangle \langle 0|$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$2^2 = 4 \text{ complex numbers}$$



Example

$$n_{\text{qubits}} = 2$$

$$|00\rangle \langle 00|$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$2^4 = 16 \text{ complex numbers}$$



Example

$$n_{\text{qubits}} = 3$$

$$|000\rangle \langle 000|$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$2^6 = 64 \text{ complex numbers}$$



Example

$$n_{\text{qubits}} = 300$$

$$|0\rangle^{\otimes 300} \langle 0|^{\otimes 300}$$

?????



Example

$$n_{\text{qubits}} = 300$$

$$|0\rangle^{\otimes 300} \langle 0|^{\otimes 300}$$

$2^{600} = 4149515568809929585124078636911611510124462322424368$
 $999956573296906528114129081463997070489471037942881978866113$
 $007891823951510754117753078868748341139636870611818034015095$
 23685376

Bigger than the number of atoms in the universe.



Reasons

The naive way to check properties of a program is to run it and **observe** its behaviour.

We **cannot observe** the state of a quantum program!

Could **simulation** on a classical machine solve this issue?

No: **exponential** space and time cost.



Reasons

The naive way to check properties of a program is to run it and **observe** its behaviour.

We **cannot observe** the state of a quantum program!

Could **simulation** on a classical machine solve this issue?

No: **exponential** space and time cost.

Solution: abstract interpretation



Ingredients

- Abstract domain
 - Abstraction function
 - Concretization function
 - Abstract operations
- Assertions



Density Matrix

Instead of dealing with a state $|\phi\rangle$ in vector form, we use its *density matrix*:

$$\rho_\phi = |\phi\rangle \langle\phi| \quad (\text{For a pure state})$$

- positive semi-definite
- $\text{Tr}(\rho) = 1$
- projection ($P = P^\dagger = P^2$)



Density Matrix

Instead of dealing with a state $|\phi\rangle$ in vector form, we use its *density matrix*:

$$\rho_\phi = |\phi\rangle \langle\phi| \quad (\text{For a pure state})$$

- positive semi-definite
- $\text{Tr}(\rho) = 1$
- projection ($P = P^\dagger = P^2$)

Example:

$$\begin{aligned} |\beta_{00}\rangle &= \frac{|00\rangle + |11\rangle}{\sqrt{2}} \\ \rho_{\beta_{00}} &= |\beta_{00}\rangle \langle\beta_{00}| = \frac{1}{2}(|00\rangle + |11\rangle)(\langle 00| + \langle 11|) \\ &= \frac{1}{2}(|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|) \\ &= \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} \end{aligned}$$



Reduced Density Matrix

Suppose we have a composite quantum system $AB = A \otimes B$, and we want to focus our attention on a state $|\phi\rangle \in AB$ with respect to the subsystem A .

$$A = \mathbb{C}^{2^n} \times \mathbb{C}^{2^n} \quad B = \mathbb{C}^{2^m} \times \mathbb{C}^{2^m}$$

$$AB = (\mathbb{C}^{2^n} \times \mathbb{C}^{2^n}) \otimes (\mathbb{C}^{2^m} \times \mathbb{C}^{2^m})$$

$$Tr_B[\rho] : AB \rightarrow A$$

$$Tr_A[\rho] : AB \rightarrow B$$

$$Tr_B[\alpha \otimes \beta] = \alpha \cdot Tr(\beta)$$


$$Tr_A[\alpha \otimes \beta] = Tr(\alpha) \cdot \beta$$

$$Tr_S[\rho + \sigma] = Tr_S[\rho] + Tr_S[\sigma] \text{ (Linearity)}$$

Alternatively:

$$Tr_B[\rho] = \sum_{v=0}^{2^m} (I_A \otimes \langle v|) \rho (I_A \otimes |v\rangle) \quad Tr_A[\rho] = \sum_{v=0}^{2^n} (\langle v| \otimes I_B) \rho (|v\rangle \otimes I_B)$$

Where v labels vectors of an orthonormal basis of the subspace we are tracing out.



Example

$$A = C^2 \times C^2 \quad B = C^2 \times C^2 \quad AB = A \otimes B$$

$$\rho_{\beta_{00}} = |\beta_{00}\rangle \langle \beta_{00}| = \frac{|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|}{2}$$

$$\text{Tr}_B[\rho_{\beta_{00}}] =$$



Example

$$A = \mathbb{C}^2 \times \mathbb{C}^2 \quad B = \mathbb{C}^2 \times \mathbb{C}^2 \quad AB = A \otimes B$$

$$\rho_{\beta_{00}} = |\beta_{00}\rangle \langle \beta_{00}| = \frac{|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|}{2}$$

$$Tr_B[\rho_{\beta_{00}}] = \frac{(Tr_B[|00\rangle \langle 00|] + Tr_B[|00\rangle \langle 11|] + Tr_B[|11\rangle \langle 00|] + Tr_B[|11\rangle \langle 11|])}{2}$$



Example

$$A = \mathcal{C}^2 \times \mathcal{C}^2 \quad B = \mathcal{C}^2 \times \mathcal{C}^2 \quad AB = A \otimes B$$

$$\rho_{\beta_{00}} = |\beta_{00}\rangle \langle \beta_{00}| = \frac{|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|}{2}$$

$$\begin{aligned} Tr_B[\rho_{\beta_{00}}] &= \frac{(Tr_B[|00\rangle \langle 00|] + Tr_B[|00\rangle \langle 11|] + Tr_B[|11\rangle \langle 00|] + Tr_B[|11\rangle \langle 11|])}{2} \\ &= \frac{(|0\rangle \langle 0| \cdot \langle 0|0\rangle) + (|0\rangle \langle 1| \cdot \langle 0|1\rangle) + (|1\rangle \langle 0| \cdot \langle 1|0\rangle) + (|1\rangle \langle 1| \cdot \langle 1|1\rangle)}{2} \end{aligned}$$



Example

$$A = \mathbb{C}^2 \times \mathbb{C}^2 \quad B = \mathbb{C}^2 \times \mathbb{C}^2 \quad AB = A \otimes B$$

$$\rho_{\beta_{00}} = |\beta_{00}\rangle \langle \beta_{00}| = \frac{|00\rangle \langle 00| + |00\rangle \langle 11| + |11\rangle \langle 00| + |11\rangle \langle 11|}{2}$$

$$\begin{aligned} Tr_B[\rho_{\beta_{00}}] &= \frac{(Tr_B[|00\rangle \langle 00|] + Tr_B[|00\rangle \langle 11|] + Tr_B[|11\rangle \langle 00|] + Tr_B[|11\rangle \langle 11|])}{2} \\ &= \frac{(|0\rangle \langle 0| \cdot \langle 0|0\rangle) + (|0\rangle \langle 1| \cdot \langle 0|1\rangle) + (|1\rangle \langle 0| \cdot \langle 1|0\rangle) + (|1\rangle \langle 1| \cdot \langle 1|1\rangle)}{2} \\ &= \frac{|0\rangle \langle 0| + |1\rangle \langle 1|}{2} \end{aligned}$$



Loss of precision

Computing a reduced density matrix **discards information!**

$$\rho_{\beta_{00}} = \frac{|00\rangle\langle 00| + |00\rangle\langle 11| + |11\rangle\langle 00| + |11\rangle\langle 11|}{2} \quad (\text{Pure state})$$

$$\rho_2 = \frac{|00\rangle\langle 00| + |01\rangle\langle 01| + |10\rangle\langle 10| + |11\rangle\langle 11|}{4} \quad (\text{Mixed state})$$



Loss of precision

Computing a reduced density matrix **discards information!**

$$\rho_{\beta_{00}} = \frac{|00\rangle\langle 00| + |00\rangle\langle 11| + |11\rangle\langle 00| + |11\rangle\langle 11|}{2} \quad (\text{Pure state})$$

$$\rho_2 = \frac{|00\rangle\langle 00| + |01\rangle\langle 01| + |10\rangle\langle 10| + |11\rangle\langle 11|}{4} \quad (\text{Mixed state})$$

$$\text{Tr}_B[\rho_{\beta_{00}}] = \frac{|0\rangle\langle 0| + |1\rangle\langle 1|}{2} = \text{Tr}_B[\rho_2]$$

The partial traces of two different initial states can be equal.

Moreover, for a state $\rho \in A \otimes B$, even if we know $\text{Tr}_B[\rho]$ and $\text{Tr}_A[\rho]$, we cannot uniquely determine ρ .



Linear Subspaces

Each projection P corresponds to a linear subspace $\{v \mid Pv = v\}$.

The support of a matrix P is the subspace orthogonal to its kernel, i.e., the set $\{v \mid Pv \neq 0\}$.



Abstract Domain

$$\mathcal{D} = \mathbb{C}^{2^n} \otimes \mathbb{C}^{2^n}, \quad S = (s_1, \dots, s_m), \quad 1 \leq m \leq 2^n, \quad s_i \subseteq [n]$$

$$AbsDom(S) = \left\{ (P_{s_1}, \dots, P_{s_m}) \mid P_{s_i} \text{ is a projection in } \mathbb{C}^{2^{|s_i|}} \otimes \mathbb{C}^{2^{|s_i|}} \right\}$$

Intuitively, given a tuple S of sets of qubits, an abstract state $\bar{\sigma} \in AbsDom(S)$ is a tuple of projections over those qubits.

Special case:

$$T = ([n]) \implies AbsDom(T) = \mathcal{D}$$



Fineness Relation

Let $S = (s_1, \dots, s_m)$ and $T = (t_1, \dots, t_m)$ (with $1 \leq m \leq 2^n$), then:

$$\underbrace{S \sqsubseteq T}_{\text{"T is finer than S"}} \triangleq \forall i \in [m]. s_i \subseteq t_i$$

T is “more concrete” than S .

Least element: $\perp = (\emptyset, \dots, \emptyset)$.

Greatest element: $\top = ([n], \dots, [n])$.

$AbsDom(\top)$ corresponds to a state so abstract that it holds no information at all.

$AbsDom(\top)$ corresponds to tuples where every projection is a concrete state.



Abstraction Function

$$S \sqsubseteq T \triangleq \forall i \in [m]. s_i \subseteq t_i$$

$$\alpha_{T \rightarrow S} : AbsDom(T) \rightarrow AbsDom(S)$$

$$\alpha_{T \rightarrow S}(Q_{t_1}, \dots, Q_{t_m}) = (P_{s_1}, \dots, P_{s_m})$$

$$P_{s_i} = \bigcap_{t_j. s_i \subseteq t_j} supp(Tr_{t_j \setminus s_i}[Q_{t_j}])$$

Given an abstract state $\bar{\tau} \in AbsDom(T) = (Q_{t_1}, \dots, Q_{t_m})$, we want to compute $\bar{\sigma} \in AbsDom(S) = (P_{s_1}, \dots, P_{s_m})$. For each $i \in [m]$:

- ① Find all Q_{t_j} s such that $s_i \subseteq t_j$. We know that at least one exists (for $j = i$), since $S \sqsubseteq T$.
- ② For each Q_{t_j} found, trace out the bits in t_j that are not in s_i .
- ③ Compute the support of the traced matrices (to preserve the structure of projections).
- ④ Compute the intersection of the supports.



Concretization Function

$$S \sqsubseteq T \triangleq \forall i \in [m]. s_i \subseteq t_i$$

$$\gamma_{S \rightarrow T} : AbsDom(S) \rightarrow AbsDom(T)$$

$$\gamma_{S \rightarrow T}(P_{s_1}, \dots, P_{s_m}) = (Q_{t_1}, \dots, Q_{t_m})$$

$$Q_{t_j} = \bigcap_{s_i. s_i \subseteq t_j} P_{s_i} \otimes I_{t_j \setminus s_i}$$

Given an abstract state $\bar{\sigma} \in AbsDom(S) = (P_{s_1}, \dots, P_{s_m})$, we want to compute $\bar{\tau} \in AbsDom(T) = (Q_{t_1}, \dots, Q_{t_m})$. For each $j \in [m]$:

- ① Find all P_{s_i} s such that $s_i \subseteq t_j$. We know at least one exists (for $i = j$), since $S \sqsubseteq T$.
- ② Extend the projection to the space of all qubits in t_j , by computing the tensor product with the identity matrix.
- ③ Compute the intersection of the extended projections.



Order Relation on Abstract States

$$1 \leq m \leq 2^n, \quad S = (s_1, \dots, s_m), \quad \forall i \in [m]. s_i \subseteq [n]$$

$$\bar{\sigma} \in \text{AbsDom}(S) = (P_{s_1}, \dots, P_{s_m}), \quad \bar{\tau} \in \text{AbsDom}(S) = (Q_{s_1}, \dots, Q_{s_m})$$

$$\bar{\sigma} \sqsubseteq \bar{\tau} \triangleq \forall i \in [m]. \underbrace{P_{s_i} \subseteq Q_{s_i}}$$

Subspace interpretation
of projections



Monotonicity

$$S \supseteq T$$
$$\forall \bar{\sigma}, \bar{\tau} \in \text{AbsDom}(T). \bar{\sigma} \sqsubseteq \bar{\tau} \implies \alpha_{T \rightarrow S}(\bar{\sigma}) \sqsubseteq \alpha$$



Computing the Projection of a Support

To compute a projection corresponding to $\text{supp}(A)$, we:

- 1 take the rows $\{r_1, \dots, r_n\}$ of A ;
- 2 extract an orthonormal set of vectors $\{b_1, \dots, b_n\}$ that span the same subspace as the rows;
- 3 create the matrix $B = \begin{pmatrix} b_1 \\ \dots \\ b_n \end{pmatrix}$;
- 4 return BB^\dagger .



Computing the Projection of an Intersection

$$\{P_1, \dots, P_k\}, \quad \forall i \in [k]. P_i = C^n \times C^n$$

$$\bigcap_{i \in [k]} P_i = I_n - \text{supp}(kl_n - \sum_{i \in [k]} P_i)$$

