

Study of spin-orbit motion in the Frozen and Quasi-Frozen rings

October 8, 2018

1 Spin decoherence effect in a perfectly aligned storage ring

We study decoherence for two major reasons:

1. it reduces the measurement time frame;
2. vertical plane decoherence is a source of systematic error. [3, p. 8]

Here we consider the decoherence resulting from the beam particles' spin tune difference.

The spin dynamics in a storage ring is governed by the FT-BMT equation, [1, p. 4] In the standard spinor formalism, spin evolution is described by a rotation matrix, whose eigenvector is the spin precession axis, \bar{n} , and eigenvalue the spin tune, ν_s . For particles traveling along the design orbit of a perfectly aligned, flat storage ring, the equilibrium direction of \bar{n} is vertical. [2, p. 1362] The spin tune expressions in the electric and magnetic fields are: [3, p. 8]

$$\nu_s^E = \left(\frac{1}{\gamma^2 - 1} - G \right) \gamma \beta^2, \quad (1)$$

$$\nu_s^B = \gamma G, \quad (2)$$

where $G = \frac{g-2}{2}$ is the magnetic moment anomaly, $\gamma = (1 - \beta^2)^{-1/2}$ is the particle's Lorentz factor.

The phase stability principle requires that orbit lengthening is accompanied by a corresponding equilibrium energy shift [4]. This means that particles traveling on different orbits will have differing spin tunes, hence decoherence. Orbit lengthening is due to two reasons: a) betatron motion, b) momentum deviation.

2 Decoherence suppression via sextupole fields

Sextupole fields influence orbit lengthening via two paths:

1. by affecting the momentum compaction factor: $\delta\alpha_1 = -\frac{S \cdot D_0^3}{L}$
2. by directly changing the orbit length: $\frac{\Delta L}{L} = \mp \frac{S \cdot D_0 \cdot \beta_{x,y} \varepsilon_{x,y}}{L}$.

3 Simulation design

4 Simulation results

References

- [1] Saleev A, Nikolaev NN, Rathmann F, Augustyniak W, Bagdasarian Z, Bai M, et al. Spin tune mapping as a novel tool to probe the spin dynamics in storage rings. Physical Review Accelerators and Beams [Internet]. 2017 Jul 7 [cited 2018 Oct 8];20(7). Available from: <http://arxiv.org/abs/1703.01295>
- [2] Barber D P, Vogt M, Hoffstätter G H. The Amplitude Dependent Spin Tune and the Invariant Spin Field in High Energy Proton Accelerators. <http://accelconf.web.cern.ch/AccelConf/e98/PAPERS/THP35G.PDF>

- [3] Yuriy Senichev. Search for the Charged Particle Electric Dipole Moments in Storage Rings. In: 25th Russian Particle Accelerator Conference (RuPAC16), St Petersburg, Russia, November 21-25, 2016 [Internet]. JACOW, Geneva, Switzerland; 2017 [cited 2017 Apr 5]. p. 610. Available from: <http://accelconf.web.cern.ch/AccelConf/rupac2016/papers/mozmh03.pdf>
- [4] Senichev Y, Zyuzin D. SPIN TUNE DECOHERENCE EFFECTS IN ELECTRO- AND MAGNETOSTATIC STRUCTURES. In: Beam Dynamics and Electromagnetic Fields [Internet]. Changhai, China: JACoW; 2013 [cited 2017 Jul 31]. p. 2579–2581. Available from: <https://accelconf.web.cern.ch/accelconf/IPAC2013/papers/wepea036.pdf>