

Proportional Size Distribution (PSD)

Alex J. Benecke

January 10, 2018

I will use a Chi-Square test to see if there is a difference between PSD between years (2013 - 2017).

Data Preparation

Load Data

```
lmbs <- read.csv("Data/Clean-Data/2012-2017_nearshore-survey-largemouth-bass_Stock_CLEAN.csv") %>%  
  filter(Year >= 2013) %>% arrange(Year, FID, Length)  
lmbs$fyrr <- as.factor(lmbs$fyrr)
```

```
unique(lmbs$Year) ### See that there is no 2012
```

```
## [1] 2013 2014 2015 2016 2017
```

View Data

```
(lmbs.LF <- xtabs(~Year+gcat,data=lmbs))
```

```
##      gcat  
## Year preferred quality stock  
## 2013      16      41     41  
## 2014      18      57     65  
## 2015      15      38     14  
## 2016      11      44     52  
## 2017       9      15     11
```

Chi-Squares Test

Is there a difference in the number of fish in each gabelhouse length category (PSD-X) during the years 2013 - 2017?

```
chisq.test(lmbs.LF)
```

```
##  
## Pearson's Chi-squared test  
##  
## data:  lmbs.LF  
## X-squared = 20.055, df = 8, p-value = 0.01013
```

This seems to suggest that the proportional stock distribution (PSD) is different for largemouth bass between years ($X^2 = 20.055$, $df = 8$, $P = 0.01013$).

Where are the differences in PSD X-Y for each year?

Creating a table of percent of fish in each gabelhouse length interval (PSD X-Y) in each year.

```
round(prop.table(lmbs.LF,margin=1)*100,0)
```

```
##          gcat
## Year    preferred quality stock
##  2013         16      42    42
##  2014         13      41    46
##  2015         22      57    21
##  2016         10      41    49
##  2017         26      43    31
```

Remarkably the percent of quality fish is the same for 2014 and 2016. the percent of quality length fish is very similar between 2013 (42%), 2014 (41%), 2016 (41%), and 2017 (43%). This trend is almost the same for stock length fish (2013 42%, 2014 46%, 2016 49%), however 2015 and 2017 have a smaller percentage of stock length fish (21% and 31% respectively). This may be explained by more variability among years for this length category, reduced sampling efficiency for smaller fish, or unstable recruitment. The year 2015 seems to break the trend with a greater percentage of preferred (22%) and quality (57%) length individuals than other years.

- 1) Could this be some sort of sampling bias?
- 2) Could this be a result of sampling different sites? where the 2015 sites more suitable for LMB?

Compare PSD-Q between years 2013 - 2017

```
lmb$ %<>% mutate(gcatQ=mapvalues(gcat,
                                from=c("stock","quality","preferred"),
                                to=c("quality-", "quality+", "quality+")),
               gcatQ=droplevels(gcatQ))

(lmb.LFQ <- xtabs(~Year+gcatQ,data = lmb$))
```

```
##          gcatQ
## Year    quality+ quality-
##  2013         57      41
##  2014         75      65
##  2015         53      14
##  2016         55      52
##  2017         24      11
```

```
chisq.test(lmb.LFQ)
```

```
##
## Pearson's Chi-squared test
##
## data:  lmb.LFQ
## X-squared = 16.815, df = 4, p-value = 0.0021
```

```
(ps.Q <- c(chisq.test(lmb.LFQ[c(1,2),])$p.value, ### 2013-2014
           chisq.test(lmb.LFQ[c(1,3),])$p.value, ### 2013-2015
           chisq.test(lmb.LFQ[c(1,4),])$p.value, ### 2013-2016
           chisq.test(lmb.LFQ[c(1,5),])$p.value, ### 2013-2017
           chisq.test(lmb.LFQ[c(2,3),])$p.value, ### 2014-2015
           chisq.test(lmb.LFQ[c(2,4),])$p.value, ### 2014-2016
           chisq.test(lmb.LFQ[c(2,5),])$p.value, ### 2014-2017
           chisq.test(lmb.LFQ[c(3,4),])$p.value, ### 2015-2016
           chisq.test(lmb.LFQ[c(3,5),])$p.value, ### 2015-2017
           chisq.test(lmb.LFQ[c(4,5),])$p.value)) ### 2016-2017
```

```
(p.val.Q <- p.adjust(ps.Q))
```

```
## [1] "13-14" "13-15" "13-16" "13-17" "14-15" "14-16" "14-17" "15-16"
## [9] "15-17" "16-17"
```

```
##      Year p-value Adjusted p
## 1  13-14  0.5694      1.0000
## 2  13-15  0.0084      0.0675
## 3  13-16  0.4060      1.0000
## 4  13-17  0.3781      1.0000
## 5  14-15  0.0007      0.0064
## 6  14-16  0.8338      1.0000
## 7  14-17  0.1583      0.9500
## 8  15-16  0.0005      0.0046
## 9  15-17  0.3515      1.0000
## 10 16-17  0.1144      0.8007
```

The PSD-Q of largemouth bass is different for at least one of the years during 2013 - 2016 (Chi-Squared, $X^2 = 16.815$, $df = 4$, $p = 0.0021$). The adjusted p-values show a *significant difference* in PSD-Q between years 2014 - 2015 ($p = 0.0064$) and 2015 - 2016 ($p = 0.0046$). The PSD-Q is not different between any other years. However 2013 and 2015 may be different ($p = 0.0675$).

Compare PSD-P between years 2013 - 2017

```
lmbs %<>% mutate(gcatP=mapvalues(gcat,
                                from=c("stock","quality","preferred"),
                                to=c("preferred-", "preferred-", "preferred+")),
              gcatP=droplevels(gcatP))

(lmb.LFP <- xtabs(~Year+gcatP,data = lmbs))
```

```
##      gcatP
## Year preferred+ preferred-
## 2013      16      82
## 2014      18     122
## 2015      15      52
## 2016      11      96
## 2017       9      26
```

```
chisq.test(lmb.LFP)
```

```
##
## Pearson's Chi-squared test
##
## data:  lmb.LFP
## X-squared = 8.2649, df = 4, p-value = 0.08234
```

```
(ps.P <- c(chisq.test(lmb.LFP[c(1,2),])$p.value, ### 2013-2014
           chisq.test(lmb.LFP[c(1,3),])$p.value, ### 2013-2015
           chisq.test(lmb.LFP[c(1,4),])$p.value, ### 2013-2016
           chisq.test(lmb.LFP[c(1,5),])$p.value, ### 2013-2017
           chisq.test(lmb.LFP[c(2,3),])$p.value, ### 2014-2015
           chisq.test(lmb.LFP[c(2,4),])$p.value, ### 2014-2016
           chisq.test(lmb.LFP[c(2,5),])$p.value, ### 2014-2017
```

```
chisq.test(lmb.LFP[c(3,4),])$p.value, ### 2015-2016
chisq.test(lmb.LFP[c(3,5),])$p.value, ### 2015-2017
chisq.test(lmb.LFP[c(4,5),])$p.value)) ### 2016-2017
```

```
## Warning in chisq.test(lmb.LFP[c(4, 5), ]): Chi-squared approximation may be
## incorrect
```

```
(p.val.P <- p.adjust(ps.P))
```

```
## [1] "13-14" "13-15" "13-16" "13-17" "14-15" "14-16" "14-17" "15-16"
## [9] "15-17" "16-17"
```

```
##      Year p-value Adjusted p
## 1  13-14  0.5724      1.0000
## 2  13-15  0.4378      1.0000
## 3  13-16  0.2837      1.0000
## 4  13-17  0.3329      1.0000
## 5  14-15  0.1212      0.8485
## 6  14-16  0.6716      1.0000
## 7  14-17  0.1048      0.8387
## 8  15-16  0.0498      0.4565
## 9  15-17  0.8964      1.0000
## 10 16-17  0.0456      0.4565
```

The PSD-P of largemouth bass is not different for any years during 2013 - 2017 (Chi-Squared, $X^2 = 8.26$, $df = 4$, $p = 0.08$). The adjusted p-values show no difference in the PSD-P between years (2013 - 2017).
T