



**A Comparison of Neuroimaging Software Procedures and
a Contour Inference Method for Group-level Task-fMRI
Analysis**

by

Alexander Bowring

St Catherine's College

Submitted to the University of Oxford

for the degree of

Doctor of Philosophy

Nuffield Department of Population Health

October 2019



Contents

Acknowledgments	iv
Declarations	v
Abstract	vi
1 Introduction	1
2 Background	2
2.1 The Study of Brain Function	2
2.2 Magnetic Resonance Imagery (MRI)	2
2.3 Task-based functional Magnetic Resonance Imagery (t-fMRI)	2
2.3.1 Overview	2
2.3.2 Pre-processing	2
2.3.3 Statistical Analysis: Subject-level	2
2.3.4 Statistical Analysis: Group-level	2
2.4 Neuroimaging Software Packages	2
2.4.1 SPM	2
2.4.2 FSL	2
2.4.3 AFNI	2
2.5 Reproducibility of fMRI Results	2
3 Exploring the Impact of Analysis Software on Task-fMRI Results	3
3.1 Data and Analysis Methods	3
3.1.1 Study Description and Data Source	3
3.1.2 Data Analyses	3
3.1.3 Comparison Methods	3
3.1.4 Permutation Test Methods	3
3.2 Results	3
3.2.1 Cross-Software Variability for Parametric Inference	3

3.2.2	Cross-Software Variability for Non-Parametric Inference . . .	3
3.2.3	Intra-Software Variability, Parametric vs Non-Parametric . . .	3
3.3	Reproducibility	3
3.3.1	Scripting of Analysis and Figures	3
3.3.2	Results Sharing	3
3.4	Discussion	3
3.4.1	Limitations	3
3.5	Conclusion	3
4	Spatial Confidence Sets for Task-fMRI Inference	4
4.1	Introduction	5
4.2	Theory	5
4.2.1	Overview	5
4.2.2	The Wild Bootstrap Method for Computation of k	5
4.3	Method	5
4.3.1	Simulations	5
4.3.2	Implementation of Contour Inference	5
4.3.3	2D Simulations	5
4.3.4	3D Simulations	5
4.3.5	Application to Human Connectome Project Data	5
4.4	Results	5
4.4.1	2D Simulations	5
4.4.2	3D Simulations	5
4.4.3	Human Connectome Project	5
4.5	Discussion	5
4.5.1	Limitations	5
4.6	Conclusion	5
4.7	Toolbox	5
5	Contour Inference for Cohen's d	6
5.1	Theory	6
5.1.1	Transforming the Residual Field	6
5.2	Method	6
5.2.1	2D Simulations	6
5.2.2	3D Simulations	6
5.2.3	Application to UK Biobank Data	6
5.3	Results	6
5.3.1	2D Simulations	6

5.3.2	3D Simulations	6
5.3.3	UK Biobank Data	6
5.3.4	Comparison to Traditional Inference Procedures	6
5.4	Discussion	6
5.4.1	Limitations	6
5.5	Conclusion	6
6	Conclusion and Future Work	7

Acknowledgments

Declarations

I, Alexander Bowring, hereby declare that except where specific reference is made to the work of others, the contents of this dissertation are original and have not been submitted in whole or in part for consideration for any other degree or qualification in these, or any other Universities. This dissertation is the result of my own work and includes nothing which is the outcome of work done in collaboration, except where specifically indicated in the text.

- The work presented in Chapter 3 has been published in the *NeuroImage* journal (Bowring et al., 2018). This work was presented at the *Organization for Human Brain Mapping* (OHBM) Annual Meetings in 2017 and 2018. At the OHBM 2018 Annual Meeting, this work was the recipient of an oral presentation and Merit Abstract Award.
- The work presented in Chapter 4 has been published in the *NeuroImage* journal (Bowring et al., 2018). This work was presented at the OHBM Annual Meeting in 2017, where it was the recipient of an oral presentation.
- The work presented in Chapter 5 is based on a pre-printed manuscript.

Alexander Bowring

September 2019

Abstract

Over the last three decades, Functional Magnetic Resonance Imaging (fMRI) has rapidly progressed to become the primary tool for human brain mapping. Recently however, considerable attention within the field has been directed towards data-sharing and open science initiatives. This has been driven by a growing apprehension about the reproducibility of findings within the neuroimaging literature, amid concerns that current inference procedures are often misused or misinterpreted causing the overall scientific conclusions to become distorted. With increasing availability to population-size neuroimaging databases, the limitations of such inference methods are also being made apparent: with sufficient statistical power, the null-hypothesis of no activation can be rejected over essentially the entire brain.

One aspect specific to neuroimaging pinpointed as a cause for poor reproducibility is the high flexibility of a typical fMRI analysis workflow. Across the pipeline, researchers can choose from a range of parameters to determine how each analysis step is carried out. In the first part of this thesis, we investigate how the choice of software package used to conduct a statistical analysis can influence the group-level results of a task-fMRI study. We use publicly shared data from three published task-fMRI studies, and reanalyze each study within the three main neuroimaging software packages, AFNI, FSL and SPM, using parametric and nonparametric inference. All information on how to process, analyze, and model each dataset we obtain from the publications. We use a variety of quantitative and qualitative comparison methods to gauge the scale of variability in our results and assess fundamental difference between each software package. While qualitatively we find broad similarities between packages, we also discover marked differences, such as Dice similarity coefficient

values ranging from 0.000 to 0.743 in comparisons of thresholded statistic maps between software. We discuss the challenges involved in our replication attempt, while also utilizing open science tools in a considerable effort to make our own research reproducible.

CHAPTER 1

Introduction

2.1 The Study of Brain Function

2.2 Magnetic Resonance Imagery (MRI)

2.3 Task-based functional Magnetic Resonance Imagery (task-fMRI)

2.3.1 Overview

2.3.2 Pre-processing

2.3.3 Statistical Analysis: Subject-level

2.3.4 Statistical Analysis: Group-level

2.4 Neuroimaging Software Packages

2.4.1 SPM

2.4.2 FSL

2.4.3 AFNI

2.5 Reproducibility of fMRI Results

Exploring the Impact of Analysis Software on Task-fMRI Results

3.1 Data and Analysis Methods

3.1.1 Study Description and Data Source

3.1.2 Data Analyses

3.1.3 Comparison Methods

3.1.4 Permutation Test Methods

3.2 Results

3.2.1 Cross-Software Variability for Parametric Inference

3.2.2 Cross-Software Variability for Non-Parametric Inference

3.2.3 Intra-Software Variability, Parametric vs Non-Parametric

3.3 Reproducibility

3.3.1 Scripting of Analysis and Figures

3.3.2 Results Sharing

3.4 Discussion

3.4.1 Limitations

3.5 Conclusion

4.1 Introduction

4.2 Theory

4.2.1 Overview

4.2.2 The Wild Bootstrap Method for Computation of k

4.3 Method

4.3.1 Simulations

4.3.2 Implementation of Contour Inference

4.3.3 2D Simulations

4.3.4 3D Simulations

4.3.5 Application to Human Connectome Project Data

4.4 Results

4.4.1 2D Simulations

4.4.2 3D Simulations

4.4.3 Human Connectome Project

4.5 Discussion

4.5.1 Limitations

4.6 Conclusion

4.7 Toolbox

5.1 Theory

5.1.1 Transforming the Residual Field

5.2 Method

5.2.1 2D Simulations

5.2.2 3D Simulations

5.2.3 Application to UK Biobank Data

5.3 Results

5.3.1 2D Simulations

5.3.2 3D Simulations

5.3.3 UK Biobank Data

5.3.4 Comparison to Traditional Inference Procedures

5.4 Discussion

5.4.1 Limitations

5.5 Conclusion

CHAPTER 6

Conclusion and Future Work

Bibliography

- A. P. Dempster, D. B. R. N. M. Laird (1977). Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society. Series B (Methodological)*, **39**(1), 1–38.
- Abney, M. (2015). Permutation testing in the presence of polygenic variation. *Genetic Epidemiology*, **39**(4), 249–258.
- Allison, D. B., Neale, M. C., Zannolli, R., Schork, N. J., Amos, C. I., and Blangero, J. (1999). Testing the robustness of the likelihood-ratio test in a variance-component quantitative-trait loci-mapping procedure. *American journal of human genetics*, **65**(2), 531–44.
- Almasy, L. and Blangero, J. (1998). Multipoint quantitative-trait linkage analysis in general pedigrees. *American Journal of Human Genetics*, **62**(5), 1198–211.
- Amemiya, T. (1977). A note on a heteroscedastic model. *Journal of Econometrics*, **6**(3), 365 – 370.
- Amos, C. I. (1994). Robust variance-components approach for assessing genetic linkage in pedigrees. *American journal of human genetics*, **54**(3), 535–43.
- Ashburner, J. and Friston, K. J. (2000). Voxel-based morphometry?the methods. *NeuroImage*, **11**(6), 805 – 821.
- B. Devlin, K. R. (1999). Genomic control for association studies. *Biometrics*, **55**(4), 997–1004.
- Balding, D. (2006). A tutorial on statistical methods for population association studies. *Nat Rev Genet*, **7**(10), 781–791.
- Basser, P., Mattiello, J., and LeBihan, D. (1994). {MR} diffusion tensor spectroscopy and imaging. *Biophysical Journal*, **66**(1), 259 – 267.

- Blangero, J. and Almasy, L. (1997). Multipoint oligogenic linkage analysis of quantitative traits. *Genetic Epidemiology*, **14**(6), 959–964.
- Blangero, J., Diego, V. P., Dyer, T. D., Almeida, M., Peralta, J., Kent, J. W., Williams, J. T., Almasy, L., and Göring, H. H. H. (2013). *A kernel of truth: statistical advances in polygenic variance component models for complex human pedigrees.*, volume 81. Academic Press.
- Blokland, G. A., McMahon, K. L., Hoffman, J., Zhu, G., Meredith, M., Martin, N. G., Thompson, P. M., de Zubicaray, G. I., and Wright, M. J. (2008). Quantifying the heritability of task-related brain activation and performance during the n-back working memory task: A twin fmri study. *Biological Psychology*, **79**(1), 70 – 79.
- Brouwer, R. M., Mandl, R. C., Peper, J. S., van Baal, G. C. M., Kahn, R. S., Boomsma, D. I., and Pol, H. E. H. (2010). Heritability of {DTI} and {MTR} in nine-year-old children. *NeuroImage*, **53**(3), 1085 – 1092.
- Burggren, A., Zeineh, M., Ekstrom, A., Braskie, M., Thompson, P., Small, G., and Bookheimer, S. (2008). Reduced cortical thickness in hippocampal subregions among cognitively normal apolipoprotein e e4 carriers. *NeuroImage*, **41**(4), 1177 – 1183.
- Buse, A. (1973). Goodness of fit in generalized least squares estimation. *The American Statistician*, **27**(3), 106–108.
- Buse, A. (1979). Goodness-of-fit in the seemingly unrelated regressions model: a generalization. *Journal of Econometrics*, **10**.
- Buse, A. (1984). Tests for additive heteroskedasticity: Goldfeld and Quandt revisited. *Empirical Economics*, **9**(4), 199–216.
- Cao, J. (1999). The size of the connected components of excursion sets of χ^2 , t and F fields. *Advances in Applied Probability*, **31**(3), 579–595.
- Cardon, L. R. and Palmer, L. J. (2003). Population stratification and spurious allelic association. *The Lancet*, **361**(9357), 598 – 604.
- Chernoff, H. (1954). On the distribution of the likelihood ratio. *The Annals of Mathematical Statistics*, **25**(3), pp. 573–578.
- Chiang, M.-C., Barysheva, M., Shattuck, D. W., Lee, A. D., Madsen, S. K., Avedissian, C., Klunder, A. D., Toga, A. W., McMahon, K. L., de Zubicaray, G. I., Wright, M. J., Srivas-

- tava, A., Balov, N., and Thompson, P. M. (2009). Genetics of brain fiber architecture and intellectual performance. *The Journal of Neuroscience*, **29**(7), 2212–2224.
- Chiang, M.-C., McMahon, K. L., de Zubicaray, G. I., Martin, N. G., Hickie, I., Toga, A. W., Wright, M. J., and Thompson, P. M. (2011). Genetics of white matter development: A {DTI} study of 705 twins and their siblings aged 12 to 29. *NeuroImage*, **54**(3), 2308 – 2317.
- Cho, Y. S., Go, M. J., Kim, Y. J., Heo, J. Y., Oh, J. H., Ban, H.-J., Yoon, D., Lee, M. H., Kim, D.-J., Park, M., *et al.* (2009). A large-scale genome-wide association study of asian populations uncovers genetic factors influencing eight quantitative traits. *Nature genetics*, **41**(5), 527–534.
- Crainiceanu, C. (2008). Likelihood ratio testing for zero variance components in linear mixed models. In D. Dunson, editor, *Random Effect and Latent Variable Model Selection*, volume 192 of *Lecture Notes in Statistics*, pages 3–17. Springer New York.
- Crainiceanu, C. M. and Ruppert, D. (2004a). Likelihood ratio tests for goodness-of-fit of a nonlinear regression model. *Journal of Multivariate Analysis*, **91**(1), 35 – 52.
- Crainiceanu, C. M. and Ruppert, D. (2004b). Likelihood ratio tests in linear mixed models with one variance component. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **66**(1), 165–185.
- Crainiceanu, C. M. and Ruppert, D. (2004c). Restricted likelihood ratio tests in non-parametric longitudinal models. *Statistica Sinica*, **14**(3), 713–730.
- den Braber, A., Bohlken, M. M., Brouwer, R. M., van 't Ent, D., Kanai, R., Kahn, R. S., de Geus, E. J. C., Hulshoff Pol, H. E., and Boomsma, D. I. (2013). Heritability of sub-cortical brain measures: A perspective for future genome-wide association studies. *NeuroImage*, **83C**, 98–102.
- Dominicus, A., Skrandal, A., Gjessing, H., Pedersen, N., and Palmgren, J. (2006). Likelihood ratio tests in behavioral genetics: Problems and solutions. *Behavior Genetics*, **36**(2), 331–340.
- Draper, N. and Stoneman, D. (1966). Testing for the Inclusion of Variables in Linear Regression by a Randomisation Technique. *Technometrics*, **8**(4), 695–699.
- Drikvandi, R., Verbeke, G., Khodadadi, A., and Partovi Nia, V. (2013). Testing multiple variance components in linear mixed-effects models. *Biostatistics (Oxford, England)*, **14**(1), 144–59.

- Ernst, M. D. (2004). Permutation Methods: A Basis for Exact Inference. *Statistical Science*, **19**(4), 676–685.
- Falconer, D. and Mackay, T. (1996). *Introduction to Quantitative Genetics*. Longman.
- Fischl, B. and Dale, A. M. (2000). Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proceedings of the National Academy of Sciences*, **97**(20), 11050–11055.
- Fisher, R. (1935). *The Design of Experiments*. Oliver and Boyd.
- Fitzmaurice, G. M., Lipsitz, S. R., and Ibrahim, J. G. (2007). A note on permutation tests for variance components in multilevel generalized linear mixed models. *Biometrics*, **63**(3), 942–6.
- Freedman, D. and Lane, D. (1983). A nonstochastic interpretation of reported significance levels. *Journal of Business and Economic Statistics*, **1**(4), 292–298.
- Freedman, D. a. (2007). How Can the Score Test Be Inconsistent? *The American Statistician*, **61**(4), 291–295.
- Friston, K., Penny, W., Phillips, C., Kiebel, S., Hinton, G., and Ashburner, J. (2002). Classical and bayesian inference in neuroimaging: Theory. *NeuroImage*, **16**(2), 465 – 483.
- Friston, K. J., Worsley, K. J., Frackowiak, R. S. J., Mazziotta, J. C., and Evans, A. C. (1994). Assessing the significance of focal activations using their spatial extent. *Human Brain Mapping*, **1**(3), 210–220.
- Ganjgahi, H., Winkler, A. M., Glahn, D. C., Blangero, J., Kochunov, P., and Nichols, T. E. (2015). Fast and powerful heritability inference for family-based neuroimaging studies. *NeuroImage*, **115**, 256 – 268.
- Genovese, C. R., Lazar, N. A., and Nichols, T. E. (2002). Thresholding of Statistical Maps in Functional Neuroimaging Using the False Discovery Rate. *NeuroImage*, **15**(4), 870–878.
- Glahn, D. C., Thompson, P. M., and Blangero, J. (2007). Neuroimaging endophenotypes: Strategies for finding genes influencing brain structure and function. *Human Brain Mapping*, **28**(6), 488–501.
- Goldfeld, S. and Quandt, R. (1965). Some Tests for Homoscedasticity. *Journal of the American Statistical ...*, **60**(310), 539–547.

- Good, P. (2005). *Permutation, Parametric and Bootstrap Tests of Hypotheses*. Springer.
- Harville, D. A. (1974). Bayesian inference for variance components using only error contrasts. *Biometrika*, **61**(2), 383–385.
- Harville, D. A. (1977). Maximum likelihood approaches to variance component estimation and to related problems. *Journal of the American Statistical Association*, **72**(358), 320–338.
- Helgason, A., Yngvadóttir, B., Hrafnkelsson, B., Gulcher, J., and Stefánsson, K. (2005). An Icelandic example of the impact of population structure on association studies. *Nature genetics*, **37**(1), 90–95.
- Hibar, D. P., Stein, J. L., Renteria, M. E., Arias-Vasquez, A., Desrivieres, S., Jahan-shad, N., Toro, R., Wittfeld, K., Abramovic, L., Andersson, M., Aribisala, B. S., Armstrong, N. J., Bernard, M., Bohlken, M. M., Boks, M. P., Bralten, J., Brown, A. A., Maller Chakravarty, M., Chen, Q., Ching, C. R. K., Cuellar-Partida, G., den Braber, A., Giddaluru, S., Goldman, A. L., Grimm, O., Guadalupe, T., Hass, J., Woldehawariat, G., Holmes, A. J., Hoogman, M., Janowitz, D., Jia, T., Kim, S., Klein, M., Kraemer, B., Lee, P. H., Olde Loohuis, L. M., Luciano, M., Macare, C., Mather, K. A., Mattheisen, M., Milaneschi, Y., Nho, K., Pampmeyer, M., Ramasamy, A., Risacher, S. L., Roiz-Santianez, R., Rose, E. J., Salami, A., Samann, P. G., Schmaal, L., Schork, A. J., Shin, J., Strike, L. T., Teumer, A., van Donkelaar, M. M. J., van Eijk, K. R., Walters, R. K., Westlye, L. T., Whelan, C. D., Winkler, A. M., Zwiers, M. P., Alhusaini, S., Athanasiu, L., Ehrlich, S., Hakobjan, M. M. H., Hartberg, C. B., Haukvik, U. K., Heister, A. J. G. A. M., Hoehn, D., Kasperaviciute, D., Liewald, D. C. M., Lopez, L. M., Makkinje, R. R. R., Matarin, M., Naber, M. A. M., Reese McKay, D., Needham, M., Nugent, A. C., Putz, B., Royle, N. A., Shen, L., Sprooten, E., Trabzuni, D., van der Marel, S. S. L., van Hulzen, K. J. E., Walton, E., Wolf, C., Almasy, L., Ames, D., Arepalli, S., Assareh, A. A., Bastin, M. E., Brodaty, H., Bulayeva, K. B., Carless, M. A., Cichon, S., Corvin, A., Curran, J. E., Czisch, M., de Zubicaray, G. I., Dillman, A., Duggirala, R., Dyer, T. D., Erk, S., Fedko, I. O., Ferrucci, L., Foroud, T. M., Fox, P. T., Fukunaga, M., Raphael Gibbs, J., Goring, H. H. H., Green, R. C., Guelfi, S., Hansell, N. K., Hartman, C. A., Hegenscheid, K., Heinz, A., Hernandez, D. G., Heslenfeld, D. J., Hoekstra, P. J., Holsboer, F., Homuth, G., Hot-tenga, J.-J., Ikeda, M., Jack Jr, C. R., Jenkinson, M., Johnson, R., Kanai, R., Keil, M., Kent Jr, J. W., Kochunov, P., Kwok, J. B., Lawrie, S. M., Liu, X., Longo, D. L., McMahon, K. L., Meisenzahl, E., Melle, I., Mohnke, S., Montgomery, G. W., Mostert, J. C., Muh-leisen, T. W., Nalls, M. A., Nichols, T. E., Nilsson, L. G., Nothen, M. M., Ohi, K., Olvera, R. L., Perez-Iglesias, R., Bruce Pike, G., Potkin, S. G., Reinvang, I., Reppermund, S., Ri-

etschel, M., Romanczuk-Seiferth, N., Rosen, G. D., Rujescu, D., Schnell, K., Schofield, P. R., Smith, C., Steen, V. M., Sussmann, J. E., Thalamuthu, A., Toga, A. W., Traynor, B. J., Troncoso, J., Turner, J. A., Valdes Hernandez, M. C., van 't Ent, D., van der Brug, M., van der Wee, N. J. A., van Tol, M.-J., Veltman, D. J., Wassink, T. H., Westman, E., Zielke, R. H., Zonderman, A. B., Ashbrook, D. G., Hager, R., Lu, L., McMahon, F. J., Morris, D. W., Williams, R. W., Brunner, H. G., Buckner, R. L., Buitelaar, J. K., Cahn, W., Calhoun, V. D., Cavalleri, G. L., Crespo-Facorro, B., Dale, A. M., Davies, G. E., Delanty, N., Depondt, C., Djurovic, S., Drevets, W. C., Espeseth, T., Gollub, R. L., Ho, B.-C., Hoffmann, W., Hosten, N., Kahn, R. S., Le Hellard, S., Meyer-Lindenberg, A., Muller-Myhsok, B., Nauck, M., Nyberg, L., Pandolfo, M., Penninx, B. W. J. H., Roffman, J. L., Sisodiya, S. M., Smoller, J. W., van Bokhoven, H., van Haren, N. E. M., Volzke, H., Walter, H., Weiner, M. W., Wen, W., White, T., Agartz, I., Andreassen, O. A., Blangero, J., Boomsma, D. I., Brouwer, R. M., Cannon, D. M., Cookson, M. R., de Geus, E. J. C., Deary, I. J., Donohoe, G., Fernandez, G., Fisher, S. E., Francks, C., Glahn, D. C., Grabe, H. J., Gruber, O., Hardy, J., Hashimoto, R., Hulshoff Pol, H. E., Jonsson, E. G., Kloszewska, I., Lovestone, S., Mattay, V. S., Mecocci, P., McDonald, C., McIntosh, A. M., Ophoff, R. A., Paus, T., Pausova, Z., Ryten, M., Sachdev, P. S., Saykin, A. J., Simmons, A., Singleton, A., Soininen, H., Wardlaw, J. M., Weale, M. E., Weinberger, D. R., Adams, H. H. H., Launer, L. J., Seiler, S., Schmidt, R., Chauhan, G., Satizabal, C. L., Becker, J. T., Yanek, L., van der Lee, S. J., Ebling, M., Fischl, B., Longstreth Jr, W. T., Greve, D., Schmidt, H., Nyquist, P., Vinke, L. N., van Duijn, C. M., Xue, L., Mazoyer, B., Bis, J. C., Gudnason, V., Seshadri, S., Ikram, M. A., Initiative, T. A. D. N., Consortium, T. C., EPIGEN, IMAGEN, SYS, Martin, N. G., Wright, M. J., Schumann, G., Franke, B., Thompson, P. M., and Medland, S. E. (2015). Common genetic variants influence human subcortical brain structures. *Nature*, **520**(7546), 224–229.

Holmes, A. P., Blair, R. C., Watson, J. D., and Ford, I. (1996). Nonparametric analysis of statistic images from functional mapping experiments. *Journal of cerebral blood flow and metabolism: official journal of the International Society of Cerebral Blood Flow and Metabolism*, **16**(1), 7–22.

Hopper, J. L. and Mathews, J. D. (1982). Extensions to multi-variate normal models for pedigree analysis. *Annals of Human Genetics*, **46**, 373–383.

Hua, X., Leow, A. D., Parikshak, N., Lee, S., Chiang, M.-C., Toga, A. W., Jr, C. R. J., Weiner, M. W., and Thompson, P. M. (2008). Tensor-based morphometry as a neuroimaging biomarker for alzheimer's disease: An {MRI} study of 676 ad, mci, and normal subjects. *NeuroImage*, **43**(3), 458 – 469.

- Jahanshad, N., Kochunov, P. V., Sprooten, E., Mandl, R. C., Nichols, T. E., Almasy, L., Blangero, J., Brouwer, R. M., Curran, J. E., de Zubicaray, G. I., Duggirala, R., Fox, P. T., Hong, L. E., Landman, B. A., Martin, N. G., McMahon, K. L., Medland, S. E., Mitchell, B. D., Olvera, R. L., Peterson, C. P., Starr, J. M., Sussmann, J. E., Toga, A. W., Wardlaw, J. M., Wright, M. J., Pol, H. E. H., Bastin, M. E., McIntosh, A. M., Deary, I. J., Thompson, P. M., and Glahn, D. C. (2013). Multi-site genetic analysis of diffusion images and voxelwise heritability analysis: A pilot project of the enigma-dti working group. *NeuroImage*, **81**(0), 455 – 469.
- Jennrich, R. and Sampson, P. (1976). Newton-raphson and related algorithms for maximum likelihood variance component estimation. *Technometrics*, **18**(1), 11–17.
- Kadri, N. K., Guldbrandtsen, B., S  rensen, P., and Sahana, G. (2014). Comparison of genome-wide association methods in analyses of admixed populations with complex familial relationships. *PLoS ONE*, **9**(3), e88926.
- Kang, H. M., Zaitlen, N. a., Wade, C. M., Kirby, A., Heckerman, D., Daly, M. J., and Eskin, E. (2008). Efficient control of population structure in model organism association mapping. **178**(3), 1709–23.
- Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. a., Kong, S.-Y., Freimer, N. B., Sabatti, C., and Eskin, E. (2010). Variance component model to account for sample structure in genome-wide association studies. *Nature genetics*, **42**(4), 348–54.
- Kochunov, P., Glahn, D., Lancaster, J., Winkler, A., Smith, S., Thompson, P., Almasy, L., Duggirala, R., Fox, P., and Blangero, J. (2010). Genetics of microstructure of cerebral white matter using diffusion tensor imaging. *NeuroImage*, **53**(3), 1109 – 1116.
- Kochunov, P., Glahn, D., Nichols, T., Winkler, A., Hong, E., Holcomb, H., Stein, J., Thompson, P., Curran, J., Carless, M., Olvera, R., Johnson, M., Cole, S., Kochunov, V., Kent, J., and Blangero, J. (2011a). Genetic analysis of cortical thickness and fractional anisotropy of water diffusion in the brain. *Frontiers in Neuroscience*, **5**(120).
- Kochunov, P., Glahn, D., Lancaster, J., Thompson, P., Kochunov, V., Rogers, B., Fox, P., Blangero, J., and Williamson, D. (2011b). Fractional anisotropy of cerebral white matter and thickness of cortical gray matter across the lifespan. *NeuroImage*, **58**(1), 41 – 49.
- Kochunov, P., Jahanshad, N., Marcus, D., Winkler, A., Sproote, E., Nichols, T., Hong, L., Behrens, T., Andersson, E. J. and Yacoub, Ugurbil, K., Brouwer, C., Landman, B.,

- Braber, A., Almasy, L., Fox, P., Olvera, R., Blangero, J., DC., G., and Van Essen, D. (2014a). Heritability of fractional anisotropy in human white matter: A comparison of human connectome project and enigma-dti data. *NeuroImage*. In Review.
- Kochunov, P., Jahanshad, N., Sprooten, E., Nichols, T. E., Mandl, R. C., Almasy, L., Booth, T., Brouwer, R. M., Curran, J. E., de Zubicaray, G. I., Dimitrova, R., Duggirala, R., Fox, P. T., Hong, L. E., Landman, B. A., Lemaitre, H., Lopez, L. M., Martin, N. G., McMahon, K. L., Mitchell, B. D., Olvera, R. L., Peterson, C. P., Starr, J. M., Sussmann, J. E., Toga, A. W., Wardlaw, J. M., Wright, M. J., Wright, S. N., Bastin, M. E., McIntosh, A. M., Boomsma, D. I., Kahn, R. S., den Braber, A., de Geus, E. J., Deary, I. J., Pol, H. E. H., Williamson, D. E., Blangero, J., van 't Ent, D., Thompson, P. M., and Glahn, D. C. (2014b). Multi-site study of additive genetic effects on fractional anisotropy of cerebral white matter: Comparing meta and megaanalytical approaches for data pooling. *NeuroImage*, **95**(0), 136 – 150.
- Koten, J. W., Wood, G., Hagoort, P., Goebel, R., Propping, P., Willmes, K., and Boomsma, D. I. (2009). Genetic contribution to variation in cognitive function: An fmri study in twins. *Science*, **323**(5922), 1737–1740.
- Kremen, W. S., Prom-Wormley, E., Panizzon, M. S., Eyler, L. T., Fischl, B., Neale, M. C., Franz, C. E., Lyons, M. J., Pacheco, J., Perry, M. E., Stevens, A., Schmitt, J. E., Grant, M. D., Seidman, L. J., Thermenos, H. W., Tsuang, M. T., Eisen, S. a., Dale, A. M., and Fennema-Notestine, C. (2010). Genetic and environmental influences on the size of specific brain regions in midlife: the vetsa mri study. *NeuroImage*, **49**(2), 1213–23.
- Lange, K. (2003). *Mathematical and statistical methods for genetic analysis*. Springer, 2nd edition.
- Lange, K., Westlake, J., and Spence, M. A. (1976). Extensions to pedigree analysis. III. variance components by the scoring method. *Annals of Human Genetics*, **39**(4), 485–91.
- Lee, O. E. and Braun, T. M. (2012). Permutation tests for random effects in linear mixed models. *Biometrics*, **68**(2), 486–493.
- Lippert, C., Listgarten, J., Liu, Y., Kadie, C. M., Davidson, R. I., and Heckerman, D. (2011a). FaST linear mixed models for genome-wide association studies. *Nature Methods*, **8**(10), 833–837.
- Lippert, C., Listgarten, J., Liu, Y., Kadie, C. M., Davidson, R. I., Heckerman, D., Lippert, C., Kadie, C. M., Davidson, R. I., Eskin, E., and Heckerman, D. (2011b). Improved

- linear mixed models for genome-wide association studies. *Nature methods*, **8**(6), 833–5.
- Listgarten, J., Lippert, C., and Heckerman, D. (2013). FaST-LMM-Select for addressing confounding from spatial structure and rare variants. *Nature Genetics*, **45**(5), 470–471.
- Longford, N. T. (1987). A fast scoring algorithm for maximum likelihood estimation in unbalanced mixed models with nested random effects. *Biometrika*, **74**(4), 817–827.
- MacCluer, J. W., Blangero, J., Dyer, T. D., and Speer, M. C. (1997). GAW10: simulated family data for a common oligogenic disease with quantitative risk factors. *Genetic epidemiology*, **14**(6), 737–42.
- Matthews, S. C., Simmons, A. N., Strigo, I., Jang, K., Stein, M. B., and Paulus, M. P. (2007). Heritability of anterior cingulate response to conflict: An fmri study in female twins. *NeuroImage*, **38**(1), 223 – 227.
- McKay, D., Knowles, E., Winkler, A., Sprooten, E., Kochunov, P., Olvera, R., Curran, J., Kent, J. JackW., Carless, M., GÁúring, H., Dyer, T., Duggirala, R., Almasy, L., Fox, P., Blangero, J., and Glahn, D. (2014). Influence of age, sex and genetic factors on the human brain. *Brain Imaging and Behavior*, **8**(2), 143–152.
- Meyer, K. (1985). Maximum likelihood estimation of variance components for a multivariate mixed model with equal design matrices. *Biometrics*, pages 153–165.
- Molenberghs, G. and Verbeke, G. (2007). Likelihood Ratio, Score, and Wald Tests in a Constrained Parameter Space. *The American Statistician*, **61**(1), 22–27.
- Morgan, B. J. T., Palmer, K. J., and Ridout, M. S. (2007). Negative Score Test Statistic. *The American Statistician*, **61**(4), 285–288.
- Mumford, J. and Nichols, T. (2006). Modeling and inference of multisubject fmri data. *Engineering in Medicine and Biology Magazine, IEEE*, **25**(2), 42–51.
- Nan M. Laird, J. H. W. (1982). Random-effects models for longitudinal data. *Biometrics*, **38**(4), 963–974.
- Neyman, J. and Pearson, E. S. (1933). On the problem of the most efficient tests of statistical hypotheses. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character*, **231**, pp. 289–337.

- Nichols, T. E. and Hayasaka, S. (2003). Controlling the familywise error rate in functional neuroimaging: a comparative review. *Statistical Methods in Medical Research*, **12**(5), 419–446.
- Nichols, T. E. and Holmes, A. P. (2001). Nonparametric permutation tests for functional neuroimaging: A primer with examples. *Human Brain Mapping*, **15**(1), 1–25.
- Nichols, T. E. and Holmes, A. P. (2002). Nonparametric permutation tests for functional neuroimaging: a primer with examples. *Human Brain Mapping*, **15**(1), 1–25.
- Ober, C., Abney, M., and McPeck, M. S. (2001). The genetic dissection of complex traits in a founder population. *The American Journal of Human Genetics*, **69**(5), 1068–1079.
- Olvera, R., Bearden, C., Velligan, D., Almasy, L., Carless, M., Curran, J., Williamson, D., Duggirala, R., Blangero, J., and Glahn, D. (2011). Common genetic influences on depression, alcohol, and substance use disorders in mexican-american families. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, **156**(5), 561–568.
- Patterson, H. D. and Thompson, R. (1971). Recovery of inter-block information when block sizes are unequal. *Biometrika*, **58**(3), 545–554.
- Pievani, M., Rasser, P., Galluzzi, S., Benussi, L., Ghidoni, R., Sabattoli, F., Bonetti, M., Binetti, G., Thompson, P., and Frisoni, G. (2009). Mapping the effect of {APOE} ?4 on gray matter loss in alzheimer’s disease in vivo. *NeuroImage*, **45**(4), 1090 – 1098.
- Pirinen, M., Donnelly, P., and Spencer, C. C. A. (2013). Efficient computation with a linear mixed model on large-scale data sets with applications to genetic studies. *Ann. Appl. Stat.*, **7**(1), 369–390.
- Polk, T. A., Park, J., Smith, M. R., and Park, D. C. (2007). Nature versus nurture in ventral visual cortex: A functional magnetic resonance imaging study of twins. *The Journal of Neuroscience*, **27**(51), 13921–13925.
- Potkin, S. G., Turner, J. A., Guffanti, G., Lakatos, A., Fallon, J. H., Nguyen, D. D., Mathalon, D., Ford, J., Lauriello, J., Macciardi, F., and FBIRN (2009a). A genome-wide association study of schizophrenia using brain activation as a quantitative phenotype. *Schizophrenia Bulletin*, **35**(1), 96–108.
- Potkin, S. G., Guffanti, G., Lakatos, A., Turner, J. A., Kruggel, F., Fallon, J. H., Saykin, A. J., Orro, A., Lupoli, S., Salvi, E., Weiner, M., Macciardi, F., and for the Alzheimer’s

Disease Neuroimaging Initiative (2009b). Hippocampal atrophy as a quantitative trait in a genome-wide association study identifying novel susceptibility genes for alzheimer's disease. *PLoS ONE*, **4**(8), 1–15.

Price, A. L., Patterson, N. J., Plenge, R. M., Weinblatt, M. E., Shadick, N. A., and Reich, D. (2006). Principal components analysis corrects for stratification in genome-wide association studies. *Nature genetics*, **38**(8), 904–909.

Price, A. L., Zaitlen, N. A., Reich, D., and Patterson, N. (2010). New approaches to population stratification in genome-wide association studies. *Nature Reviews*, **11**(June), 459–463.

Pritchard, J. K., Stephens, M., Rosenberg, N. A., and Donnelly, P. (2000). Association mapping in structured populations. *The American Journal of Human Genetics*, **67**(1), 170 – 181.

Rao, C. R. (2008). *Linear Statistical Inference and its Applications*. John Wiley & Sons, Inc.

Reich, D. E., Cargill, M., Bolk, S., Ireland, J., Sabeti, P. C., Richter, D. J., Lavery, T., Kouyoumjian, R., Farhadian, S. F., Ward, R., and Lander, E. S. (2001). Linkage disequilibrium in the human genome. *Nature*, **411**, 199–204.

Rimol, L. M., Panizzon, M. S., Fennema-Notestine, C., Eyler, L. T., Fischl, B., Franz, C. E., Hagler, D. J., Lyons, M. J., Neale, M. C., Pacheco, J., Perry, M. E., Schmitt, J. E., Grant, M. D., Seidman, L. J., Thermenos, H. W., Tsuang, M. T., Eisen, S. a., Kremen, W. S., and Dale, A. M. (2010). Cortical thickness is influenced by regionally specific genetic factors. *Biological psychiatry*, **67**(5), 493–9.

Sabatti, C., Service, S. K., Hartikainen, A.-L., Pouta, A., Ripatti, S., Brodsky, J., Jones, C. G., Zaitlen, N. A., Varilo, T., Kaakinen, M., *et al.* (2009). Genome-wide association analysis of metabolic traits in a birth cohort from a founder population. *Nature genetics*, **41**(1), 35–46.

Salimi-Khorshidi, G., Smith, S. M., and Nichols, T. E. (2010). Adjusting the effect of nonstationarity in cluster-based and TFCE inference. *NeuroImage*, **54**(3), 2006–2019.

Samuh, M. H., Grilli, L., Rampichini, C., Salmaso, L., and Lunardon, N. (2012). The use of permutation tests for variance components in linear mixed models. *Communications in Statistics - Theory and Methods*, **41**(16-17), 3020–3029.

- Searle, S. (1971). *Linear Models*. Number v. 1 in Wiley Publication in Mathematical Statistics. John Wiley & Sons.
- Searle, S. R., Casella, G., and McCulloch, C. E. (2009). *Variance components*, volume 391. John Wiley & Sons.
- Self, S. G. and Liang, K.-Y. (1987). Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. *Journal of the American Statistical Association*, **82**(398), pp. 605–610.
- Servin, B. and Stephens, M. (2007). Imputation-based analysis of association studies: Candidate regions and quantitative traits. *PLoS Genet*, **3**(7).
- Shephard, N. (1993). Maximum likelihood estimation of regression models with stochastic trend components. *Journal of the American Statistical Association*, **88**(422), pp. 590–595.
- Shephard, N. G. and Harvey, A. C. (1990). On the probability of estimating a deterministic component in the local level model. *Journal of Time Series Analysis*, **11**(4), 339–347.
- Silvapulle, M. J. (1992). Robust wald-type tests of one-sided hypotheses in the linear model. *Journal of the American Statistical Association*, **87**(417), 156–161.
- Silvapulle, M. J. and Silvapulle, P. (1995). A score test against one-sided alternatives. *Journal of the American Statistical Association*, **90**(429), 342–349.
- Smith, S. and Nichols, T. (2009). Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage*, **44**(1), 83–98.
- Smith, S. M., Jenkinson, M., Johansen-Berg, H., Rueckert, D., Nichols, T. E., Mackay, C. E., Watkins, K. E., Ciccarelli, O., Cader, M. Z., Matthews, P. M., and Behrens, T. E. (2006). Tract-based spatial statistics: Voxelwise analysis of multi-subject diffusion data. *NeuroImage*, **31**(4), 1487 – 1505.
- Stein, J. L., Hua, X., Morra, J. H., Lee, S., Hibar, D. P., Ho, A. J., Leow, A. D., Toga, A. W., Sul, J. H., Kang, H. M., Eskin, E., Saykin, A. J., Shen, L., Foroud, T., Pankratz, N., Huentelman, M. J., Craig, D. W., Gerber, J. D., Allen, A. N., Corneveaux, J. J., Stephan, D. A., Webster, J., DeChairo, B. M., Potkin, S. G., Jr., C. R. J., Weiner, M. W., and Thompson, P. M. (2010a). Genome-wide analysis reveals novel genes influencing temporal lobe structure with relevance to neurodegeneration in alzheimer's disease. *NeuroImage*, **51**(2), 542 – 554.

Stein, J. L., Hua, X., Lee, S., Ho, A. J., Leow, A. D., Toga, A. W., Saykin, A. J., Shen, L., Foroud, T., Pankratz, N., Huentelman, M. J., Craig, D. W., Gerber, J. D., Allen, A. N., Corneveaux, J. J., DeChairo, B. M., Potkin, S. G., Weiner, M. W., and Thompson, P. M. (2010b). Voxelwise genome-wide association study (vgwas). *NeuroImage*, **53**(3), 1160 – 1174. Imaging Genetics.

Stein, J. L., Medland, S. E., Vasquez, A. A., Hibar, D. P., Senstad, R. E., Winkler, A. M., Toro, R., Appel, K., Barteczek, R., Bergmann, O., Bernard, M., Brown, A. A., Cannon, D. M., Chakravarty, M. M., Christoforou, A., Domin, M., Grimm, O., Hollinshead, M., Holmes, A. J., Homuth, G., Hottenga, J.-J., Langan, C., Lopez, L. M., Hansell, N. K., Hwang, K. S., Kim, S., Laje, G., Lee, P. H., Liu, X., Loth, E., Lourdasamy, A., Mattingsdal, M., Mohnke, S., Maniega, S. M., Nho, K., Nugent, A. C., O'Brien, C., Papmeyer, M., Putz, B., Ramasamy, A., Rasmussen, J., Rijpkema, M., Risacher, S. L., Roddey, J. C., Rose, E. J., Ryten, M., Shen, L., Sprooten, E., Strengman, E., Teumer, A., Trabzuni, D., Turner, J., van Eijk, K., van Erp, T. G. M., van Tol, M.-J., Wittfeld, K., Wolf, C., Woudstra, S., Aleman, A., Alhusaini, S., Almasy, L., Binder, E. B., Brohawn, D. G., Cantor, R. M., Carless, M. A., Corvin, A., Czisch, M., Curran, J. E., Davies, G., de Almeida, M. A. A., Delanty, N., Depondt, C., Duggirala, R., Dyer, T. D., Erk, S., Fagerness, J., Fox, P. T., Freimer, N. B., Gill, M., Goring, H. H. H., Hagler, D. J., Hoehn, D., Holsboer, F., Hoogman, M., Hosten, N., Jahanshad, N., Johnson, M. P., Kasperaviciute, D., Kent, J. W., Kochunov, P., Lancaster, J. L., Lawrie, S. M., Liewald, D. C., Mandl, R., Matarin, M., Mattheisen, M., Meisenzahl, E., Melle, I., Moses, E. K., Muhleisen, T. W., Nauck, M., Nothen, M. M., Olvera, R. L., Pandolfo, M., Pike, G. B., Puls, R., Reinvang, I., Renteria, M. E., Rietschel, M., Roffman, J. L., Royle, N. A., Rujescu, D., Savitz, J., Schnack, H. G., Schnell, K., Seiferth, N., Smith, C., Steen, V. M., Valdes Hernandez, M. C., Van den Heuvel, M., van der Wee, N. J., Van Haren, N. E. M., Veltman, J. A., Volzke, H., Walker, R., Westlye, L. T., Whelan, C. D., Agartz, I., Boomsma, D. I., Cavalleri, G. L., Dale, A. M., Djurovic, S., Drevets, W. C., Hagoort, P., Hall, J., Heinz, A., Jack, C. R., Foroud, T. M., Le Hellard, S., Macciardi, F., Montgomery, G. W., Poline, J. B., Porteous, D. J., Sisodiya, S. M., Starr, J. M., Sussmann, J., Toga, A. W., Veltman, D. J., Walter, H., Weiner, M. W., Bis, J. C., Ikram, M. A., Smith, A. V., Gudnason, V., Tzourio, C., Vernooij, M. W., Launer, L. J., DeCarli, C., and Seshadri, S. (2012). Identification of common variants associated with human hippocampal and intracranial volumes. *Nat Genet*, **44**(5), 552–561.

Stram, D. O. and Lee, J. W. (1994). Variance components testing in the longitudinal mixed effects model. *Biometrics*, **50**(4), pp. 1171–1177.

Svishcheva, G. R., Axenovich, T. I., Belonogova, N. M., van Duijn, C. M., and Aulchenko,

- Y. S. (2012). Rapid variance components–based method for whole-genome association analysis. *Nature Genetics*, **44**(10), 1166–1170.
- ter Braak, C. J. (1992). *Permutation Versus Bootstrap Significance Tests in Multiple Regression and Anova*, volume 376 of *Lecture Notes in Economics and Mathematical Systems*. Springer Berlin Heidelberg.
- Verbeke, G. and Molenberghs, G. (2003). The use of score tests for inference on variance components. *Biometrics*, **59**(2), pp. 254–262.
- Verbeke, G. and Molenberghs, G. (2007). What Can Go Wrong With the Score Test? *The American Statistician*, **61**(4), 289–290.
- Voight, B. F. and Pritchard, J. K. (2005). Confounding from cryptic relatedness in case-control association studies. *PLoS Genet*, **1**(3).
- Weir, B. S., Anderson, A. D., and Hepler, A. B. (2006). Genetic relatedness analysis: modern data and new challenges. *Nature Reviews Genetics*, **7**(10), 771–780.
- Widmer, C., Lippert, C., Weissbrod, O., Fusi, N., Kadie, C., Davidson, R., Listgarten, J., and Heckerman, D. (2014). Further improvements to linear mixed models for genome-wide association studies. *Scientific reports*, **4**, 6874.
- Winkler, A. M., Kochunov, P., Blangero, J., Almasy, L., Zilles, K., Fox, P. T., Duggirala, R., and Glahn, D. C. (2010). Cortical thickness or grey matter volume? the importance of selecting the phenotype for imaging genetics studies. *NeuroImage*, **53**(3), 1135–1146.
- Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., and Nichols, T. E. (2014). Permutation inference for the general linear model. *NeuroImage*, **92C**, 381–397.
- Worsley, K. J., Evans, A. C., Marrett, S., and Neelin, P. (1992). A three-dimensional statistical analysis for CBF activation studies in human brain. *Journal of Cerebral Blood Flow and Metabolism*, **12**(6), 900–918.
- Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M., and Price, A. L. (2014). Advantages and pitfalls in the application of mixed-model association methods. *Nature genetics*, **46**(2), 100–6.
- Yu, J., Pressoir, G., Briggs, W. H., Vroh Bi, I., Yamasaki, M., Doebley, J. F., McMullen, M. D., Gaut, B. S., Nielsen, D. M., Holland, J. B., Kresovich, S., and Buckler, E. S. (2006). A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nature genetics*, **38**(2), 203–8.

- Zhang, Z., Ersoz, E., Lai, C.-Q., Todhunter, R. J., Tiwari, H. K., Gore, M. a., Bradbury, P. J., Yu, J., Arnett, D. K., Ordovas, J. M., and Buckler, E. S. (2010). Mixed linear model approach adapted for genome-wide association studies. *Nature genetics*, **42**(4), 355–360.
- Zhou, X. and Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association studies. *Nature genetics*, **44**(7), 821–4.