

Alex Caramanico

3-16-22

1.

```
. summarize mathscore if aide == 1
```

Variable	Obs	Mean	Std. dev.	Min	Max
mathscore	2,043	482.9187	45.94009	339	626

```
. summarize mathscore if aide == 0
```

Variable	Obs	Mean	Std. dev.	Min	Max
mathscore	3,743	487.062	48.56821	320	626

Classes with teaching aides score slightly less on average (about 4 points).

2.

```
. regress mathscore aide
```

Source	SS	df	MS	Number of obs	=	5,786
Model	22687.6225	1	22687.6225	F(1, 5784)	=	9.99
Residual	13136518.1	5,784	2271.18225	Prob > F	=	0.0016
				R-squared	=	0.0017
				Adj R-squared	=	0.0016
Total	13159205.8	5,785	2274.71145	Root MSE	=	47.657

mathscore	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
aide	-4.143235	1.310905	-3.16	0.002	-6.713099	-1.573372
_cons	487.062	.7789617	625.27	0.000	485.5349	488.589

3.

. regress mathscore if aide == 0

Source	SS	df	MS	Number of obs	=	3,743
Model	0	0	.	F(0, 3742)	=	0.00
Residual	8826893.62	3,742	2358.87056	Prob > F	=	.
Total	8826893.62	3,742	2358.87056	R-squared	=	0.0000
				Adj R-squared	=	0.0000
				Root MSE	=	48.568

mathscore	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
_cons	487.062	.7938567	613.54	0.000	485.5055	488.6184

. regress mathscore if aide == 1

Source	SS	df	MS	Number of obs	=	2,043
Model	0	0	.	F(0, 2042)	=	0.00
Residual	4309624.51	2,042	2110.49193	Prob > F	=	.
Total	4309624.51	2,042	2110.49193	R-squared	=	0.0000
				Adj R-squared	=	0.0000
				Root MSE	=	45.94

mathscore	Coefficient	Std. err.	t	P> t	[95% conf. interval]	
_cons	482.9187	1.016384	475.13	0.000	480.9255	484.912

.