

ETC3550

Applied forecasting for business and economics

Ch12. Some practical forecasting
issues

QTexts.org/fnp3/



Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Models for different frequencies

Models for annual data

- ETS, ARIMA, Dynamic regression

Models for different frequencies

Models for annual data

- ETS, ARIMA, Dynamic regression

Models for quarterly data

- ETS, ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA

Models for different frequencies

Models for annual data

- ETS, ARIMA, Dynamic regression

Models for quarterly data

- ETS, ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA

Models for monthly data

- ETS, ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA

Models for different frequencies

Models for weekly data

- ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA, TBATS

Models for different frequencies

Models for weekly data

- ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA, TBATS

Models for daily, hourly and other sub-daily data


- ARIMA/SARIMA, Dynamic regression, Dynamic harmonic regression, STL+ETS, STL+ARIMA, TBATS

Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Positive forecasts

```
recent_prices <- prices %>% filter(!is.na(eggs))  
recent_prices %>%  
  model(ETS(log(eggs) ~ error("A") + trend("A") + season("N"))) %>%  
  forecast(h=50) %>%  
  autoplot(recent_prices)
```



Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Forecast combinations

Clemen (1989)


“The results have been virtually unanimous: combining multiple forecasts leads to increased forecast accuracy. ... In many cases one can make dramatic performance improvements by simply averaging the forecasts.”

Forecast combinations

```
aus_cafe <- aus_retail %>%  
  filter(Industry == "Cafes, restaurants and catering services") %>%  
  summarise(Turnover = sum(Turnover))  
fc <- aus_cafe %>%  
  filter(Month <= yearmonth("2013 Sep")) %>%  
  model(  
    ETS = ETS(Turnover),  
    ARIMA = ARIMA(Turnover)  
  ) %>%  
  mutate(  
    Combination = (ETS + ARIMA)/2  
  ) %>%  
  forecast(h = "5 years")
```

Forecast combinations

```
fc %>% autoplot(aus_cafe, level = NULL) +  
  labs(x = "Year", y = "$ billion",  
       title = "Australian monthly expenditure on eating out")
```



Forecast combinations

```
fc %>% accuracy(aus_cafe)
```

```
## # A tibble: 3 x 10
```

| ## | .model | .type | ME | RMSE | MAE | MPE | MAPE | MASE | RMSSE | ACF1 |
|------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| ## | <chr> | <chr> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> |
| ## 1 | ARIMA | Test | 112. | 122. | 112. | 5.44 | 5.44 | 1.80 | 1.50 | 0.510 |
| ## 2 | Combination | Test | 120. | 125. | 120. | 5.81 | 5.81 | 1.93 | 1.55 | 0.382 |
| ## 3 | ETS | Test | 128. | 133. | 128. | 6.18 | 6.18 | 2.06 | 1.64 | 0.324 |

Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Missing values

Functions which can handle missing values

- `ARIMA()`
- `TSLM()`
- `NNETAR()`
- `VAR()`
- `FASSTER()`

Models which cannot handle missing values

- `ETS()`
- `STL()`
- `TBATS()`

Missing values

Functions which can handle missing values

- `ARIMA()`
- `TSLM()`
- `NNETAR()`
- `VAR()`
- `FASSTER()`

Models which cannot handle missing values


- `ETS()`
- `STL()`
- `TBATS()`

What to do?

- 1 Model section of data after last missing value.


Missing values

```
gold <- as_tsibble(forecast::gold)  
gold %>% autoplot(value)
```



Missing values

```
gold_complete <- gold %>%  
  model(ARIMA(value)) %>%  
  interpolate(gold)  
gold_complete %>%  
  autoplot(value, colour = "red") +  
  autolayer(gold, value)
```



Outline

- 1 Models for different frequencies
- 2 Ensuring forecasts stay within limits
- 3 Forecast combinations
- 4 Missing values
- 5 Outliers

Outliers

```
fit <- gold %>%  
  model(ARIMA(value))  
augment(fit) %>%  
  mutate(stdres = .resid/sd(.resid, na.rm=TRUE)) %>%  
  filter(abs(stdres) > 10)
```

```
## # A tsibble: 2 x 7 [1]  
## # Key:           .model [1]  
##   .model      index value .fitted .resid .innov stdres  
##   <chr>      <dbl> <dbl>   <dbl> <dbl> <dbl> <dbl>  
## 1 ARIMA(value)  770  594.   499.   94.7  94.7  16.4  
## 2 ARIMA(value)  771  487.   562. -74.8 -74.8 -12.9
```