

# ECE 6560 Final Project - Image Smoothing

Alexander Domagala

Spring 2024

## 1 Problem Description

Although image processing techniques began to appear around the 1960s, the proliferation of inexpensive digital computing power has greatly widened the application pool. Image processing techniques can now be found in areas such as digital photography, medical imaging, and object detection/tracking.

All these application areas can be affected by a very common problem: image noise. Noise can render further processing ineffective, thus there exist a variety of approaches by which we can attempt to smooth and denoise our images. Traditional image processing techniques may rely on fixed kernels that can be swept over an image in order to smooth it. However, this type of approach will result in uniform smoothing can blur the edges of an image.

We can instead leverage PDEs to describe how an image should be updated depending on its local characteristics. More specifically, we will examine Anisotropic Diffusion. This technique can be used to reduce image noise while lessening the blurring that is done to edges. This project will attempt to develop PDEs that can be used to reduce noise in high/low contrast images.

We need more information about the actual high/low contrast problem!

## 2 Mathematical Modeling

Before explicitly developing our PDEs, we must first understand what behavior we want our system to have. The calculus of variations can be used to minimize an energy functional. The choice of setup for the energy functional will determine the system behavior.

Let's begin by defining our image as

$$I(x, y)$$

Note that the derivations in sections (2) and (3) are being performed exclusively in continuous space:  $x, y \in \mathbb{R}$ .

### 3 Derivation of PDE

Now, let's introduce the Euler-Lagrange equation

$$\begin{aligned} L_f - \frac{\partial}{\partial x} L_{f'} &= 0 \quad (1\text{-D}) \\ L_I - \frac{\partial}{\partial x} L_{I_x} - \frac{\partial}{\partial y} L_{I_y} &= 0 \quad (2\text{-D}) \end{aligned}$$

We can begin working towards obtaining our PDE by setting up a gradient descent

$$\begin{aligned} I_t &= -\nabla_I E \\ I_t &= -L_I + \frac{\partial}{\partial x} L_{I_x} + \frac{\partial}{\partial y} L_{I_y} \end{aligned}$$

We will now have to compute terms  $L_I$ ,  $L_{I_x}$ ,  $L_{I_y}$  using the previously obtained energy functional

$$\begin{aligned} L(I, I_x, I_y, x, y) &= \frac{\lambda}{1+e^\alpha}, \text{ where } \alpha = -\frac{1}{\beta}(\|\nabla_I\|) \\ L_I &= \frac{\partial}{\partial I}(L) \\ L_I &= 0 \\ L_{I_x} &= \frac{\partial}{\partial I_x}(L) \\ L_{I_x} &= \frac{\lambda}{\beta} \frac{e^\alpha}{(1+e^\alpha)^2} \frac{I_x}{\sqrt{I_x^2 + I_y^2}} \\ L_{I_y} &= \frac{\partial}{\partial I_y}(L) \\ L_{I_y} &= \frac{\lambda}{\beta} \frac{e^\alpha}{(1+e^\alpha)^2} \frac{I_y}{\sqrt{I_x^2 + I_y^2}} \end{aligned}$$

Now that we have obtained our expressions for  $L_{I_x}$  and  $L_{I_y}$ , we must compute their partial derivatives as shown by the Euler-Lagrange equation. This will be shown for  $\frac{\partial}{\partial x} L_{I_x}$ .  $\frac{\partial}{\partial y} L_{I_y}$  will be obtained by examining the expression of  $\frac{\partial}{\partial x} L_{I_x}$ .

Let  $\phi$  denote  $\frac{e^\alpha}{(1+e^\alpha)^2}$  and let  $\gamma$  denote  $\frac{I_x}{\sqrt{I_x^2 + I_y^2}}$ . We can begin finding  $\frac{\partial}{\partial x} L_{I_x}$  by using the product-rule  $\frac{\partial}{\partial x}(\gamma)\phi + \frac{\partial}{\partial x}(\phi)\gamma$ . We will start with the left side of the sum. Note that we must include  $\frac{\lambda}{\beta}$  in the final expression.

$$\begin{aligned} &\frac{\partial}{\partial x}(\gamma)\phi \\ &\frac{\partial}{\partial x}\left(\frac{I_x}{\sqrt{I_x^2 + I_y^2}}\right)\phi \\ &\left(\frac{I_{xx}}{(I_x^2 + I_y^2)^{\frac{1}{2}}} + \frac{I_x}{(I_x^2 + I_y^2)^{\frac{3}{2}}}(I_x I_{xx} + I_y I_{xy})\right)\phi \end{aligned}$$

We can now examine the right side of  $\frac{\partial}{\partial x}(\gamma)\phi + \frac{\partial}{\partial x}(\phi)\gamma$ .

$$\begin{aligned} & \frac{\partial}{\partial x}(\phi)\gamma \\ & \frac{\partial}{\partial x}\left(\frac{e^\alpha}{(1+e^\alpha)^2}\right)\gamma \\ & \frac{\partial}{\partial x}((e^\alpha)(1+e^\alpha)^{-2})\gamma \end{aligned}$$

We see that we will need to again perform the product-rule between  $(e^\alpha)$  and  $(1+e^\alpha)^{-2}$ . Taking the partial derivative of  $(e^\alpha)$

$$-\frac{1}{\beta}e^\alpha \frac{1}{(I_x^2+I_y^2)^{\frac{1}{2}}}(I_x I_{xx} + I_y I_{xy})$$

Taking the partial derivative of  $(1+e^\alpha)^{-2}$

$$-2(1+e^\alpha)^{-3}\left(-\frac{1}{\beta}e^\alpha \frac{1}{(I_x^2+I_y^2)^{\frac{1}{2}}}(I_x I_{xx} + I_y I_{xy})\right)$$

Thus, after factoring common terms,  $\frac{\partial}{\partial x}((e^\alpha)(1+e^\alpha)^{-2})$  yields

$$\left[-\frac{1}{\beta}(e^\alpha)\left(\frac{1}{(I_x^2+I_y^2)^{\frac{1}{2}}}\right)(I_x I_{xx} + I_y I_{xy})\right]\left[(1+e^\alpha)^{-2} + (e^\alpha)(-2(1+e^\alpha)^{-3})\right]$$

We have reached the final expression for  $\frac{\partial}{\partial x}L_{I_x}$

$$\begin{aligned} \frac{\partial}{\partial x}L_{I_x} = & \frac{\lambda}{\beta}\left[\left(\frac{I_{xx}}{(I_x^2+I_y^2)^{\frac{1}{2}}}\right) - \left(\frac{I_x}{(I_x^2+I_y^2)^{\frac{3}{2}}}\right)(I_x I_{xx} + I_y I_{xy})\left(\frac{e^\alpha}{(1+e^\alpha)^2}\right) + \right. \\ & \left. \left(\frac{I_x}{(I_x^2+I_y^2)^{\frac{1}{2}}}\right)\left[-\frac{1}{\beta}(e^\alpha)\left(\frac{1}{(I_x^2+I_y^2)^{\frac{1}{2}}}\right)(I_x I_{xx} + I_y I_{xy})\right]\left[(1+e^\alpha)^{-2} + (e^\alpha)(-2(1+e^\alpha)^{-3})\right]\right] \end{aligned}$$

$\frac{\partial}{\partial y}L_{I_y}$  can be obtained by modifying  $\frac{\partial}{\partial x}L_{I_x}$

$$\begin{aligned} \frac{\partial}{\partial y}L_{I_y} = & \frac{\lambda}{\beta}\left[\left(\frac{I_{yy}}{(I_x^2+I_y^2)^{\frac{1}{2}}}\right) - \left(\frac{I_y}{(I_x^2+I_y^2)^{\frac{3}{2}}}\right)(I_x I_{xy} + I_y I_{yy})\left(\frac{e^\alpha}{(1+e^\alpha)^2}\right) + \right. \\ & \left. \left(\frac{I_y}{(I_x^2+I_y^2)^{\frac{1}{2}}}\right)\left[-\frac{1}{\beta}(e^\alpha)\left(\frac{1}{(I_x^2+I_y^2)^{\frac{1}{2}}}\right)(I_x I_{xy} + I_y I_{yy})\right]\left[(1+e^\alpha)^{-2} + (e^\alpha)(-2(1+e^\alpha)^{-3})\right]\right] \end{aligned}$$

Our final gradient-descent PDE is

$$\begin{aligned}
I_t = & \frac{\lambda}{\beta} \left[ \left( \frac{I_{xx}}{(I_x^2 + I_y^2)^{\frac{1}{2}}} \right) - \left( \frac{I_x}{(I_x^2 + I_y^2)^{\frac{3}{2}}} \right) (I_x I_{xx} + I_y I_{xy}) \left( \frac{e^\alpha}{(1+e^\alpha)^2} \right) + \right. \\
& \left( \frac{I_x}{(I_x^2 + I_y^2)^{\frac{1}{2}}} \right) \left[ -\frac{1}{\beta} (e^\alpha) \left( \frac{1}{(I_x^2 + I_y^2)^{\frac{1}{2}}} \right) (I_x I_{xx} + I_y I_{xy}) \right] [(1+e^\alpha)^{-2} + (e^\alpha)(-2(1+e^\alpha)^{-3})] + \\
& \left( \frac{I_{yy}}{(I_x^2 + I_y^2)^{\frac{1}{2}}} \right) - \left( \frac{I_y}{(I_x^2 + I_y^2)^{\frac{3}{2}}} \right) (I_x I_{xy} + I_y I_{yy}) \left( \frac{e^\alpha}{(1+e^\alpha)^2} \right) + \\
& \left. \left( \frac{I_y}{(I_x^2 + I_y^2)^{\frac{1}{2}}} \right) \left[ -\frac{1}{\beta} (e^\alpha) \left( \frac{1}{(I_x^2 + I_y^2)^{\frac{1}{2}}} \right) (I_x I_{xy} + I_y I_{yy}) \right] [(1+e^\alpha)^{-2} + (e^\alpha)(-2(1+e^\alpha)^{-3})] \right]
\end{aligned}$$

Where  $\alpha = -\frac{1}{\beta} (\|\nabla I\|)$

## 4 Discretization and Implementation

The PDE that was obtained in the previous section is only applicable for continuous time and space variables. We must discretize the PDE so that it can actually be implemented in software.

Before explicitly discretizing the PDE, we must also obtain numeric approximations for its partial derivatives. Beginning with the left side of the PDE, let's approximate  $I_t$ . It is important to understand that the PDE is defining how the image will be smoothed over successive iterations. In other words, it specifies how the image is updated as time increases. Thus, it is appropriate to use a forward difference.

The forward difference for a single (space) variable can be obtained using the numeric differentiation method shown in class

$$\begin{aligned}f(x, t) &= f(x, t) \\f(x, t + \Delta t) &= f(x, t) + \Delta t f'(x, t) + O(\Delta t^2)\end{aligned}$$

After eliminating  $f(x, t)$  terms on the right side of the above equations, we obtain our approximation. Note that the approximation includes an error term  $O(\Delta t)$ .

$$f'(x, t) = \frac{f(x, t + \Delta t) - f(x, t)}{\Delta t}$$

Expanding this process to two (space) dimensions yields

$$I_t(x, y, t) = \frac{I(x, y, t + \Delta t) - I(x, y, t)}{\Delta t}$$

Continuing with the right side of the PDE, approximations are needed for the following partial derivatives:  $I_x$ ,  $I_y$ ,  $I_{xx}$ ,  $I_{yy}$ , and  $I_{xy}$ . Since these approximations capture how the image is changing with respect to space, it is more appropriate to use a central difference rather than a forward difference.

Suppose we are trying to approximate  $I_x$ . A forward difference would introduce some bias because the value of  $I_x$  at a certain point is dependent on an adjacent image value in the positive x-direction only. The central difference will balance out the value of  $I_x$  since both directions are considered. This will make the approximation more robust to variations in image content.

The central difference for a single (space) variable can be obtained using the numeric differentiation method shown in class

$$\begin{aligned} f(x, t) &= f(x, t) \\ f(x + \Delta x, t) &= f(x, t) + \Delta x f'(x, t) + O(\Delta x^2) \\ f(x - \Delta x, t) &= f(x, t) - \Delta x f'(x, t) + O(\Delta x^2) \end{aligned}$$

After eliminating  $f(x, t)$  terms on the right side of the above equations, we obtain our approximation. Note that the approximation includes an error term  $O(\Delta x)$ .

$$f'(x, t) = \frac{f(x + \Delta x, t) - f(x - \Delta x, t)}{2\Delta x}$$

Expanding this process to two (space) dimensions yields

$$I_x(x, y, t) = \frac{I(x + \Delta x, y, t) - I(x - \Delta x, y, t)}{2\Delta x}$$

$$I_y(x, y, t) = \frac{I(x, y + \Delta y, t) - I(x, y - \Delta y, t)}{2\Delta y}$$

In order to obtain approximations for  $I_{xx}$  and  $I_{yy}$ , we can simply add additional terms to the Taylor Series expansion and then repeat the elimination process

$$I_{xx}(x, y, t) = \frac{I(x + \Delta x, y, t) - 2I(x, y, t) + I(x - \Delta x, y, t)}{\Delta x^2}$$

$$I_{yy}(x, y, t) = \frac{I(x, y + \Delta y, t) - 2I(x, y, t) + I(x, y - \Delta y, t)}{\Delta y^2}$$

Finally, we need to obtain an approximation for the mixed partial derivative  $I_{xy}$ . This is achieved by taking the central difference approximation for  $I_x$  and applying another central difference with respect to  $y$

$$I_x(x, y, t) = \frac{I(x + \Delta x, y, t)}{2\Delta x} - \frac{I(x - \Delta x, y, t)}{2\Delta x}$$

$$I_{xy}(x, y, t) = \frac{\partial}{\partial y} \left( \frac{I(x + \Delta x, y, t)}{2\Delta x} \right) - \frac{\partial}{\partial y} \left( \frac{I(x - \Delta x, y, t)}{2\Delta x} \right)$$

$$I_{xy}(x, y, t) = \frac{1}{2\Delta y} \left( \frac{I(x + \Delta x, y + \Delta y, t)}{2\Delta x} - \frac{I(x + \Delta x, y - \Delta y, t)}{2\Delta x} \right) - \frac{1}{2\Delta y} \left( \frac{I(x - \Delta x, y + \Delta y, t)}{2\Delta x} - \frac{I(x - \Delta x, y - \Delta y, t)}{2\Delta x} \right)$$

$$I_{xy}(x, y, t) = \frac{I(x + \Delta x, y + \Delta y, t) - I(x + \Delta x, y - \Delta y, t) - I(x - \Delta x, y + \Delta y, t) + I(x - \Delta x, y - \Delta y, t)}{4\Delta x \Delta y}$$

## 5 Experimental Results

## 6 Summary