First- and Second-Order Coherence of a Point Source Through Spacetime Foam

Author One
Department of Physics, University X, City, Country
(Dated: April 22, 2026)

We derive the first- and second-order coherence functions for light from point and extended sources propagating through spacetime foam parametrized by the Ng–Perlman exponent α. In the narrow-band limit, foam-induced arrival-time jitter suppresses the observed first-order coherence by $e^{-\sigma_\phi^2[1-C_\alpha(\tau)]}$ (where $\sigma_\phi = 2\pi\sigma_\ell(D)/\lambda$) while leaving the second-order coherence unchanged, producing a Siegert violation $\mathcal{S}(\tau) = |g^{(1)}_{\rm src}|^2(1-e^{-2\sigma_\phi^2[1-C_\alpha(\tau)]}) \geq 0$ that encodes α and D independently of the source spectrum. For extended sources, photon pairs from directions separated by more than the transverse foam coherence length $r_\perp$ accumulate independent jitters, suppressing the zero-lag bunching peak by $\delta g^{(2)} = 1 - [1 + (\sigma_0/\tau_c)^2]^{-1/2}$ in a way that grows with source angular size and directly probes $r_\perp$. The fringe-erasure condition of Lieu & Hillman and the Strehl-ratio suppression of Ng & Perlman are recovered as limiting cases, with temporal HBT less sensitive than imaging by a factor $\Delta\lambda/\lambda$. Joint measurements of the Siegert discrepancy and bunching suppression across sources of varying redshift and angular size constrain $\{\alpha, \sigma_\ell(D), r_\perp\}$ simultaneously.

I. MODEL

A. Arrival-Time Jitter as a Time Reparameterization

We model the detected field as a stochastically time-warped version of the intrinsic source field:

$$E_{\rm obs}(t) = E_{\rm src}(t - \delta t(t))\,, \tag{1}$$

where $E_{\rm src}(t)$ is a stationary zero-mean complex Gaussian process (a single chaotic/thermal mode) and $\delta t(t)$ is the foam-induced arrival-time jitter, a stationary stochastic process statistically independent of $E_{\rm src}$. The jitter arises from cumulative path-length fluctuations along the line of sight of length $\ell^{(D)}$:

$$\delta t(t) = \frac{1}{c}\,\Delta\ell^{(D)}_{\rm tot}(t). \tag{2}$$

B. Ng–Perlman Variance and Temporal Covariance

Following Ng and collaborators [1–3], the single-ray path-length variance obeys the power-law scaling

$$\sigma^2_\ell(D) = A_\alpha\,[\ell^{(D)}]^{2(1-\alpha)}\,\ell_{\rm P}^{2\alpha}, \tag{3}$$

where $\ell_{\rm P} = \sqrt{\hbar G/c^3} \simeq 1.616\times10^{-35}$ m is the Planck length, α parametrizes the foam model ($\alpha = 1$: conservative; $\alpha = 2/3$: holographic; $\alpha = 1/2$: random walk), and A_α is a dimensionless coefficient of order unity. The temporal covariance of the aggregate path-length fluctuation along a single ray takes the factorized form

$$\big\langle \Delta\ell^{(D)}_{\rm tot}(t)\,\Delta\ell^{(D)}_{\rm tot}(t+\tau)\big\rangle = \sigma^2_\ell(D)\,C_\alpha(\tau; D), \tag{4}$$

where the temporal decorrelation function $C_\alpha(\tau; D)$ satisfies $C_\alpha(0; D) = 1$ and decays to zero as $\tau \to \infty$.

C. Differential Jitter and Its Distribution

Define the differential jitter between two detection times separated by lag τ:

$$\Delta(\tau) \equiv \delta t(t+\tau) - \delta t(t). \tag{5}$$

By stationarity, the statistics of $\Delta(\tau)$ depend only on τ. Its variance follows directly from Eq. (4):

$$\begin{aligned}\sigma^2_\Delta(\tau) &\equiv \langle\Delta(\tau)^2\rangle \\ &= \frac{1}{c^2}\big\langle[\Delta\ell_{\rm tot}(t+\tau) - \Delta\ell_{\rm tot}(t)]^2\big\rangle \\ &= \frac{2\sigma^2_\ell(D)}{c^2}\big[1 - C_\alpha(\tau; D)\big].\end{aligned} \tag{6}$$

Because $\Delta(\tau)$ is a sum of contributions from many weakly correlated foam cells, the central limit theorem implies it is Gaussian with zero mean. We therefore write its probability density as

$$p(\Delta;\tau) = \frac{1}{\sqrt{2\pi}\,\sigma_\Delta(\tau)} \exp\left(-\frac{\Delta^2}{2\sigma_\Delta^2(\tau)}\right). \tag{7}$$

Note the two limits: $\sigma_\Delta^2(0) = 0$ (no jitter between simultaneous arrivals) and $\sigma_\Delta^2(\infty) = 2\sigma_\ell^2(D)/c^2$ (maximal decorrelation at large lag).

II. FIRST-ORDER COHERENCE

A. General Result

The observed first-order coherence is

$$g^{(1)}_{\rm obs}(\tau) = \frac{\langle E^*_{\rm obs}(t)\, E_{\rm obs}(t+\tau)\rangle}{\langle |E_{\rm obs}(t)|^2\rangle}. \tag{8}$$

Substituting Eq. (1) and letting $u = t - \delta t(t)$, $v = t + \tau - \delta t(t+\tau)$, so that $v - u = \tau - \Delta(\tau)$, stationarity of $E_{\rm src}$ gives

$$\langle E^*_{\rm src}(u)\, E_{\rm src}(v)\rangle = \langle |E_{\rm src}|^2\rangle\, g^{(1)}_{\rm src}(\tau - \Delta(\tau)). \tag{9}$$

Since δt is independent of $E_{\rm src}$, averaging over the jitter yields

$$\boxed{g^{(1)}_{\rm obs}(\tau) = \left\langle g^{(1)}_{\rm src}(\tau - \Delta(\tau))\right\rangle_{\delta t} = \int_{-\infty}^{\infty} p(\Delta;\tau)\, g^{(1)}_{\rm src}(\tau - \Delta)\, d\Delta.} \tag{10}$$

The observed first-order coherence is the intrinsic coherence function convolved with the Gaussian differential-jitter kernel.

B. Closed Form for a Lorentzian Source

A thermal or AGN source typically has a Lorentzian power spectrum, so

$$g^{(1)}_{\rm src}(\tau) = e^{-|\tau|/\tau_c}, \tag{11}$$

where τ_c is the coherence time. Evaluating Eq. (10) by completing the square and splitting on the sign of $\tau - \Delta$ gives, for $\tau > 0$:

$$g^{(1)}_{\rm obs}(\tau) = e^{\sigma_\Delta^2/2\tau_c^2} \cdot \frac{1}{2}\left[e^{-\tau/\tau_c}\,\mathrm{erfc}\left(\frac{\sigma_\Delta/\tau_c - \tau/\sigma_\Delta}{\sqrt{2}}\right) + e^{\tau/\tau_c}\,\mathrm{erfc}\left(\frac{\sigma_\Delta/\tau_c + \tau/\sigma_\Delta}{\sqrt{2}}\right)\right], \tag{12}$$

where $\sigma_\Delta \equiv \sigma_\Delta(\tau)$. In the weak-jitter limit $\sigma_\Delta \ll \tau_c$, expanding the erfc gives

$$g^{(1)}_{\rm obs}(\tau) \approx e^{-|\tau|/\tau_c}\left[1 - \frac{\sigma_\Delta^2(\tau)}{\tau_c}\,|\tau| + \mathcal{O}(\sigma_\Delta^4)\right]. \tag{13}$$

The normalization $g^{(1)}_{\rm obs}(0) = 1$ is preserved exactly, since $\sigma_\Delta(0) = 0$.

C. Narrow-Band Limit

For a quasi-monochromatic source with center frequency ω_0 and bandwidth $\Delta\omega \ll \omega_0$, write $E_{\rm src}(t) = A(t)\, e^{-i\omega_0 t}$ where $A(t)$ varies on timescale $\tau_c = 1/\Delta\omega$. For jitter $|\delta t| \ll \tau_c$, the envelope is unaffected, $A(t - \delta t) \approx A(t)$, and the jitter acts purely as a carrier phase modulation:

$$E_{\rm obs}(t) \approx A(t)\, e^{-i\omega_0 t}\, e^{i\omega_0 \delta t(t)}. \tag{14}$$

Since $A(t)$ and $\delta t(t)$ are independent, the first-order coherence factorizes. The jitter contribution is the characteristic function of the Gaussian $\Delta(\tau)$ evaluated at ω_0:

$$\left\langle e^{i\omega_0 \Delta(\tau)} \right\rangle_{\delta t} = e^{-\omega_0^2 \sigma_\Delta^2(\tau)/2}. \tag{15}$$

Substituting $\omega_0 = 2\pi c/\lambda$ and Eq. (6):

$$\boxed{g^{(1)}_{\rm obs}(\tau) = g^{(1)}_{\rm src}(\tau) \cdot \exp\!\left(-\sigma_\phi^2 \left[1 - C_\alpha(\tau; D)\right]\right),} \tag{16}$$

where we define the rms accumulated phase

$$\sigma_\phi \equiv \frac{2\pi\, \sigma_\ell(D)}{\lambda} = \frac{2\pi\, \sqrt{A_\alpha}\, [\ell^{(D)}]^{1-\alpha}\, \ell_{\rm P}^{\alpha}}{\lambda}. \tag{17}$$

We write the foam suppression factor as

$$\mathcal{F}_{\rm foam}(\tau) \equiv e^{-\sigma_\phi^2 [1 - C_\alpha(\tau; D)]}, \tag{18}$$

so $g^{(1)}_{\rm obs}(\tau) = g^{(1)}_{\rm src}(\tau) \cdot \mathcal{F}_{\rm foam}(\tau)$. At $\tau = 0$: $\mathcal{F}_{\rm foam}(0) = 1$. At $\tau \to \infty$: $\mathcal{F}_{\rm foam}(\infty) = e^{-\sigma_\phi^2}$.

III. CONNECTION TO FRINGE VISIBILITY AND STREHL RATIO

A. Fringe Visibility and the Lieu–Hillman Condition

The fringe visibility measured in a stellar interferometer at time delay τ is $V(\tau) = |g^{(1)}_{\rm obs}(\tau)|$. From Eq. (16):

$$V(\tau) = V_{\rm src}(\tau) \cdot \mathcal{F}_{\rm foam}(\tau), \tag{19}$$

where $V_{\rm src}(\tau) = |g^{(1)}_{\rm src}(\tau)|$ is the intrinsic source visibility. Lieu & Hillman [4] argue that fringes are erased when the accumulated phase randomizes over $[0, 2\pi)$, i.e. $\sigma_\phi \gtrsim 1$. In our language this is the limit $\mathcal{F}_{\rm foam} \to 0$: the foam suppression factor collapses and $V \to 0$. The Lieu–Hillman condition $\Delta\phi \geq 2\pi$, where $\Delta\phi \simeq 2\pi\sigma_\ell(D)/\lambda$ [1], is precisely $\sigma_\phi \geq 1$. Our result extends this binary criterion to a continuous suppression curve $V(\tau)$ whose shape encodes the temporal decorrelation $C_\alpha(\tau; D)$.

B. Strehl Ratio

For spatially uncorrelated phase noise of rms σ_ϕ across the telescope aperture, the Strehl ratio (ratio of the observed PSF peak to the diffraction-limited peak) is [1]

$$S = e^{-\sigma_\phi^2}. \tag{20}$$

Comparing with Eq. (18), the Strehl ratio equals the long-lag limit of the foam suppression factor:

$$S = \mathcal{F}_{\rm foam}(\tau \to \infty). \tag{21}$$

Both the Strehl and the Lieu–Hillman fringe visibility measure the same underlying quantity $\sigma_\phi^2 = (2\pi\sigma_\ell/\lambda)^2$ — the total accumulated phase variance. The difference is observational channel: Strehl and fringes probe the *spatial* phase variation across the aperture, while the temporal $g^{(1)}$ probes the *temporal* phase variation along a single ray as a function of lag.

C. Why Temporal HBT is Less Sensitive

In the narrow-band limit the relevant sensitivity parameter for the temporal measurement is $\sigma_\Delta(\tau)/\tau_c$, not σ_ϕ directly. Using $c\tau_c = \lambda^2/\Delta\lambda$ for fixed fractional bandwidth:

$$\frac{\sigma_\Delta(\tau)}{\tau_c} = \frac{\sqrt{2[1 - C_\alpha(\tau)]}\, \sigma_\ell(D)}{c\tau_c} = \sigma_\phi \cdot \frac{\sqrt{2[1 - C_\alpha(\tau)]}}{\omega_0 \tau_c} \sim \sigma_\phi \cdot \frac{\Delta\lambda}{\lambda} \cdot \sqrt{2[1 - C_\alpha]}. \tag{22}$$

TABLE I. Comparison of foam observables at $D = 1$ Gpc, $\lambda = 500$ nm, $\Delta\lambda/\lambda = 10^{-3}$, $A_\alpha = 1$.

Method	Probe	Sensitivity
Strehl / Lieu–Hillman	$\sigma_\phi = 2\pi\sigma_\ell/\lambda$	Direct
Temporal $g^{(1)}$ at lag τ	$\sigma_\phi\sqrt{1 - C_\alpha(\tau)} \cdot (\Delta\lambda/\lambda)$	Suppressed by $\Delta\lambda/\lambda$

The suppression factor $\Delta\lambda/\lambda \ll 1$ means that even when $\sigma_\phi \gg 1$ (fringes erased), the temporal HBT ratio can be of order unity or smaller.

For $\alpha = 1/2$ at $D = 1$ Gpc: $\sigma_\phi \approx 4\pi \times 10^9$ (fringes completely erased), while $\sigma_\Delta/\tau_c \approx 0.59$ (marginally accessible to HBT). This reflects the fact that phase-coherence constraints already decisively rule out $\alpha = 1/2$ at optical wavelengths [5], whereas the temporal HBT signature for the same model is only at the edge of detectability.

IV. SECOND-ORDER COHERENCE

A. Conditional Siegert Relation

Fix a realization of the jitter process $\delta t(\cdot)$. Conditioned on it, $E_{\rm obs}(t)$ is a deterministic time-warp of a complex Gaussian field, hence is itself complex Gaussian. For any complex Gaussian field the Siegert relation holds exactly, so

$$g^{(2)}_{\rm obs}(\tau \mid \delta t) - 1 = \left| g^{(1)}_{\rm obs}(\tau \mid \delta t) \right|^2 . \tag{23}$$

Given a fixed δt, the differential jitter $\Delta(\tau)$ is a deterministic number, so $g^{(1)}_{\rm obs}(\tau \mid \delta t) = g^{(1)}_{\rm src}(\tau - \Delta(\tau))$. Using the intrinsic Siegert relation $g^{(2)}_{\rm src} - 1 = |g^{(1)}_{\rm src}|^2$:

$$g^{(2)}_{\rm obs}(\tau \mid \delta t) - 1 = \left| g^{(1)}_{\rm src}(\tau - \Delta(\tau)) \right|^2 = g^{(2)}_{\rm src}(\tau - \Delta(\tau)) - 1. \tag{24}$$

B. Unconditional Second-Order Coherence

Averaging over the jitter distribution:

$$\boxed{g^{(2)}_{\rm obs}(\tau) - 1 = \left\langle g^{(2)}_{\rm src}(\tau - \Delta(\tau)) - 1 \right\rangle_{\delta t} = \int_{-\infty}^{\infty} p(\Delta; \tau) \left[g^{(2)}_{\rm src}(\tau - \Delta) - 1 \right] d\Delta.} \tag{25}$$

The structure is identical to Eq. (10): both $g^{(1)}_{\rm obs}$ and $g^{(2)}_{\rm obs}$ are convolutions of their intrinsic counterparts with the *same* Gaussian jitter kernel.

C. Narrow-Band Limit: $g^{(2)}$ is Unchanged

In the narrow-band limit, the jitter $\delta t \ll \tau_c$ does not affect the amplitude envelope $A(t)$. Since intensity $I_{\rm obs}(t) = |E_{\rm obs}(t)|^2 = |A(t)|^2 = I_{\rm src}(t)$, the intensity statistics are identical to the intrinsic source:

$$g^{(2)}_{\rm obs}(\tau) = g^{(2)}_{\rm src}(\tau) \qquad \text{(narrow-band limit, exactly).} \tag{26}$$

The physical reason is that intensity is blind to the carrier phase $e^{i\omega_0 \delta t(t)}$: a timing jitter that shifts the wavepacket by $\delta t \ll \tau_c$ does not change the envelope shape, only the phase of the underlying oscillation. By contrast, $g^{(1)}$ *is* sensitive to this phase because $\omega_0 \delta t \sim \omega_0 \sigma_\ell / c = \sigma_\phi$ can be large even when $\delta t \ll \tau_c$: for $\alpha = 1/2$ at $D = 1$ Gpc and $\lambda = 500$ nm, $\sigma_\phi \sim 4\pi \times 10^9$.

V. SIEGERT RELATION VIOLATION

A. General Bound

Define the Siegert discrepancy:

$$\mathcal{S}(\tau) \equiv \left[g^{(2)}_{\rm obs}(\tau) - 1\right] - \left|g^{(1)}_{\rm obs}(\tau)\right|^2 . \tag{27}$$

Let $X \equiv g^{(1)}_{\rm src}(\tau - \Delta(\tau))$, a real random variable. From Eqs. (10) and (25):

$$g^{(1)}_{\rm obs}(\tau) = \langle X \rangle_{\delta t}, \qquad g^{(2)}_{\rm obs}(\tau) - 1 = \langle X^2 \rangle_{\delta t}. \tag{28}$$

Therefore

$$\mathcal{S}(\tau) = \langle X^2 \rangle - \langle X \rangle^2 = {\rm Var}(X) \geq 0, \tag{29}$$

by the Cauchy–Schwarz inequality, with equality only when $\Delta(\tau)$ has zero variance (i.e., at $\tau = 0$ or in the absence of foam). The Siegert relation is violated for all $\tau \neq 0$ whenever foam is present.

B. Explicit Form in the Narrow-Band Limit

In the narrow-band limit, $g^{(2)}_{\rm obs} = g^{(2)}_{\rm src}$ exactly (Eq. 26), while $g^{(1)}_{\rm obs}$ is suppressed by $\mathcal{F}_{\rm foam}(\tau)$ (Eq. 16). Using the intrinsic Siegert relation $g^{(2)}_{\rm src}(\tau) - 1 = |g^{(1)}_{\rm src}(\tau)|^2$:

$$\boxed{\mathcal{S}(\tau) = \left|g^{(1)}_{\rm src}(\tau)\right|^2 \left(1 - e^{-2\sigma_\phi^2[1 - C_\alpha(\tau;D)]}\right).} \tag{30}$$

C. Properties

The key properties of $\mathcal{S}(\tau)$ are summarized in Table II.

TABLE II. Properties of the Siegert discrepancy $\mathcal{S}(\tau)$ in the narrow-band limit.

Limit	$\mathcal{S}(\tau)$	Reason
$\tau = 0$	0	$\Delta(0) = 0$; both coherences agree
$\sigma_\phi \ll 1$ (weak jitter)	$\approx 2\sigma_\phi^2[1 - C_\alpha]\,\lvert g^{(1)}_{\rm src}\rvert^2$	Linear in phase variance
$\sigma_\phi \gg 1$ (strong jitter)	$\approx \lvert g^{(1)}_{\rm src}(\tau)\rvert^2$	$g^{(1)}_{\rm obs} \to 0$; full violation
$\tau \to \infty$	$\to 0$	$\lvert g^{(1)}_{\rm src}\rvert \to 0$ faster

D. Physical Interpretation

In the narrow-band limit foam acts as a *phase screen*: it modulates the carrier phase $\omega_0 \delta t(t)$ randomly without touching the intensity envelope. This asymmetry is the root cause of the Siegert violation. The discrepancy $\mathcal{S}(\tau)$ is a direct observable — both $g^{(1)}_{\rm obs}$ and $g^{(2)}_{\rm obs}$ are measurable from the same dataset — and carries a specific distance dependence $\sigma_\phi^2 \propto [\ell^{(D)}]^{2(1-\alpha)}$ that distinguishes foam from instrumental effects, which would not scale with source distance. It is also independent of the unknown intrinsic source spectrum: whatever $g^{(1)}_{\rm src}(\tau)$, the violation is controlled solely by α, D, and $C_\alpha(\tau; D)$.

VI. EXTENDED SOURCES: TRANSVERSE FOAM COHERENCE

A. Field Model and Spatiotemporal Covariance

We now generalize to an extended incoherent source whose angular brightness distribution is $W(\boldsymbol{\theta})$ (normalized to unity). The detected field is a weighted superposition over rays labelled by direction offset $\boldsymbol{\theta}$:

$$E_{\rm obs}(t) = \int d\boldsymbol{\theta}\, W^{1/2}(\boldsymbol{\theta})\, E_{\boldsymbol{\theta}}(t - \delta t(t;\boldsymbol{\theta}))\,, \tag{31}$$

where the $E_{\boldsymbol{\theta}}$ are independent, identically distributed complex Gaussian processes (one per sky direction), and each ray carries its own foam jitter $\delta t(t;\boldsymbol{\theta})$. The spatiotemporal covariance of the path-length aggregate is

$$\big\langle \Delta\ell_{\rm tot}(t;\boldsymbol{\theta})\, \Delta\ell_{\rm tot}(t+\tau;\boldsymbol{\theta}')\big\rangle = \sigma_\ell^2(D)\, C_\alpha(\tau;D)\, \Phi\!\left(\frac{b(\boldsymbol{\theta},\boldsymbol{\theta}')}{r_\perp}\right), \tag{32}$$

where $b(\boldsymbol{\theta},\boldsymbol{\theta}') \simeq \ell^{(D)}|\boldsymbol{\theta}-\boldsymbol{\theta}'|$ is the transverse separation between rays at distance $\ell^{(D)}$, $r_\perp$ is the foam transverse coherence length, and Φ is the transverse coherence function satisfying $\Phi(0) = 1$ and $\Phi(b/r_\perp) \to 0$ for $b \gg r_\perp$.

B. First-Order Coherence is Unchanged

The numerator of $g^{(1)}_{\rm obs}(\tau)$ involves cross terms $\langle E^*_{\boldsymbol{\theta}}(\ldots)\, E_{\boldsymbol{\theta}'}(\ldots)\rangle$. Because different source directions are spatially incoherent,

$$\langle E^*_{\boldsymbol{\theta}}(s)\, E_{\boldsymbol{\theta}'}(s')\rangle \propto \delta(\boldsymbol{\theta}-\boldsymbol{\theta}'), \tag{33}$$

so only the diagonal $\boldsymbol{\theta} = \boldsymbol{\theta}'$ contributes. Each diagonal term reproduces the single-ray result of Section II C, and the angular integral over W gives unity. Therefore:

$$g^{(1)}_{\rm obs}(\tau) = g^{(1)}_{\rm src}(\tau) \cdot e^{-\sigma_\phi^2[1-C_\alpha(\tau;D)]}, \tag{34}$$

identical to the point-source result Eq. (16). The first-order coherence probes only single-ray statistics; the transverse foam structure leaves no imprint on $g^{(1)}$.

C. Cross-Ray Differential Jitter

For $g^{(2)}$, the Wick contraction in the Gaussian moment theorem enforces pairings between rays $\boldsymbol{\theta}$ and $\boldsymbol{\theta}'$ that need not be equal. The relevant quantity is the *cross-ray differential jitter*:

$$\Delta_{\boldsymbol{\theta},\boldsymbol{\theta}'}(\tau) \equiv \delta t(t+\tau;\boldsymbol{\theta}) - \delta t(t;\boldsymbol{\theta}'), \tag{35}$$

the time offset between the "second photon" arriving along $\boldsymbol{\theta}$ at $t+\tau$ and the "first photon" along $\boldsymbol{\theta}'$ at t. Computing its variance from Eq. (32):

$$\begin{aligned}
\sigma^2_\Delta(\tau;\boldsymbol{\theta},\boldsymbol{\theta}') &= \frac{1}{c^2}\big\langle[\Delta\ell_{\rm tot}(t+\tau;\boldsymbol{\theta}) - \Delta\ell_{\rm tot}(t;\boldsymbol{\theta}')]^2\big\rangle \\
&= \frac{1}{c^2}\left[\sigma_\ell^2 + \sigma_\ell^2 - 2\sigma_\ell^2\, C_\alpha(\tau)\, \Phi\!\left(\frac{b}{r_\perp}\right)\right] \\
&= \frac{2\sigma_\ell^2(D)}{c^2}\left[1 - C_\alpha(\tau;D)\, \Phi\!\left(\frac{b(\boldsymbol{\theta},\boldsymbol{\theta}')}{r_\perp}\right)\right].
\end{aligned} \tag{36}$$

Setting $\boldsymbol{\theta} = \boldsymbol{\theta}'$ (and hence $\Phi(0) = 1$) recovers the single-ray result Eq. (6) exactly. For $b \gg r_\perp$, $\Phi \to 0$ and the variance becomes $2\sigma_\ell^2/c^2$ *at all lags, including* $\tau = 0$ — a new effect absent in the point-source case.

D. General Second-Order Coherence

By the Gaussian moment theorem, the unconditional $g^{(2)}$ for an extended source is an angularly weighted mixture over all ray pairs:

$$\boxed{g^{(2)}_{\rm obs}(\tau) - 1 = \iint d\boldsymbol{\theta}\, d\boldsymbol{\theta}'\; W(\boldsymbol{\theta})W(\boldsymbol{\theta}') \int_{-\infty}^{\infty} p_{\boldsymbol{\theta},\boldsymbol{\theta}'}(\Delta;\tau)\left[g^{(2)}_{\rm src}(\tau-\Delta) - 1\right] d\Delta,} \tag{37}$$

where $p_{\boldsymbol{\theta},\boldsymbol{\theta}'}(\Delta;\tau)$ is Gaussian with variance $\sigma^2_\Delta(\tau;\boldsymbol{\theta},\boldsymbol{\theta}')$ from Eq. (36). In the limit $\Phi \equiv 1$ (all rays transversely coherent), every kernel reduces to the single-ray $p_{\rm 1D}$ and Eq. (37) reduces to the point-source result Eq. (25).

E. Effective Transverse Coherence Factor

To reduce Eq. (37) to a tractable form, define the effective transverse coherence factor:

$$\Phi_{\rm eff} \equiv \iint d\boldsymbol{\theta}\, d\boldsymbol{\theta}'\; W(\boldsymbol{\theta})W(\boldsymbol{\theta}')\, \Phi\!\left(\frac{b(\boldsymbol{\theta},\boldsymbol{\theta}')}{r_\perp}\right). \tag{38}$$

This replaces the ray-dependent differential variance with a single effective variance:

$$\sigma^2_{\tau,\rm eff}(\tau) \equiv \frac{2\sigma^2_\ell(D)}{c^2}\Big[1 - C_\alpha(\tau;D)\,\Phi_{\rm eff}\Big], \tag{39}$$

and allows Eq. (37) to be approximated by the simpler 1D convolution of Eq. (25) with σ^2_Δ replaced by $\sigma^2_{\tau,\rm eff}$. The two limits are: $\Phi_{\rm eff} \to 1$ (point-like source or $r_\perp \gg \ell^{(D)}\Delta\theta$) recovers the single-ray result; $\Phi_{\rm eff} \to 0$ (fully incoherent, $r_\perp \ll \ell^{(D)}\Delta\theta$) maximizes the effective variance.

F. Zero-Lag Bunching Peak Suppression

At $\tau = 0$, $C_\alpha(0;D) = 1$, so $\sigma^2_{\tau,\rm eff}(0)$ reduces to

$$\sigma^2_0 \equiv \frac{2\sigma^2_\ell(D)}{c^2}(1 - \Phi_{\rm eff}), \tag{40}$$

which is strictly positive whenever $\Phi_{\rm eff} < 1$. Evaluating the zero-lag convolution in Eq. (37) for a Lorentzian $g^{(2)}_{\rm src}(\tau) = 1 + e^{-|\tau|/\tau_c}$:

$$\boxed{g^{(2)}_{\rm obs}(0) = 1 + \frac{1}{\sqrt{1 + (\sigma_0/\tau_c)^2}},} \tag{41}$$

so the zero-lag bunching peak is suppressed below its ideal value of 2 by

$$\delta g^{(2)} \equiv 2 - g^{(2)}_{\rm obs}(0) = 1 - \frac{1}{\sqrt{1 + (\sigma_0/\tau_c)^2}}. \tag{42}$$

In the point-source limit $\Phi_{\rm eff} \to 1$, $\sigma_0 \to 0$ and $\delta g^{(2)} \to 0$ as expected. Since $\sigma^2_0 \propto (1 - \Phi_{\rm eff})$ and $\Phi_{\rm eff}$ decreases as the source angular diameter increases at fixed D and $r_\perp$, more extended sources experience greater suppression of the bunching peak.

The physical origin is distinct from the temporal phase-scrambling effect. Even at $\tau = 0$, photon pairs from *different* sky directions $\boldsymbol{\theta} \neq \boldsymbol{\theta}'$ accumulate independent instantaneous jitters $\delta t(t;\boldsymbol{\theta}) \neq \delta t(t;\boldsymbol{\theta}')$ when $b \gg r_\perp$. Their relative offset $\delta t(t;\boldsymbol{\theta}) - \delta t(t;\boldsymbol{\theta}')$ blurs the $g^{(2)}(0)$ measurement even though $\tau = 0$. This is a purely *spatial* effect with no analogue in the point-source case.

G. Extended-Source Siegert Discrepancy

In the narrow-band limit, $g^{(2)}_{\rm obs}(\tau) - 1$ for the extended source is obtained from Eq. (37) with the Gaussian carrier-phase substitution. The result is

$$g^{(2)}_{\rm obs}(\tau) - 1 = \left|g^{(1)}_{\rm src}(\tau)\right|^2 \mathcal{G}(\tau), \tag{43}$$

where

$$\mathcal{G}(\tau) = \iint d\boldsymbol{\theta}\, d\boldsymbol{\theta}'\, W(\boldsymbol{\theta}) W(\boldsymbol{\theta}') \exp\!\big(-2\sigma_\phi^2 \big[1 - C_\alpha(\tau)\big]\big[1 - \Phi(b/r_\perp)\big]\big)\,. \tag{44}$$

Since $g^{(1)}_{\rm obs}(\tau) = g^{(1)}_{\rm src}(\tau) \cdot e^{-\sigma_\phi^2[1-C_\alpha]}$ (Eq. 34), the Siegert discrepancy is:

$$\boxed{\mathcal{S}(\tau) = \left|g^{(1)}_{\rm src}(\tau)\right|^2 \left[\mathcal{G}(\tau) - e^{-2\sigma_\phi^2[1-C_\alpha(\tau;D)]}\right].} \tag{45}$$

For a point source ($\Phi \equiv 1$, so $\mathcal{G} \to 1$), this reduces to the point-source result Eq. (30).

H. Two Distinct Signatures

The two observable signatures of foam for extended sources are summarized in Table III.

TABLE III. Two foam signatures for an extended source in the narrow-band limit.

Signature	Origin	Non-zero at $\tau = 0$?	Depends on $r_\perp$?
Siegert discrepancy $\mathcal{S}(\tau)$	Temporal phase scrambling (single ray)	No	Only via $\mathcal{G}(\tau)$
Bunching suppression $\delta g^{(2)}$	Transverse spatial decorrelation	Yes	Yes — direct probe of $r_\perp$

The two signatures are complementary. The Siegert discrepancy $\mathcal{S}(\tau)$ is controlled by σ_ϕ^2 and $C_\alpha(\tau; D)$, probing α and D independently of $r_\perp$. The bunching suppression $\delta g^{(2)}$ is controlled by $\sigma_0^2 \propto \sigma_\ell^2(D)(1 - \Phi_{\rm eff})$, providing a direct handle on the transverse coherence length $r_\perp$ through the dependence of $\Phi_{\rm eff}$ on source angular size. Joint measurements of point-like and extended sources at the same redshift therefore allow $\{\alpha, \sigma_\ell(D), r_\perp\}$ to be constrained simultaneously and largely independently of the unknown intrinsic source spectrum.

ACKNOWLEDGMENTS

[Acknowledgments to be added.]

[1] Y. J. Ng and E. S. Perlman, Universe **8**, 382 (2022).
[2] Y. J. Ng, Modern Physics Letters A **18**, 1073 (2003).
[3] Y. J. Ng, Physical Review Letters **86**, 2946 (2001).
[4] R. Lieu and L. W. Hillman, The Astrophysical Journal **585**, L77 (2003).
[5] E. S. Perlman, S. A. Rappaport, W. A. Christiansen, Y. J. Ng, J. DeVore, and D. Pooley, The Astrophysical Journal **805**, 10 (2015).