

Class groups are essential in our life

Alex Gélin

Laboratoire de Mathématiques de Versailles
Paris-Saclay – UVSQ – CNRS

ANTS Summer School – *at home* ☹️

2020/06/25

Why do we study class groups?

Why do we study class groups?



Why do we study class groups?

- Because Gauss did!

Why do we study class groups?

- Because Gauss did!
- Class groups are everywhere

Why do we study class groups?

- Because Gauss did!
- Class groups are everywhere
- Beautiful mathematical challenge

Why do we study class groups?

- Because Gauss did!
- Class groups are everywhere
- Beautiful mathematical challenge
- Finite group \implies Applications in Cryptology

Why do we study class groups?

- Because Gauss did!
- Class groups are everywhere
- Beautiful mathematical challenge
- Finite group \implies Applications in Cryptology
- Make use of trendy structures

FROM ALEX GELIN'S
MATHEMATICAL WORLD

CLASS GROUPS

AND WHERE
TO FIND THEM



Number fields

K number field \Rightarrow finite-degree extension of **Q**

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

Number fields

\mathbf{K} number field \Rightarrow finite-degree extension of \mathbf{Q}

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

T is a **defining polynomial** of \mathbf{K} and there exist infinitely many of them

Number fields

\mathbf{K} number field \Rightarrow finite-degree extension of \mathbf{Q}

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

T is a **defining polynomial** of \mathbf{K} and there exist infinitely many of them

$\mathcal{O}_{\mathbf{K}}$ denotes the ring of integers of \mathbf{K}

Number fields

\mathbf{K} number field \Rightarrow finite-degree extension of \mathbf{Q}

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

T is a **defining polynomial** of \mathbf{K} and there exist infinitely many of them

$\mathcal{O}_{\mathbf{K}}$ denotes the ring of integers of \mathbf{K}

Two manners for representing elements:

Number fields

\mathbf{K} number field \Rightarrow finite-degree extension of \mathbf{Q}

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

T is a **defining polynomial** of \mathbf{K} and there exist infinitely many of them

$\mathcal{O}_{\mathbf{K}}$ denotes the ring of integers of \mathbf{K}

Two manners for representing elements:

- The minimal polynomial

Number fields

\mathbf{K} number field \Rightarrow finite-degree extension of \mathbf{Q}

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

T is a **defining polynomial** of \mathbf{K} and there exist infinitely many of them

$\mathcal{O}_{\mathbf{K}}$ denotes the ring of integers of \mathbf{K}

Two manners for representing elements:

- The minimal polynomial

A polynomial modulo T

Number fields

\mathbf{K} number field \Rightarrow finite-degree extension of \mathbf{Q}

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

T is a **defining polynomial** of \mathbf{K} and there exist infinitely many of them

$\mathcal{O}_{\mathbf{K}}$ denotes the ring of integers of \mathbf{K}

Two manners for representing elements:

- The minimal polynomial

A polynomial modulo T

- The conjugates

Number fields

\mathbf{K} number field \Rightarrow finite-degree extension of \mathbf{Q}

Primitive Element Theorem $\Rightarrow \exists T \in \mathbf{Z}[X]$ monic such that

$$\mathbf{K} \simeq \mathbf{Q}[X] / \langle T \rangle$$

T is a **defining polynomial** of \mathbf{K} and there exist infinitely many of them

$\mathcal{O}_{\mathbf{K}}$ denotes the ring of integers of \mathbf{K}

Two manners for representing elements:

- The minimal polynomial

A polynomial modulo T

- The conjugates

Vector of $[\mathbf{K}:\mathbf{Q}]$ complex coordinates \Rightarrow Vector of $[\mathbf{K}:\mathbf{Q}]$ real coordinates

Two interesting structures in number fields:

Two interesting structures in number fields:

- Group of fractional ideals

Two interesting structures in number fields:

- Group of fractional ideals \longrightarrow contains subgroup of principal ideals

Two interesting structures in number fields:

- Group of fractional ideals \longrightarrow contains subgroup of principal ideals

Quotient \implies class group $Cl(\mathbf{K})$

Two interesting structures in number fields:

- Group of fractional ideals \longrightarrow contains subgroup of principal ideals

Quotient \implies class group $Cl(\mathbf{K})$

$$1 \longrightarrow P(\mathbf{K}) \longrightarrow I(\mathbf{K}) \longrightarrow Cl(\mathbf{K}) \longrightarrow 1$$

Two interesting structures in number fields:

- Group of fractional ideals \longrightarrow contains subgroup of principal ideals

Quotient \implies class group $Cl(\mathbf{K})$

$$1 \longrightarrow P(\mathbf{K}) \longrightarrow I(\mathbf{K}) \longrightarrow Cl(\mathbf{K}) \longrightarrow 1$$

- Group of units

Two interesting structures in number fields:

- Group of fractional ideals \longrightarrow contains subgroup of principal ideals

Quotient \implies class group $Cl(\mathbf{K})$

$$1 \longrightarrow P(\mathbf{K}) \longrightarrow I(\mathbf{K}) \longrightarrow Cl(\mathbf{K}) \longrightarrow 1$$

- Group of units \longrightarrow finitely generated

Two interesting structures in number fields:

- Group of fractional ideals \longrightarrow contains subgroup of principal ideals

Quotient \implies class group $Cl(\mathbf{K})$

$$1 \longrightarrow P(\mathbf{K}) \longrightarrow I(\mathbf{K}) \longrightarrow Cl(\mathbf{K}) \longrightarrow 1$$

- Group of units \longrightarrow finitely generated

Generators \implies fundamental units

Two interesting structures in number fields:

- Group of fractional ideals \longrightarrow contains subgroup of principal ideals

Quotient \implies class group $Cl(\mathbf{K})$

$$1 \longrightarrow P(\mathbf{K}) \longrightarrow I(\mathbf{K}) \longrightarrow Cl(\mathbf{K}) \longrightarrow 1$$

- Group of units \longrightarrow finitely generated

Generators \implies fundamental units

$$1 \longrightarrow U(\mathbf{K}) \longrightarrow \mathbf{K}^* \longrightarrow P(\mathbf{K}) \longrightarrow 1$$

$$\begin{array}{ccccccc}
 & & 1 & & 1 & & 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & U(\mathbf{K}) & \longrightarrow & J_{S_\infty}(\mathbf{K}) & \longrightarrow & C_{S_\infty}(\mathbf{K}) \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & \mathbf{K}^* & \longrightarrow & J(\mathbf{K}) & \longrightarrow & C(\mathbf{K}) \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & P(\mathbf{K}) & \longrightarrow & I(\mathbf{K}) & \longrightarrow & Cl(\mathbf{K}) \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 1 & & 1 & & 1
 \end{array}$$

Number fields

Number-field *size* is given by the **discriminant** Δ_K

$$\Delta_K = \det(\sigma_i(\omega_j))^2$$

Number fields

Number-field *size* is given by the **discriminant** $\Delta_{\mathbf{K}}$ ($H(T) = \max |t_k|$ and $n = [\mathbf{K} : \mathbf{Q}]$)

$$\Delta_{\mathbf{K}} = \det(\sigma_i(\omega_j))^2 \quad \text{and} \quad |\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Number fields

Number-field *size* is given by the **discriminant** $\Delta_{\mathbf{K}}$ ($H(T) = \max |t_k|$ and $n = [\mathbf{K} : \mathbf{Q}]$)

$$\Delta_{\mathbf{K}} = \det(\sigma_i(\omega_j))^2 \quad \text{and} \quad |\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Algorithm complexities are given according to the size of $|\Delta_{\mathbf{K}}|$

Number fields

Number-field *size* is given by the **discriminant** $\Delta_{\mathbf{K}}$ ($H(T) = \max |t_k|$ and $n = [\mathbf{K} : \mathbf{Q}]$)

$$\Delta_{\mathbf{K}} = \det(\sigma_i(\omega_j))^2 \quad \text{and} \quad |\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Algorithm complexities are given according to the size of $|\Delta_{\mathbf{K}}|$

- Polynomial: $(\log |\Delta_{\mathbf{K}}|)^c$ operations

Number fields

Number-field *size* is given by the **discriminant** $\Delta_{\mathbf{K}}$ ($H(T) = \max |t_k|$ and $n = [\mathbf{K} : \mathbf{Q}]$)

$$\Delta_{\mathbf{K}} = \det(\sigma_i(\omega_j))^2 \quad \text{and} \quad |\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Algorithm complexities are given according to the size of $|\Delta_{\mathbf{K}}|$

- Polynomial: $(\log |\Delta_{\mathbf{K}}|)^c$ operations
- Exponential: $|\Delta_{\mathbf{K}}|^c$ operations

Number fields

Number-field size is given by the **discriminant** $\Delta_{\mathbf{K}}$ ($H(T) = \max |t_k|$ and $n = [\mathbf{K} : \mathbf{Q}]$)

$$\Delta_{\mathbf{K}} = \det(\sigma_i(\omega_j))^2 \quad \text{and} \quad |\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Algorithm complexities are given according to the size of $|\Delta_{\mathbf{K}}|$

- Polynomial: $(\log |\Delta_{\mathbf{K}}|)^c$ operations
- Exponential: $|\Delta_{\mathbf{K}}|^c$ operations
- Between?

Number fields

Number-field size is given by the **discriminant** $\Delta_{\mathbf{K}}$ ($H(T) = \max |t_k|$ and $n = [\mathbf{K} : \mathbf{Q}]$)

$$\Delta_{\mathbf{K}} = \det(\sigma_i(\omega_j))^2 \quad \text{and} \quad |\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Algorithm complexities are given according to the size of $|\Delta_{\mathbf{K}}|$

- Polynomial: $(\log |\Delta_{\mathbf{K}}|)^c$ operations
- Exponential: $|\Delta_{\mathbf{K}}|^c$ operations
- Between?

Subexponential L -notation :

$$L_N(\alpha, c) = \exp((c + o(1))(\log N)^\alpha (\log \log N)^{1-\alpha})$$

Number fields

Number-field size is given by the **discriminant** $\Delta_{\mathbf{K}}$ ($H(T) = \max |t_k|$ and $n = [\mathbf{K} : \mathbf{Q}]$)

$$\Delta_{\mathbf{K}} = \det(\sigma_i(\omega_j))^2 \quad \text{and} \quad |\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Algorithm complexities are given according to the size of $|\Delta_{\mathbf{K}}|$

- Polynomial: $(\log |\Delta_{\mathbf{K}}|)^c$ operations
- Exponential: $|\Delta_{\mathbf{K}}|^c$ operations
- Between?

Subexponential L -notation :

$$L_{|\Delta_{\mathbf{K}}|}(0, c) \approx (\log |\Delta_{\mathbf{K}}|)^c \mid L_{|\Delta_{\mathbf{K}}|}(1, c) \approx |\Delta_{\mathbf{K}}|^c$$

$$L_N(\alpha, c) = \exp((c + o(1))(\log N)^\alpha (\log \log N)^{1-\alpha})$$

1969 Shanks: quadratic number fields in $O(|\Delta_{\mathbf{K}}|^{\frac{1}{5}})$

1989 Hafner and McCurley: imaginary quadratic number fields in $L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2}, \sqrt{2})$

1990 Buchmann: all number fields with fixed degree in $L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2}, 1.7)$

2014 Biasse and Fieker: all number fields in $L_{|\Delta_{\mathbf{K}}|}(\frac{2}{3} + \varepsilon)$ and $L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2})$ if $n \leq (\log |\Delta_{\mathbf{K}}|)^{\frac{3}{4} - \varepsilon}$

2014 Biasse and Fieker: number fields defined by a *good* polynomial in $L_{|\Delta_{\mathbf{K}}|}(a)$, $\frac{1}{3} \leq a < \frac{1}{2}$



how i compute class groups

JA

1969 Shanks: quadratic number fields in $O(|\Delta_K|^{\frac{1}{5}})$

1989 Hafner and McCurley: imaginary quadratic number fields in $L_{|\Delta_K|}(\frac{1}{2}, \sqrt{2})$

1990 Buchmann: all number fields with fixed degree in $L_{|\Delta_K|}(\frac{1}{2}, 1.7)$

2014 Biasse and Fieker: all number fields in $L_{|\Delta_K|}(\frac{2}{3} + \varepsilon)$ and $L_{|\Delta_K|}(\frac{1}{2})$ if $n \leq (\log |\Delta_K|)^{\frac{3}{4} - \varepsilon}$

2014 Biasse and Fieker: number fields defined by a *good* polynomial in $L_{|\Delta_K|}(a)$, $\frac{1}{3} \leq a < \frac{1}{2}$

Well-known, used for discrete logarithms

❶ **Factor base**

Fix a factor base composed of small elements

❷ **Relation collection**

Collect some relations between those small elements, corresponding to linear equations

❸ **Linear algebra**

Deduce the sought result performing linear algebra on the system built

The factor base

$$\mathcal{B} = \{\text{prime ideals in } \mathcal{O}_{\mathbf{K}} \text{ of norm below } B\}$$

B is determined such that \mathcal{B} generates the whole class group

The factor base

$$\mathcal{B} = \{\text{prime ideals in } \mathcal{O}_{\mathbf{K}} \text{ of norm below } B\}$$

B is determined such that \mathcal{B} generates the whole class group

Minkowski's bound: every class contains an ideal of norm smaller than

$$M_{\mathbf{K}} = \sqrt{|\Delta_{\mathbf{K}}|} \left(\frac{4}{\pi}\right)^{r_2} \frac{n!}{n^n}$$

The factor base

$$\mathcal{B} = \{\text{prime ideals in } \mathcal{O}_{\mathbf{K}} \text{ of norm below } B\}$$

B is determined such that \mathcal{B} generates the whole class group

Minkowski's bound: every class contains an ideal of norm smaller than

$$M_{\mathbf{K}} = \sqrt{|\Delta_{\mathbf{K}}|} \left(\frac{4}{\pi}\right)^{r_2} \frac{n!}{n^n}$$

Bach's bound: assuming ERH, classes of ideals of norm less than $12(\log|\Delta_{\mathbf{K}}|)^2$ generate the class group

The factor base

$$\mathcal{B} = \{\text{prime ideals in } \mathcal{O}_{\mathbf{K}} \text{ of norm below } B\}$$

B is determined such that \mathcal{B} generates the whole class group

Minkowski's bound: every class contains an ideal of norm smaller than

$$M_{\mathbf{K}} = \sqrt{|\Delta_{\mathbf{K}}|} \left(\frac{4}{\pi}\right)^{r_2} \frac{n!}{n^n}$$

Bach's bound: assuming ERH, classes of ideals of norm less than $12(\log|\Delta_{\mathbf{K}}|)^2$ generate the class group

Practically

$$B = L_{|\Delta_{\mathbf{K}}|}(\beta, c_b)$$

$$\mathcal{B} = (\mathfrak{p}_1, \dots, \mathfrak{p}_N)$$

Surjective morphism:

$$\begin{array}{ccccc} \mathbf{Z}^N & \longrightarrow & I & \longrightarrow & Cl(K) \\ (e_1, \dots, e_N) & \longmapsto & \prod_i \mathfrak{p}_i^{e_i} & \longmapsto & \left[\prod_i \mathfrak{p}_i^{e_i} \right] \end{array}$$

$$Cl(\mathbf{K}) \simeq \mathbf{Z}^N / \{(e_1, \dots, e_N) \in \mathbf{Z}^N \mid \prod \mathfrak{p}_i^{e_i} = (\alpha) \mathcal{O}_{\mathbf{K}}\}$$

$$\mathcal{B} = (\mathfrak{p}_1, \dots, \mathfrak{p}_N)$$

Surjective morphism:

$$\begin{array}{ccccc} \mathbf{Z}^N & \longrightarrow & I & \longrightarrow & Cl(K) \\ (e_1, \dots, e_N) & \longmapsto & \prod_i \mathfrak{p}_i^{e_i} & \longmapsto & \left[\prod_i \mathfrak{p}_i^{e_i} \right] \end{array}$$

$$Cl(\mathbf{K}) \simeq \mathbf{Z}^N / \{(e_1, \dots, e_N) \in \mathbf{Z}^N \mid \prod \mathfrak{p}_i^{e_i} = (\alpha) \mathcal{O}_{\mathbf{K}}\}$$

Idea:

- 1 Pick at random $\mathfrak{a} = \prod \mathfrak{p}_i^{a_i}$
- 2 Find a *reduced* ideal \mathfrak{b} in the same class
- 3 If \mathfrak{b} splits over \mathcal{B} ($\iff \mathfrak{b} = \prod \mathfrak{p}_i^{b_i}$) then

$$\mathfrak{a} \cdot \mathfrak{b}^{-1} = \prod \mathfrak{p}_i^{a_i - b_i} \quad \text{is principal}$$

Correspondence between ideals and lattices:

$$\mathfrak{a} \longleftrightarrow \sigma(\mathfrak{a}) = (\sigma_i(\mathfrak{a}_j))_{i,j}$$

Correspondence between ideals and lattices:

ideal-lattices

$$\mathfrak{a} \longleftrightarrow \sigma(\mathfrak{a}) = (\sigma_i(\mathfrak{a}_j))_{i,j}$$

Correspondence between ideals and lattices:

ideal-lattices

$$\mathfrak{a} \longleftrightarrow \sigma(\mathfrak{a}) = (\sigma_i(\mathfrak{a}_j))_{i,j}$$

Buchmann's reduction:

- Shortest Vector
- Exponential in the extension degree

Correspondence between ideals and lattices:

ideal-lattices

$$\mathfrak{a} \longleftrightarrow \sigma(\mathfrak{a}) = (\sigma_i(\mathfrak{a}_j))_{i,j}$$

Buchmann's reduction:

- Shortest Vector
- Exponential in the extension degree

Biasse-Fieker's reduction:

- BKZ
- Trade-off between time spent and approximation factor
- Subexponential complexity

- Relations stored in a matrix of size about $N \times N$
- Structure of the class group given by the *Smith Normal Form* of the matrix
- First compute *Hermite Normal Form* with a premultiplier because we need kernel vectors
- Storjohann and Labahn algorithm, runtime in $N^{\omega+1}$
($2 \leq \omega \leq 3$ exponent of matrix multiplication)

We find a tentative class group H , but the class group $Cl(\mathbf{K})$ may be only a quotient of H
 \Rightarrow Need an approximation of the class number $h_{\mathbf{K}} = |Cl(\mathbf{K})|$

We find a tentative class group H , but the class group $Cl(\mathbf{K})$ may be only a quotient of H
 \Rightarrow Need an approximation of the class number $h_{\mathbf{K}} = |Cl(\mathbf{K})|$

Class Number Formula + Euler Product:

$$h_{\mathbf{K}} \cdot \text{Reg}_{\mathbf{K}} \approx EP \cdot \frac{w_{\mathbf{K}} \cdot \sqrt{|\Delta_{\mathbf{K}}|}}{2^{r_1} \cdot (2\pi)^{r_2}}$$


We find a tentative class group H , but the class group $Cl(\mathbf{K})$ may be only a quotient of H
 \Rightarrow Need an approximation of the class number $h_{\mathbf{K}} = |Cl(\mathbf{K})|$

Class Number Formula + Euler Product:

$$h_{\mathbf{K}} \cdot \text{Reg}_{\mathbf{K}} \approx EP \cdot \frac{w_{\mathbf{K}} \cdot \sqrt{|\Delta_{\mathbf{K}}|}}{2^{r_1} \cdot (2\pi)^{r_2}}$$


From the relations, we can also deduce a candidate for an approximation of $\text{Reg}_{\mathbf{K}}$ and perform the verification step

General case:




First general subexponential algorithm

Special case:




Only if \mathbf{K} is defined by T such that $H(T) = L_{|\Delta_K|}(1 - \alpha)$

General case:




Refinement on the complexity analysis

General case:



Improvement through a better parameters choice

General case:



Improvement using special lattice-reduction algorithm

Complexity history

1969 Shanks: quadratic number fields in $O(|\Delta_{\mathbf{K}}|^{\frac{1}{5}})$

1989 Hafner and McCurley: imaginary quadratic number fields in $L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2}, \sqrt{2})$

1990 Buchmann: all number fields with fixed degree in $L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2}, 1.7)$

2014 Biasse and Fieker: all number fields in $L_{|\Delta_{\mathbf{K}}|}(\frac{2}{3} + \varepsilon)$ and $L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2})$ if $n \leq (\log |\Delta_{\mathbf{K}}|)^{\frac{3}{4} - \varepsilon}$

2014 Biasse and Fieker: number fields defined by a *good* polynomial in $L_{|\Delta_{\mathbf{K}}|}(a)$, $\frac{1}{3} \leq a < \frac{1}{2}$

What is a *good* polynomial?

We want a polynomial that defines a fixed number field:

- The degree is fixed
- We want the coefficients as small as possible

What is a *good* polynomial?

We want a polynomial that defines a fixed number field:

- The degree is fixed
- We want the coefficients as small as possible

For $T = \sum a_k X^k \in \mathbf{Z}[X]$, the **height** is the maximal norm of its coefficients

$$H(T) = \max_k |a_k|$$

What is a *good* polynomial?

We want a polynomial that defines a fixed number field:

- The degree is fixed
- We want the coefficients as small as possible

For $T = \sum a_k X^k \in \mathbf{Z}[X]$, the **height** is the maximal norm of its coefficients

$$H(T) = \max_k |a_k|$$

For every degree- n number field \mathbf{K} and every defining polynomial T of \mathbf{K} ,

$$|\Delta_{\mathbf{K}}| \leq n^{2n} H(T)^{2n-2}$$

Why minimizing the height?

Idea: Look at small algebraic integers $x \in \mathcal{O}_{\mathbf{K}}$ such that $x = A(\theta)$ with $\deg A \leq c_d$ and $H(A) \leq C_H$.

Why minimizing the height?

Idea: Look at small algebraic integers $x \in \mathcal{O}_{\mathbf{K}}$ such that $x = A(\theta)$ with $\deg A \leq c_d$ and $H(A) \leq C_H$. Then

$$\mathcal{N}(\langle x \rangle) \leq f(n, c_d) \cdot C_H^n \cdot H(T)^{c_d}$$

Why minimizing the height?

Idea: Look at small algebraic integers $x \in \mathcal{O}_{\mathbf{K}}$ such that $x = A(\theta)$ with $\deg A \leq c_d$ and $H(A) \leq C_H$. Then

$$\mathcal{N}(\langle x \rangle) \leq f(n, c_d) \cdot C_H^n \cdot H(T)^{c_d}$$

How: A tricky reduction of the lattice $\sigma(\mathcal{O}_{\mathbf{K}})$.

- Classic reduction: same magnitude for all the coordinates
- Tricky reduction: one dominant coordinate

Why minimizing the height?

Idea: Look at small algebraic integers $x \in \mathcal{O}_{\mathbf{K}}$ such that $x = A(\theta)$ with $\deg A \leq c_d$ and $H(A) \leq C_H$. Then

$$\mathcal{N}(\langle x \rangle) \leq f(n, c_d) \cdot C_H^n \cdot H(T)^{c_d}$$

How: A tricky reduction of the lattice $\sigma(\mathcal{O}_{\mathbf{K}})$.


- Classic reduction: same magnitude for all the coordinates
- Tricky reduction: one dominant coordinate

Example: Number field defined by $T = x^5 - 5843635x^4 + 931633x^2 + 6577x - 8570$

- Rounded conjugates: $[-0.38, -0.10, 0.095, 0.39, 5843634.999999997]$
- Rounded shortest vector: $[-84411, -23707, -1315, 20616, 88819]$


$$x^5 - 2x^4 - 8001397580x^3 - 31542753393650x^2 + 3636653302451131875x + 4818547529425280067500$$

Special case:




Only if \mathbf{K} is defined by T such that $H(T) = L_{|\Delta_K|}(1 - \alpha)$

General case:



Without any condition

Special case:




Only when it is better than the method based on ideal reductions

CLASS GROUPS AND CRYPTOLOGY


pl

private key
Bob

Public Key Cryptography




Public Key Cryptography



- Everyone uses the **public key** to encrypt

Public Key Cryptography



- Everyone uses the **public key** to encrypt
- Only Bob can decrypt thanks to his **private key**

The Principal Ideal Problem

Definition

The *Principal Ideal Problem* (PIP) consists in finding a generator of an ideal, assuming it is principal.

The Principal Ideal Problem

Definition

The *Short Principal Ideal Problem* (SPIP) consists in finding a *short* generator of an ideal, assuming it is principal.

The Principal Ideal Problem

Definition

The *Short Principal Ideal Problem* (SPIP) consists in finding a *short* generator of an ideal, assuming it is principal.

- Base of several cryptographical schemes ([SV10],[GGH13])

The Principal Ideal Problem

Definition

The *Short Principal Ideal Problem* (SPIP) consists in finding a **short** generator of an ideal, assuming it is principal.

- Base of several cryptographical schemes ([SV10],[GGH13])
- Two distinct phases:
 - 1 Given the \mathbf{Z} -basis of the ideal $\mathfrak{a} = \langle \mathbf{g} \rangle$, find a — not necessarily short — generator $\mathbf{g}' = \mathbf{g} \cdot \mathbf{u}$ for a unit \mathbf{u}
 - 2 From \mathbf{g}' , find a short generator of the ideal

The Principal Ideal Problem

Definition

The *Short Principal Ideal Problem* (SPIP) consists in finding a *short* generator of an ideal, assuming it is principal.

- Base of several cryptographical schemes ([SV10],[GGH13])
- Two distinct phases:
 - 1 Given the \mathbf{Z} -basis of the ideal $\mathfrak{a} = \langle \mathbf{g} \rangle$, find a — not necessarily short — generator $\mathbf{g}' = \mathbf{g} \cdot \mathbf{u}$ for a unit \mathbf{u}
 - 2 From \mathbf{g}' , find a short generator of the ideal

2014 - Campbell, Groves, and Sheperd:

Reduction in polynomial time for power-of-two cyclotomic fields

2016 - Cramer, Ducas, Peikert, and Regev:

Proof and extension to prime-power cyclotomic fields

Key Generation:

- 1 Fix the security parameter $N = 2^n$
- 2 Let $F(X) = X^N + 1$ be the polynomial defining the cyclotomic field $\mathbf{K} = \mathbf{Q}(\zeta_{2N})$
- 3 Set $G(X) = 1 + 2 \cdot S(X)$,
for $S(X)$ of degree $N - 1$ with coefficients in $[-2^{\sqrt{N}}, 2^{\sqrt{N}}]$,
such that the norm $\mathcal{N}(\langle G(\zeta_{2N}) \rangle)$ is prime
- 4 Set $\mathbf{g} = G(\zeta_{2N}) \in \mathcal{O}_{\mathbf{K}}$
- 5 Return the **private key** $\text{sk} = \mathbf{g}$ and the **public key** $\text{pk} = \text{HNF}(\langle \mathbf{g} \rangle)$

Key Generation:

- 1 Fix the security parameter $N = 2^n$
- 2 Let $F(X) = X^N + 1$ be the polynomial defining the cyclotomic field $\mathbf{K} = \mathbf{Q}(\zeta_{2N})$
- 3 Set $G(X) = 1 + 2 \cdot S(X)$,
for $S(X)$ of degree $N - 1$ with coefficients in $[-2^{\sqrt{N}}, 2^{\sqrt{N}}]$,
such that the norm $\mathcal{N}(\langle G(\zeta_{2N}) \rangle)$ is prime
- 4 Set $\mathbf{g} = G(\zeta_{2N}) \in \mathcal{O}_{\mathbf{K}}$
- 5 Return the **private key** $\text{sk} = \mathbf{g}$ and the **public key** $\text{pk} = \text{HNF}(\langle \mathbf{g} \rangle)$

Goal: Recover the private key from the public key

- ① Perform a reduction from the cyclotomic field to its totally real subfield, allowing to work in smaller dimension
- ② Then a descent makes the sizes of involved ideals decrease
- ③ Collect relations and run linear algebra to construct small ideals and a generator
- ④ Eventually run the derivation of the small generator from a bigger one

- 1 Perform a reduction from the cyclotomic field to its totally real subfield, allowing to work in smaller dimension
- 2 Then a descent makes the sizes of involved ideals decrease
- 3 Collect relations and run linear algebra to construct small ideals and a generator
- 4 Eventually run the derivation of the small generator from a bigger one

All the complexities are expressed as a function of the field discriminant $\Delta_{\mathbf{Q}(\zeta_{2N})} = N^N$, for $N = 2^n$. For instance,

$$L_{|\Delta_{\mathbf{K}}|}(\alpha) = 2^{N^{\alpha+o(1)}}$$

1. Reduction to the totally real subfield

Goal: Halve the dimension of the ambient field

1. Reduction to the totally real subfield

Goal: Halve the dimension of the ambient field

- Based on the algorithm of Gentry and Szydło

1. Reduction to the totally real subfield

Goal: Halve the dimension of the ambient field

- Based on the algorithm of Gentry and Szydło
- Polynomial complexity

1. Reduction to the totally real subfield

Goal: Halve the dimension of the ambient field


- Based on the algorithm of Gentry and Szydło
- Polynomial complexity
- **Input:** a \mathbf{Z} -basis of $\mathfrak{a} = \langle \mathbf{g} \rangle$

1. Reduction to the totally real subfield

Goal: Halve the dimension of the ambient field


- Based on the algorithm of Gentry and Szydło
- Polynomial complexity
- **Input:** a \mathbf{Z} -basis of $\mathfrak{a} = \langle \mathbf{g} \rangle$
- **Output:** a \mathbf{Z} -basis of $\mathfrak{a}^+ = \langle \mathbf{g} + \bar{\mathbf{g}} \rangle \subset \mathbf{Q}(\zeta + \zeta^{-1})$ and $\mathbf{g} \cdot \bar{\mathbf{g}}^{-1}$ to recover \mathbf{g} from $\mathbf{g} + \bar{\mathbf{g}}$

2. The descent



Input ideal – Norm arbitrary large


2. The descent



Input ideal – Norm arbitrary large

Initial reduction – Norm: $L_{|\Delta_K|}(\frac{3}{2})$


2. The descent



Input ideal – Norm arbitrary large

Initial reduction – $L_{|\Delta_{\mathbf{K}}|}(1)$ -smooth

2. The descent




Input ideal – Norm arbitrary large

Initial reduction – $L_{|\Delta_{\mathbf{K}}|}(1)$ -smooth

First step – Norm: $L_{|\Delta_{\mathbf{K}}|}\left(\frac{5}{4}\right)$

2. The descent




Input ideal – Norm arbitrary large

Initial reduction – $L_{|\Delta_K|}(1)$ -smooth

First step – $L_{|\Delta_K|}(\frac{3}{4})$ -smooth

2. The descent




Input ideal – Norm arbitrary large

Initial reduction – $L_{|\Delta_{\mathbf{K}}|}(1)$ -smooth

First step – $L_{|\Delta_{\mathbf{K}}|}(\frac{3}{4})$ -smooth

Second step – Norm: $L_{|\Delta_{\mathbf{K}}|}(\frac{9}{8})$

2. The descent




Input ideal – Norm arbitrary large

Initial reduction – $L_{|\Delta_K|}(1)$ -smooth

First step – $L_{|\Delta_K|}(\frac{3}{4})$ -smooth

Second step – $L_{|\Delta_K|}(\frac{5}{8})$ -smooth

2. The descent



Input ideal – Norm arbitrary large


Initial reduction – $L_{|\Delta_K|}(1)$ -smooth

First step – $L_{|\Delta_K|}(\frac{3}{4})$ -smooth

Second step – $L_{|\Delta_K|}(\frac{5}{8})$ -smooth

Last but one step – Norm: $\approx L_{|\Delta_K|}(1)$

2. The descent



Input ideal – Norm arbitrary large


Initial reduction – $L_{|\Delta_{\mathbf{K}}|}(1)$ -smooth

First step – $L_{|\Delta_{\mathbf{K}}|}(\frac{3}{4})$ -smooth

Second step – $L_{|\Delta_{\mathbf{K}}|}(\frac{5}{8})$ -smooth

Last but one step – $\approx L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2})$ -smooth

2. The descent



Input ideal – Norm arbitrary large

Initial reduction – $L_{|\Delta_{\mathbf{K}}|}(1)$ -smooth


First step – $L_{|\Delta_{\mathbf{K}}|}(\frac{3}{4})$ -smooth

Second step – $L_{|\Delta_{\mathbf{K}}|}(\frac{5}{8})$ -smooth

Last but one step – $\approx L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2})$ -smooth

Last step – Norm: $L_{|\Delta_{\mathbf{K}}|}(1)$

2. The descent



Input ideal – Norm arbitrary large

Initial reduction – $L_{|\Delta_{\mathbf{K}}|}(1)$ -smooth

First step – $L_{|\Delta_{\mathbf{K}}|}(\frac{3}{4})$ -smooth

Second step – $L_{|\Delta_{\mathbf{K}}|}(\frac{5}{8})$ -smooth

Last but one step – $\approx L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2})$ -smooth

Last step – $L_{|\Delta_{\mathbf{K}}|}(\frac{1}{2})$ -smooth

3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

Index Calculus Method:

- **Factor base:** set of all prime ideals with norm below B

3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

Index Calculus Method:

- **Factor base:** set of all prime ideals with norm below B
- **Relation collection:** construction of a full-rank matrix M

3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

Index Calculus Method:

- **Factor base:** set of all prime ideals with norm below B
- **Relation collection:** construction of a full-rank matrix M

Relation: principal ideal that splits on the factor base.

Test ideals generated by $\mathbf{v} = \sum v_i(\zeta^i + \zeta^{-i})$ for $|v_i| \leq \log|\Delta_K|$

3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

Index Calculus Method:

- **Factor base:** set of all prime ideals with norm below B
- **Relation collection:** construction of a full-rank matrix M

Relation: principal ideal that splits on the factor base.

Test ideals generated by $\mathbf{v} = \sum v_i(\zeta^i + \zeta^{-i})$ for $|v_i| \leq \log|\Delta_K|$

Norm below $L_{|\Delta_K|}(1) \implies L_{|\Delta_K|}(\frac{1}{2})$ -smooth ideals in $L_{|\Delta_K|}(\frac{1}{2})$

3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

Index Calculus Method:

- **Factor base:** set of all prime ideals with norm below B
- **Relation collection:** construction of a full-rank matrix M

$$\begin{pmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \\ \vdots \\ \mathbf{v}_{Q|\mathcal{B}|} \end{pmatrix} \rightarrow \begin{pmatrix} M_{1,1} & \cdots & M_{1,|\mathcal{B}|} \\ M_{2,1} & \cdots & M_{2,|\mathcal{B}|} \\ \vdots & & \vdots \\ M_{Q|\mathcal{B}|,1} & \cdots & M_{Q|\mathcal{B}|,|\mathcal{B}|} \end{pmatrix} \Rightarrow \forall i, \langle \mathbf{v}_i \rangle = \prod_{j=1}^{|\mathcal{B}|} \mathfrak{p}_j^{M_{i,j}}$$

3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

Index Calculus Method:

- **Factor base:** set of all prime ideals with norm below B
- **Relation collection:** construction of a full-rank matrix M
- A N -dimensional vector Y including all the valuations of the smooth ideals in the \mathfrak{p}_i


3. Solution for smooth ideals

Input: Bunch of prime ideals of norm below $B = L_{|\Delta_K|}(\frac{1}{2})$

Index Calculus Method:


- **Factor base:** set of all prime ideals with norm below B
- **Relation collection:** construction of a full-rank matrix M
- A N -dimensional vector Y including all the valuations of the smooth ideals in the \mathfrak{p}_i
- A solution of $MX = Y$ provides a generator of the product of the $L_{|\Delta_K|}(\frac{1}{2})$ -smooth ideals

4. The backtracking




A generator for the product

4. The backtracking



A generator for the product

4. The backtracking




A generator for the initial ideal



A generator for the product

4. The backtracking



A generator for the initial ideal



A generator for the product

Thanks

Kia ora

Illustrations by Alexia R. (@a_draw_r)