

OLLSCOIL NA hÉIREANN, CORCAIGH
THE NATIONAL UNIVERSITY OF IRELAND, CORK

COLÁISTE NA hOLLSCOILE, CORCAIGH
UNIVERSITY COLLEGE, CORK

AUTUMN EXAMINATIONS, 2011

B.E. (ELECTRICAL)

DIGITAL SIGNAL PROCESSING
EE4008

Dr L. Seed
Dr. M. Creed
Dr. W.P. Marnane

Time Allowed: *3 hours*

Answer *five* questions.

All questions carry equal marks.

The use of departmental approved non-programmable calculators is permitted

1. (a) Starting with the ideal frequency response $H_d(\omega)$, describe the windows method of designing a band pass filter. [10 marks]
(b) Determine the filter length M and the coefficients $h(1)$ and $h(\frac{M-1}{2})$ using the “Windows” method of the band pass filter design that meets the following specification:
 - Passband:- 150 – 250Hz
 - Transition Width:- 50Hz
 - Passband Ripple:- 0.01dB
 - Stopband attenuation:- > 60 dB
 - Sampling frequency:- 1kHz

The parameters of common window functions are given in the Appendix. [10 marks]

2. (a) A First Order Lowpass IIR Digital Filter has a transfer function

$$H_{LP}(z) = G \frac{1 + z^{-1}}{1 - \alpha z^{-1}}$$

Determine the Gain factor G in terms of α . [5 marks]

- (b) Derive an expression for α in terms of the 3dB cutoff frequency ω_c . [5 marks]

- (c) Determine the transfer function $H_1(z)$, of a first-order low pass filter with a 3-dB cutoff Frequency of 0.6π . Draw the pole/zero Plot of $H_1(z)$ and sketch the magnitude response of the filter for $0 \leq \omega \leq \pi$, clearly identifying the 3dB cutoff frequency ω_c [5 marks]

- (d) A comb filter with a transfer function $G_1(z)$ is formed by taking the transfer function $H_1(z)$ and replacing each delay by M delays, such that:

$$G_1(z) = H_1(z^M)$$

Draw the pole/zero plot for $M = 2$ and sketch the magnitude response for the same value of M , for $0 \leq \omega \leq \pi$. [5 marks]

3. Consider the z-transform

$$H(z) = \frac{z(z + 0.5)}{z^2 - 0.65z + 0.1}$$

- (a) Draw the Pole-Zero plot of $H(z)$ and identify the three possible regions of convergence. [6 marks]

- (b) Use the partial fractions method to determine the inverse z-transform $h(n)$ where
 i. $h(n)$ is a causal sequence
 ii. $h(n)$ is an anti-causal sequence
 iii. $h(n)$ is a two sided sequence. [10 marks]

- (c) Determine the first three values of the causal $h(n)$ sequence, using the long division method of inverting the z-transform. [4 marks]

4. (a) Let $x(n)$ be a Wide Sense Stationary random process with mean m_X , autocorrelation $\phi_{XX}(k)$ and power spectral density $P_{XX}(\omega)$. $x(n)$ is filtered by a Stable Linear Time Invariant System with impulse response $h(n)$ to produce output $y(n)$. Determine the mean m_Y , autocorrelation $\phi_{YY}(k)$ and power spectral density $P_{YY}(\omega)$ of $y(n)$. [10 marks]

- (b) Unit variance white noise is filtered by a LTI filter with impulse response:

$$h(n) = \frac{1}{5} \left(-\frac{1}{5}\right)^n u(n) + \frac{1}{5} \left(-\frac{1}{5}\right)^{n-3} u(n-3)$$

Determine the mean and the power spectral density of the filter output in trigonometric form.

[10 marks]

5. (a) Show how the N point DFT

$$X(k) = \sum_{n=0}^{N-1} x(n)e^{(-j2\pi nk/N)} \quad k = 0, 1, \dots, N-1$$

can be reduced to two $\frac{N}{2}$ point DFTs of the odd and even indexed values of $x(n)$.

[6 marks]

- (b) Hence show that the computational complexity of a radix 2 decimation in time FFT algorithm is

$$O\left(\frac{N}{2} \log_2 N\right)$$

[4 marks]

- (c) 1024 data values of the Wide Sense Stationary Random Process

$$x(n) = A \sin(n\omega_1 + \phi_1) + A \sin(n\omega_2 + \phi_2) + w(n)$$

are recorded, where A , ω_1 and ω_2 are fixed constants; ϕ_1 and ϕ_2 are random variables uniformly distributed over the interval $-\pi, \pi$ and $w(n)$ is Gaussian White Noise with variance σ_w . For $\Delta\omega = (\omega_2 - \omega_1) = 0.015\pi$ compare the performance of the following 4 methods of spectral estimation in terms of resolution, variance reduction and computation complexity.

- i. Periodogram
- ii. Modified Periodogram using a Hamming window
- iii. Bartlett method
- iv. Welch method with 50% overlap and a Hanning window

You may assume that the FFT is used in each of the spectral estimation methods and parameters of the window function are given in the Appendix.

[10 marks]

6. (a) Starting with the Yule Walker Equation for an ARMA process of order (p, q)

$$\phi_{XX}(k) = - \sum_{l=1}^p a(l)\phi_{XX}(k-l) + \sigma_v^2 \sum_{l=0}^{q-k} b(l+k)h^*(l)$$

derive the Yule Walker Equations for an

- i. AR Process of order p
- ii. MA Process of order q

[4 marks]

- (b) Compare the following AR methods of parametric spectral estimation

- i. Autocorrelation
- ii. Covariance
- iii. Modified Covariance

[9 marks]

- (c) In the covariance method of parametric spectral estimation the value $\hat{c}(j, k)$, where

$$\hat{c}(j, k) = \frac{1}{N-p} \sum_{n=p}^{N-1} x^*(n-j)x(n-k)$$

is used as an estimate of the autocorrelation values $\phi_{XX}(j-k)$. Given

$$\begin{aligned} \hat{c}(0, 0) &= 1 \\ \hat{c}(0, 1) = \hat{c}(1, 0) &= 0.66 \\ \hat{c}(0, 2) = \hat{c}(2, 0) &= 0.25 \\ \hat{c}(1, 1) &= 0.95 \\ \hat{c}(1, 2) = \hat{c}(2, 1) &= 0.65 \\ \hat{c}(2, 2) &= 0.94 \end{aligned}$$

Estimate the spectrum using the covariance method of AR parametric spectral estimation for an AR(2) process.

[7 marks]

Appendix of Equations

- Window Functions

Window $w(n)$	Sidelobe	Δf	Stopband Attenuation	Passband Ripple	$\Delta\omega_{3db}$
Rectangular	-13db	$\frac{0.9}{N}$	21db	0.7416db	$0.89\frac{2\pi}{N}$
$w(n) = \begin{cases} 1 & 0 \leq n \leq N-1 \\ 0 & \text{otherwise} \end{cases}$					
Hanning	-31db	$\frac{3.1}{N}$	44db	0.0546db	$1.44\frac{2\pi}{N}$
$w(n) = \begin{cases} 0.5 - 0.5 \cos\left(\frac{2\pi n}{N-1}\right) & 0 \leq n \leq N-1 \\ 0 & \text{otherwise} \end{cases}$					
Hamming	-41db	$\frac{3.3}{N}$	53db	0.0194db	$1.30\frac{2\pi}{N}$
$w(n) = \begin{cases} 0.54 - 0.46 \cos\left(\frac{2\pi n}{N-1}\right) & 0 \leq n \leq N-1 \\ 0 & \text{otherwise} \end{cases}$					

- Table of Z-Transforms

Signal	Z-Transform	ROC
$x(n)$	$X(z)$	
$u(n)$	$\frac{1}{1-z^{-1}}$	$ z > 1$
$a^n u(n)$	$\frac{1}{1-az^{-1}}$	$ z > a$
$-a^n u(-n-1)$	$\frac{1}{1-az^{-1}}$	$ z < a$