

Notes on curved-sky quadratic estimation

Julien Carron

This document supplements the release of the Planck 2018 CMB lensing [1] pipeline. It collects the formulae relevant to curved-sky quadratic estimators in the spin-weight, position-space formalism, including in particular estimator cross-responses and Gaussian noise biases between arbitrary pairs of quadratic estimators. **JC: Document to be included with the pipeline release after submission of the revised L08. in progress**

CONTENTS

I. Spin-weight estimators	1
II. Gaussian covariance calculations	2
III. Response and cross-responses calculations	2
IV. Derivation of optimal QE weights	4
V. Examples	4
A. Lensing	4
B. Modulation	4
C. Polarization rotation	4
D. Point sources	4
E. Noise variance map anisotropies	4
References	5

Prior to projection onto gradient and curl modes, and prior to proper normalization, separable quadratic estimators can be written as a (sum of) products of two position-space maps. Let \hat{q} be such an unnormalized estimator:

$${}_{s_o+t_o}\hat{q}(\hat{n}) \equiv \left(\sum_{\ell m} w_{\ell}^{s_o s_i} {}_s\bar{X}_{\ell m} {}_{s_o}Y_{\ell m}(\hat{n}) \right) \cdot \left(\sum_{\ell m} w_{\ell}^{t_o t_i} {}_{t_i}\bar{X}_{\ell m} {}_{t_o}Y_{\ell m}(\hat{n}) \right) \quad (1.5)$$

where s_i, t_i are input spins, s_o, t_o outputs spins, and $w_{\ell}^{s_o s_i}, w_{\ell}^{t_o t_i}$ associated weights. For simplicity we use the same symbol w for the first and second leg even if they may differ in general for the same spin indices. By consistency with ${}_{-s_o-t_o}\hat{q}(\hat{n}) = {}_{s_o+t_o}\hat{q}^*(\hat{n})$ the weights have symmetry $w_{\ell}^{-s_o-s_i} = (-1)^{s_o+s_i} w_{\ell}^{*s_o s_i}$.

The maps ${}_s\bar{X}_{\ell m}$ are the inverse variance filtered CMB maps; the filtered scalar temperature

$${}_0\bar{X}_{\ell m} = \bar{T}_{\ell m} \quad (1.6)$$

and filtered spin ± 2 Stokes polarization ${}_{\pm 2}\bar{P} = \bar{Q} \pm i\bar{U}$,

$${}_{\pm 2}\bar{X}_{\ell m} = {}_{\pm 2}\bar{P}_{\ell m} = -(\bar{E}_{\ell m} \pm i\bar{B}_{\ell m}), \quad (1.7)$$

(for the purposes of the analytical calculations in this document) are isotropically related to the (beam-deconvolved) data maps ${}_sX^{\text{dat}}$ through a matrix F ,

$${}_s\bar{X}_{\ell m} \equiv \sum_{s_2=0,2,-2} F_{\ell}^{ss_2} {}_{s_2}X_{\ell m} \quad (1.8)$$

(isotropic approximation of $\bar{X} = \mathcal{B}^{\dagger} \text{Cov}^{-1} X^{\text{dat}}$ in the notation of Ref. [1]) Correlation functions:

$$\begin{aligned} \xi_{+}^{st}(\beta) &\equiv \left\langle e^{-is\alpha} {}_sX(\hat{n}_1) ({}_tX(\hat{n}_2)e^{-it\gamma})^{*} \right\rangle \\ \xi_{-}^{st}(\beta) &\equiv \left\langle (e^{-is\alpha} {}_sX(\hat{n}_1))^{*} ({}_tX(\hat{n}_2)e^{-it\gamma})^{*} \right\rangle \end{aligned} \quad (1.9)$$

γ is the angle at \hat{n}_1 that aligns the local x -axis to the geodesic connecting \hat{n}_1 and \hat{n}_2 (with the axis pointing towards \hat{n}_2), β the angle between \hat{n}_1 and \hat{n}_2 , and α is defined just as γ but at \hat{n}_2 . [2, 3] **JC: Mixups with n1 and n2 defs...fix this!**

I. SPIN-WEIGHT ESTIMATORS

JC: too many α s A complex spin- s field ${}_sf(\hat{n})$ is defined by the condition that it transforms under a rotation of angle ψ of the local axes through ${}_sf(\hat{n}) \rightarrow e^{is\psi} {}_sf(\hat{n})$. We use further the notation ${}_{-s}f(\hat{n}) = {}_sf^{*}(\hat{n})$.

The gradient (g) and curl (c) harmonic modes of definite parity of ${}_sf$ are then defined as follows

$$g_{\ell m}^s = -\frac{1}{2} (|s|f_{\ell m} + (-1)^s {}_{-|s|}f_{\ell m}) \quad (1.1)$$

$$c_{\ell m}^s = -\frac{1}{2i} (|s|f_{\ell m} - (-1)^s {}_{-|s|}f_{\ell m}) \quad (1.2)$$

where ${}_{\pm s}f_{\ell m} \equiv \int d^2n {}_{\pm s}f(\hat{n}) {}_{\pm s}Y_{\ell m}^{*}(\hat{n})$. Spin-0 fields are real and pure gradients. With these conventions, we have in particular for the CMB polarization ${}_{\pm P}$

$${}_{\pm 2}P_{\ell m} = -(E_{\ell m} \pm iB_{\ell m}). \quad (1.3)$$

where E and B are the gradient and curl mode. Note that the spin-0 intensity ${}_0T$ gradient is $-T_{\ell m}$ and not $T_{\ell m}$. Our polarization conventions are such that ${}_{\pm 2}P = Q \pm iU$, with the local x and y axes at each point \hat{n} pointing south and east (following e.g. the healpix software conventions). The relation inverse to Eqs. (1.1) and (1.2) is

$${}_{\pm |s|}f_{\ell m} = -(\pm)^s (g_{\ell m}^s \pm ic_{\ell m}^s). \quad (1.4)$$

$$\begin{aligned}\xi_+^{st}(\beta) &= (+1)^s \sum_L \left(\frac{2L+1}{4\pi} \right) \left[C_L^{g^s g^t} + C_L^{c^s c^t} - i \left(C_L^{g^s c^t} + C_L^{c^s g^t} \right) \right] d_{st}^L(\beta) \\ \xi_-^{st}(\beta) &= (-1)^s \sum_L \left(\frac{2L+1}{4\pi} \right) \left[C_L^{g^s g^t} - C_L^{c^s c^t} - i \left(C_L^{g^s c^t} - C_L^{c^s g^t} \right) \right] d_{-st}^L(\beta)\end{aligned}\quad (1.10)$$

(we adopt the convention, standard in CMB lensing, to write quadratic estimator multipoles with L, M and use

ℓ, m for the CMB fields from which they are built)

The formulae exposed in this document can be derived through simple application of this formal relation,

$$\begin{aligned}& \sum_{m_1, m_2} \int d^2 n_1 \, {}_{s_1} Y_{\ell_1 m_1}(\hat{n}_1) \, {}_{s_2} Y_{\ell_2 m_2}(\hat{n}_1) \, {}_{r_1} Y_{LM}(\hat{n}_1) \int d^2 n_2 \, {}_{t_1} Y_{\ell_1 m_1}(\hat{n}_2) \, {}_{t_2} Y_{\ell_2 m_2}(\hat{n}_2) \, {}_{r_2} Y_{L'M'}(\hat{n}_2) \\ &= \delta_{LL'} \delta_{MM'} \frac{2\ell_1+1}{4\pi} \frac{2\ell_2+1}{4\pi} 2\pi \int_{-1}^1 d\beta \, d_{s_1 t_1}^{\ell_1}(\beta) d_{s_2 t_2}^{\ell_2}(\beta) d_{r_1 r_2}^L(\beta) \quad (\text{whenever } s_1 + s_2 + r_1 = 0 = t_1 + t_2 + r_2).\end{aligned}\quad (1.11)$$

where $d_{mm'}^\ell$ are Wigner small d-matrices.

curl modes results in

$$\begin{aligned}\left\langle \hat{g}_{LM}^r \hat{g}_{L'M'}^{*,r'} \right\rangle_{\text{G.}} &= \delta_{LL'} \delta_{MM'} \frac{1}{2} \Re \left[C_L^{rr'} + (-1)^r C_L^{-rr'} \right] \\ \left\langle \hat{c}_{LM}^r \hat{c}_{L'M'}^{*,r'} \right\rangle_{\text{G.}} &= \delta_{LL'} \delta_{MM'} \frac{1}{2} \Re \left[C_L^{rr'} - (-1)^r C_L^{-rr'} \right] \\ \left\langle \hat{g}_{LM}^r \hat{c}_{L'M'}^{*,r'} \right\rangle_{\text{G.}} &= \delta_{LL'} \delta_{MM'} \frac{1}{2} \Im \left[-C_L^{rr'} - (-1)^r C_L^{-rr'} \right] \\ \left\langle \hat{c}_{LM}^r \hat{g}_{L'M'}^{*,r'} \right\rangle_{\text{G.}} &= \delta_{LL'} \delta_{MM'} \frac{1}{2} \Im \left[C_L^{rr'} - (-1)^r C_L^{-rr'} \right]\end{aligned}\quad (2.3)$$

where

$$C_L^{\pm rr'} \equiv 2\pi \int_{-1}^1 d\mu \, d_{\pm rr'}^L(\mu) \xi_{\pm}^{rr'}(\beta) \quad (2.4)$$

(\Re and \Im stands for real and imaginary parts respectively). Ref. [1] calculates the covariance matrix based on these equations using the empirical, realisation dependent power spectra $\bar{C}_\ell^{s_i t_i}$. A gradient-curl mode cross-covariance may be sourced by gradient-curl couplings in the inverse-variance filtered CMB fields (i.e., non-zero $C_\ell^{\bar{T}\bar{B}}$ or $C_\ell^{\bar{E}\bar{B}}$). In most relevant situations there is no such couplings and the gradient to curl and curl to gradient covariance vanish.

III. RESPONSE AND CROSS-RESPONSES CALCULATIONS

We now turn to the calculation of the response of the estimator to a source of anisotropy. Anisotropy can sometimes be parametrized at the level of the CMB maps, (for example for lensing), with

$${}_s \delta X(\hat{n}) = \sum_{a=\pm r} {}_a \alpha(\hat{n}) \left(\sum_{\ell m} R_\ell^{a,s} {}_s X_{\ell m} {}_{s-a} Y_{\ell m}(\hat{n}) \right) \quad (3.1)$$

for response kernel functions $R_\ell^{r,s}$. More generally, let the covariance of the CMB data respond as follows to a spin- r anisotropy source α :

II. GAUSSIAN COVARIANCE CALCULATIONS

Q.E. noise covariance can be evaluated with a series of one-dimensional integrals as was first demonstrated by Ref. [1]. For two generic estimators as defined in Eq. (1.5), we now obtain their gradient (g) and curl (c) covariances with four integrals as follows. **JC: describe here the matrix** For an isotropy estimator ${}_r \hat{\alpha}$ let $s = (s_i, s_o, w^{s_i s_o})$ collectively describes the in and out spins and weight function of the left leg, and similarly with t for the right leg (with $s_o + t_o = r$). In the same way, let u and v describes another estimator ${}_{r'} \hat{\alpha}$ (with $u_o + v_o = r'$). Then, their Gaussian correlation functions are

$$\xi_{\pm}^{rr'}(\beta) = \xi^{\pm s, u}(\beta) \xi^{\pm t, v}(\beta) + \xi^{\pm s, v}(\beta) \xi^{\pm t, u}(\beta), \quad (2.1)$$

where $\xi^{s,t}$ is

$$\xi^{s,t}(\beta) \equiv \sum_{\ell} \left(\frac{2\ell+1}{4\pi} \right) w_\ell^{s_o s_i} w_\ell^{* t_o t_i} \bar{C}_\ell^{s_i t_i} d_{s_o t_o}^\ell(\beta) \quad (2.2)$$

and $\bar{C}_\ell^{s_i t_i} \equiv \langle {}_{s_i} \bar{X}_{\ell m} {}_{t_i} \bar{X}_{\ell m}^* \rangle$. Projecting onto gradient and




FIG. 1. The geometry and angles in Eq. (1.9), with the local axes in green. It holds $\alpha(\hat{n}_2, \hat{n}_1) = \pi - \gamma(\hat{n}_1, \hat{n}_2)$ and $\gamma(\hat{n}_2, \hat{n}_1) = \pi - \alpha(\hat{n}_1, \hat{n}_2)$. Figure originally from Wayne Hu tutorials, <http://background.uchicago.edu/~whu/tamm/webversion/node5.html>.

$$\delta \langle {}_s X(\hat{n}_1) {}_t X^*(\hat{n}_2) \rangle = \sum_{\ell m, a=\pm r} {}_a \alpha(\hat{n}_1) W_\ell^{a, st} {}_{s-a} Y_{\ell m}(\hat{n}_1) {}_t Y_{\ell m}^*(\hat{n}_2) + W_\ell^{*a, ts} {}_s Y_{\ell m}(\hat{n}_1) {}_{t-a} Y_{\ell m}^*(\hat{n}_2) {}_a \alpha^*(\hat{n}_2) \quad (3.2)$$

for some weights functions $W_\ell^{a, st}$. For map-level descriptions in Eq. (3.1) then holds

$$W_\ell^{a, st} = R^{a, s} C_\ell^{st}. \quad (3.3)$$

However, Eq. (3.2) is more general. Section ?? lists weights functions of some relevant cases.

Let as before s, t denote collectively the QE spins and weight functions for an estimator ${}_r \hat{\alpha}(\hat{n})$ of spin $r = s_o + t_o$, and let r' be the spin of anisotropy source ${}_{r'} \beta(\hat{n})$ with covariance response kernel $W^{r'}$ as above. Let $\mathcal{R}_L^{g_r g_{r'}} \delta_{LL'} \delta_{MM'}$ be defined as the response of the gradi-

ent mode of α_{LM} to the gradient mode of $\beta_{L'M'}$, and similarly for the curl. It holds:

$$\begin{aligned} \mathcal{R}_L^{g_r g_{r'}} &= \Re \left[R_L^{st, r'} + (-1)^{r'} R_L^{st, -r'} \right] \\ \mathcal{R}_L^{c_r c_{r'}} &= \Re \left[R_L^{st, r'} - (-1)^{r'} R_L^{st, -r'} \right] \\ \mathcal{R}_L^{g_r c_{r'}} &= \Im \left[-R_L^{st, r'} + (-1)^{r'} R_L^{st, -r'} \right] \\ \mathcal{R}_L^{c_r g_{r'}} &= \Im \left[R_L^{st, r'} + (-1)^{r'} R_L^{st, -r'} \right] \end{aligned} \quad (3.4)$$

where

$$R_L^{st, r'} = 2\pi \int_{-1}^1 d\mu \, d_{rr'}^L(\mu) \sum_{\tilde{s}_i, \tilde{t}_i=0,2,-2} \left[\xi^{s_o s_i \tilde{s}_i}(\mu) \psi^{t_o t_i \tilde{t}_i \tilde{s}_i, r'}(\mu) + \xi^{t_o t_i \tilde{t}_i}(\mu) \psi^{s_o s_i \tilde{s}_i \tilde{t}_i, r'}(\mu) \right] \quad (3.5)$$

and

$$\begin{aligned}\xi^{s_o s_i \tilde{s}_i}(\mu) &\equiv \sum_{\ell} \left(\frac{2\ell+1}{4\pi} \right) w_{\ell}^{s_o s_i} F_{\ell}^{s_i \tilde{s}_i} d_{s_o, \tilde{s}_i}^{\ell}(\mu) \\ \psi^{s_o s_i \tilde{s}_i \tilde{t}_i, r'}(\mu) &\equiv \sum_{\ell} \left(\frac{2\ell+1}{4\pi} \right) w_{\ell}^{s_o s_i} F_{\ell}^{s_i \tilde{s}_i} W_{\ell}^{*-r', -\tilde{t}_i \tilde{s}_i} d_{s_o, -\tilde{t}_i + r'}^{\ell}(\mu)\end{aligned}\quad (3.6)$$

Again, in most relevant cases, the gradient to curl and curl to gradient responses do vanish. If there is a unique source of anisotropy, properly normalized gradient and curl estimators are then given by $\hat{g}_{LM}^r/\mathcal{R}_L^{g_r g_r}$ and $\hat{c}_{LM}^r/\mathcal{R}_L^{c_r c_r}$.

IV. DERIVATION OF OPTIMAL QE WEIGHTS

Optimal (in the sense of minimal Gaussian variance) QE weights are easily gained from the representation in Eq. 3.2 of the anisotropy. Let the CMB likelihood gradients be

$$\pm_r \hat{\alpha}(\hat{n}) = \frac{\delta}{\delta_{\mp r} \alpha(\hat{n})} - \frac{1}{2} {}_{s_1} X \text{Cov}_{s_1 s_2}^{-1} {}_{s_2} X \Big|_{\alpha \equiv 0} \quad (4.1)$$

where $\text{Cov}_{s_1 s_2}(\hat{n}, \hat{n}') \equiv \langle {}_{s_1} X(\hat{n}) {}_{s_2} X(\hat{n}') \rangle$, and where ${}_r \alpha(\hat{n})$ and ${}_{-r} \alpha(\hat{n})$ are treated as independent variables. Using Eq. (3.2) and comparing to Eq. (1.5), we find

$$\boxed{w_{\ell}^{st} = \delta_{st} \text{ (1st leg)} \quad w_{\ell}^{-s+r, t} = 2W_{\ell}^{-r, -st} \text{ (2nd leg)}} \quad (4.2)$$

JC: why 2 again? JC: FIXME: The right expression is

$${}_r \hat{g}(\hat{n}) = \sum_s -s \bar{X}(\hat{n}) \cdot (2W_{\ell}^{-r, st} {}_t \bar{X}_{\ell m} {}_{s+r} Y_{\ell m}(\hat{n})) \quad (4.3)$$

where \bar{X} has the $(0, 2, -2)$ elements (note the additional factor of 2! in pol w.r.t. to naive spin defs.)

$$\begin{pmatrix} \bar{T} \\ -\frac{1}{2} (\bar{E} + i\bar{B}) \\ -\frac{1}{2} (\bar{E} - i\bar{B}) \end{pmatrix} \quad (4.4)$$

Factor of 2 in front of W comes from $2\delta/\delta {}_{-r} \alpha(\hat{n})$ to get $d/dre + d/dim$ (?).

V. EXAMPLES

Examples include:

A. Lensing

The source of anisotropy is the spin-1 deflection field ${}_1 \alpha(\hat{n})$, with linear response (see Ref. [4]) $\delta_s X(\hat{n}) = -\frac{1}{2} \alpha_1(\hat{n}) \bar{\eth}_s X(\hat{n}) - \frac{1}{2} \alpha_{-1}(\hat{n}) \eth_s X(\hat{n})$ where \eth and $\bar{\eth}$ are the spin raising and spin lowering operator respectively. From the action of these operator on the spherical harmonics follow immediately

$$\begin{aligned}R_{\ell}^{-1, s} &= -\frac{1}{2} \sqrt{(l-s)(l+s+1)} \\ R_{\ell}^{1, s} &= +\frac{1}{2} \sqrt{(l+s)(l-s+1)}\end{aligned}\quad (5.1)$$

B. Modulation

The anisotropy source is a scalar, with response $\delta_s X(\hat{n}) = {}_0 \alpha(\hat{n}) {}_s X(\hat{n})$, hence

$$R_{\ell}^{0, s} = 1 \quad (5.2)$$

C. Polarization rotation

This is relevant in the case of systematic polarization angle miscalibration, or within more speculative ideas including cosmic birefringence. The observed polarization is rotated according to $\pm_2 P$ is $e^{\mp 2i {}_0 \alpha} \pm_2 P$. Hence,

$$R_{\ell}^{0, \pm 2} = \mp 2i \quad (5.3)$$

D. Point sources

Point sources in temperature (S^2 , see Ref. [5]): here anisotropy is sought of the form $\delta \langle T(\hat{n}) T(\hat{n}') \rangle = \delta_{\hat{n} \hat{n}'} S^2(\hat{n})$. Hence,

$$W_{\ell}^{r, st} = \frac{1}{4} \delta_{r0} \delta_{s0} \delta_{t0} \quad (5.4)$$

E. Noise variance map anisotropies

This is conceptually the same as point sources but acting on beam-convolved maps

$$W_{\ell}^{r, st} = \frac{1}{4} \delta_{r0} \delta_{s0} \delta_{t0} \frac{1}{b_{\ell}^2} \quad (5.5)$$




FIG. 2. Lensing gradient and curl reconstruction noise levels for a *Planck*-like experiment.

-
- [1] N. Aghanim *et al.* (Planck), “Planck 2018 results. VIII. Gravitational lensing,” (2018), arXiv:1807.06210 [astro-ph.CO].
 - [2] Gayoung Chon, Anthony Challinor, Simon Prunet, Eric Hivon, and Istvan Szapudi, “Fast estimation of polarization power spectra using correlation functions,” *Mon. Not. Roy. Astron. Soc.* **350**, 914 (2004), arXiv:astro-ph/0303414 [astro-ph].
 - [3] Anthony Challinor and Antony Lewis, “Lensed CMB power spectra from all-sky correlation functions,” *Phys. Rev. D* **71**, 103010 (2005), arXiv:astro-ph/0502425 [astro-ph].
 - [4] Anthony Challinor and Gayoung Chon, “Geometry of weak lensing of CMB polarization,” *Phys. Rev. D* **66**, 127301 (2002), arXiv:astro-ph/0301064 [astro-ph].
 - [5] Stephen J. Osborne, Duncan Hanson, and Olivier Doré, “Extragalactic Foreground Contamination in Temperature-based CMB Lens Reconstruction,” *JCAP* **1403**, 024 (2014), arXiv:1310.7547 [astro-ph.CO].