

## Lecture 2. Basic Probability Rules

**YULIA R. GEL**

**CS/SE/STAT 3341 Probability and Statistics  
in Computer Science and Software Engineering**

January 17, 2017

# 1 Operations of set theory


# Basics

It is useful to interpret (and visualize!) events via algebra of sets. First, we start from basic notations:

- The statement that a possible outcome of an experiment  $s$  is a member of  $S$  is denoted symbolically by the relation  $s \in S$ .
- Any event  $E$  can be regarded as a certain subset of a sample space  $S$ , i.e.  $E \subset S$  for all events  $E$ .
- An event  $A$  is said to be contained in another event  $B$  if every outcome that belongs to the subset defining the event  $A$  also belongs to the subset  $B$ , i.e.  $A \subset B$ .
- Some events are impossible! E.g., roll of a die cannot produce a negative number. Hence, such event of observing a negative number contains no outcomes. This subset of  $S$  is called the empty set and is denoted by the symbol  $\emptyset$ . Note that  $\emptyset \subset A \subset S$  for all events  $A$ .

# Unions

If  $A$  and  $B$  are any two events then the union of  $A$  and  $B$  is defined to be the event containing all outcomes that belong to  $A$  alone, to  $B$  alone or to both  $A$  and  $B$ . We denote the union of  $A$  and  $B$  by  $A \cup B$ .



# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cup B =$

# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cup B = B \cup A$

# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cup B = B \cup A$

②  $A \cup \emptyset =$

# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cup B = B \cup A$

②  $A \cup \emptyset = A$



# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $A \cup B = B \cup A$
- 2  $A \cup \emptyset = A$
- 3  $A \cup S =$

# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $A \cup B = B \cup A$
- 2  $A \cup \emptyset = A$
- 3  $A \cup S = S$

# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $A \cup B = B \cup A$
- 2  $A \cup \emptyset = A$
- 3  $A \cup S = S$
- 4  $A \cup A =$

# Properties of a Union

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- ①  $A \cup B = B \cup A$
- ②  $A \cup \emptyset = A$
- ③  $A \cup S = S$
- ④  $A \cup A = A$

# Properties of a Union


For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $A \cup B = B \cup A$
- 2  $A \cup \emptyset = A$
- 3  $A \cup S = S$
- 4  $A \cup A = A$

The union of  $n$  events  $E_1, E_2, \dots, E_n$  is defined to be the event that contains all outcomes which belong to at least one of the  $n$  events, and is denoted by  $\cup_{i=1}^n E_i$ .

# Intersection

If  $A$  and  $B$  are any two events then the intersection of  $A$  and  $B$  is defined to be the event containing all outcomes that belong both to  $A$  and  $B$ . We denote the intersection of  $A$  and  $B$  by  $A \cap B$ .



# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cap B =$

# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cap B = B \cap A$



# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cap B = B \cap A$

②  $A \cap \emptyset =$

# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $A \cap B = B \cap A$

②  $A \cap \emptyset = \emptyset$

# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- ①  $A \cap B = B \cap A$
- ②  $A \cap \emptyset = \emptyset$
- ③  $A \cap S =$

# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $A \cap B = B \cap A$
- 2  $A \cap \emptyset = \emptyset$
- 3  $A \cap S = A$

# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- ①  $A \cap B = B \cap A$
- ②  $A \cap \emptyset = \emptyset$
- ③  $A \cap S = A$
- ④  $A \cap A =$

# Properties of an Intersection

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- ①  $A \cap B = B \cap A$
- ②  $A \cap \emptyset = \emptyset$
- ③  $A \cap S = A$
- ④  $A \cap A = A$

# Properties of an Intersection


For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- ①  $A \cap B = B \cap A$
- ②  $A \cap \emptyset = \emptyset$
- ③  $A \cap S = A$
- ④  $A \cap A = A$

The intersection of  $n$  events  $E_1, E_2, \dots, E_n$  is defined to be the event that contains all outcomes which belong to all of the  $n$  events, and is denoted by  $\cap_{i=1}^n E_i$ .

# Complement

The complement of an event  $A$  is defined to be event that contains all outcomes in the sample space  $S$  which do not belong to  $A$ . The complement of  $A$  is denoted as  $A^c$ .





# Properties of a Compliment

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $(A^c)^c =$

# Properties of a Compliment

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $(A^c)^c = A$

# Properties of a Complement

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

①  $(A^c)^c = A$

②  $\emptyset^c =$

# Properties of a Compliment

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $(A^c)^c = A$
- 2  $\emptyset^c = S$

# Properties of a Compliment

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $(A^c)^c = A$
- 2  $\emptyset^c = S$
- 3  $A \cup A^c =$

# Properties of a Complement

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $(A^c)^c = A$
- 2  $\emptyset^c = S$
- 3  $A \cup A^c = S$

# Properties of a Compliment

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $(A^c)^c = A$
- 2  $\emptyset^c = S$
- 3  $A \cup A^c = S$
- 4  $A \cap A^c =$

# Properties of a Compliment

For any events  $A$  and  $B$  such that  $A \subset S$  and  $B \subset S$ , the following properties hold:

- 1  $(A^c)^c = A$
- 2  $\emptyset^c = S$
- 3  $A \cup A^c = S$
- 4  $A \cap A^c = \emptyset$



# Disjoint and Exhaustive Events

It is said that two events  $E_1$  and  $E_2$  are **disjoint** or **mutually exclusive** if  $E_1$  and  $E_2$  have no outcomes in common. It follows that  $E_1$  and  $E_2$  are disjoint if and only if  $E_1 \cap E_2 = \emptyset$ .

Events  $E_1, E_2, \dots$  are called **exhaustive** if  $\cup_i E_i = S$  and at least one  $E_i$  occurs for sure.

Example 1.  $E$  and  $E^c$  are disjoint and exhaustive. For instance, gender of a child, i.e. male and female are disjoint and exhaustive.

# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

①  $(A \cup B)^c =$

# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

$$\textcircled{1} (A \cup B)^c = A^c \cap B^c$$

# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

$$\textcircled{1} (A \cup B)^c = A^c \cap B^c$$

$$\textcircled{2} (A \cap B)^c =$$

# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

$$\textcircled{1} \quad (A \cup B)^c = A^c \cap B^c$$

$$\textcircled{2} \quad (A \cap B)^c = A^c \cup B^c$$

# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

$$\textcircled{1} (A \cup B)^c = A^c \cap B^c$$

$$\textcircled{2} (A \cap B)^c = A^c \cup B^c$$

$$\textcircled{3} A \cap (B \cup C) =$$

# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

$$\textcircled{1} (A \cup B)^c = A^c \cap B^c$$

$$\textcircled{2} (A \cap B)^c = A^c \cup B^c$$

$$\textcircled{3} A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

- 1  $(A \cup B)^c = A^c \cap B^c$
- 2  $(A \cap B)^c = A^c \cup B^c$
- 3  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- 4  $A \cup (B \cap C) =$



# DeMorgan's Laws

Let  $A$ ,  $B$  and  $C$  be subsets of  $S$ . Then,

- 1  $(A \cup B)^c = A^c \cap B^c$
- 2  $(A \cap B)^c = A^c \cup B^c$
- 3  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- 4  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$