

1. Convexity of Sets

Definition. A set C is convex if and only if the line segment between any two points in C lies in C :

$$C \text{ is convex} \iff \forall \vec{x}_1, \vec{x}_2 \in C, \forall \theta \in [0, 1], \theta \vec{x}_1 + (1 - \theta) \vec{x}_2 \in C \quad (1)$$

(a) Show that the **intersection of convex sets is convex**:

$$C_1, C_2 \text{ are convex} \implies C = C_1 \cap C_2 \text{ is convex} \quad (2)$$

(b) Show that the following sets are convex:

i. **(OPTIONAL)** A vector subspace of \mathbb{R}^n .

ii. **(OPTIONAL)** A hyperplane, $\mathcal{L} = \{\vec{x} \mid \vec{a}^\top \vec{x} = b\}$.

iii. A halfspace, $\mathcal{H} = \{\vec{x} \mid \vec{a}^\top \vec{x} \leq b\}$.

Definition. A function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is affine if it is the sum of a linear function and a constant,

$$f(\vec{x}) = A\vec{x} + \vec{b}, \quad (3)$$

for $A \in \mathbb{R}^{m \times n}$ and $\vec{b} \in \mathbb{R}^m$.

(c) **(OPTIONAL)** Prove that if $S \subseteq \mathbb{R}^n$ is convex, then the image of S under an affine function f ,

$$f(S) = \{f(\vec{x}) \mid \vec{x} \in S\}, \quad (4)$$

is convex.

2. Convexity of Functions

Definition. A function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is convex if $\text{dom}(f)$ is a convex set and if for all $\vec{x}, \vec{y} \in \text{dom}(f)$ and $\theta \in [0, 1]$, we have,

$$f(\theta\vec{x} + (1 - \theta)\vec{y}) \leq \theta f(\vec{x}) + (1 - \theta)f(\vec{y}). \quad (5)$$

The function f is strictly convex if the inequality is strict.

Definition. A function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is concave if $\text{dom}(f)$ is a convex set and if for all $\vec{x}, \vec{y} \in \text{dom}(f)$ and θ with $0 \leq \theta \leq 1$, we have,

$$f(\theta\vec{x} + (1 - \theta)\vec{y}) \geq \theta f(\vec{x}) + (1 - \theta)f(\vec{y}). \quad (6)$$

The function f is strictly concave if the inequality is strict.

Property. A function f is concave if and only if $-f$ is convex. An affine function is both convex and concave.

Property: Jensen's inequality. The inequality in Equation (5) is known as **Jensen's Inequality**. This can be extended to convex combinations of more than one point. If f is convex, and $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k \in \text{dom}(f)$, and $\theta_1, \theta_2, \dots, \theta_k \geq 0$ with $\sum_{i=1}^k \theta_i = 1$ then,

$$f(\theta_1\vec{x}_1 + \theta_2\vec{x}_2 + \dots + \theta_k\vec{x}_k) \leq \theta_1 f(\vec{x}_1) + \theta_2 f(\vec{x}_2) + \dots + \theta_k f(\vec{x}_k). \quad (7)$$

Property: first order condition. Suppose f is differentiable. Then f is convex if and only if $\text{dom}(f)$ is convex and

$$f(\vec{y}) \geq f(\vec{x}) + \nabla f(\vec{x})^\top (\vec{y} - \vec{x}), \quad (8)$$

for all $\vec{x}, \vec{y} \in \text{dom}(f)$.

Property: Second order condition. Suppose f is twice differentiable. Then f is convex if and only if, $\text{dom}(f)$ is convex and the Hessian of f , $\nabla^2 f(\vec{x})$, is positive semi-definite for all $\vec{x} \in \text{dom}(f)$.

(a) Under what condition on $A \in \mathbb{R}^{n \times n}$, where A is symmetric, is the function $f: \vec{x} \rightarrow \vec{x}^\top A \vec{x}$ convex?

(b) (OPTIONAL) Restriction to a line.

Show that a function f is convex if and only if for all $\vec{x} \in \text{dom}(f)$ and all \vec{v} , the function $g : \text{dom}(g) \rightarrow \mathbb{R}$ given by $g(t) = f(\vec{x} + t\vec{v})$ is convex for $\text{dom}(g) = \{t \in \mathbb{R} \mid \vec{x} + t\vec{v} \in \text{dom}(f)\}$.

(c) (OPTIONAL) Non-negative weighted sum.

Show that the non-negative weighted sum of convex functions is convex: i.e. if f_1, \dots, f_n are n convex functions from \mathbb{R}^n to \mathbb{R} and $w_1, \dots, w_n \in \mathbb{R}_+$ are n positive scalars, then the function:

$$f = \sum_{i=1}^n w_i f_i \quad (9)$$

is convex. To make the question easier, you can assume that the functions f_1, \dots, f_n are twice-differentiable.

(d) Point-wise maximum.

Show that if f_1 and f_2 are convex functions then their pointwise maximum f , defined by

$$f(\vec{x}) = \max(f_1(\vec{x}), f_2(\vec{x})), \quad (10)$$

with $\text{dom}(f) = \text{dom}(f_1) \cap \text{dom}(f_2)$, is also convex.

(e) Show that a piece-wise linear function that can be written as,

$$f(\vec{x}) = \max(\vec{a}_1^\top \vec{x} + \vec{b}_1, \vec{a}_2^\top \vec{x} + \vec{b}_2, \dots, \vec{a}_m^\top \vec{x} + \vec{b}_m), \quad (11)$$

is convex.

3. Convexity and composition of functions

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$. Define the composition of f with g as $h = f \circ g : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $h(\vec{x}) = f(g(\vec{x}))$.

- (a) Show that if f is convex and non decreasing and g is convex, then h is convex.
- (b) Show that there exists f non decreasing and g convex, such that $h = f \circ g$ is not convex.
- (c) Show that there exists f convex and g convex such that $h = f \circ g$ is not convex.