

# Potential Predictive Value of *TP53* and *KRAS* Mutation Status for Response to PD-1 Blockade Immunotherapy in Lung Adenocarcinoma

Zhong-Yi Dong<sup>1,2</sup>, Wen-Zhao Zhong<sup>1</sup>, Xu-Chao Zhang<sup>1</sup>, Jian Su<sup>1</sup>, Zhi Xie<sup>1</sup>, Si-Yang Liu<sup>1</sup>, Hai-Yan Tu<sup>1</sup>, Hua-Jun Chen<sup>1</sup>, Yue-Li Sun<sup>1</sup>, Qing Zhou<sup>1</sup>, Jin-Ji Yang<sup>1</sup>, Xue-Ning Yang<sup>1</sup>, Jia-Xin Lin<sup>1</sup>, Hong-Hong Yan<sup>1</sup>, Hao-Ran Zhai<sup>1,2</sup>, Li-Xu Yan<sup>3</sup>, Ri-Qiang Liao<sup>1</sup>, Si-Pei Wu<sup>1</sup>, and Yi-Long Wu<sup>1,2</sup>

## Abstract

**Purpose:** Although clinical studies have shown promise for targeting programmed cell death protein-1 (PD-1) and ligand (PD-L1) signaling in non-small cell lung cancer (NSCLC), the factors that predict which subtype patients will be responsive to checkpoint blockade are not fully understood.

**Experimental Design:** We performed an integrated analysis on the multiple-dimensional data types including genomic, transcriptomic, proteomic, and clinical data from cohorts of lung adenocarcinoma public (discovery set) and internal (validation set) database and immunotherapeutic patients. Gene set enrichment analysis (GSEA) was used to determine potentially relevant gene expression signatures between specific subgroups.

**Results:** We observed that *TP53* mutation significantly increased expression of immune checkpoints and activated T-effector and interferon- $\gamma$  signature. More importantly, the *TP53/KRAS* comutated subgroup manifested exclusive increased expression

of PD-L1 and a highest proportion of *PD-L1*<sup>+</sup>/*CD8A*<sup>+</sup>. Meanwhile, *TP53*- or *KRAS*-mutated tumors showed prominently increased mutation burden and specifically enriched in the transversion-high (TH) cohort. Further analysis focused on the potential molecular mechanism revealed that *TP53* or *KRAS* mutation altered a group of genes involved in cell-cycle regulating, DNA replication and damage repair. Finally, immunotherapeutic analysis from public clinical trial and prospective observation in our center were further confirmed that *TP53* or *KRAS* mutation patients, especially those with co-occurring *TP53/KRAS* mutations, showed remarkable clinical benefit to PD-1 inhibitors.

**Conclusions:** This work provides evidence that *TP53* and *KRAS* mutation in lung adenocarcinoma may be served as a pair of potential predictive factors in guiding anti-PD-1/PD-L1 immunotherapy. *Clin Cancer Res*; 23(12); 3012–24. ©2016 AACR.

## Introduction

Recent clinical trials with anti-programmed cell death 1 (PD-1) and its ligand PD-1 ligand (PD-L1) therapies have shown unprecedented durable responses in patients with non-small cell lung cancer (NSCLC; refs. 1, 2). Unfortunately, only a minority of the total of treated patients respond to the current immunotherapy (3). The factors that determine which patients will be drug sensitive or resistant are not fully understood. Therefore, it has become a primary priority to identify the biomarkers that determine the responsiveness to checkpoint blockade, and to develop

strategies that could potentially increase the patient response rates. Encouragingly, recent studies had demonstrated that tumor mutational load (4–6), DNA mismatch repair (MMR) deficiency (7), the intensity of *CD8*<sup>+</sup> T cell infiltrates (8, 9) and intratumoral PD-L1 expression (10, 11) have each been proposed as distinct biomarkers of response to anti-PD-1/PD-L1 therapies. Meanwhile, these factors are functionally interrelated and are often found coordinately in individual tumor specimens (12). This raises the question of whether there exist some other variables simultaneously affect two or more of these above factors so as to provide stronger predictive value for therapeutic outcomes.

The identification of subsets of lung adenocarcinoma with oncogenic drivers has transformed the treatment of NSCLC, particularly for patients whose tumors harbor activating mutations in *EGFR*. However, the goal of developing specific therapeutic strategies for those bearing activating mutations in *KRAS* has thus far proven elusive. Meanwhile, mutations in tumor suppressor genes *TP53* and *STK11* are also common in lung adenocarcinoma and frequently co-occur with *KRAS* mutations (13–15). Given that activation of specific oncogenic pathways can have broad effects on gene expression, it is reasonable to imagine that the genetic make-up of cancer cells could have major effects on the immune tumor microenvironment (TME), by driving specific immune-related pathways. This could be through induction of immune checkpoints, secretion

<sup>1</sup>Guangdong Lung Cancer Institute, Guangdong General Hospital and Guangdong Academy of Medical Sciences, Guangzhou, China. <sup>2</sup>Southern Medical University, Guangzhou, China. <sup>3</sup>Department of Pathology and Laboratory Medicine, Guangdong General Hospital and Guangdong Academy of Medical Sciences, Guangzhou, China.

**Note:** Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

**Corresponding Author:** Yi-Long Wu, Guangdong General Hospital and Guangdong Academy of Medical Sciences, 106 Zhongshan Er Road, Guangzhou 510080, China. Phone: 8620-83877855; Fax: 8620-83827712; E-mail: [syylwu@live.cn](mailto:syylwu@live.cn)

**doi:** 10.1158/1078-0432.CCR-16-2554

©2016 American Association for Cancer Research.

### Translational Relevance

Programmed cell death ligand 1 (PD-L1) expression, tumor mutational load, and the intensity of CD8<sup>+</sup> T-cell infiltrates have recently been proposed as predictive biomarkers for response to PD-1 blockade immunotherapy. However, there are still many treatment responses beyond the explanation of these factors. It is increased need for more effective biomarkers for PD-1 blockade. We demonstrated *TP53* and *KRAS* mutation had remarkable effects on increasing PD-L1 expression, facilitating T-cell infiltration and augmenting tumor immunogenicity. More important, we confirmed that patients with *TP53* and/or *KRAS* mutation showed sensitivity to PD-1 blockade. These findings represent the first demonstration of potential predictive value of *TP53* and *KRAS* mutation for response to PD-1 blockade immunotherapy in lung adenocarcinoma.

of specific cytokines or production of chemokines that recruit specific cell types (12). Recent studies had shown *TP53* or *KRAS* mutant NSCLC expressed higher levels of PD-L1 protein compared with corresponding wild-type tumors (16, 17). Meanwhile, it has been demonstrated that loss of *TP53* function decreased genomic stability and was associated with defects in DNA damage repair, indicating a higher mutational burden might occur in *TP53* mutational tumor (18). Therefore, we speculate that common mutations as *TP53* and *KRAS* in lung adenocarcinoma may be served as effective predictive factors in guiding anti-PD-1/PD-L1 immunotherapy.

Here, in order to systematically address the potential mechanism that *TP53*/*KRAS* mutation mediate immune response to lung adenocarcinoma, we describe an integrative analysis that incorporates tumor mutational load, DNA MMR deficiency, intratumoral PD-L1 expression, and content of CD8<sup>+</sup> T-cell infiltrates from cohorts of both lung adenocarcinoma repository database analysis and clinical immunotherapeutic patients. Significantly, we uncover *TP53*/*KRAS* mutation as a superiority group to anti-PD-1/PD-L1 therapies and highlight a new insight into common mutations in guiding immunotherapy.

## Materials and Methods

### Clinical cohorts

The Cancer Genome Atlas (TCGA), GSE72094 and Broad cohorts were retrieved from online data repository. A total of 462 patients were included in the TCGA cohort with mRNA expression profiling and gene mutation data. The GSE72094 cohort recruited 442 patients with detailed mRNA expression data and *EGFR*/*KRAS*/*TP53*/*STK11* sanger sequencing analysis (19). The Broad cohort contained 183 lung adenocarcinomas and matched normal tissues with detail information about mutation load and mutation spectrum (20). Most of the patients enrolled in the three cohorts were early-stage lung adenocarcinomas. A total of 85 lung adenocarcinomas from the Guangdong Lung Cancer Institute (GLCI), Guangdong General Hospital (GGH) were underwent whole genome sequencing (WGS). Key variables including demographic and clinical information are provided in Supplementary Table S1.

### Immunotherapeutic patients

Clinical and mutation data for 34 NSCLC [29 adenocarcinoma (ADC)] patients were retrieved from cBioPortal ([http://www.cbioportal.org/study.do?cancer\\_study\\_id=luad\\_mskcc\\_2015](http://www.cbioportal.org/study.do?cancer_study_id=luad_mskcc_2015)). All patients treated with pembrolizumab (anti-PD-1) from 2012 to 2013 followed the protocol NCT01295827 (KEYNOTE-001). Objective response to pembrolizumab was assessed by investigator-assessed immune-related response criteria (irRC) by a study radiologist (5).

Another group consisted of 20 NSCLC (15 ADC) patients were collected prospectively in the GLCI from August 2015 to August 2016. Eleven of them were treated with pembrolizumab and nine patients were treated with nivolumab. Tumor specimens were obtained for Sanger sequencing and IHC analysis. This study was approved by the Institutional Review Board of GLCI of GGH, and all patients provided specimens with written informed consent. Clinicopathologic and molecular information are provided in Supplementary Table S2.

### mRNA expression profiling and reverse phase protein array (RPPA) analysis

For lung adenocarcinomas included in the TCGA cohort, experimental procedures regarding RNA extraction from tumors, mRNA library preparation, sequencing (on the Illumina HiSeq platform), quality control, and subsequent data processing for quantification of gene expression have been previously reported (21). Gene expression data for the GSE72094 lung adenocarcinomas have been deposited in the GEO repository (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE72094>). Tumors from the GSE72094 cohort were profiled using a custom Affymetrix GeneChip. The gene expression cutoff value was chosen as median over the entire dataset (TCGA and GSE72094) to ensure all analyses of each gene were based on the same cutoff value (22, 23).

Proteomic analysis was based on RPPA from the TCGA database. The RPPA methodology and data analysis pipeline have been previously described (21). For TCGA, level 3 data were downloaded directly from the TCGA portal and utilized in subsequent analyses.

### Mutation data analysis

For the discovery set, somatic mutation data (level 2) of the 462 lung adenocarcinomas were retrieved from the TCGA data portal (<https://gdc.cancer.gov/>). To assess the mutational load, the number of mutated genes carrying at least one nonsynonymous mutation in the coding region was computed for each tumor. Somatic mutation data of 183 lung adenocarcinomas in Broad cohort was retrieved from cBioPortal ([http://www.cbioportal.org/study.do?cancer\\_study\\_id=luad\\_broad](http://www.cbioportal.org/study.do?cancer_study_id=luad_broad)). Somatic substitutions and covered bases within their trinucleotide sequence context were analyzed to characterize the mutation spectrum of 183 lung adenocarcinoma. Mutation spectrum for each sample was calculated as the percentage of each of six possible single nucleotide changes (AT>CG, AT>GC, AT>TA, GC>AT, GC>CG, GC>TA) among all single-nucleotide substitutions. The most frequent mutation signatures were C→T transitions and C→A transversions.

For the validation set (GLCI), we conducted whole-exome sequencing of DNA from tumors and matched normal blood from 85 lung adenocarcinoma patients. Enriched exome libraries were sequenced on the HiSeq 2000 platform (Illumina) to >100× coverage. Alignment, base-quality score recalibration and duplicate-read removal were performed, germline variants were

excluded, mutations annotated and indels evaluated as previously described (4, 5, 24). Mutations between clinical groups were compared using the Mann-Whitney test.

### Gene set enrichment analysis (GSEA)

For GSEA (25), the javaGSEA Desktop Application was downloaded from <http://software.broadinstitute.org/gsea/index.jsp>. GSEA was used to associate the gene signature with the *TP53* or *KRAS* mutation status (*TP53*-mut vs. *TP53*-wt; *KRAS*-mut vs. *KRAS*-wt). The genes identified to be on the leading edge of the enrichment profile were subject to pathway analysis. Fold-change values were exported for all genes and analyzed with version 2.2.0 of GSEA, using the GseaPreranked module. The normalized enrichment score (NES) is the primary statistic for examining gene set enrichment results. The nominal *P* value estimates the statistical significance of the enrichment score. A gene set with nominal  $P \leq 0.05$  was considered to be significantly enriched in genes.

### Immunohistochemistry

Tumor sections were assessed immunohistochemically using PD-L1 (clone: SP142, Spring Bioscience, Inc) and CD8 (clone: C8/144B, Gene Tech (Shanghai) Co. Ltd). The IHC-stained tissue sections were scored separately by two pathologists blinded to the clinical parameters.

PD-L1 expression on tumor cells and immune cell was evaluated using a three-tiered grading system. Strong:  $\geq 50\%$  for tumor cell (TC) or  $\geq 10\%$  for immune cell (IC); weak: 5%–49% for TC or 5%–9% for IC; negative:  $< 5\%$  for TC or IC.

The percentages of CD8<sup>+</sup> lymphocytes compared with the total amount of nucleated cells in the stromal compartments were assessed. Scoring cutoff points at 25% or 50% for each core according to the degree of cell densities: low density:  $< 25\%$ ; intermediate density: 25% to 49%; high density:  $\geq 50\%$ .

### Sanger sequencing

Genomic DNA from each sample was used for sequence analysis of *EGFR* exons 18–21, *KRAS* exons 2–3 and *TP53* exons 2–11. These exons were amplified by the polymerase chain reaction (PCR) as previously described (19, 26), and the resulting PCR products were purified and labeled for sequencing using the Big Dye 3.1 Kit (Applied Biosystems) according to the manufacturer's protocol.

### Statistical analyses

Statistical analyses were conducted using GraphPad Prism (version 7.01) and SPSS version 22.0 (SPSS, Inc.). Scatter dot plot and Box and whisker plots indicate median and 95% confidence interval (CI). Statistical tests were used to analyze the clinical and genomic data, including the Mann-Whitney *U*,  $\chi^2$ , Fisher exact, and Kruskal-Wallis. Kaplan-Meier curves analysis of progression-free survival (PFS) were compared using the log-rank test. All reported *P* values are two-tailed, and for all analyses,  $P \leq 0.05$  is considered statistically significant, unless otherwise specified.

## Results

### Correlation between *TP53* and *KRAS* mutation and PD-L1 expression in lung adenocarcinoma

To investigate the correlation between common mutations (*TP53*, *KRAS*, *EGFR*, and *STK11*) and immune checkpoint status

in lung adenocarcinoma, we thus initially interrogated RNA sequencing (RNA-Seq) expression data from a repository database including 462 lung adenocarcinomas from The Cancer Genome Atlas (TCGA) and 442 lung adenocarcinomas from GEO repository (GSE72094). Both the TCGA and GEO databases showed significantly increased *PD-L1* mRNA expression in the *TP53* mutation subgroup than in other gene mutation. Specifically, the *TP53* and *KRAS* comutated group manifested prominent higher *PD-L1* expression than other comutation types (Fig. 1A).


We next sought to explore the impact of *TP53* and *KRAS* mutation on PD-L1 expression in both PD-L1 mRNA expression profiling and RPPA analysis based on the TCGA database. The results demonstrated that it was *TP53* mutation but not *KRAS* mutation that boosted PD-L1 expression (Supplementary Fig. S1A and S1B). Significantly, those with co-occurring mutations in *TP53* and *KRAS* revealed the highest PD-L1 expression (both mRNA and protein level) than single gene mutation or wild-type tumors, indicating potential synergistic effect on activating PD-L1 expression (Fig. 1B). To confirm the association between *TP53/KRAS* mutation and PD-L1 expression as repository data demonstrated, we detected 93 lung adenocarcinoma surgical specimens using an IHC analysis (Fig. 1C; Supplementary Fig. S1C and Table S3) and immunostaining shows *TP53/KRAS* comutated specimens the strongest staining for the PD-L1 protein (Fig. 1D).

Next, we further analyzed the association between *TP53* or *KRAS* mutation and other non-PD-L1 immune checkpoints. A heatmap depicted the expression level of key immune checkpoints to three groups (*TP53*, *KRAS*, and *TP53/KRAS*; Fig. 1E). The results displayed remarkable increased expression of most checkpoints in the *TP53* mutation group while decreased expression in the *KRAS* mutation group. More interestingly, the *TP53/KRAS* comutated subgroup manifested exclusive increased expression of PD-L1; however, it showed decreased expression of some other non-PD-L1 immune inhibitory checkpoints, such as Lymphocyte Activating 3 (LAG3) and V-Set Domain Containing T Cell Activation Inhibitor 1 (VTCN1; ref. 27), implying a potential candidate population for anti-PD-1/PD-L1 immunotherapy (Fig. 1F).

### *TP53* mutation facilitates CD8<sup>+</sup> T-cell infiltration and activates T-effector and interferon- $\gamma$ (IFN $\gamma$ ) associated gene signature

The presence of tumor-infiltrating lymphocytes (TIL) is an important biomarker for predicting responses to PD-L1 blockade therapy. We continue to analyze the correlation between these above common mutations and CD8<sup>+</sup> TIL contents in lung adenocarcinoma based on the TCGA database. Our results revealed significantly increased expression of *CD8A* in *TP53* mutation and *TP53/KRAS* comutated than other groups (Fig. 2A). It has been proposed that four different types of immune tumor microenvironments (TME) exist based on the presence or absence of TIL and PD-L1 expression. To further explore whether *TP53* or *KRAS* mutation would influence the TME, we analyzed the correlation between *TP53* or *KRAS* mutation and TME immune types classified based on *PD-L1* and *CD8A* expression as previously described (28, 29). Positive *PD-L1* and *CD8A* were defined as above-median expression. We identified that the *TP53* mutation group displayed a higher proportion of dual positive *PD-L1* and *CD8A* (*PD-L1*<sup>+</sup>/*CD8A*<sup>+</sup>) than the *TP53* wild-type group, while there was no difference between *KRAS* mutation and wild-type (Fig. 2B and C),



**Figure 1.**

Correlation of *TP53* and *KRAS* mutation with PD-L1 expression in patients with lung adenocarcinoma. **A**, Correlations between common mutations (*TP53/KRAS/EGFR/STK11*) and PD-L1 mRNA expression in lung adenocarcinoma patients based on the analysis of the TCGA and GEO repository (GSE72094) database. **B**, Quantitative analysis of PD-L1 mRNA and protein expression based on *TP53* and *KRAS* mutation status. **C**, Comparison of PD-L1 IHC H-score between *TP53* or *KRAS* mutation and corresponding wild-type tumors in a cohort of 93 lung adenocarcinomas. **D**, Representative images of PD-L1 immunostaining in lung adenocarcinoma tissues with indicated gene mutation. Scale bar, 200  $\mu$ m. **E**, Heatmap representation of relative mRNA expression levels of selected immune inhibitory checkpoints. **F**, Quantitative analysis of two typical inhibitory checkpoints (*LAG3* and *VTCN1*) on the base of *TP53* and *KRAS* mutation status. Mut, mutation; wt, wild-type; \*\*\*,  $P < 0.001$ ; \*\*,  $P < 0.01$ ; \*,  $P < 0.05$ .


suggesting an adaptive immune resistance TME existed in the *TP53* mutation population. More importantly, the *TP53/KRAS* comutated subgroup showed the highest proportion of *PD-L1*<sup>+</sup>/*CD8A*<sup>+</sup> than the *TP53* or *KRAS* single mutation and wild-type group (Fig. 2D). These observations were further confirmed by our IHC analysis that *TP53/KRAS* comutated patients manifested a strong staining of PD-L1 and high intensity of *CD8*<sup>+</sup>TILs (Fig. 2E).

Given that *TP53* mutation had effects on the TME in lung adenocarcinoma, we subsequently sought to assess the relationship between *TP53* mutation and T-effector and IFN $\gamma$ -associated gene signature, which have previously been associated with activated T cells, immune cytolytic activity, and IFN $\gamma$  release (30, 31). An integrated heatmap depicting expression levels of T-effector and IFN $\gamma$  associated gene signature in tumors with *TP53* mutation compared with *TP53* wild-type. We identified significant increased expression of both T-effector and IFN $\gamma$ -associated genes in the *TP53* mutation group, while there were no differences between *KRAS* mutation and wild-type, indicating preexisting immunity within *TP53* mutation tumor tissue (Fig. 2F and G).

### *TP53* and *KRAS* mutation shows increased mutation burden and distinct mutation spectrum

Recent studies have highlighted the relevance of tumor mutational loads and response to PD-1 blockade (5). We next speculate whether there are some common mutations in lung adenocarcinoma that affect the whole tumor mutational profile and change the tumor antigenicity. We first analyzed the TCGA and Broad databases as discovery set. The TCGA analysis showed significantly increased mutational loads in the *TP53* mutation group (median, 325), followed by *KRAS* (median, 179) and *STK11* (median, 132) mutation, *EGFR* (median, 60) mutation tumor had the lowest mutational loads. Meanwhile, the *TP53/KRAS* comutated subgroup showed significantly higher mutational loads (median, 358) than other comutated subgroup (Fig. 3A). We then tested these findings using another dataset (Broad), which consisted of 183 lung adenocarcinomas with detailed somatic mutation data, and confirmed that *TP53* and *KRAS* mutation and the *TP53/KRAS* comutated group had higher mutational loads than other groups (Fig. 3A). To further verify these findings, a total of 85 lung adenocarcinomas from GLCI detected by whole genome sequencing were defined as the validation set. GLCI data manifested the

Dong et al.


**Figure 2.**

*TP53* mutation facilitates  $CD8^+$  T-cell infiltration and activates T-effector and IFN $\gamma$ -associated gene signature. **A**, Association between common mutations (*TP53*/*KRAS*/*EGFR*/*STK11*) and *CD8A* mRNA expression in lung adenocarcinoma patients based on analysis of the TCGA dataset. **B-D**, The correlation between *TP53* or *KRAS* mutation status and TME immune types classified based on *PD-L1* and *CD8A* expression. Positive *PD-L1* and *CD8A* were defined as above-median expression. **E**, Representative images of *PD-L1* and *CD8* immunostaining in different subgroups according to *TP53* and *KRAS* mutation status. **F**, Heatmap depicting mRNA expression levels of T-effector and IFN $\gamma$ -associated gene signature. **G**, Quantitative analysis of four key genes (*GZMB*, *CXCL9*, *STAT1*, and *IFN $\gamma$* ) in T-effector and IFN $\gamma$  gene signature on the base of *TP53* and *KRAS* mutation status. Mut, mutation; wt, wild-type; \*\*\*,  $P < 0.001$ ; \*\*,  $P < 0.01$ ; \*,  $P < 0.05$ .

similar results with the discovery set that *TP53* mutation and the *TP53*/*KRAS* comutated group had higher mutational loads than other groups (Fig. 3B). It is well known that tobacco exposure was responsible for much of the mutagenesis in NSCLC. Multivariate linear regression analysis of mutation count in patients stratified by smoking status manifested that *TP53* mutation was an independent factor responsible for increased mutation burden regardless of smoking status, while *KRAS* mutation showed increased mutation burden only in nonsmokers (Supplementary Table S4).

Previous studies have established the notion that somatic mutations are primarily GC>TA transversions (32). We next investigated whether these above common mutations could affect

tumor mutation spectrum by using a TCGA cohort. Transversion-high (TH) and transversion-low (TL) was based on smoking history and GC>AT, GC>TA frequency as previously described (5, 21). We can identify *KRAS* mutations were significantly enriched in the TH cohort, while *EGFR* mutations were significantly enriched in the TL group (Supplementary Fig. S2). Consistent with TCGA results, the Broad dataset showed a high rate of transversion/transition (Tv/Ti) in *KRAS* mutation and the *TP53*/*KRAS* comutated group while the lowest rate Tv/Ti in *EGFR* mutation (Fig. 3C). Notably, *TP53* and *KRAS* mutation was significantly correlated with high somatic mutations, high rate of Tv/Ti and C>A transversion and high smoking index (pack-years; Fig. 3D).

**Figure 3.**

*TP53* and *KRAS* mutation augments tumor antigenicity by transforming the mutational profile. **A**, Different tumor mutational burden driving by a specific mutation gene analyzed on the base of the discovery set (TCGA and Broad database). **B**, Different tumor mutational burden driving by specific mutation gene analyzed on the base of the validation set (GLIC data). **C**, Box plot represents the proportion of Tv/Ti according to indicated mutation subgroups. **D**, Heatmap displays integrated relationship between mutation burden, mutation spectrum, smoking, and 4 common mutations status based on analysis of the Broad database. \*\*\*,  $P < 0.001$ ; \*\*,  $P < 0.01$ ; \*,  $P < 0.05$ .

### Impact of *TP53* and *KRAS* mutation on the cell cycle, DNA replication, and damage repair-related genes

We sought to determine whether alterations in DNA replication and damage repair-related genes resulted from *TP53* or *KRAS* mutation could account for differential mutation burden and mutation spectrum. GSEA reveals prominent enrichment of signatures relating to cell cycle, DNA replication and DNA repair in both the *TP53* and *KRAS* mutation groups. However, there were distinct differences between these two groups. *TP53* mutation predominantly led to acceleration of cell-cycle and DNA replication, which potentially increased mutation probability, for unrepaired DNA damages that do not kill the cell by blocking replication would tend to cause replication errors and thus mutation. *KRAS* mutation manifested various defects of DNA repair including MMR, nucleotide excision repair (NER), and base excision repair (BER) that greatly enhanced point mutation (Fig. 4A).

Recent studies showed that *POLE* mutation is associated with disruption of the exonuclease activity required for DNA proof-reading and results in a high mutational burden or an "ultra-mutator" phenotype (33, 34). We identified significantly increased mutation frequencies of *POLE* in the *TP53* mutation group ( $P = 0.002$ ) while decreased mutation frequencies of *POLE* in the *EGFR* and *STK11* mutation groups compare with their


corresponding wild-type group, indicating *TP53* mutation tends to cause DNA replication errors (Fig. 4B).

We next determined the correlation between these common mutations and DNA damage repair-related genes. DNA double-strand breaks (DSB) elicit that DNA damage response largely relies on the activity of ataxia telangiectasia mutated (ATM), which have been found to be mutated in human disorders associated with genome instability (35, 36). The results had revealed that a high frequency of ATM mutation was found predominantly in the *KRAS* and *STK11* mutation groups, and ATM protein analysis further confirmed that waning expression of ATM protein was specifically found in the *KRAS* and *STK11* mutation groups (Fig. 4B and C).

MMR-deficient tumors were recently shown susceptibility to checkpoint blockade immunotherapy (7). Our former GSEA identified that *KRAS* mutation was negatively correlated with MMR-related gene expression, and we next verified whether *KRAS* or other genes mutation affected the mutation status and protein expression of MMR-related genes. Four primary MMR-related genes, including *MSH2*, *MSH6*, *MLH1*, and *PMS2*, were co-analyzed. Consistent with GSEA, high mutation frequency of MMR-related genes was exclusively identified in the *KRAS* mutation group. Furthermore, the protein of *MSH2* and *MSH6* was significantly decreased in tumors with *KRAS* mutation; however, it was



Dong et al.

**Figure 4.**


Impact of *TP53* and *KRAS* mutation on the cell cycle, DNA replication and damage repair-related gene signatures. **A**, GSEA reveals acceleration of cell-cycle and DNA replication-related gene signatures as prominent modules in the *TP53* mutation group and impaired of DNA damage repair-related gene signatures, including MMR, NER, and BER in the *KRAS* mutation group compared with wild-type. **B**, Estimated proportion representation of *POLE*, *ATM* and MMR-related gene mutations in four groups according to indicated gene mutational status. **C**, Box plot representation of MMR-related proteins (MSH2 and MSH6) and ATM protein in four groups according to indicated genes' mutational status. NSE, normalized enrichment score; mut, mutation; wt, wild-type; \*\*\*,  $P < 0.001$ ; \*\*,  $P < 0.01$ ; \*,  $P < 0.05$ .

increased in tumors with *TP53* mutation, suggesting that *KRAS* mutation might be a potential driver agent to induce MMR deficiency and in consequence produce more neoantigens (Fig. 4B and C).

#### Patients with *TP53* or *KRAS* mutation, especially co-occurring *TP53/KRAS* mutations, show favorable clinical benefit to anti-PD-1 treatment

*TP53* and *KRAS* mutation showed remarkable effects on regulating PD-L1 expression, facilitating T-cell infiltration and

augmenting tumor immunogenicity. We presumed that patients with these two mutations probably had increased sensitivity to PD-1 blockade immunotherapy. In support of this hypothesis, publicly available trial data (MSKCC, KEYNOTE-001) were re-analyzed. A total of 34 advanced NSCLC (29 ADC) patients were prescribed pembrolizumab from 2012 to 2013 following the NCT01295827 protocol. All tumor tissues underwent whole-exome sequencing. We observed significantly increased nonsynonymous mutation and candidate neoantigen burden in the *TP53* or *KRAS* mutation group compared with the wild-type

**Figure 5.**

The correlation between *TP53*/*KRAS* mutation and clinical response to PD-1 blockade. Comparison of nonsynonymous mutation (**A**) and candidate neoantigen (**B**) burden in *TP53* or *KRAS* mutation and wild-type group. **C**, Proportion representation of transversion dominant mutation in the indicated group based on *TP53* or *KRAS* mutation. **D** and **E**, Kaplan-Meier survival curves estimates of PFS compared *TP53* or *KRAS* mutation with the wild-type group in patients treated with pembrolizumab. **F**, Proportional representation of clinical benefit of pembrolizumab in the indicated group based on *TP53* or *KRAS* mutation. **G**, Individual PFS of 34 NSCLC patients coupled with their mutational status of *TP53* and *KRAS*, pathology, PD-L1 expression, mutation burden, mutation spectrum and clinical benefit to pembrolizumab in each patient. TH, transversion high; TL, transversion low; DCB, durable clinical benefit; NDB, no durable benefit; ADC, adenocarcinoma; mut, mutation; H, High nonsynonymous burden (mutation  $\geq 209$ ); L, low nonsynonymous burden (mutation  $< 209$ ); PD-L1 expression: S, strong ( $\geq 50\%$ ); W, weak ( $1\sim 49\%$ ); N, negative ( $< 1\%$ ); NA, not available;  $***$ ,  $P < 0.001$ ;  $*$ ,  $P < 0.05$ .



Dong et al.

group (Fig. 5A and B). Consistent with the analysis from mutation burden, there was a strikingly high proportion of TH in the *TP53* and *KRAS* mutation group (Fig. 5C).

Notably, *TP53* or *KRAS* mutation patients obtained a significantly prolonged progression-free survival (PFS) compared with wild-type patients who underwent pembrolizumab treatment (median PFS, *TP53*-mut vs. *KRAS*-mut vs. wild-type: 14.5 vs. 14.7 vs. 3.5 months,  $P = 0.012$ ; Fig. 5D and E). Most of *TP53* or *KRAS* mutation patients enjoyed a durable clinical benefit during treatment, while most of wild-type patients showed no durable benefit (Fig. 5F). More importantly, 4 lung adenocarcinoma patients who concomitantly harbored *TP53* and *KRAS* mutation manifested superior PFS, and all of them had a durable clinical benefit. Meanwhile, these patients displayed a high mutation burden, high rate of transversion, and strong staining of PD-L1 (Fig. 5G).

To further confirm these observations from a public database, we prospectively collected 20 NSCLC (15 ADC) patients who were treated with pembrolizumab ( $n = 11$ ) or nivolumab ( $n = 9$ ) from August 2015 to August 2016 in our center (GLCI). All of the patients underwent at least one assessment after baseline. Patients' tissues were used for DNA sequencing for *EGFR*, *KRAS*, and *TP53*, and paraffin-embedded specimens were detected for IHC analysis of PD-L1 and CD8 (Table 1). Eight of the patients showed *TP53* mutation and 3 patients showed *KRAS* mutation. Up to August 25, 2016, 6 patients experienced partial response (PR). Two of them concomitantly harbored *TP53* and *KRAS* mutation, 3 patients with single *KRAS* or *TP53* mutation, and 1 patient without common mutation. Besides, 6 patients were evaluated as progression disease (PD) after 2 or 3 cycles of immunotherapy and 2 of them harbored *EGFR* mutation (Fig. 6A and Table 1). Patients with *TP53* and/or *KRAS* mutation showed prolonged PFS than both genes negative patients treated with PD-1 inhibitors. Six patients were assessed as PR and five of them had an ongoing response (Fig. 6B).

Next, we focused on 1 patient who had a durable clinical benefit (DCB) with *TP53* and *KRAS* comutation (Fig. 6C). IHC was used for analysis of PD-L1, CD8, and MMR-related genes, including *MSH2*, *MSH6*, *MLH1*, and *PMS2*. Consistent with our expectation, the patient showed strong staining for both PD-L1 and CD8. Meanwhile, four MMR-related genes displayed different intensity of immunostaining: weak positive for *MLH1*, moderate positive for *MSH6* and *PMS2*, and strong positive for *MSH2*, suggesting that a potential possibility of MMR deficiency existed in this tumor (Fig. 6D).


## Discussion

Although the expression of PD-L1 on the surface of tumor cells, as measured by IHC, is recommended as a predictive factor to identify patients who would benefit from PD-1 blockade, not all PD-L1-positive patients respond well (10, 37). The underlying biology of such limitations has not been clearly understood until recent studies, which showed that the presence of TILs and mutational burden correlated with T-effector signature and immunogenic features that supported the response to anti-PD-1/PD-L1 therapy (5, 8, 12, 38, 39). Here, we first identified a group of oncogenic driver (*EGFR* and *KRAS*) and tumor suppressor (*TP53* and *STK11*) mutations of lung adenocarcinoma that distinctively affected immune checkpoints expression, T-cell infiltration, and tumor immunogenicity. Specifically, our findings revealed that *TP53* mutation remarkably increased PD-L1

**Table 1.** Molecular and demographic characteristics and efficacy of PD-1 inhibitors in 20 non-small cell lung cancer

Characteristic	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	P13	P14	P15	P16	P17	P18	P19	P20
Age	66	54	52	53	56	58	58	58	51	53	62	50	57	64	58	75	59	70	50	53
Gender	Male	Male	Male	Male	Male	Female	Female	Male	Male	Male	Male	Male	Female	Male	Female	Male	Male	Male	Male	Male
Smoking history (page years)	40	60	30	40	70	0	0	60	30	30	120	48	0	60	0	10	0	75	20	35
ECOG PS	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2	1	1	1	1	1
Pathology	ADC	ADC	ADC	SCC	ADC	ADC	ADC	LELC	ADC	ADC	SCC	NEC	ADC	ADC	ADC	ADC	SCC	ADC	ADC	ADC
Clinical staging	IV	IIIA	IV	IIIA	IV	IV	IIIB	IIIB	IV	IV	IV	IV	IIIB	IV	IV	IV	IV	IV	IV	IV
EGFR	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT
KRAS	G12D	WT	G12V	WT	G12C	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT	WT
TP53	WT	WT	Q167fs	Y234H	Y126_Splice	WT	G187_Splice	WT	V151F	P87L	NA	WT	NA	H214R	WT	WT	WT	WT	WT	WT
PD-L1	Weak	NA	NA	Weak	Strong	NA	Negative	Weak	Negative	Strong	Negative	NA	NA	Strong	NA	Weak	Strong	Negative	Negative	Negative
CD8+TILs	Negative	NA	NA	Weak	Strong	NA	Negative	Weak	NA	Strong	Negative	NA	NA	Strong	NA	Weak	Negative	Negative	Negative	Negative
Immunotherapy	Pembro	Pembro	Nivo	Pembro	Pembro	Nivo	Nivo	Nivo	Nivo	Nivo	Nivo	Pembro	Pembro	Pembro	Pembro	Pembro	Pembro	Pembro	Pembro	Nivo
Best response	PR	PR	PR	PR	PR	PR	PR	PR	PR	SD	SD	SD	SD	SD	SD	SD	SD	SD	SD	SD
PFS (months)	11 m	9.5 m	9.5 m	8 m	7 m	7.5 m	6.0 m	5.5 m	5.0 m	4.5 m	4.5 m	3.5 m	3.5 m	3.5 m	2 m	2 m	2.5 m	2.0 m	1.5 m	1.5 m

Abbreviations: ADC, adenocarcinoma; SCC, squamous carcinoma; CY, cycles; LELC, lymphoepithelioma-like carcinoma; NA, not available; NEC, neuroendocrine carcinoma; Nivo, nivolumab; PD, progression disease; Pembro, pembrolizumab; PFS, progression-free survival; PR, partial response; SD, stable disease; WT, wild-type.

**Figure 6.**

Antitumor activity and biomarkers analysis of PD-1 blockade in patients with NSCLC. **A**, Best tumor burden change from baseline in target lesions in 20 NSCLC (15 ADC) patients who received nivolumab or pembrolizumab. The presence of mutation genes in each patient was indicated. **B**, Time to progression and duration of response in individual patients, as defined by Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1. **C**, PET/CT scan shows typical imaging alteration in a patient after 4 cycles of pembrolizumab treatment. **D**, Biomarker analysis of *TP53* and *KRAS* mutation status and protein expression of PD-L1, CD8, MSH2, MSH6, MLH1, and PMS2 detected by DNA sequencing and IHC.

expression and facilitated CD8<sup>+</sup> T-cell infiltration, and accompanied with a higher proportion of dual positive *PD-L1* and *CD8A* than other mutation groups. Furthermore, *TP53*-mutated tumors showed prominently increased somatic mutation burden and specifically enriched in the TH subset. Previous studies have classified the TME into four groups on the basis of *PD-L1* expression and TIL recruitment. These include type I (*PD-L1* positive with TILs driving adaptive immune resistance), type II (*PD-L1* negative with no TIL indicating immune ignorance), type III (*PD-L1* positive with no TIL indicating intrinsic induction), and type IV (*PD-L1* negative with TIL indicating the role of other suppressors in promoting immune tolerance; refs. 28, 29). Significantly, type I is lately thought to be associated with a high mutational burden, *PD-L1* amplification, and oncogenic viral infection, which defines a subtype sensitivity to PD-1 blockade (29). These notions, to some extent, support our findings that *TP53* mutation represents a state of adaptive immune resistance and a high immunogenicity, which contributes to a probable sensitivity to PD-1 blockade (40). Nevertheless, we could also discover a fact that *TP53* mutation equally enhanced some other non-*PD-L1* immune-inhibitory checkpoints expression, such as *LAG3* and *VTCN1*, which might serve as potential primary resistance to *TP53* mutation patients treated with anti-PD-1/*PD-L1* (41).

Recent studies based on a phase III clinical trial have identified that patients who harbored an *EGFR* mutation displayed unfavorable response to PD-1 blockade than those with a wild-type *EGFR* (42–44). This may be the first finding in which driver mutation of NSCLC was involved in altering sensitivity to immunotherapy. The most likely explanation is that patients with *EGFR* mutation were prone to produce a weak immunogenic tumor and an immunosuppressive TME. These perspectives were also confirmed in our study that *EGFR* mutation showed the lowest mutation burden and lowest rate of Tv/Ti than other mutations. Besides, *EGFR* mutation did not increase the expression of *PD-L1*, like others reported, but with a relatively lower expression than *TP53* and *KRAS* mutation (45), which further supported their hyposensitivity to PD-1 blockade. *KRAS* mutation was the second important oncogenic driver mutation in lung adenocarcinoma. The development of more effective treatment strategies for patients with *KRAS* mutation is hampered by the biologic and phenotypic heterogeneity of *KRAS*-mutant tumors. More recently, some studies suggested that patients with activating mutations in *KRAS* may probably benefit from PD-1 blockade, but the underlying mechanisms remained elusive and most of the researchers attributed this predilection to the association between smoking and the presence of *KRAS* mutations (5, 42). In this study, we uncovered potential mechanisms that account for this correlation. We discovered a significant increase of mutation load in *KRAS*-mutant tumors. Particularly, a predominant higher proportion of Tv/Ti was also found in this subgroup. Furthermore, we observed that *KRAS* mutations defected DNA repair, especially in MMR, which supported the notion that MMR deficiency acted as a favorable agent for PD-1 blockade (7).

It is well known that smoking-related lung cancers were characterized by greater mutation burden, higher rate of transversion, and more frequent *KRAS* mutation than that occurred in never smokers (21, 32, 46, 47). More recently, studies have demonstrated the association of *PD-L1* expression with significant smoking history (48). In our study, we discovered that *TP53* mutation, especially *TP53*/*KRAS* comutation, showed increased *PD-L1* expression and augmented tumor immunogenicity. To confirm

whether these correlations are more related to tobacco exposure, a multivariate linear regression analysis of mutation count and *PD-L1* expression stratified by smoking status was performed. We demonstrated that *TP53* mutation was responsible for increased mutation burden and *PD-L1* expression independent of smoking status (Supplementary Tables S4 and S5). Recent studies based on subgroup analysis demonstrated those with a history of current or ever smoking showed much better benefits of PD-1 blockade than non-smokers. So we can imagine current or ever-smoker patients with *TP53* and/or *KRAS* mutation may be the optimal population for PD-1 blockade immunotherapy.

Co-occurring mutations in *TP53* and *KRAS* have recently been defined as a specific cluster associated with activation of antitumor immunity and immune tolerance/escape (49). Interestingly, our study identified *TP53* and *KRAS* comutant tumors manifested exclusive increased expression of *PD-L1* and a highest proportion of *PD-L1*<sup>+</sup>/*CD8A*<sup>+</sup> than *TP53* or *KRAS* single mutation. Meanwhile, *TP53*/*KRAS* dual mutation showed predominant increased mutation burden and enriched in the TH subset. Consistent with these preclinical predictions, the clinical analysis on the base of MSKCC and our center database had further confirmed that those with co-occurring mutations in *TP53* and *KRAS* showed remarkable clinical benefit from pembrolizumab. These results implicated a possibility that *TP53* and *KRAS* mutation played a role with synergistic and complementary in regulating immune biomarkers, which gave rise to a responsive TME with adaptive immune resistance and high immunogenicity. However, these findings were established in a relatively small cohort and even fewer patients with *TP53* and *KRAS* comutation. Based on the preliminary evidence, a prospective study with a larger sample size of *TP53*/*KRAS* mutation and *PD-L1* expression for response to PD-1 blockade is warranted in the future.

Taken together, the results of this study provided an insight into immune regulation driving by some common mutations of lung adenocarcinoma. We discovered a prominent significance of *TP53* and *KRAS* mutation in boosting *PD-L1* expression, facilitating T-cell infiltration, and augmenting tumor immunogenicity. This work provided evidence that *TP53* and *KRAS* mutation in lung adenocarcinoma might be served as a pair of potential predictive factors in guiding PD-1 blockade immunotherapy.

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Authors' Contributions

Conception and design: Z.-Y. Dong, Y.-L. Wu

Development of methodology: Z.-Y. Dong, J. Su, Z. Xie, H.-H. Yan, S.-P. Wu, Y.-L. Wu

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): Z.-Y. Dong, W.-Z. Zhong, X.-C. Zhang, J. Su, Z. Xie, H.-Y. Tu, H.-J. Chen, Y.-L. Sun, Q. Zhou, X.-N. Yang, J.-X. Lin, H.-R. Zhai, L.-X. Yan, R.-Q. Liao

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): Z.-Y. Dong, W.-Z. Zhong, X.-C. Zhang, J. Su, Z. Xie, S.-Y. Liu, H.-Y. Tu, H.-J. Chen, Y.-L. Sun, Q. Zhou, H.-H. Yan, L.-X. Yan

Writing, review, and/or revision of the manuscript: Z.-Y. Dong, W.-Z. Zhong, J. Su, S.-Y. Liu, H.-Y. Tu, H.-J. Chen, Q. Zhou, X.-N. Yang, J.-X. Lin, H.-H. Yan, L.-X. Yan, Y.-L. Wu

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): Z.-Y. Dong, J. Su, Z. Xie, J.-J. Yang, H.-R. Zhai  
Study supervision: Z.-Y. Dong, W.-Z. Zhong, X.-C. Zhang, Y.-L. Wu



## Acknowledgments

The authors thank Professor Li Liu and Professor De-Hua Wu's team in Nanfang Hospital for providing lung adenocarcinoma-related public databases (TCGA, Broad and GEO databases) and helping with bioinformatic analysis. The authors also appreciate support from doctors and patients.

## Grant Support

This study was supported by the Guangzhou Science and Technology Bureau (grant nos. 2014Y2-00050 and 2014Y2-00545), The National Key Research and Development Program of China (grant no. 2016YFC1303800), the Special Fund for Research in the Public Interest from the National Health and Family

Planning Commission of the People's Republic of China (grant 201402031), the Project of National Natural Science Foundation (grant no. 81372285), and the Guangdong Provincial Applied Science and Technology Research and Development Program (grant no. 2016B020237006).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received October 11, 2016; revised December 1, 2016; accepted December 22, 2016; published OnlineFirst December 30, 2016.

## References

- Brahmer JR, Tykodi SS, Chow LQ, Hwu WJ, Topalian SL, Hwu P, et al. Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med* 2012;366:2455–65.
- O'Sullivan Coyne G, Madan RA, Gulley JL. Nivolumab: promising survival signal coupled with limited toxicity raises expectations. *J Clin Oncol* 2014;32:986–8.
- Carbognin L, Pilotto S, Milella M, Vaccaro V, Brunelli M, Calio A, et al. Differential activity of nivolumab, pembrolizumab and MPDL3280A according to the tumor expression of programmed death-ligand-1 (PD-L1): sensitivity analysis of trials in melanoma, lung and genitourinary cancers. *PLoS One* 2015;10:e0130142.
- Snyder A, Makarov V, Merghoub T, Yuan J, Zaretsky JM, Desrichard A, et al. Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N Engl J Med* 2014;371:2189–99.
- Rizvi NA, Hellmann MD, Snyder A, Kvistborg P, Makarov V, Havel JJ, et al. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* 2015;348:124–8.
- Blank CU, Haanen JB, Ribas A, Schumacher TN. Cancer Immunology. The "cancer immunogram". *Science* 2016;352:658–60.
- Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509–20.
- Tang H, Wang Y, Chlewicki LK, Zhang Y, Guo J, Liang W, et al. Facilitating T cell infiltration in tumor microenvironment overcomes resistance to PD-L1 blockade. *Cancer Cell* 2016;29:285–96.
- Dong ZY, Wu SP, Liao RQ, Huang SM, Wu YL. Potential biomarker for checkpoint blockade immunotherapy and treatment strategy. *Tumour Biol* 2016;37:4251–61.
- Herbst RS, Soria JC, Kowanetz M, Fine GD, Hamid O, Gordon MS, et al. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature* 2014;515:563–7.
- Shukuya T, Carbone DP. Predictive markers for the efficacy of anti-PD-1/PD-L1 antibodies in lung cancer. *J Thorac Oncol* 2016;11:976–88.
- Topalian SL, Taube JM, Anders RA, Pardoll DM. Mechanism-driven biomarkers to guide immune checkpoint blockade in cancer therapy. *Nat Rev Cancer* 2016;16:275–87.
- Pao W, Girard N. New driver mutations in non-small-cell lung cancer. *Lancet Oncol* 2011;12:175–80.
- Ding L, Getz G, Wheeler DA, Mardis ER, McLellan MD, Cibulskis K, et al. Somatic mutations affect key pathways in lung adenocarcinoma. *Nature* 2008;455:1069–75.
- Gibbons DL, Byers LA, Kurie JM. Smoking, p53 mutation, and lung cancer. *Mol Cancer Res* 2014;12:3–13.
- Cortez MA, Ivan C, Valdecana D, Wang X, Peltier HJ, Ye Y, et al. PDL1 regulation by p53 via miR-34. *J Natl Cancer Inst* 2016;108:djv303.
- Ji M, Liu Y, Li Q, Li X, Ning Z, Zhao W, et al. PD-1/PD-L1 expression in non-small-cell lung cancer and its correlation with EGFR/KRAS mutations. *Cancer Biol Ther* 2016;17:407–13.
- Soussi T, Beroud C. Assessing TP53 status in human tumours to evaluate clinical outcome. *Nat Rev Cancer* 2001;1:233–40.
- Schabath MB, Welsh EA, Fulp WJ, Chen L, Teer JK, Thompson ZJ, et al. Differential association of STK11 and TP53 with KRAS mutation-associated gene expression, proliferation and immune surveillance in lung adenocarcinoma. *Oncogene* 2016;35:3209–16.
- Imielinski M, Berger AH, Hammerman PS, Hernandez B, Pugh TJ, Hodis E, et al. Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* 2012;150:1107–20.
- Cancer Genome Atlas Research N. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* 2014;511:543–50.
- Deng L, Gyorffy B, Na F, Chen B, Lan J, Xue J, et al. Association of PDCD1 and CTLA-4 gene expression with clinicopathological factors and survival in non-small-cell lung cancer: results from a large and pooled microarray database. *J Thorac Oncol* 2015;10:1020–6.
- Choi YE, Meghani K, Brault ME, Leclerc L, He YJ, Day TA, et al. Platinum and PARP inhibitor resistance due to overexpression of MicroRNA-622 in BRCA1-mutant ovarian cancer. *Cell Rep* 2016;14:429–39.
- Ho AS, Kannan K, Roy DM, Morris LG, Ganly I, Katabi N, et al. The mutational landscape of adenoid cystic carcinoma. *Nat Genet* 2013;45:791–8.
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 2005;102:15545–50.
- Silwal-Pandit L, Vollen HK, Chin SF, Rueda OM, McKinney S, Osako T, et al. TP53 mutation spectrum in breast cancer is subtype specific and has distinct prognostic relevance. *Clin Cancer Res* 2014;20:3569–80.
- Anderson AC, Joller N, Kuchroo VK. Lag-3, Tim-3, and TIGIT: Co-inhibitory receptors with specialized functions in immune regulation. *Immunity* 2016;44:989–1004.
- Teng MW, Ngiew SF, Ribas A, Smyth MJ. Classifying cancers based on T-cell infiltration and PD-L1. *Cancer Res* 2015;75:2139–45.
- Ock CY, Keam B, Kim S, Lee JS, Kim M, Kim TM, et al. Pan-cancer immunogenomic perspective on the tumor microenvironment based on PD-L1 and CD8 T-cell infiltration. *Clin Cancer Res* 2016;22:2261–70.
- Fehrenbacher L, Spira A, Ballinger M, Kowanetz M, Vansteenkiste J, Mazieres J, et al. Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial. *Lancet* 2016;387:1837–46.
- Rooney MS, Shukla SA, Wu CJ, Getz G, Hacohen N. Molecular and genetic properties of tumors associated with local immune cytolytic activity. *Cell* 2015;160:48–61.
- Lee W, Jiang Z, Liu J, Haverty PM, Guan Y, Stinson J, et al. The mutation spectrum revealed by paired genome sequences from a lung cancer patient. *Nature* 2010;465:473–7.
- Mehnert JM, Panda A, Zhong H, Hirshfield K, Damare S, Lane K, et al. Immune activation and response to pembrolizumab in POLE-mutant endometrial cancer. *J Clin Invest* 2016;126:2334–40.
- van Gool IC, Eggink FA, Freeman-Mills L, Stelloo E, Marchi E, de Bruyn M, et al. POLE proofreading mutations elicit an antitumor immune response in endometrial cancer. *Clin Cancer Res* 2015;21:3347–55.
- Caron P, Choudhary J, Clouaire T, Bugler B, Daburon V, Aguirrebengoa M, et al. Non-redundant functions of ATM and DNA-PKcs in response to DNA double-strand breaks. *Cell Rep* 2015;13:1598–609.
- Lima M, Bouzid H, Soares DG, Selle F, Morel C, Galmarini CM, et al. Dual inhibition of ATR and ATM potentiates the activity of trabectedin and lurbectedin by perturbing the DNA damage response and homologous recombination repair. *Oncotarget* 2016;7:25885–901.

Dong et al.

37. Garon EB, Rizvi NA, Hui R, Leighl N, Balmanoukian AS, Eder JP, et al. Pembrolizumab for the treatment of non-small-cell lung cancer. *N Engl J Med* 2015;372:2018–28.
38. Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, et al. Signatures of mutational processes in human cancer. *Nature* 2013;500:415–21.
39. Schumacher TN, Schreiber RD. Neoantigens in cancer immunotherapy. *Science* 2015;348:69–74.
40. Tumei PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJ, Robert L, et al. PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature* 2014;515:568–71.
41. Koyama S, Akbay EA, Li YY, Herter-Sprie GS, Buczkowski KA, Richards WG, et al. Adaptive resistance to therapeutic PD-1 blockade is associated with upregulation of alternative immune checkpoints. *Nat Commun* 2016;7:10501.
42. Borghaei H, Paz-Ares L, Horn L, Spigel DR, Steins M, Ready NE, et al. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015;373:1627–39.
43. Herbst RS, Baas P, Kim DW, Felip E, Perez-Gracia JL, Han JY, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet* 2016;387:1540–50.
44. Gainor JF, Shaw AT, Sequist LV, Fu X, Azzoli CG, Piotrowska Z, et al. EGFR mutations and ALK rearrangements are associated with low response rates to PD-1 pathway blockade in non-small cell lung cancer: a retrospective analysis. *Clin Cancer Res* 2016;22:4585–93.
45. Chen N, Fang W, Zhan J, Hong S, Tang Y, Kang S, et al. Upregulation of PD-L1 by EGFR activation mediates the immune escape in EGFR-driven NSCLC: implication for optional immune targeted therapy for NSCLC patients with EGFR mutation. *J Thorac Oncol* 2015;10:910–23.
46. Govindan R, Ding L, Griffith M, Subramanian J, Dees ND, Kanchi KL, et al. Genomic landscape of non-small cell lung cancer in smokers and never-smokers. *Cell* 2012;150:1121–34.
47. Pfeifer GP, Denissenko MF, Olivier M, Tretyakova N, Hecht SS, Hainaut P. Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers. *Oncogene* 2002;21:7435–51.
48. Huynh TG, Morales-Oyarvide V, Campo MJ, Gainor JF, Bozkurtlar E, Uruga H, et al. Programmed cell death ligand 1 expression in resected lung adenocarcinomas: association with immune microenvironment. *J Thorac Oncol* 2016;11:1869–78.
49. Skoulidis F, Byers LA, Diao L, Papadimitrakopoulou VA, Tong P, Izzo J, et al. Co-occurring genomic alterations define major subsets of KRAS-mutant lung adenocarcinoma with distinct biology, immune profiles, and therapeutic vulnerabilities. *Cancer Discov* 2015;5:860–77.

# Clinical Cancer Research

## Potential Predictive Value of *TP53* and *KRAS* Mutation Status for Response to PD-1 Blockade Immunotherapy in Lung Adenocarcinoma

Zhong-Yi Dong, Wen-Zhao Zhong, Xu-Chao Zhang, et al.

*Clin Cancer Res* 2017;23:3012-3024. Published OnlineFirst December 30, 2016.

**Updated version** Access the most recent version of this article at:  
doi:[10.1158/1078-0432.CCR-16-2554](https://doi.org/10.1158/1078-0432.CCR-16-2554)

**Supplementary Material** Access the most recent supplemental material at:  
<http://clincancerres.aacrjournals.org/content/suppl/2016/12/30/1078-0432.CCR-16-2554.DC1>

**Cited articles** This article cites 49 articles, 12 of which you can access for free at:  
<http://clincancerres.aacrjournals.org/content/23/12/3012.full#ref-list-1>

**Citing articles** This article has been cited by 24 HighWire-hosted articles. Access the articles at:  
<http://clincancerres.aacrjournals.org/content/23/12/3012.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://clincancerres.aacrjournals.org/content/23/12/3012>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.